

HETEROCYCLES, Vol. 81, No. 12, 2010, pp. 2817 - 2830. © The Japan Institute of Heterocyclic Chemistry
Received, 21st September, 2010, Accepted, 21st October, 2010, Published online, 25th October, 2010
DOI: 10.3987/COM-10-12069

**SYNTHESIS OF TWO ESTRADIOL-IMIDAZOLE C-RIBONUCLEOSIDE
HYBRID COMPOUNDS EXHIBITING INHIBITORY EFFECTS
AGAINST TYPE 1 17 β -HYDROXYSTEROID DEHYDROGENASE**

**Shinya Harusawa,* Chihiro Kojima, Kensuke Fujii, Yuusaku Yamashita,
Tomoya Tanaka, Lisa Araki, Toshinobu Yoshimura, Minoru Sakaguchi,
Yoshihide Usami, and Masanori Takaoka**

Osaka University of Pharmaceutical Sciences, 4-20-1 Nasahara, Takatsuki, Osaka
569-1094, Japan. e-mail: harusawa@gly.oups.ac.jp

Abstract – Novel estradiol-imidazole C-nucleoside hybrid compounds **4a** and **4b**, which have C4-linked C₀- and C₂-imidazole ribonucleosides as adenosine mimics and amide bond linkers, were designed and synthesized based on EM-1745, an inhibitor of type 1 17 β -hydroxysteroid dehydrogenase (17 β -HSD1). Compounds **4a** and **4b** were also tested as enzyme inhibitors.

INTRODUCTION

Breast cancer is one of the most common cancers diagnosed among women and approximately 60% of breast cancers are hormone-responsive.¹ Estradiol (E₂), the most potent female sex hormone, stimulates the growth of mammary tumors² and endometriosis³ by activating the estrogen receptor. One treatment approach for this type of cancer is to decrease the level of E₂ by inhibiting one of the enzymes involved in its biosynthesis.^{4,5} Among those enzymes, type 1 17 β -hydroxysteroid dehydrogenase (17 β -HSD1) catalyzes the last step in the process of biosynthesis of E₂, as illustrated in Figure 1.^{6,7} As this enzyme, utilizing the cofactor NAD(P)H, reduces the C17 ketone of estrone (E₁) into E₂,⁸ 17 β -HSD1 inhibitors are regarded as promising new agents for estrogen-induced diseases.⁹ Poirier and co-workers recently developed E₂-adenosine hybrid compounds as a new type of 17 β -HSD1 inhibitor (Figure 2).^{10,11} The compounds were designed to exhibit affinity for both substrate (E₁ or E₂) and cofactor [NAD(P)H] binding domains of the enzyme.^{11a,12} The most potent hybrid inhibitor is EM-1745 (**1**: IC₅₀ = 52 nM),¹⁰ in which E₂ is linked to the adenosine moiety via a 16 β -oriented eight-CH₂ ester spacer. Crystal structure analysis of a complex of EM-1745 and 17 β -HSD1 led to the identification of a series of hydrogen bonds

demonstrated the importance of particular adenine and guanine bases in the catalytic mechanism of ribozymes, showing that the imidazole-substituted ribozymes are active in both cleavage and ligation. The results suggested that the *C*-nucleoside **3a** and its C_2 -elongated homologue **3b** could be used as structural mimics of adenosine or guanosine having purine bases.¹⁶




Figure 3. Imidazole C_0 - and C_2 -nucleosides **3a** and **3b** as purine-base mimics

In this context, we envisioned that the adenosine moiety and ester linkage in EM-1745, which were susceptible to hydrolysis, could be replaced with imidazole *C*-nucleoside and amide bond, respectively.¹⁹ Further, the endocyclic amine function of the imidazole had possibility to form hydrogen bonds with Asp65 of 17 β -HSD1, similar to the 6-amino group of adenine in EM-1745 (Figure 2). We herein report the synthesis of new E_2 -imidazole *C*-nucleoside hybrid compounds (**4a** and **4b**) and the preliminary evaluations for 17 β -HSD1 (Figure 4).





Figure 4. New E_2 -imidazole *C*-nucleoside hybrid compounds

RESULTS AND DISCUSSION

The synthesis of 5'-amino imidazole C_0 - and C_2 -nucleosides **10a** and **10b**, each of which constitutes the right half of the target molecules **4a** and **4b**, is shown in Scheme 1. Tribenzylated β -ribofuranosyl imidazole **6** was prepared through the three steps from commercially available 2,3,5-tri-*O*-benzyl-D-ribose **5**²⁰ according to our previous procedures.²¹ The imidazole-*N* of **6** was

protected by a [2-(trimethylsilyl)ethoxy]methyl (SEM) group to give the 3:1 isomeric mixture **7** (75%) at the endocyclic *N* functions of the imidazole. Debenzylation of **7** with Pd(OH)₂/C and cyclohexene and subsequent Mitsunobu reaction [DEAD/Ph₃P/phthalimide] afforded selectively desired 5'-substituted phthalimide **9a** in 36% yield from **7**.²² Deprotection of **9a** with hydrazine hydrate afforded 5'-amino derivative **10a** in 60% yield. By the way, we have recently reported a method to generate vinylimidazole **11** (*E/Z* = 2/1) from the starting material **5** by the six-step route in 52% yield.¹⁶ Then, for the synthesis of C₂- amino homologue **10b**, vinylimidazole **11** was used as the key intermediate.¹⁶ The introduction of a




Scheme 1. Synthesis of **10a** and **10b**

Reagents and conditions: (a) NaH, SEMCl, THF, rt; (b) cyclohexene, 20% Pd(OH)₂/C, EtOH, reflux; (c) phthalimide, Ph₃P, DEAD, rt; d) NH₂NH₂·H₂O, EtOH, reflux.

SEM group at the ^{im}N position of **11** followed by debenzylation and reduction of the double bond produced *N*-SEM-imidazole C₂-ribonucleoside **8b** (73%). Conversion of **8b** into phthalimide **9b** (63%) and deprotection with hydrazine hydrate provided 5'-amino imidazole C₂-nucleoside **10b** (79%).


Carboxylic acid **16** as the left half was synthesized as shown in Scheme 2 and the starting protected allyl-E₂ **13** was synthesized from E₁ according to the literature.²³ Poirier *et al.* recently reported the synthesis of **16** by a cross-metathesis (CM)¹³ between allyl compound **13** and 7-octenal followed by oxidation of CM product.^{11b} On the other hand, as ruthenium olefin metathesis catalysts were tolerant against even the protonic functionalities like carboxylic acids and other functional groups,²⁴ we tried to use the commercially available 7-octenoic acid (**14**) on **13** for CM. When a mixture of allyl compound **13** and carboxylic acid **14** was refluxed in dichloromethane for 20 h in the presence of Hoveyda-Grubbs second generation catalyst (HG-2, 5 mol%), which was a powerful catalyst for CM,²⁵ a coupling product **15** could be obtained in 63% yield. Catalytic hydrogenation of the double bond of **15** yielded the saturated carboxylic acid **16**^{11b} (88%). Direct use of carboxylic acid **14** made four steps of Poirier procedure unnecessary: three-step preparation of 7-octenal from 6-bromo-1-hexanol and oxidation of CM product.^{11b}



Scheme 2. Synthesis of left half **16**

With amines **10a** and **10b** and carboxylic acid **16**, amide bond was formed in the presence of diethyl phosphorocyanidate (DEPC)²⁶ and triethylamine to give amide **17a** (*n* = 0, 67%), as shown in Scheme 3. The SEM and *tert*-butyldimethylsilyl (TBDMS) groups with tetra-*n*-butylammonium fluoride under co-existing ethylenediamine²⁷ were removed in refluxing THF for 2 h to give unsubstituted imidazole **18a** in 89% yield. Finally, cleavage of the tetrahydropyranyl (THP) group of **18a** with *p*-toluenesulfonic acid (*p*-TSA) successfully afforded the target C₀-compound **4a** in 86% yield. Similarly, amide compound **17b**

($n = 2$), prepared from carboxylic acid **16** and C₂-amino compound **10b**, was converted into the desired C₂-hybrid compound **4b**.



Scheme 3. Synthesis of **4a** and **4b**.

Reagents and conditions: (a) DEPC, Et₃N, DMF, rt, 22 h; (b) Bu₄NF, NH₂(CH₂)₂NH₂, THF, reflux, 3.5 h; (c) *p*-TSA, MeOH, rt, 2 h.

Compounds **4a** and **4b** were evaluated for their ability to inhibit the *in vitro* transformation of E₁ into E₂ by a human recombinant 17 β -HSD1.²⁸ Preliminary results were that **4b** (IC₅₀: 3.5 μ M) showed more

potent 17 β -HSD1 inhibitory effect than **4a** (IC₅₀: > 10 μ M). In case of the ribose-(CH₂)_n-imidazole moiety of **4a** and **4b**, insertion of the two-methylene spacer (**4b**, n = 2) remarkably increased the inhibitory effect compared to the case of n = 0 (**4a**). Although the inhibitory potency of compound **4b** is even much lower (about 1 / 70) than EM-1745 (IC₅₀: 52 nM), such kinds of hybrid compounds as E₂-imidazole C-nucleoside lead to the synthesis of several analogues for a structure-activity relationship study. Further work on application of the imidazole C-nucleosides toward biofunctional molecules is under way and will be published in due course.

EXPERIMENTAL

Optical rotation measurements were recorded with a DIP-1000 digital polarimeter (JASCO). IR spectra were recorded on an IR-435 spectrometer (Shimadzu). ¹H- and ¹³C-NMR spectra were measured with tetramethylsilane as the internal standard on Gemini-200, Mercury-300, and UNITY INOVA-500 spectrometers (Varian). Low-resolution MS and high-resolution MS were obtained on a JMS-700(2) (JEOL). Reactions with air- and moisture-sensitive compounds were carried out under the argon atmosphere. Unless otherwise noted, all extracts were dried over Na₂SO₄ and the solvent was removed in a rotary evaporator under reduced pressure. BW-127ZH and Chromatorex NH-DM 1020 [(NH-silica gel), Fuji Silysia] were used for column chromatography. Dehydrated THF was purchased (Wako). TLC was performed on the pre-coated TLC plates with 60F₂₅₄ (silica gel, Merck).

4(5)-(2,3,5-Tri-*O*-benzyl- β -D-ribofuranosyl)-1-[2-(trimethylsilyl)ethoxymethyl]imidazole (7) To suspension of NaH (60%, 54 mg, 1.34 mmol) in mineral oil in THF (5 mL) was added a solution of **6**²¹ (420 mg, 0.89 mmol) in THF (3 mL). The mixture was stirred at room temperature (rt) for 1.5 h to stop the evolution of hydrogen. A solution of SEMCl (237 mg, 1.34 mmol) in THF (6 mL) was added to the resulting mixture. After stirring at rt for 1 h, saturated aqueous ammonium chloride was added and the whole was evaporated to give a residue, which was subsequently distributed between CH₂Cl₂ and water. After the CH₂Cl₂ layer was separated, the aqueous layer was further extracted with CH₂Cl₂. The combined CH₂Cl₂ layer was washed with water and brine, dried over anhydrous MgSO₄, filtered and evaporated. The crude product was purified by column chromatography on silica gel using hexane, followed by EtOAc/hexane (7/3) to give **7** (402 mg, 75%) as a pale yellow oil. The 3:1 isomeric mixture of **7** was assigned on the basis of the following ¹H-NMR data: ¹H-NMR (CDCl₃) δ : 0.21 (9H, s), 1.00–1.20 (2H, m), 3.57–3.68 (2H, m), 3.68–4.00 (2H, m), 4.23–4.60 (3H, m), 4.63–4.94 (6H, m), 5.23–5.60 (3H, m), 7.18 (0.25H, s), 7.23 (0.75H, s), 7.40–7.63 (15H, m), 7.77 (1H, s). HR-MS: *m/z*: 601.3103 [Calcd for C₃₅H₄₅N₂O₅Si (M+H)⁺: 601.3095].

4(5)-(5-Deoxy-5-phthaloylamino- β -D-ribofuranosyl)-1-[2-(trimethylsilyl)ethoxymethyl]imidazole (9a)

A mixture of **7** (177 mg, 0.30 mmol), 20% Pd(OH)₂/C (106 mg), and cyclohexene (0.9 mL, 8.85 mmol) in EtOH (8.5 mL) was refluxed for 28 h. After filtration through Celite, the filtrate was evaporated to give 4(5)-(5-deoxy-5-phthaloylamino- β -D-ribofuranosyl)-1-[2-(trimethylsilyl)ethoxymethyl]imidazole (**8a**, 96 mg). ¹H-NMR (CD₃OD) δ : 0.00 (9H, s), 0.91 (2H, t, J = 7.2 Hz), 3.50-3.82 (4H, m), 3.93-4.03 (1H, m), 4.05-4.18 (2H, m), 4.75 (1H, d, J = 6.0 Hz), 5.43 (1.6H, s), 5.63 (0.4 H, q, J = 7.0 Hz), 7.48 (1H, s), 8.35 (1H, s). HR-MS: m/z : 331.1685 [Calcd for C₁₄H₂₇N₂O₅Si (M+H)⁺: 331.1687]. Phthalimide (48 mg, 0.32 mmol) and Ph₃P (270 mg, 1.03 mmol) were dissolved in a solution of **8a** (96 mg) in CH₂Cl₂ (5 mL).^{22,29} To this mixture, a solution of DEAD (40%, 0.47 mL, 1.03 mmol) in toluene was added slowly with stirring. After the reaction mixture was stirred for 2 h at rt, the reaction was quenched with MeOH (0.5 mL). The whole was evaporated to give a residue that was purified by column chromatography on silica gel using hexane, 50% EtOAc/hexane, EtOAc, and 10% MeOH/EtOAc to give **9a** (49 mg, 36%) as an oil. ¹H-NMR (CD₃OD) δ : -0.20 (9H, s), 0.78-1.00 (2H, m), 3.40-4.33 (7H, m), 4.76 (1H, d, J = 4.0 Hz), 5.30 (1.2H, s), 5.40 (0.8H, s), 6.90 (0.4H, s), 7.28 (0.6H, s), 7.60-7.90 (5H, m). EI-MS: m/z : 460 (M⁺+H). HR-MS: m/z : 460.1901 [Calcd for C₂₂H₃₀N₃O₆Si (M+H)⁺: 460.1904].

4(5)-(5-Amino-5-deoxy- β -D-ribofuranosyl)-1-[2-(trimethylsilyl)ethoxymethyl]imidazole (10a)

A solution of **9a** (93 mg, 0.20 mmol) and NH₂NH₂·H₂O (0.06 mL, 1.02 mmol) in EtOH (10 mL) was refluxed for 4 h and then cooled. A heaping spatula of 10% Pd/C was added to the solution and the reaction mixture was further refluxed for 20 min. After removal of the catalyst by filtration through a Celite pad, NH-silica gel (1 g) was added to the filtrate. The solvent was evaporated to give a coated silica gel, which was subsequently placed in a column (NH-silica gel, 4 g). Chromatography using 5%, 10%, and 15% MeOH in EtOAc as the eluent gave **10a** (40 mg, 60%) as an oil. ¹H-NMR (CD₃OD) δ : 0.00 (9H, s), 0.80-1.00 (2H, m), 2.67-3.00 (2H, m), 3.53 (2H, t, J = 8.0 Hz), 3.82-4.00 (2H, m), 4.15 (0.6H, t, J = 6.0 Hz), 4.23 (0.4H, t, J = 6.0 Hz), 4.72 (0.6H, d, J = 6.0 Hz), 4.88 (0.4H, d, J = 6.0 Hz), 5.33 (1.2H, s), 5.44 (0.8H, s), 7.04 (0.4H, s), 7.25 (0.6H, s), 7.80 (1H, s). ¹³C-NMR (CD₃OD) δ : -1.4 (-1.40), -1.4 (-1.37), 18.5, 45.0, 45.1, 67.0, 67.4, 73.7, 73.8, 75.7, 76.6, 77.1, 77.2, 80.8, 85.3, 86.3, 119.1, 128.0, 132.1, 139.5, 140.9, 142.0. HR-MS: m/z : 330.1845 [Calcd for C₁₄H₂₈N₃O₄Si (M+H)⁺: 330.1849].

4(5)-[(E,Z)-2-(2,3,5-Tri-O-benzyl- β -D-ribofuranos-1-yl)vinyl]-1-[2-(trimethylsilyl)ethoxymethyl]imidazole (12)

Using the same procedure as that for the preparation of **7**, **11**¹⁶ (E/Z = 2/1, 1.30 g, 2.5 mmol) was converted into the isomeric mixture **12** (1.10 g, 70%) as an oil. ¹H-NMR (CDCl₃) δ : -0.05 (2H, s), -0.03 (3H, s), -0.01 (4H, s), 0.86 (0.4H, t, J = 8.1 Hz), 0.88 (0.7H, t, J = 8.1 Hz), 0.90

(0.9H, t, $J = 8.1$ Hz), 3.40-3.50 (2H, m), 3.52-3.65 (2H, m), 3.74 (0.2H, t, $J = 5.3$ Hz), 3.82 (0.5H, t, $J = 5.3$ Hz), 3.86 (0.3H, t, $J = 5.3$ Hz), 3.93-4.04 (1H, m), 4.22-4.30 (1H, m), 4.47-4.76 (7H, m), 5.14 (0.4H, s), 5.17 (0.6H, s), 5.20 (1H, s), 5.57 (0.3H, dd, $J = 11.7, 8.3$ Hz), 6.04 (0.2H, dd, $J = 16.7, 6.7$ Hz), 6.34 (0.5H, dd, $J = 16.7, 6.7$ Hz), 6.49 (0.3H, d, $J = 11.7$ Hz), 6.60 (0.5H, d, $J = 16.7$ Hz), 6.62 (0.2H, d, $J = 16.7$ Hz), 6.79 (0.3H, s), 7.12 (0.2H, s), 7.20-7.40 (15.5H, m), 7.55 and 7.58 (1H, 2s). HR-MS: m/z : 627.3252 [Calcd for $C_{37}H_{47}N_2O_5Si$ ($M+H$) $^+$: 627.3254].

4(5)-[2-(β -D-Ribofuranos-1-yl)ethyl]-1-[2-(trimethylsilyl)ethoxymethyl]imidazole (8b) Using the same procedure as that for the preparation of **8a**, **12** (84 mg, 0.13 mmol) was converted into **8b** (35 mg, 73%) as an oil. 1H -NMR (CD_3OD) δ : -0.02 (9H, s), 0.89 and 0.90 (2H, 2t, $J = 8.2$ Hz), 1.75-2.04 (2H, m), 2.57-2.94 (2H, m), 3.50-3.61 (4H, m), 3.64-3.82 (4H, m), 5.32 (1.3H, s), 5.38 (0.7H, s), 6.90 (0.3H, s), 7.05 (0.7H, s), 7.89 (0.7H, s), 7.93 (0.3H, s). HR-MS: m/z : 358.1919 [Calcd for $C_{16}H_{30}N_2O_5Si$ (M^+): 358.1922].

4(5)-[2-(5-Deoxy-5-phthaloylamino- β -D-ribofuranos-1-yl)ethyl]-1-[2-(trimethylsilyl)ethoxymethyl]imidazole (9b) Phthalimide (21 mg, 0.14 mmol) and Ph_3P (89 mg, 0.34 mmol) were dissolved in a solution of **8b** (35 mg, 0.097 mmol) in THF (4 mL). To this mixture, a toluene solution of DEAD (40%, 0.15 mL, 0.34 mmol) was added slowly with stirring. The reaction mixture was stirred for 16 h at rt and then the reaction was quenched with a small amount of water. The whole mixture was evaporated to give a residue, which was subsequently purified by column chromatography on silica gel using 10% MeOH/EtOAc to give **9b** (30 mg, 63%) as a colorless oil. 1H -NMR (CD_3OD) δ : -0.04 (9H, s), 0.87 (2H, t, $J = 8.3$ Hz), 1.70-1.99 (2H, m), 2.56-2.81 (2H, m), 3.42 (2H, t, $J = 8.3$ Hz), 3.72-4.13 (6H, m), 5.15 (1.3H, s), 5.17 (0.7H, s), 6.74 (0.3H, s), 6.77 (0.7H, s), 7.49 (1H, m), 7.64-7.82 (4H, m). HR-MS: m/z : 487.2126 [Calcd for $C_{24}H_{33}N_3O_6Si$ (M^+): 487.2137].

4(5)-[2-(5-Amino-5-deoxy- β -D-ribofuranos-1-yl)ethyl]-1-[2-(trimethylsilyl)ethoxymethyl]imidazole (10b) Using the same procedure as that for the preparation of **10a**, **9b** (29 mg, 0.06 mmol) was converted into **10b** (17 mg, 79%) as a colorless oil. 1H -NMR (CD_3OD) δ : -0.01 (9H, s), 0.84 (2H, t, $J = 8.1$ Hz), 1.72-2.06 (2H, m), 2.57-2.91 (4H, m), 3.51 and 3.53 (2H, 2t, $J = 8.1$ Hz), 3.68-3.82 (4H, m), 5.28 (0.7H, s), 5.34 (0.3H, s), 6.78 (0.3H, s), 6.96 (0.7H, s), 7.67 (1H, s). HR-MS: m/z : 357.2089 [Calcd for $C_{16}H_{31}N_3O_4Si$ (M^+): 357.2082].

(*E,Z*)-9-[3-(*tert*-Butyldimethylsilyloxy)-17 β -(tetrahydro-2*H*-pyran-2-yl-oxy)-estra-

1,3,5(10)-trien-16 β -yl]-7-nonenoic acid (15) A mixture of protected 16 β -allyl E₂ **13**²³ (114 mg, 0.22 mmol), 7-octenoic acid **14** (71 mg, 0.48 mmol), and HG-2 (10 mg, 0.015 mmol) in dry CH₂Cl₂ (7 mL) was refluxed for 20 h. The solvent was evaporated to give a residue, which was subsequently chromatographed with EtOAc/hexane (10 to 30%) as eluent to give **15** (87 mg, 63%) as a foam. It was dissolved by visual bubbling in saturated aqueous sodium bicarbonate. IR (film) cm⁻¹: 1640-1810 (br, C=O). ¹H-NMR (CDCl₃) δ : 0.19 (6H, s), 0.80 and 0.84 (3H, 2s), 0.97 (9H, s), 1.15-2.28 (30H, m), 2.35 (2H, t, J = 11.4 Hz), 2.72-2.86 (2H, m), 3.43-3.59 (1H, m), 3.68-3.85 (1H, m), 3.85-4.02 (1H, m), 4.61-4.79 (1H, m), 5.32-5.44 (2H, m), 6.54 (1H, d, J = 3.8 Hz), 6.60 (1H, dd, J = 12.2, 3.8 Hz), 7.10 and 7.12 (1H, 2d, J = 12.2 Hz). Selected ¹³C-NMR (CDCl₃) δ : 117.1, 119.9, 126.1, 133.2, 137.8, 153.2, 179.5 (C=O). HR-MS: m/z : 624.4205 [Calcd for C₃₈H₆₀O₅Si (M⁺): 624.4207].

9-[3-(*tert*-Butyldimethylsilyloxy)-17 β -(tetrahydro-2H-pyran-2-yl-oxy)-estra-1,3,5(10)-trien-16 β -yl]nonanoic acid (16) A solution of **15** (35 mg, 0.056 mmol) in EtOH (5 mL) was hydrogenated over 10% Pd on carbon (21 mg) at 3.0 kg/cm² for 3 h. After filtration through Celite, a small amount of silica gel was added to the filtrate. The solvent was evaporated to give a coated silica gel, which was subsequently placed in a column. Chromatography using EtOAc/hexane (5:95) as eluent gave **16**^{11b} (31 mg, 88%) as a colorless oil. It was dissolved by visual bubbling in saturated aqueous sodium bicarbonate. ¹H-NMR (CDCl₃) δ : 0.18 (6H, s), 0.79 and 0.83 (3H, 2s), 0.97 (9H, s), 1.03-2.30 (32H, m), 2.35 (2H, t, J = 7.0 Hz), 2.72-2.85 (2H, m), 3.44-3.56 (1H, m), 3.71 and 3.78 (1H, 2d, J = 9.6 Hz), 3.84-4.04 (1H, m), 4.58-4.77 (1H, m), 6.54 (1H, d, J = 1.9 Hz), 6.60 (1H, dd, J = 8.7, 1.9 Hz), 7.09 and 7.11 (1H, 2d, J = 8.7 Hz). HR-MS: m/z : 626.4359 [Calcd for C₃₈H₆₂O₅Si (M⁺): 626.4363].

N-[1-(Imidazol-4-yl)-5-deoxy- β -D-ribofuranos-5-yl]-9-[3-hydroxy-17 β -(tetrahydro-2H-pyran-2-yloxy)-estra-1,3,5(10)-trien-16 β -yl]nonanamide (18a) To a solution of **16** (89 mg, 0.14 mmol) in DMF (3 mL) were added a solution of **10a** (47 mg, 0.14 mmol) in DMF (2 mL), a solution of DEPC (90%, 38 mg, 0.21 mmol) in DMF (2 mL), and Et₃N (58 μ L, 0.42 mmol) in turn. The mixture was stirred at rt for 22 h and then diluted with EtOAc-hexane (3:1). The whole mixture was washed with H₂O, saturated aq. NaHCO₃, and brine, dried, filtered and evaporated to give a crude oil, which was purified by column chromatography (100% EtOAc to 5% MeOH in EtOAc) to give **17a** (87 mg, 67%) as an oil. ¹H-NMR (CD₃OD) δ : 0.00 (9H, s), 0.17 (6H, s), 0.75-0.95 (5H, m), 0.99 (9H, s), 1.03-2.33 (35H, m), 2.70-2.88 (2H, br), 3.39-4.26 (10H, m), 4.57-4.66 (0.5H, br), 4.70 (1H, d, J = 7.2 Hz), 4.70-4.78 (0.5H, br), 5.33 (1.3H, s), 5.45 (0.7H, s), 6.50 (1H, s), 6.57 (1H, d, J = 8.0 Hz), 7.03 (0.3H, s), 7.07-7.14 (1H, m), 7.26 (0.7H, s), 7.79 (0.3H, s), 7.81 (0.7H, s). HR-MS: m/z : 938.6116 [Calcd for C₅₂H₈₈N₃O₈Si₂ (M+H)⁺:

938.6110]. Next, a 1 M solution of tetra-*n*-butylammonium fluoride (0.46 mL, 0.46 mmol) in THF and ethylenediamine (0.05 mL, 0.74 mmol) were added to a solution of **17a** (87 mg, 0.09 mmol) in THF (6 mL). The resulting mixture was refluxed for 3.5 h and then THF was evaporated to give a residue. The residue was subjected to chromatography [NH-silica gel, MeOH/CHCl₃ (10:90 to 20:80)] to give a pale yellow oil. As ¹H-NMR measurement of the oil indicated the presence of remaining tetra-*n*-butylammonium hydroxide, it was removed by filtration on C18 silica gel [MeOH/H₂O (70:30 to 100:0)] to give **18a** (57 mg, 89%) as a colorless oil. ¹H-NMR (CD₃OD) δ : 0.77 (1.2H, s), 0.82 (1.8H, s), 0.86-2.30 (35H, m), 2.72-2.80 (2H, br), 3.18-4.12 (8H, m), 4.57-4.62 (0.5H, br), 4.70-4.75 (0.5H, br), 4.75 (1H, d, J = 7.2 Hz), 6.46 (1H, s), 6.52 (1H, d, J = 8.0 Hz), 7.02-7.08 (1H, m), 7.10 (1H, s), 7.71 (1H, s). HR-MS: m/z : 694.4428 [Calcd for C₄₀H₆₀N₃O₇ (M+H)⁺: 694.4431].

***N*-[1-(Imidazol-4-yl)-5-deoxy- β -D-ribofuranos-5-yl]-9-[3,17 β -dihydroxy-estra-**

1,3,5(10)-trien-16 β -yl]nonanamide (4a) *p*-TsOH (15 mg, 0.09 mmol) was added to a solution of **18a** (56 mg, 0.08 mmol) in MeOH (3.5 mL). After stirring at rt for 2 h, the reaction mixture was neutralized with saturated aq. NaHCO₃ and evaporated. The residue was dissolved in EtOAc and the solution was washed with saturated aq. NaHCO₃ and brine, dried, filtered, and evaporated. The residual oil was purified on N-H silica gel using MeOH-CHCl₃ (5:95 to 15:85, v/v) to give **4a** (42 mg, 86%) as a colorless oil. R_f = 0.33 (30% MeOH in CHCl₃). $[\alpha]_D^{25}$ = +26.2° (c = 1.85, MeOH). ¹H-NMR (CD₃OD) δ : 0.76 (3H, s), 0.93-2.30 (29H, m), 2.71-2.84 (2H, m), 3.20-3.70 (3H, m), 3.93-3.99 (2H, m), 4.06-4.11 (1H, m), 4.73 (1H, d, J = 6.1 Hz), 6.47 (1H, d, J = 2.7 Hz), 6.53 (1H, dd, J = 8.5, 2.7 Hz), 7.06 (1H, d, J = 8.5 Hz), 7.09 (1H, s), 7.68 (1H, s). ¹³C-NMR (CD₃OD) δ : 13.3, 27.1, 27.6, 28.8, 29.9, 30.4, 30.5, 30.7, 30.8, 31.0, 33.0, 33.7, 37.1, 39.1, 40.0, 41.7, 42.5, 45.2, 45.5, 49.9, 50.0, 62.3, 73.5, 76.8, 80.2, 83.4, 83.8, 113.7, 116.1, 127.2, 132.7, 137.0, 138.8, 155.9, 176.7. HR-MS: m/z : 610.3860 [Calcd for C₃₅H₅₂N₃O₆ (M+H)⁺: 610.3856].

***N*-{1-[2-(Imidazol-4-yl)ethyl]-5-deoxy- β -D-ribofuranos-5-yl]-9-[3-hydroxy-17 β -**

(tetrahydro-2H-pyran-2-yloxy)-estra-1,3,5(10)-trien-16 β -yl]nonanamide (18b) A mixture of **16** (67 mg, 0.11 mmol), **10b** (41 mg, 0.12 mmol), DEPC (29 mg, 0.16 mmol), and Et₃N (0.05 mL, 0.32 mmol) in DMF (10 mL) was stirred at rt for 13 h to give **17b** (64 mg, 62%) as a colorless oil using the same procedure as that for the preparation of **17a**. ¹H-NMR (CD₃OD) δ : -0.08 (9H, s), 0.17 (6H, s), 0.78 and 0.82 (3H, 2s), 0.84-0.93 (2H, m), 0.96 (9H, s), 1.02-2.06 (33H, m), 2.08-2.30 (2H, m), 2.18 (2H, t, J = 6.9 Hz), 2.62-2.88 (4H, m), 3.36-3.58 (5H, m), 3.66-3.88 (5H, m), 3.89-4.15 (1H, m), 4.57-4.75 (1H, m), 5.18 (0.7H, s), 5.21 (0.3H, s), 6.50 (1H, d, J = 2.3 Hz), 6.55 (1H, dd, J = 9.2, 2.3 Hz), 6.78 (0.3H, s), 6.94

(0.7H, s), 7.08 and 7.10 (1H, 2d, $J = 9.2$ Hz), 7.52 (1H, s). HR-MS: m/z : 966.6422 [Calcd for $C_{54}H_{92}N_3O_8Si_2$ (M+H) $^+$: 966.6418]. A mixture of **17b** (18 mg, 0.02 mmol), 1 M THF solution of tetra-*n*-butylammonium fluoride (0.09 mL, 0.09 mmol) and ethylenediamine (0.01 mL, 0.15 mmol) in THF (1.5 mL) was refluxed for 2 h to give **18b** (9 mg, 69%) as a colorless oil using the same procedure as that for the preparation of **18a**. 1H -NMR (CD_3OD) δ : 0.78 and 0.82 (3H, 2s), 1.08-2.06 (37H, m), 2.06-2.83 (2H, m), 2.20 (2H, t, $J = 8.2$ Hz), 2.60-2.83 (4H, m), 3.26-3.56 (2H, m), 3.64-3.82 (5H, m), 3.84-4.03 (1H, m), 4.56-4.73 (1H, m), 5.32-5.43 (2H, m), 6.46 (1H, d, $J = 3.0$ Hz), 6.52 (1H, dd, $J = 9.0, 3.0$ Hz), 6.77 (1H, s), 7.04 and 7.05 (1H, 2d, $J = 9.0$ Hz), 7.53 (1H, s). HR-MS: m/z : 722.4739 [Calcd for $C_{42}H_{64}N_3O_7$ (M+H) $^+$: 722.4740].

***N*-{1-[2-(Imidazol-4-yl)ethyl]-5-deoxy- β -D-ribofuranos-5-yl}-9-[3,17 β -dihydroxy-estra-1,3,5(10)-trien-16 β -yl]nonanamide (4b)** A mixture of **18b** (20 mg, 0.03 mmol) and *p*-TsOH (5 mg, 0.03 mmol) in MeOH (1 mL) was stirred at rt for 4 h to give **4b** (12 mg, 70%) as a colorless oil using the same procedure as that for the preparation of **4a**. $R_f = 0.32$ (20% MeOH in $CHCl_3$). $[\alpha]_D^{25} = +25.5^\circ$ ($c = 1.65$, MeOH). 1H -NMR (CD_3OD) δ : 0.76 (3H, s), 1.22-2.34 (32H, m), 2.20 (2H, t, $J = 7.5$ Hz), 2.60-2.87 (4H, m), 3.64-3.83 (5H, m), 6.46 (1H, d, $J = 3.2$ Hz), 6.52 (1H, dd, $J = 7.9, 3.2$ Hz), 6.78 (1H, s), 7.06 (1H, d, $J = 7.9$ Hz), 7.56 (1H, s). ^{13}C -NMR (CD_3OD) δ : 13.5, 24.1, 27.2, 27.7, 28.9, 30.0, 30.5, 30.6, 30.8, 30.9, 31.2, 33.1, 33.7, 34.9, 37.1, 39.1, 40.1, 41.7, 42.8, 45.2, 45.5, 50.0, 73.8, 76.0, 83.1, 83.3, 113.2, 115.5, 126.6, 132.1, 135.1, 138.1, 155.1, 175.6. HR-MS: m/z : 638.4165 [Calcd for $C_{37}H_{56}N_3O_6$ (M+H) $^+$: 638.4166].

Biological Assay

Human 17 β -HSD1 was overexpressed in the BL21-AI strain of *Escherichia coli* containing the pET-41 Ek/LIC-17 β -HSD1 vector, basically according to the method of Chang *et al.*²⁸ The human recombinant 17 β -HSD1 was purified by using a Glutathione Sepharose 4B column (GE Healthcare) and a Mono Q 5/50 GL column (GE Healthcare). The purified enzyme was incubated with substrate E_1 and cofactor NADH in the presence or absence of inhibitors at pH 7.4. The residual E_1 and newly produced E_2 in the incubation mixture were extracted with hexane. The organic phase was evaporated, the residue containing E_1 and E_2 was dissolved in acetonitrile solution and then measured by reverse-phase high-performance liquid chromatography with an amperometric detector (HTEC-500, EICOM).

ACKNOWLEDGEMENTS

We are grateful to Dr. K. Minoura and Ms. M. Fujitake of Osaka University of Pharmaceutical Sciences

for NMR and MS measurements, respectively. This work was supported in part by a Grant-in-Aid for Scientific Research [No. 21590130 (S. H.)] from JSPS and a Grant-in-Aid for “High Technology Research Center from MEXT (Ministry of Education, Sciences, Sports and Culture), 2009-2013, Japan.

REFERENCES AND NOTES

1. A. J. Theobald, *Int. J. Clin. Pract.*, 2000, **54**, 665.
2. R. C. Travis and T. J. Key, *Breast Cancer Res.*, 2003, **5**, 239.
3. G. S. Dizerega, D. L. Barber, and G. D. Hodgen, *Fertil. Steril.*, 1980, **33**, 649.
4. F. Labrie, V. Luu-The, S. X. Lin, J. Simard, C. Labrie, M. Elalfy, G. Pelletier, and A. Bélanger, *J. Mol. Endocrinol.*, 2000, **25**, 1.
5. H. J. Smith, P. J. Nicholls, C. Simons, and R. Le Latin, *Exp. Opin. Ther. Patents*, 2001, **11**, 789.
6. V. Luu-The, Y. Zhang, D. Poirier, and F. Labrie, *J. Steroid Biochem. Mol. Biol.*, 1995, **55**, 581.
7. V. Luu-The, *J. Steroid Biochem. Mol. Biol.*, 2001, **76**, 143.
8. R. Breton, D. Housset, C. Mazza, and J. C. Fontecilla-Camps, *Structure*, 1996, **4**, 905.
9. J. M. Day, H. J. Tutill, A. Purohit, and M. J. Reed, *Endocr. Relat. Cancer.*, 2008, **15**, 665 and references therein.
10. D. Poirier, R. P. Boivin, M. R. Tremblay, M. Bérubé, W. Qiu, and S.-X. Lin, *J. Med. Chem.*, 2005, **48**, 8134.
11. (a) D. Fournier, D. Poirier, M. Mazumdar, and S. -X. Lin, *Eur. J. Med. Chem.*, 2008, **43**, 2298; (b) M. Bérubé and D. Poirier, *J. Enzyme Inhib. Med. Chem.*, 2009, **24**, 832.
12. W. Qiu, R. L. Campbell, A. Gangloff, P. Dupuis, R. P. Boivin, M. R. Tremblay, D. Poirier, and S.-X. Lin, *FASEB.*, 2002, **16**, 1829.
13. (a) M. Bérubé and D. Poirier, *Org. Lett.*, 2004, **6**, 3127; (b) M. Bérubé and D. Poirier, *Can. J. Chem.*, 2009, **87**, 1180.
14. (a) L. Araki, S. Harusawa, M. Yamaguchi, S. Yonezawa, N. Taniguchi, D. M. J. Lilley, Z. Zhao, and T. Kurihara, *Tetrahedron Lett.*, 2004, **45**, 2657; (b) L. Araki, S. Harusawa, M. Yamaguchi, S. Yonezawa, N. Taniguchi, D. M. J. Lilley, Z. Zhao, and T. Kurihara, *Tetrahedron*, 2005, **61**, 11976.
15. L. Araki, Z. Zhao, D. M. J. Lilley, and S. Harusawa, *Heterocycles*, 2010, **81**, 1861.
16. L. Araki, K. Morita, M. Yamaguchi, Z. Zhao, T. J. Wilson, D. M. J. Lilley, and S. Harusawa, *J. Org. Chem.*, 2009, **74**, 2350.
17. Z. Zhao, A. Mcleod, S. Harusawa, L. Araki, M. Yamaguchi, T. Kurihara, and D. M. J. Lilley, *J. Am. Chem. Soc.*, 2005, **127**, 5026.
18. T. J. Wilson, J. Ouellet, Z. Zhao, S. Harusawa, L. Araki, T. Kurihara, and D. M. J. Lilley, *RNA*, 2006, **12**, 980.

19. D. E. Levy and C. Tang, 'Tetrahedron Organic Chemistry Series: The Chemistry of C-Glycosides,' Vol. 13, Chap. 1, ed. by J. E. Baldwin and P. D. Magnus, Pergamon, New York, 1995, pp. 4-27.
20. 2,3,5-Tri-*O*-benzyl-D-ribofuranose **5** was purchased from Carbosynth Ltd.
21. S. Harusawa, M. Murai, H. Moriyama, T. Imazu, H. Ohishi, R. Yoneda, and T. Kurihara, *J. Org. Chem.*, 1996, **61**, 4405.
22. S. Harusawa, H. Moriyama, Y. Murai, T. Imazu, H. Ohishi, R. Yoneda, T. Kurihara, H. Hata, and Y. Sakamoto, *Chem Pharm. Bull.*, 1997, **45**, 53.
23. M. R. Tremblay and D. Poirier, *J. Comb. Chem.*, 2000, **2**, 48.
24. (a) S. J. Connon and S. Blechert, *Angew. Chem. Int. Ed.*, 2003, **42**, 1900; (b) A. Fürstner, *Angew. Chem. Int. Ed.*, 2000, **39**, 3012.
25. S. Randl, S. Gessler, H. Wakamatsu, and S. Blechert, *Synlett*, 2001, 430.
26. T. Shioiri, Y. Yokoyama, Y. Kasai, and S. Yamada, *Tetrahedron*, 1976, **32**, 2211.
27. J. M. Muchowski and D. R. Solas, *J. Org. Chem.*, 1984, **49**, 203.
28. Y. H. Chang, Y. L. Wang, J. Y. Lin, L. Y. Chuang, and C. C. Hwang, *Mol. Biotechnol.*, 2010, **44**, 133.
29. Since triol **8a** was insoluble in THF, dichloromethane was used as a solvent for the reaction.