

ChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: S. Martinez-Erro, A. Bermejo Gómez, A. Vázquez-Romero, E. Erbing and B. Martín-Matute, *Chem. Commun.*, 2017, DOI: 10.1039/C7CC04823H.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Chemical Communications

COMMUNICATION

Received 00th January 20xx,

2,2-Diiododimedone: a mild electrophilic iodinating agent for the selective synthesis of α -iodoketones from allylic alcohols

Accepted 00th January 20xx

Samuel Martinez-Erro,[†] Antonio Bermejo Gómez,[†] Ana Vázquez-Romero, Elis Erbing, and Belén Martín-Matute*

DOI: 10.1039/x0xx00000x

www.rsc.org/


2,2-Diiodo-5,5-dimethylcyclohexane-1,3-dione is reported as a new electrophilic iodinating agent that selectively iodineates electron-rich aromatics. In contrast to other common electrophilic iodinating reagents, its mild nature allows it to be used for the selective synthesis of α -iodinated carbonyl compounds from allylic alcohols through a 1,3-hydrogen shift / iodination process catalyzed by iridium(III) complexes.

Halogenated organic compounds are ubiquitous, not only as natural products,¹ but also as synthetic compounds, and they have applications in medicinal chemistry,² agrochemistry,³ and materials science.⁴ The halogen functional group is very versatile. It can participate in many synthetic transformations, including nucleophilic substitutions, eliminations, dehydrohalogenations, and oxidative additions to metal centers. The nature of the halogen gives each carbon–halogen bond unique properties,⁵ resulting in numerous applications in synthetic organic chemistry.

Significant progress has been made in the development of synthetic methods for the construction of C–F bonds, by both electrophilic and nucleophilic approaches.⁶ There also exist many synthetic methods for the nucleophilic and electrophilic chlorination⁷ and bromination⁸ of organic molecules. When it comes to electrophilic methods for the construction of C–I bonds reagents such as I₂, NIS (*N*-iodosuccinimide), IPy₂BF₄ (Barluenga's reagent), and ICl have been widely used in organic synthesis.⁹ Also, methods based on the conversion of iodide into the corresponding iodonium species through oxidation have also been reported.¹⁰ These methods have some drawbacks, however, such as the high reactivity of the electrophilic iodinating agents, or the need for strong oxidizing agents, which result in low functional-group tolerance and low

chemo- or regioselectivity. Thus, the development of mild and selective electrophilic iodination methods remains a challenging and desirable goal for synthetic chemists.

We have previously reported the use of iridium catalysts for the conversion of allylic alcohols (**1**) into α -functionalized carbonyl compounds.^{11,7c,8c,12} Carbon and halogen (i.e., F, Cl, and Br) functional groups could be introduced through a process involving a 1,3-hydrogen shift mediated by the metal complex. This process allows the selective formation of either the α or α' constitutional isomer, as the new bond is constructed exclusively at the carbon that was originally part of the olefin. Attempts to introduce an iodide failed in our hands, mainly due to the high reactivity of the available electrophilic iodinating agents, but also due to the instability of the starting allylic alcohols and/or the final iodoketones under the reaction conditions. In this paper, we report the development of a new mild iodinating agent (Scheme 1, top) and its application in electrophilic iodination of aromatic compounds, and also in the synthesis of α -iodoketones (**3**) from allylic alcohols mediated by iridium catalysts (Scheme 1, bottom). This methodology allows for the first time the one-step synthesis of α -iodoketones as single constitutional isomers from allylic alcohols in a one-step procedure. The utility of α -iodoketones for the preparation of valuable



Scheme 1 Synthesis of 2,2-diiododimedone (**2**, top) and of α -iodocarbonyl compounds from allylic alcohols.

Department of Organic Chemistry, Arrhenius Laboratory,
Stockholm University, 10691 Stockholm, Sweden.
E-mail: belen.martin.matute@su.se

[†] These authors contribute equally to this work.


Electronic Supplementary Information (ESI) available: detailed synthetic procedures, characterization, NMR spectra, optimization tables and deuterium labelling studies. See DOI: 10.1039/x0xx00000x

COMMUNICATION

Journal Name

building blocks is also presented.


The synthesis of the new iodinating reagent, 2,2-diiododimedone (**2**), was easily accomplished by treatment of dimedone with iodine monochloride in 1,4-dioxane at room temperature. The synthesis of **2** can be carried out on a gram scale, and purification by simple filtration gives the product in excellent yield. We initially tested **2** as an iodinating agent for aromatic compounds (Scheme 2). In optimization studies, the best results were obtained using a catalytic amount of FeCl₃, and hexafluoroisopropanol (HFIP) as the solvent. The need to use these conditions reflects the mildness of **2**, which is important for increasing the functional group tolerance in electrophilic halogenation reactions. After a short optimization (see ESI), we found that anisole **4a** could be iodinated under an air atmosphere in just 2 h. Two other substrates were also iodinated using these reaction conditions; both *N,N*-dimethylaniline (**4b**) and *N*-pivaloyl indole (**4c**) were converted into the corresponding iodinated analogues (**5b** and **5c**) in very good yields after short reaction times and with excellent regioselectivity.



Scheme 2 Iodination of electron-rich aromatics using **2**.

Next, we evaluated the performance of **2** in the iridium-catalyzed isomerization / α -iodination of allylic alcohols. For a successful outcome, the reaction must take place under mild conditions, and the starting alcohols (**1**) and final products (**3**) should not undergo undesired iodination. Oct-1-en-3-ol (**1a**) was chosen as a model substrate for the optimization studies, and several Ir(III) catalysts and solvents were evaluated (Table 1). When [Cp*IrCl₂]₂ (**I**, 2 mol% [Ir]) was used as the catalyst in a mixture of acetone and H₂O (1:1), full conversion was observed, but only 34% of the desired product **3a** was formed, together with 64% of α,β -unsaturated ketone **6a** (entry 1). This solvent mixture gave, on the contrary, excellent results in the bromination of allylic alcohols.^{8c} The use of THF / H₂O (1:1) resulted in a clear decrease in the formation of **6a** (entry 2). This effect was even clearer when THF was replaced by 2-methyltetrahydrofuran (2-MeTHF). In this case, the desired α -iodinated compound (**3a**) was obtained in 85% yield and only 11% of **6a** was formed (entry 3). The amount of **6a** could be decreased further by using [Cp*Ir(H₂O)₃]SO₄ (**II**) or [(Cp*Ir)₂(OH)₃]OH·11H₂O (**III**) as catalysts (entries 4 and 5). When the proportion of H₂O in the solvent mixture was increased (i.e., 1:2), higher amounts of the undesired **6a** were formed (entries 6 and 7); when the proportion of 2-MeTHF was increased, the yield of **6a** dropped to just 5-6% (entries 8


Table 1 Optimization.^a



| Entry | Catalyst | Solvent (v / v) | Conv. [%] ^b | 3a/6a [%] ^{b,c} |
|-----------------|----------|--------------------------------|------------------------|--------------------------|
| 1 | I | Acetone/H ₂ O (1:1) | >99 | 36/64 |
| 2 | I | THF/H ₂ O (1:1) | 86 | 58/28 |
| 3 | I | 2-MeTHF/H ₂ O (1:1) | 96 | 85/11 |
| 4 | II | 2-MeTHF/H ₂ O (1:1) | 99 | 93/6 |
| 5 | III | 2-MeTHF/H ₂ O (1:1) | >99 | 92/8 |
| 6 | II | 2-MeTHF/H ₂ O (1:2) | >99 | 91/9 |
| 7 | III | 2-MeTHF/H ₂ O (1:2) | >99 | 90/10 |
| 8 | II | 2-MeTHF/H ₂ O (2:1) | 98 | 93/5 |
| 9 | III | 2-MeTHF/H ₂ O (2:1) | >99 | 94/6 |
| 10 ^d | II | 2-MeTHF/H ₂ O (2:1) | 86 | 79/7 |
| 11 ^d | III | 2-MeTHF/H ₂ O (2:1) | 92 | 88/4 |

^a Unless otherwise noted: **1a** (0.5 mmol, 0.1 M) and 2 mol% of [Ir] was used. ^b Determined by ¹H NMR spectroscopy. ^c Octan-3-one was not formed. ^d 1 mol% of [Ir].

Table 2 Evaluation of a variety of iodinating agents.^a



| Entry | Reagent | Conv. [%] ^b | 3a/6a [%] ^b |
|-------|---|------------------------|------------------------|
| 1 | 2 | >99 | 94/6 |
| 2 | I ₂ | 66 | -/3 ^c |
| 3 | ICl | 66 | -/1 ^c |
| 4 | NIS | >99 | 17/- |
| 5 | IPy ₂ BF ₄ | >99 | -/- |
| 6 | - | <5 | - |
| 7 | 2 ^d | 4 | - |
| 8 | I ₂ ^d | 25 | - ^e |
| 9 | ICl ^d | 95 | - ^e |
| 10 | NIS ^d | 90 | - ^e |
| 11 | IPy ₂ BF ₄ ^d | 76 | - ^e |

^a Unless otherwise noted: **1a** (0.5 mmol, 0.1 M). ^b Determined by ¹H NMR spectroscopy. ^c 28% (entry 2) and 21% (entry 3) of octan-3-one was formed. ^d Without [Ir] catalyst. ^e A mixture of unknown by-products was observed.

and 9), while a very good yield for **3a** was maintained. Increasing the proportion of 2-MeTHF further did not prevent the formation of **6a**, and also compromised the conversion (See ESI for a full description of solvent effects). When the catalyst loading was decreased to 1 mol% [Ir], catalyst **III** was better than **II** (entries 10 and 11). When 0.6 and 0.2 equiv. of **2** were used, yields of 70 and 27%, respectively, were observed (see ESI). This result suggests that more than one iodo unit can be donated by the reagent. All reactions were carried out under an atmosphere of air.

When compared to other electrophilic iodinating reagents, the superiority of **2** was clearly seen (Table 2). None of the other reagents tested were able to give good yields of **3a**, mainly due to decomposition of the starting allylic alcohol **1a** (entries 1-5). This is not surprising, since it is well known that double bonds promptly react with these reagents to give cyclic iodonium ions. This undesired reaction was not observed when 2,2-diiododimedone (**2**) was used. A series of control experiments was also carried out (Table 2, entries 6-11). Catalyst **III** was not able alone to react with the allylic alcohol in the absence of **2** (entry 6). We recently reported that the presence of a halogen ligand on Ir is important for the activity of [Cp*Ir] complexes of this type, and described the *in situ* formation of such an Ir-halogen bond through reaction of the Ir precatalyst with the halogenating agent.¹³ In another control experiment, we observed that **2** does not react with allylic alcohol **1a** in the absence of the iridium catalyst after 16 h (entry 7). It can therefore be concluded that the mildness of iodinating agent **2** is key to the success of this tandem isomerization / halogenation reaction. Mechanistic investigations were carried out with deuterium-labelled alcohols, and the results suggest that an intramolecular 1,3-H shift is operating under these reaction conditions (see ESI).

Having established these optimized reaction conditions, we went on to study the scope of the reaction (Table 3). Alcohols **1a-1f** and **1k-1m** bearing a terminal double bond gave the corresponding α -iodocarbonyl compounds (**3a-3f** and **3k-3m**) in high yields within 16 h. Allylic alcohols with internal double bonds (**1g-1j**) were also converted into the corresponding α -iodinated products in good yields. Additional double bonds in the molecule were unaffected (**1f** and **1g**); only the allylic alcohol functionality reacted. The reaction is not restricted to secondary alcohols; primary alcohols gave the corresponding α -iodoaldehydes (**3n-3r**) in moderate to good yields. The stability of the products during the purification process accounts for the difference between the yields estimated against an internal standard and the isolated yields.


The utility of α -iodoketones as versatile intermediates for organic synthesis was demonstrated in a series of transformations (Schemes 3 and 4). First, the synthesis of imidazole **7** was easily accomplished by allowing an amidine to react with α -iodoketone **3e**. In a one-pot procedure starting from **1e**, iodohydrin **8** was synthesized in 59% yield with excellent diastereoselectivity (93:7). Subsequent basic treatment gave the corresponding epoxide **9** with excellent diastereoselectivity. α -Cyanoepoxide **10** was also prepared in good yield and with a good diastereomeric ratio in one step by the reaction of **3e** with KCN.

The synthetic utility of α -iodocarbonyl compounds was also demonstrated by the preparation of valuable α -aminoketones. This structural motif is found in many compounds with important biological activities.¹⁴ Although there are several procedures for the synthesis of compounds containing this motif, their regioselective preparation when two enolizable positions are present remains a challenge.^{14b-d} We found that secondary amines react with α -iodocarbonyl derivatives — prepared from allylic alcohols as described in this paper — to


give α -amino ketones as single constitutional isomers (Scheme 4).

In conclusion, we present a new mild and selective

Table 3 Substrate scope.^a




^aReactions carried out on a 1 mmol scale (0.1 M). Yield determined by ¹H NMR spectroscopy. Isolated yields in parentheses.



Scheme 3 α -Iodoketones as versatile synthetic intermediates.

electrophilic iodinating reagent, 2,2-diiododimedone (**2**). Using this reagent, we have been able to carry out the first selective synthesis of α -iodocarbonyl compounds as single constitutional isomers. The method relies on an iridium-catalyzed 1,3-hydrogen shift mediated by iridium(III) complexes. The utility of α -iodocarbonyl compounds has been demonstrated by their conversion into other important organic structures for which selective synthetic procedures have not previously been available.



Scheme 4 Regioselective synthesis of α -amino ketones.

Acknowledgements


This project was supported by the Swedish Research Council (VR), the Swedish Research Council Formas, and the Knut and Alice Wallenberg Foundation. We also thank the Swedish Governmental Agency for Innovation Systems (VINNOVA) for a VINNMER grant to B.M.-M.

Notes and references



- 1 a) M. M. Haeggblom and I. D. Bossert, *Dehalogenation*, Kluwer Academic Publishers, New York, 2003, pp 6-29; b) G. W. Gribble, *Naturally occurring organohalogen compounds - A comprehensive update*; Springer: Wien, 2010; c) G. W. Gribble, *Progress in Heterocyclic Chemistry*, 2003, **15**, 58.
- 2 a) S. Purser, P. R. Moore, S. Swallow and V. Gouverneur, *Chem. Soc. Rev.* 2008, **37**, 320; b) M. Z. Hernandez, S. M. T. Cavalcanti, D. R. M. Moreira, W. F., Jr. de Azeved and A. C. L. Leite, *Curr. Drug Targets*, 2010, **11**, 303; c) F. Adasme-Carreno, C. Muñoz-Gutierrez and J. H. Alzate-Morales, *RSC Advances*, 2016, **6**, 61837; d) S. Jiang, L. Zhang, D. Cui, Z. Yao, B. Gao, J. Lin and D. Wei, *Scientific Reports*, 2016, **6**, 34750.
- 3 P. Jeschke, W. Kraimer, U. Schirmer and M. Witschel, *Modern Methods in Crop Protection Research*, Wiley-VCH: Weinheim, 2013. Chapter 4, pp 73-119.
- 4 a) M. L. Tang and Z. Bao, *Chem. Mater.* 2011, **23**, 446; b) F. Yang, C. Li, W. Lai, A. Zhang, H. Huang and W. Li, *Mater. Chem. Front.*, 2017, Advance Article.
- 5 G. Cavallo, P. Metrengolo, R. Milani, T. Pilati, A. Priimagi, G. Resnati and G. Terraneo, *Chem. Rev.*, 2016, **116**, 2478.
- 6 a) C. Hollingworth and V. Gouverneur, *Chem. Commun.* 2012, **48**, 2929; b) T. Umemoto, *J. Fluor. Chem.* 2014, **167**, 3; c) M. G. Campbell and T. Ritter, *Chem. Rev.* 2015, **115**, 612; d) J.-D. Yang, Y. Wang, X.-S. Xue and J.-P. Cheng, *J. Org. Chem.* 2017, **82**, 4129.
- 7 a) I. Lengyel, V. Cesare and R. Stephani, *Synthetic Communications*, 1998, **28**, 1891; b) X. Wan, Z. Ma, B. Li, K. Zhang, S. Cao, S. Zhang and Z. Shi, *J. Am. Chem. Soc.* 2006, **128**, 7416; c) N. Ahlsten, A. Bermejo Gómez and B. Martín-Matute, *Angew. Chem. Int. Ed.* 2013, **52**, 6273; d) R. A. Rodriguez, C. -M. Pan, Y. Yabe, Y. Kawamata, M. D. Eastgate and P. S. Baran, *J. Am. Chem. Soc.* 2014, **136**, 6908; e) S. M. Maddox, C. J. Nalbandian, D. E. Smith and J. L. Gustafson, *Org. Lett.* 2015, **17**, 1042; f) M. Wang, Y. Zhang, T. Wang, C. Wang, D. Xue, J. Xiao, *Org. Lett.*, 2016, **18**, 1976.
- 8 a) O. A. Attanasi, S. Beretta, G. Favi, P. Filippone, G. Mele, G. Moscatelli and R. Saladino, *Org. Lett.* 2006, **8**, 4291; b) F. Mo, J. M. Yan, D. Qiu, F. Li, Y. Zhang and J. Wang, *Angew. Chem. Int. Ed.* 2010, **49**, 2028; c) A. Bermejo Gómez, E. Erbing, M. Batuecas, A. Vázquez-Romero and B. Martín-Matute, *Chem. Eur. J.* 2014, **20**, 10703; d) I. Saikia, A. J. Borah and P. Phukan, *Chem. Rev.* 2016, **116**, 6837; e) M. H. Shinde and U. A. Kshirsagar, *Green Chem.*, 2016, **18**, 1455.
- 9 a) J. Barluenga, *Pure Appl. Chem.* 1999, **71**, 431 and references therein; b) J. Barluenga, J. M. González, M. A. G. Martín, P. J. Campos and G. Asensio, *J. Org. Chem.* 1993, **58**, 2058; c) R. Johnsson, A. Meijer and U. Ellervik, *Tetrahedron*, 2005, **61**, 11657; d) X.-C. Wang, Y. Hu, S. Bonacorsi, Y. Hong, R. Burrell and J.-Q. Yu, *J. Am. Chem. Soc.* 2013, **135**, 10326; e) J. He, J. Duan, H. Shi, J. Huang, J. Huang, L. Yu, M. Zeller, A. D. Hunter, and Z. Xu, *J. Inorg. Chem.* 2014, **53**, 6837; f) D. T. Racys, S. A. I. Sharif, S. L. Pimlott and A. Sutherland, *J. Org. Chem.* 2016, **81**, 772; g) Y. Yang, L. Zhang, G.-J. Deng and H. Gong, *Scientific Reports*, 2017, **7**, 40430.
- 10 a) J. Barluenga, M. Marco-Arias, F. González-Bobes, A. Balleteros, J. M. González, *Chem. Comm.* 2004, **22**, 2616; b) S. Stavber, M. Jereb and M. Zupan, *Synthesis*, 2008, **10**, 1487; c) R. Singha, S. Dhara and J. K. Ray, *RSC Advances*, 2013, **3**, 23989. d) S. Song, X. Sun, X. Li, Y. Yuan and N. Jiao, *Org. Lett.* 2015, **17**, 2886; e) J. Iskra and S. S. Murphree, *Tetrahedron Lett.* 2017, **58**, 645.
- 11 a) N. Ahlsten, B. Martín-Matute, *Chem. Commun.* 2011, **47**, 8331; b) A. Bartoszewicz, M. Livendahl, B. Martín-Matute, *Chem. Eur. J.* 2008, **14**, 10547; c) N. Ahlsten, B. Martín-Matute, *Adv. Synth. Catal.* 2009, **351**, 2657.
- 12 Base-catalyzed isomerization; S. Martinez-Erro, A. Sanz-Marco, A. Bermejo Gómez, A. Vázquez-Romero, M. S. G. Ahlquist, B. Martín-Matute, *J. Am. Chem. Soc.* 2016, **138**, 13408.
- 13 E. Erbing, A. Vázquez-Romero, A. Bermejo Gómez, A. E. Platero-Prats, F. Carson, X. Zou, P. Tolstoy and Martín-Matute, *Chem. Eur. J.* 2016, **22**, 15659.
- 14 a) P. C. Meltzer, D. Butler, J. R. Deschamps, B. K. Madras, *J. Med. Chem.* 2006, **49**, 1420; b) R. W. Evans, J. R. Zbieg, S. Wu, W. Li, D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2013, **135**, 16074; c) S. Guha, V. Rajeshkumar, S. S. Kotha, G. Sekar, *Org. Lett.* 2015, **17**, 406; d) S. Liang, C.-C. Zeng, H.-Y. Tian, B.-G. Sun, X.-G. Luo, F.-Z. Ren, *J. Org. Chem.* 2016, **81**, 11565.



- New iodinating agent
- Easy synthesis at gram-scale
- Mild and high functional group tolerance



$[\text{Cp}^*\text{Ir}]_{\text{cat}}$



18 examples
Versatile building blocks