

The template effect of solvents on high yield synthesis, co-cyclization of pillar[6]arenes and interconversion between pillar[5]- and pillar[6]arenes†

Cite this: *Chem. Commun.*, 2014, 50, 5774Received 17th March 2014,
Accepted 7th April 2014

DOI: 10.1039/c4cc01968g

www.rsc.org/chemcomm

Tomoki Ogoshi,^{a,b} Naosuke Ueshima,^a Tomohiro Akutsu,^a Daiki Yamafuji,^a Takuya Furuta,^a Fumiyasu Sakakibara^a and Tada-aki Yamagishi^a

We have synthesized a pillar[6]arene in high yield and a co-pillar[6]arene using chlorocyclohexane as a solvent.

The properties of host–guest complexes based on macrocycles such as cyclodextrins,¹ crown ethers,² calixarenes³ and cucurbiturils⁴ have been studied extensively. The development of methods for the high yield synthesis of macrocyclic host homologues possessing different cavity sizes has extended this field of chemistry considerably, because of different host–guest properties depending on their cavity sizes. The selective synthesis of the macrocyclic homologues, however, is generally difficult by the kinetically controlled cyclization process. The use of template compounds (thermodynamically controlled cyclization process) is therefore necessary for the selective synthesis of particular homologues.

Pillar[*n*]arenes, which were first reported by our group in 2008,⁵ are some of the new macrocyclic compounds at the core of supramolecular chemistry. A cyclic pentamer, pillar[5]arene, was prepared in high yields by the condensation reaction of 1,4-dialkoxybenzene with paraformaldehyde and BF₃·OEt₂ in 1,2-dichloroethane (DCE) because DCE acts as a templating solvent. The high yield synthesis of pillar[5]arenes has led to the rapid development of the chemistry of pillar[5]arenes.^{5–7} Pillar[6]arenes, which are cyclic hexamers, have been synthesized, but their yields were not high (<45%) even under the optimized conditions because the cyclization process is kinetically controlled.^{8–10} The low yields encountered in the synthesis of pillar[6]arenes compared to pillar[5]arenes represent a bottleneck in terms of further expanding the field of pillar[6]arene chemistry. Herein, we described our research towards developing a thorough understanding of the exact role of the template effect of solvents on the synthesis of a pillar[6]arene. By the thermodynamically controlled cyclization process, we successfully synthesized




Fig. 1 (a) Synthesis of pillar[5]- and pillar[6]arenes from dialkoxybenzene monomers (M1–M4). (b) Pillar[n]arenes with tri(ethylene oxide) groups (TEO[5] *n* = 5; TEO[6] *n* = 6).

a pillar[6]arene in high yield, and a co-pillar[6]arene by the co-cyclization approach for the first time.

DCE is a suitable solvent for the synthesis of pillar[5]arenes. Pillar[5]arene P1[5] (Fig. 1) was obtained in high yield (71%) when DCE was used as the solvent with 1,4-dimethoxybenzene (M1) as the monomer (Table 1, entry 1). The cavity size of pillar[5]arenes is ca. 5.5 Å and generally fits linear alkanes containing electron-withdrawing groups.⁶ The linear dihaloalkane DCE therefore fits well into the pillar[5]arene cavity and acts as the template for the synthesis of the pillar[5]arene, indicating a thermodynamically controlled cyclization process. In contrast, the reaction of 1,4-diethoxybenzene (M2) with paraformaldehyde in the presence of BF₃·OEt₂ in chloroform (Table 1, entry 2) resulted in a mixture of pillar[5–10]arenes,¹⁰ with P2[5] and P2[6] being isolated in yields of 20 and 15%, which prevented the pillar[5]arene from forming stable complexes.⁶ This indicates that the cyclization is carried out under kinetic control. In this study, several other reaction solvents capable of fitting the pillar[6]arene cavity were also investigated for

^a Graduate School of Natural Science and Technology, Kanazawa University, Kakuma-machi, Kanazawa, 920-1192, Japan. E-mail: ogoshi@se.kanazawa-u.ac.jp
^b JST, PRESTO, 4-1-8 Honcho, Kawaguchi, Saitama, 332-0012, Japan
 † Electronic supplementary information (ESI) available: Experimental section, characterization data, DFT calculations and ¹H NMR spectra. See DOI: 10.1039/c4cc01968g

Table 1 Solvent effects

Entry	Monomer	Solvent	Pillar[n]arenes (yield%)	
			<i>n</i> = 5	<i>n</i> = 6
1 ^{6b}	M1	DCE	P1[5] (71)	P1[6] (0)
2 ⁹	M2	Chloroform	P2[5] (20)	P2[6] (15)
3 ^a	M3	DCE	P3[5] (3)	P3[6] (0)
4 ^b	M3	Chloroform	P3[5] (10)	P3[6] (8)
5 ^c	M3	CyC6	P3[5] (0)	P3[6] (0)
6 ^c	M3	Cl-CyC6	P3[5] (3)	P3[6] (87)

^a The same reaction conditions as in ref. 6b. ^b The same reaction conditions as in ref. 9. ^c Detailed experimental conditions are provided in the ESI.

the synthesis of pillar[6]arenes. The cavity size of pillar[6]arenes is greater than that of pillar[5]arenes, which is *ca.* 7.5 Å, and this large cavity size allows pillar[6]arenes to bind various bulky hydrocarbons.⁹ Based on the host-guest chemistry of pillar[6]arenes, it was envisaged that bulky hydrocarbon solvents would be ideal solvents for the synthesis of pillar[6]arenes. Thus, we evaluated the use of cyclohexane (CyC6), chlorocyclohexane (Cl-CyC6), cyclooctane and decaline as solvents. 1,4-Bis(methylcyclohexyl ether)benzene (**M3**) was used as a monomer in these reactions to overcome the poor solubility of 1,4-dialkoxybenzenes in the hydrocarbon solvents because monomer **M3** contained bulky lipophilic moieties that would enhance its solubility in the hydrocarbon solvents. We initially examined the reaction of **M3** in DCE under the same conditions used for the high yield synthesis of pillar[5]arenes (Table 1, entry 1). Although the conditions were suitable for the synthesis of pillar[5]arenes, the yield of **P3[5]** was much lower, about 3% (Table 1, entry 3). This low yield was attributed to steric hindrance of the bulky cyclohexylmethyl groups, which would have prevented the formation of **P3[5]** and also **P3[6]**. When **M3** was reacted in chloroform, **P3[5]** and **P3[6]** were formed with yields of 10 and 8%, respectively (Table 1, entry 4). Chloroform does not act as a template for the synthesis of both pillar[5]- and pillar[6]arenes,⁶ thus the reaction in chloroform resulted in the formation of **P3[5]** and **P3[6]**. This result therefore displayed a similar trend to that of the reaction of **M2** in chloroform (Table 1, entry 2). Pleasingly, when Cl-CyC6 was used as the solvent with **M3**, the pillar[6]arene **P3[6]** was isolated as the major product in 87% yield (Table 1, entry 6). This high yield represents a significant improvement over those previously reported for the synthesis of pillar[6]arenes (35–45%).⁸ This method therefore allowed for the selective and high yield synthesis of a pillar[6]arene using Cl-CyC6 as a bulky hydrocarbon solvent. Several other bulky hydrocarbon solvents were also investigated, including CyC6 (Table 1, entry 5), but did not afford any of the desired pillar[n]arene products. The use of halogenated solvents is also suitable for the synthesis of pillar[n]arenes, because the cavities of pillar[n]arenes are electron-rich spaces and can therefore readily accommodate guest molecules containing electron-withdrawing groups such as cyano and halogen moieties.⁶ The haloalkane Cl-CyC6 would therefore be included in the pillar[6]arene cavity where it would act as a template for the synthesis of a pillar[6]arene. As with pillar[6]arenes, **P3[6]** also forms a host-guest complex with a DABCO cation⁹ (Fig. S2, ESI†) and deprotection of cyclohexylmethyl groups by BBr₃ afforded per-hydroxylated pillar[6]arene in high yield (yield 95%).




Fig. 2 ¹H NMR spectra of (a) DCE, and 1:1 mixtures of DCE with (b) **TEO[5]** or (c) **TEO[6]**, (d) Cl-CyC6, and 1:1 mixtures of Cl-CyC6 with (e) **TEO[5]** or (f) **TEO[6]** at 25 °C.

In an attempt to examine the solvent template effect on the selective formation of pillar[5]- and pillar[6]arenes, we recorded the ¹H NMR spectra of mixtures of DCE and Cl-CyC6 with **P3[5]** and **P3[6]** in CDCl₃ (Fig. S3 and S4, ESI†). It was not possible, however, to detect clear peak shifts in these spectra because the complexation was too weak in CDCl₃. We recently found that pillar[5]- and pillar[6]arenes carrying tri(ethylene oxide) chains (Fig. 1b, **TEO[5]** and **TEO[6]**) exist as liquids at room temperature.¹¹ We proceeded to evaluate the bulk state host-guest complexation process by ¹H NMR with guest molecules using **TEO[5]** and **TEO[6]** as solvents. The proton signal from the ethylene moiety of DCE in the bulk state was observed at 3.64 ppm (Fig. 2a). When DCE was mixed with **TEO[5]** (Fig. 2b), the signal of the ethylene moiety underwent a significant upfield shift of 2.53 ppm. In contrast, no upfield shift was observed for this peak in a 1:1 mixture of DCE and **TEO[6]** (Fig. 2c). These results indicated that DCE formed a strong complex with **TEO[5]** in the bulk state but not with **TEO[6]**. In contrast, when Cl-CyC6 and **TEO[5]** were mixed (Fig. 2e), only small up-field shifts were observed in the proton signals belonging to Cl-CyC6. In the case of a mixture of Cl-CyC6 and **TEO[6]** (Fig. 2f), large upfield shifts were observed in the signals belonging to Cl-CyC6. Cl-CyC6 can form much stronger interaction with **TEO[6]** than that with **TEO[5]**. From these results, it can be inferred that DCE acts as a template for pillar[5]arenes but not for pillar[6]arenes, and Cl-CyC6 behaves as a template in the synthesis of pillar[6]arenes but not in the synthesis of pillar[5]arenes. DFT calculations (Fig. S5, ESI†) also supported these experimental data.

To examine the mechanism of the reaction leading to the pillar[6]arene using Cl-CyC6 (Table 1, entry 6), we monitored the size-exclusion chromatography (SEC) profiles of the products obtained from the reaction over time (Fig. 3a). After 1 min, peaks of **P3[6]** and **P3[5]** were observed with retention times of 19.0 and 19.5 min. Peaks of the corresponding oligomers were also observed in the




Fig. 3 Monitoring the reaction of entry 6 (Table 1). (a) SEC traces, (b) partial ^1H NMR spectra (CDCl_3 , 25°C) and (c) ratios of **M3** (green diamonds), pillar[5]arene **P3[5]** (blue circles), pillar[6]arene **P3[6]** (red squares) and oligomers (black triangles) determined by ^1H NMR of the reaction mixtures over time.

range of 17–20 min. As the reaction proceeded, the sizes of the peaks corresponding to **P3[5]** and the oligomers decreased, whereas the peak corresponding to **P3[6]** increased in size. After 1.5 min, we detected peaks of **P3[6]** and the oligomers. Beyond this time point, the trace remained largely unchanged. The reaction was also monitored using ^1H NMR (Fig. 3b). After 20 s, the proton signal of **M3** (green peak, 6.80 ppm) and multiple other proton signals (6.60–6.73 ppm) of the oligomers were observed. After 1 min, the spectrum contained proton signals of **P3[5]** (blue peak, 6.84 ppm), **P3[6]** (red peak, 6.75 ppm) and the oligomers. After 3 min, the proton signal of **M3** completely disappeared, with the main signal in the spectrum belonging to **P3[6]**. The integration ratios of the different species were investigated over time (Fig. 3c). At the beginning of the reaction, **M3** was converted to linear oligomers. Following the formation of these oligomers, there was a reduction in the amount of oligomers and an increase in the amount of **P3[6]** as the reaction progressed. This reaction behaviour indicated that pillar[*n*]arenes are constructed by a reversible reaction between the oligomer and the macrocyclic compounds and the reaction proceeds under thermodynamic control.

Based on the solvent template effect and the reversible reaction behaviour, we examined the “dynamic” interconversion between the pillar[5]arene and pillar[6]arene with $\text{BF}_3\cdot\text{OEt}_2$. Firstly, we examined conversion of **P3[6]** to **P3[5]** in DCE. The reaction was monitored by ^1H NMR measurements (Fig. 4a). At 25°C , the ring-opening reaction of **P3[6]** did not proceed, but, when the mixture was heated at 80°C , new peaks corresponding to **P3[5]** and the oligomers were observed. It was necessary to heat the reaction to provide the energy required to overcome the kinetic barrier to the ring-opening reaction. A decrease in the ratio of **P3[6]** together with an increase in the ratio of **P3[5]** was




Fig. 4 Interconversion between pillar[5]arene **P3[5]** and pillar[6]arene **P3[6]**. Ratios of **P3[5]** (blue circles), **P3[6]** (red squares) and the oligomers (black triangles) by cleaving (a) **P3[6]** in DCE and (b) **P3[5]** in Cl-CyC6 at 80°C , which were monitored by ^1H NMR analysis of the reaction mixtures over time.

subsequently observed as the reaction progressed, indicating that **P3[6]** was converted to **P3[5]**. We also examined the conversion of **P3[5]** to **P3[6]** in Cl-CyC6 (Fig. 4b). Heating at 80°C was also required for the cleavage of the **P3[5]** ring. As the reaction proceeded, we observed a decrease in the amount of **P3[5]** together with a simultaneous increase in the amount of **P3[6]**, even though the ratio of oligomers in the reaction mixture was high. This result indicated that **P3[5]** was converted to **P3[6]**.¹² By contrast, conversions did not take place (Fig. S5, ESI[†]) in combinations of host-guest complexes (**P3[5]** in DCE or **P3[6]** in Cl-CyC6), meaning “freezing” by inclusion of a template solvent. We can reversibly carry out interconversion using different template solvents, in which **P3[6]** can be converted to **P3[5]** using DCE, and **P3[5]** can be converted back to **P3[6]** using Cl-CyC6.

The co-cyclization method, which involves the co-cyclization of different monomers, represents a useful approach for the synthesis of functionalized pillar[5]arenes. This method, however, cannot be applied to the synthesis of functionalized pillar[6]arenes because pillar[5]arenes are formed as the major products when DCE is used as the solvent.¹³ In this study, however, we found that a pillar[6]arene was formed as the major product using Cl-CyC6. We investigated the development of a co-cyclization method for the synthesis of a co-pillar[6]arene (Fig. 5). A large amount of **M4** is found to be best




Fig. 5 Synthesis of co-pillar[6]arene by the co-cyclization of different monomers in Cl-CyC6.

for the synthesis of co-pillar[6]arene **P3[5]P4[1]**.¹⁴ A mixture of **M3** and **M4** (**M3**:**M4** = 1:16) with paraformaldehyde and BF₃·OEt₂ was stirred in Cl-CyC6 at room temperature. After purification by column chromatography, co-pillar[6]arene **P3[5]P4[1]** was isolated in 9% yield, which was higher than the yields observed for homopillar[*n*]arenes **P3[5]** (1%) and **P3[6]** (8%). The use of Cl-CyC6 as a solvent therefore allowed for the synthesis of **P3[5]P4[1]** by the co-cyclization method for the first time.

In conclusion, we demonstrated the template effect of different solvents on the formation of pillar[5]- and pillar[6]arenes, and discovered that Cl-CyC6 is a good solvent for synthesis of pillar[6]arenes. Cl-CyC6 is an inexpensive and commercially available solvent, and we therefore believe that our new procedure could be used to provide better access to pillar[6]arenes. We have also demonstrated for the first time the synthesis of co-pillar[6]arene using the co-cyclization method. The success of this solvent-templated synthesis in the construction of pillar[6]arenes suggests that a similar strategy could be applied to the selective synthesis of the other pillar[*n*]arene homologues with larger ring sizes. This is currently underway in our laboratory.

Notes and references

- 1 S. A. Nepogodiev and J. F. Stoddart, *Chem. Rev.*, 1998, **98**, 1959; A. Harada, A. Hashidzume, H. Yamaguchi and Y. Takashima, *Chem. Rev.*, 2009, **109**, 5974.
- 2 C. J. Pedersen, *J. Am. Chem. Soc.*, 1967, **89**, 7017.
- 3 C. D. Gutsche, *Acc. Chem. Res.*, 1983, **16**, 161; A. Ikeda and S. Shinkai, *Chem. Rev.*, 1997, **97**, 1713; J. Rebek Jr., *Acc. Chem. Res.*, 1999, **32**, 278.
- 4 J. W. Lee, S. Samal, N. Selvapalam, H. J. Kim and K. Kim, *Acc. Chem. Res.*, 2000, **36**, 621; J. Lagona, P. Mukhopadhyay, S. Chakrabarti and L. Isaacs, *Angew. Chem., Int. Ed.*, 2005, **44**, 4844.
- 5 T. Ogoshi, S. Kanai, S. Fujinami, T. Yamagishi and Y. Nakamoto, *J. Am. Chem. Soc.*, 2008, **130**, 5022.
- 6 (a) X. Shu, S. Chen, J. Li, Z. Chen, L. Weng, X. Jia and C. Li, *Chem. Commun.*, 2012, **48**, 2967; (b) T. Boinski and A. Szumna, *Tetrahedron*, 2012, **68**, 9419.
- 7 T. Ogoshi, *J. Inclusion Phenom. Macrocyclic Chem.*, 2012, **72**, 247; T. Ogoshi and T. Yamagishi, *Chem. Commun.*, 2014, **50**, 4776; P. J. Cragg and K. Sharma, *Chem. Soc. Rev.*, 2012, **41**, 597; M. Xue, Y. Yang, X. Chi, Z. Zhang and F. Huang, *Acc. Chem. Res.*, 2012, **45**, 1294; T. Ogoshi and T. Yamagishi, *Eur. J. Org. Chem.*, 2013, 2961.
- 8 D. Cao, Y. Kou, J. Liang, Z. Chen, L. Wang and H. Meier, *Angew. Chem., Int. Ed.*, 2009, **48**, 9721; H. Tao, D. Cao, L. Liu, Y. Kou, L. Wang and H. Meier, *Sci. China: Chem.*, 2012, **55**, 223.
- 9 T. Ogoshi, H. Kayama, D. Yamafuji, T. Aoki and T. Yamagishi, *Chem. Sci.*, 2012, **3**, 3221; W. Xia, X. Y. Hu, Y. Chen, C. Lin and L. Wang, *Chem. Commun.*, 2013, **49**, 5085.
- 10 X. B. Hu, Z. Chen, L. Chen, L. Zhang, J. L. Hou and Z. T. Li, *Chem. Commun.*, 2012, **48**, 10999.
- 11 T. Ogoshi, T. Aoki, R. Shiga, R. Iizuka, S. Ueda, K. Demachi, D. Yamafuji, H. Kayama and T. Yamagishi, *J. Am. Chem. Soc.*, 2012, **134**, 20322; T. Ogoshi, K. Kida and T. Yamagishi, *J. Am. Chem. Soc.*, 2012, **134**, 20146.
- 12 Due to the weak association constant between Cl-CyC6 and **P3[6]** at 80 °C, the efficiency of the conversion from **P3[5]** to **P3[6]** should be low.
- 13 Z. Zhang, B. Xia, C. Han, Y. Yu and F. Huang, *Org. Lett.*, 2010, **12**, 3285.
- 14 This trend is the same for the synthesis of co-pillar[5]arenes; L. Liu, D. Cao, Y. Jin, H. Tao, Y. Kou and H. Meier, *Org. Biomol. Chem.*, 2011, **9**, 7007.