ELSEVIER

Contents lists available at ScienceDirect

Journal of Molecular Structure

journal homepage: www.elsevier.com/locate/molstr

Novel molecules containing structural features of NSAIDs and 1,2,3-triazole ring: Design, synthesis and evaluation as potential cytotoxic agents

Jyoti Mareddy^a, Kazi Amirul Hossain^b, N. Sudhakar Yadav^c, Venkanna Banothu^d, Jaya Shree Anireddy^e, Sarbani Pal^{a,*}

^a Department of Chemistry, MNR Degree & PG College, Kukatpally, Hyderabad 500085, India

^b Department of Physical Chemistry, Gdansk University of Technology, Gdansk, Poland

^c VNR Vignana Jyothi Institute of Engineering and Technology, Vignana Jyothi Nagar, Pragathi Nagar, Nizampet (S.O.), Hyderabad, Telangana 500090, India

^d Department of Biotechnology, Institute of Science and Technology, Jawaharlal Nehru Technological University Hyderabad, Hyderabad 500085, India

^e Center for Chemical Sciences and Technology, Institute of Science and Technology, Jawaharlal Nehru Technological University Hyderabad, Hyderabad 500085, India

ARTICLE INFO

Article history: Received 13 May 2021 Revised 25 July 2021 Accepted 30 July 2021 Available online 2 August 2021

Keywords: NSAID

Triazole CuAAC Cytotoxicity In silico study

ABSTRACT

For the first time the template containing structural features of more than one NSAIDs and the 1,2,3triazole ring was explored for the identification of potential cytotoxic agents. These new and complex molecules were predicted to be effective inhibitors of PDE4B by molecular modelling studies *in silico*. The multi-step synthesis of these compounds were carried out starting from the well-known drug nimesulide and involved the use of copper-catalyzed azide-alkyne cycloaddition (CuAAC) approach as the key step. Mainly two types of compounds e.g. 1-aryl-1*H*-1,2,3-triazoles and *N*-aryl-2-(1*H*-1,2,3-triazol-1yl)acetamide derivatives were synthesized by using this method in good yields. The *in vitro* screening of these compounds against two cancer cell lines e.g. HCT-15 (human colon cancer cell line) and NCI-H226 (human lung cancer cell line) using a colorimetric MTT assay allowed identification of two preliminary hit molecules i.e. **8a** and **8f**. The SAR (Structure Activity Relationship) analysis indicated that the presence of an amide linker between the aryl ring and the 1,2,3-triazole moiety was favorable for the activities. The compound **8a** and **8f** showed significant inhibition of PDE4B *in vitro* and good interactions with this protein *in silico* suggesting PDE4B as their potential target. The usefulness and concerns of these molecules in the light of computational ADME prediction were analyzed. Overall, novel molecules were identified as potential cytotoxic agents for further study.

© 2021 Elsevier B.V. All rights reserved.

1. Introduction

Assembly of two or more frameworks in a single molecular entity (molecular hybridization) has become an interesting strategy in the identification of promising new chemical entities (NCEs) or bioactive agents in various therapeutic areas including cancer. Triazole being bio-isosters of amide, ester and carboxylic acid in addition to its ability to mimic various amino acids has been explored widely for this purpose [1]. Till date numerous polyfunctionalized or densely substituted molecules containing the 1,2,3-triazole framework has been reported for the identification of various bioactive compounds including the antiproliferative agents. For example, molecules based on C-14 dehydroa-

* Corresponding author. *E-mail address:* sarbani277@yahoo.com (S. Pal). bietic acid (DHAA)-1,2,3-triazole hybrid (A, Fig. 1) were synthesized and evaluated against SK-OV-3 (ovary), PC-3 (prostate), MDA-MB-231 (breast) and MCF-7 (breast) cancer cell lines [2]. Similarly, molecules based on diphenyl-1H-pyrazole-1,2,3-triazole hybrid linked through acrylate moiety (**B**, Fig. 1) were explored as apoptosis inducing cytotoxic and anti-inflammatory agents [3]. Molecules based on benzosuberone-1,2,3-triazole hybrid (C, Fig. 1) were evaluated for their anti-proliferative activities [4]. Notably, besides the presence of 1,2,3-triazole ring all these molecules contain either an amide or ester as a linker that allowed successful hybridization of different frameworks. Nevertheless, the molecular hybridization of more than two frameworks is rather uncommon in the literature. On the other hand, due to our interest in the identification of potential anti-proliferative agents we have explored the nimesulide-1,2,3-triazole hybrid (e.g. D, Fig. 2) previously [5]. In further continuation of this research we also reported the synthe-





Fig. 1. Example of polyfunctionalized or densely substituted molecules containing the 1,2,3-triazole framework.



Fig. 2. Design of new compounds E and F from the known agent D.

sis and biological evaluation of complex molecules [6-8] containing 1,2,3-triazole ring along with the framework of certain NSAIDs (Nonsteroidal anti-inflammatory drugs). Indeed, the PDE4B (Phosphodiesterase 4B) inhibition along with cytotoxic properties of a series of 2,2,4-trimethyl-1,2-dihydroquinolinyl substituted 1,2,3triazole derivatives were evaluated previously [9]. Prompted by the earlier report on dual cytotoxic and anti-inflammatory properties [3] including the PDE4 inhibition [5, 9] of 1,2,3-triazole based hybrid molecules we devoted our efforts on constructing a library of similar hybrid molecules represented by E and F as shown in Fig. 2. Indeed, the 1-aryl-1H-1,2,3-triazole based template E was designed by modifying the C-4 substituent of the 1,2,3-triazole ring of **D** whereas the design of N-aryl-2-(1H-1,2,3triazol-1-yl)acetamide based template F involved incorporation of a linker between the 1,2,3-triazole ring and the substituted benzene moiety. The group Z was mostly designed via incorporating the structural features of a range of known drugs mostly NSAIDs or analgesics individually and separately. These include mefenamic acid, naproxen, ibuprofen, tolfenamic acid, acetylsalicylic acid (aspirin), nimesulide (all NSAIDs) and acetaminophen (paracetamol) (an analgesic). We anticipated that this strategy would introduce more drug-like structural diversity into the template to afford new molecules for pharmacological studies towards the identification of potential cytotoxic agents that may inhibit PDE4. Notably, the inhibition of PDE4 (that is widely expressed in cancer cells) provides a new approach for targeting certain types of cancer. This is supported by the beneficial effects shown by PDE4 inhibitors in (i) inhibiting the brain tumor cell growth [10,11], (ii) reducing the proliferation and angiogenesis of lung cancer cell lines [12] and (iii)



Fig. 3. The 2D (H-bond presented in magenta and pi-pi in green color) and 3D interaction diagrams (where H-bond and pi-pi stacking are shown in orange and cyan dashed lines, respectively) of molecule **G** with PDE4B (PDB ID: 4MYQ) prepared using Maestro visualizer (Schrödinger, LLC).

causing selective apoptosis of malignant cells without affecting the normal cells [13]. Nevertheless, while molecules containing structural features of single NSAID and the 1,2,3-triazole ring have been studied earlier however to the best of our knowledge the template like **E** / **F** containing structural features of more than one NSAIDs and the 1,2,3-triazole ring was not explored for the identification of potential cytotoxic agents previously.

In order to gain some preliminary idea regarding the PDE4 inhibitory potential of molecules derived from **F** the *in silico* docking studies were performed using a representative molecule **G**. The protein PDE4B (PDB ID: 4MYQ) was collected from the Protein Data Bank whereas MarvinSketch [14], AutoDock tool [15] and AutoDock Vina [16] were used for the preparation of studied ligands and performing the docking study. The 2D and 3D interaction diagrams of molecule **G** with PDE4B (Fig. 3, see also Fig S-1 in suppl data) suggested that the molecule interacted well with the target protein with the docking score -10.7. Indeed, the molecule formed a H-bond with SER614 through its amidic NH group though at a distance of 2.8 Å. Additionally, it also participated in (i) a pipi interaction with HIS406 and a pi-pi stacking with PHE586 and



Scheme 1. Synthesis of 1-aryl-1H-1,2,3-triazole derivatives 5 based on E (Fig 2).



Scheme 2. Synthesis of N-aryl-2-(1H-1,2,3-triazol-1-yl)acetamide derivatives 8 based on F (Fig 2).

(ii) several hydrophobic interactions with residues ILE582, PHE678, MET603, LEU674 etc within the distance of 4 Å. All these results prompted us to undertake a straightforward synthesis of compounds based on **E** and **F**.

2. Results and discussion

The synthesis of both class of compounds i.e. compounds based on E (or the 1-aryl-1H-1,2,3-triazole derivatives 5) and that based on **F** [or *N*-aryl-2-(1*H*-1,2,3-triazol-1-yl)acetamide derivatives **8**] were involved the use of a common strategy (Schemes 1 and 2). It is well known that the construction of a 1,2,3-triazole ring is commonly performed by using the click chemistry approach that involves copper (I)-catalyzed 1,3-cycloaddition reaction between a terminal alkyne and an azide [17]. Being known as coppercatalyzed azide-alkyne cycloaddition (CuAAC) this approach has found enormous applications in organic synthesis and medicinal chemistry [1-9]. This CuAAC based approach has been used as a key step in the current synthesis of target compounds. The required azide 3 and 7 were synthesized from nimesulide (1). Thus the aniline derivative (2) obtained from 1 via reduction of its nitro group [18] was transformed into the azide **3** following the known and traditional method (Scheme 1). In another effort the aniline 2 was N-acylated using chloroacetyl chloride [19] to give the compound $\mathbf{6}$ that on treatment with KI followed by NaN₃ afforded the azide 7 (Scheme 2). These azides were separately coupled with an appropriate terminal alkyne (4) under CuAAC condi-



Scheme 3. Preparation of alkyne 4.

tions. A mixture of $CuSO_4 \cdot 5H_2O$ and sodium ascorbate was used as a pre-catalyst system in this coupling reaction to generate the Cu(I) species *in situ* that actually catalyzed the reaction. The required terminal alkynes (**4a-i**) were prepared *via* the reaction of propargyl bromide with various drugs e.g. mefenamic acid, naproxen, ibuprofen, tolfenamic acid, acetylsalicylic acid (aspirin), nimesulide, and acetaminophen (paracetamol) or compound **2** or a reduced product of naproxen in DMF at 90 °C (Scheme 3). All the terminal alkynes obtained were employed to react with the azide **3** or **7** under the CuAAC conditions separately and individually. The duration of the reaction was 5-15 min depending on the nature and type of terminal alkynes used. The reaction proceeded well in all these cases affording the desired product in good yield (Table 1). Thus a variety of compound **5** and **8** were prepared using the method described in Schemes 1 and 2.

All the synthesized compounds were characterized by spectral (NMR, IR and MS) data. For example, ¹H NMR spectra of a representative compound **5c** (Fig. 4A) showed (i) a doublet at δ 0.87 (for 2CH₃) and a quintet at δ 1.81 (for CH) due to the *i*-propyl group, (ii) a doublet at δ 2.41 due to the CH₂ protons between the *i*-propyl and phenyl group, (iii) a doublet at δ 1.48 and a quartet at δ 3.71 due to the CH₃ group attached to the benzylic carbon and the benzylic proton, (iv) one singlet at δ 3.08 due to the -SO₂CH₃ protons and (v) one singlet at δ 5.24 and δ 7.01 due to the CH₂ group attached to the triazole ring and the triazole ring proton. In the ¹³C NMR spectra, the Me of *i*-propyl group and the methanesulfonamide moiety and that attached to the benzylic carbon appeared at 22.4, 39.9 and 18.3 ppm. Further, the CH₂, OCH₂, and -O-C=O group appeared at 44.9, 57.8 and 174.6 ppm, respectively. A similar ¹H and ¹³C NMR spectral analysis can be presented for another compound 8c (Fig. 4B).

Next, the synthesized compounds were tested *in vitro* against two cancer cell lines e.g. HCT-15 (human colon cancer cell line)

Table 1

List of 1-aryl-1H-1,2,3-triazole derivatives 5 and N-aryl-2-(1H-1,2,3-triazol-1-yl)acetamide derivatives 8 synthesized.



*Figure within the bracket represents %yield obtained from the final (CuAAC) step.

and NCI-H226 (human lung cancer cell line) using a colorimetric MTT assay. The well-known anticancer agent doxorubicin was used as a reference compound in this assay. The results of active compounds that showed moderate to good effects are presented in Table 2. Among them the compound 8a was found to be most active against HCT-15 whereas the compound 8f was identified as the best active agent against NCI-H226 cell line. Notably, 8a was not so effective against NCI-H226 whereas 8f was least effective against HCT-15 indicating their selectivity towards the particular cancer cell line. Among the rest of the compounds, 5d and 5e



Fig. 4. Partial representation of 1 H and 13 C NMR spectral data of compound (A) 5c and (B) 8c.

Table 2									
The %	control g	rowth of	cancer	cells	at				
10^{-4} M concentration of test compounds.									
	1	11075.45							

Compound	HCT-15	NCI-H226	
5c	53.6	57.9	
5d	50.7	59.0	
5e	57.7	51.6	
8a	44.1	52.3	
8b	55.2	69.1	
8f	65.2	28.3	
Doxorubicin	23.1	27.6	

Values are average of three independent determinations

showed some effects against HCT-15 and NCI-H226, respectively. Form the viewpoint of SAR (Structure Activity Relationship) the Naryl-2-(1H-1,2,3-triazol-1-yl)acetamide derivatives (8) appeared as a better active class of compounds over 1-aryl-1H-1,2,3-triazole derivatives (5) suggesting that the presence of amide linker between the aryl ring and the 1,2,3-triazole moiety was favorable for the observed activities. In a preliminary assay both 8a and 8f showed good inhibition of PDE4B e.g. 67 and 78%, respectively at 10 µM concentration when tested in vitro [20] whereas the reference compound rolipram (a well-known inhibitor of PDE4) showed 90% inhibition at 10 µM in the same assay. While interaction of compound 8a (or compound G) with PDE4B in silico has been presented earlier (see Fig. 3) we performed docking studies using the compound 8f (Fig. 5, see also Fig S-2 in suppl data) to understand its interaction with this protein. The molecule 8f showed strong interactions with PDE4B which is reflected by its binding affinity (i.e. docking score -11.0). The molecule participated in several noncovalent interactions that include (i) a H-bonding with GLN456 at 2.68Å and (ii) pi-pi interactions with TYR405, HIS406, and HIS450. The molecule also participated in several van der Waals / hy-



Fig. 5. The 2D (H-bond presented in magenta and pi-pi in green color) and 3D interaction diagrams (where H-bond and pi-pi stacking are shown in orange and cyan dashed lines, respectively) of molecule **8f** with PDE4B (PDB ID: 4MYQ) prepared using Maestro visualizer (Schrödinger, LLC).

drophobic contacts within the cut-off distance of 4 Å among which the contact with PHE678 (a key residue in the CR3 region) is mention worthy. The other noticeable hydrophobic contacts of **8f** were observed (i) with the hydrophobic residues like ILE582, PHE586, PHE618, MET603, LEU674, MET519 etc and (ii) with the hydrophobic regions of polar residues e.g. SER454, ASN455, SER520, ASN567 etc. Notably, the reference compound rolipram participated in two H-bonds with GLN615 and HIS406 at a distance of 2.9 and 3.09 Å (the second one being somewhat weaker H-bond), and two aromatic pi interactions with PHE618 and PHE586 (Fig. 6, see also Fig S-3 in suppl data). Additionally, it participated in hydrophobic contacts (at a distance cut-off 4 Å) mainly with residues PHE678, TYR405, MET519, THR579, ASN567, etc.

Having identified the compound **8a** and **8f** as potential hits it was necessary to conduct some initial assessment about ADME (absorption, distribution, metabolism, and excretion) or pharmacokinetic properties of these compounds and compare with that of rolipram. Thus the computational ADME prediction of these compounds was carried out using Swiss ADME web-tool [21] and results are summarized in Table 3 (among the various descriptors only notable one are listed in the table). Because of relatively high molecular weight it was expected that unlike rolipram these molecules would not show favorable drug likeness and high GI absorption. However, this type of concerns could be found for a number of drugs like macrolide antibiotics or anticancer drug Taxol

Table 3

Computational ADME prediction of 8a, 8f and the reference compound rolipram.

Properties	Molecules		
(i) Physicochemical	8a	8f	rolipram
Molecular Weight (g/mol)	640.71	550.59	275.34
Consensus Log P ^a	4.28	2.08	2.44
Log S (ESOL) ^b	-6.75 (poorly soluble)	-4.00 (moderately soluble)	-2.90 (soluble)
(ii) Pharmacokinetics			
GI ^c absorption	Low	Low	High
BBB ^d penetration	No	No	Yes
P-gp ^e substrate	No	Yes	Yes
(iii) Drug likeness			
Lipinski rule	No; 2 violations: $MW^f > 500$, N or $O^g > 10$	No; 2 violations: Rotors ^h > 10, TPSA ⁱ > 140	No violation
Veber rule	No; 2 violations: MW>500, N or 0 > 10	No; 2 violations: Rotors $>$ 10, TPSA $>$ 140	No violation

^a Log P: Lipophilicity.

^b Log S (ESOL): water solubility, calculated by ESOL method which is a Quantitative Structure-Property Relationship (QSPR) based model.

^c GI: Gastrointestinal.

^d BBB: Blood Brain Barrier

^e P-gp: permeability glycoprotein.

f MW: molecular weight,

g N or O: Nitrogen/Oxygen indicates number of H-bond acceptor,

^h Rotors: total number of rotatable bonds,

ⁱ TPSA: Total polar surface area.





Fig. 6. 2D followed by 3D interaction diagram of reference compound rolipram with PDE4B (PDB ID: 4MYQ).

that still entered into the market and are currently in patient's use. Moreover, both **8a** and **8f** were predicted to be a non-penetrant of BBB. Finally, a non-Pgp substrate was predicted for **8a** though not for **8f**.

3. Conclusions

In conclusion, novel molecules possessing structural features of more than one NSAIDs and the 1.2.3-triazole ring was explored for the identification of potential cytotoxic agents. These new and complex molecules were predicted to be effective inhibitors of PDE4B by molecular modelling studies in silico. The multi-step synthesis of these compounds were carried out starting from the wellknown anti-inflammatory drug nimesulide and involved the use of copper-catalyzed azide-alkyne cycloaddition (CuAAC) approach as the key step. Mainly two types of compounds e.g. 1-aryl-1H-1,2,3triazoles and N-aryl-2-(1H-1,2,3-triazol-1-yl)acetamide derivatives were synthesized by using this method in good yields. The in vitro screening of these compounds against two cancer cell lines e.g. HCT-15 (human colon cancer cell line) and NCI-H226 (human lung cancer cell line) using a colorimetric MTT assay allowed identification of two preliminary hit molecules i.e. 8a and 8f. The SAR (Structure Activity Relationship) analysis indicated that the presence of an amide linker between the aryl ring and the 1,2,3-triazole moiety was favorable for the activities. The compound 8a and 8f showed significant inhibition of PDE4B in vitro and good interactions with this protein in silico suggesting PDE4B as their potential target. The usefulness and concerns of these molecules in the light of computational ADME prediction were analysed and discussed. Overall, the current research involving the synthesis and identification of novel molecules as potential cytotoxic agents for further study would attract interest.

Author statement

Jyoti Mareddy was involved in the preparation, isolation, purification and characterization of all the target compounds presented in the current manuscript.

Kazi Amirul Hossain, N. Sudhakar Yadav and Venkanna Banothu were involved in performing all the *in silico* studies as well as *in vitro* assays.

Jaya Shree Anireddy and Sarbani Pal were responsible for conceptualization, coordination and overall supervision of the entire work presented in the submitted manuscript.

Supplementary data

Supplementary data associated with this article can be found, in the on line version, at xxxxxxxx

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgement

The authors thank the management of MNR Degree and PG College, Hyderabad, India for continuous support and encouragement.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.molstruc.2021.131222.

References

- S. Kumar, B. Sharma, V. Mehra, V. Kumar, Recent accomplishments on the synthetic/biological facets of pharmacologically active 1*H*-1,2,3-triazoles, Eur. J. Med. Chem. 212 (2021) 113069.
- [2] W. Hou, Z. Luo, G. Zhang, D. Cao, D. Li, H. Ruan, B.H. Ruan, L. Su, H. Xu, Click chemistry-based synthesis and anticancer activity evaluation of novel C-14 1,2,3-triazole dehydroabietic acid hybrids, Eur. J. Med. Chem. 138 (2017) 1042–1052.
- [3] M.F. Khan, T. Anwer, A. Bakht, G. Verma, W. Akhtar, M.M. Alam, M.A. Rizvi, M. Akhter, M. Shaquiquzzam, Unveiling novel diphenyl-1H-pyrazole based acrylates tethered to 1,2,3-triazole as promising apoptosis inducing cytotoxic and anti-inflammatory agents, Bioorg. Chem. 87 (2019) 667–678.
- [4] C.B.P Devi, K Vijay, B.H Babu, S Farooq Adil, M.M Alam, M Vijjulatha, M.B Ansari, CuSO₄/sodium ascorbate catalysed synthesis of benzosuberone and 1,2,3-triazole conjugates: design, synthesis and *in vitro* anti-proliferative activity, J. Saudi Chem. Soc. 23 (2019) 980–991.
- [5] J. Mareddy, N. Suresh, C.G. Kumar, R.K. Kapavarapu, A. Jayasree, S Pal, 1,2,3-Triazole-nimesulide hybrid: their design, synthesis and evaluation as potential anticancer agents, Bioorg. Med. Chem. Lett 27 (2017) 518–523.
- [6] S. Srinivas, P. Neeraja, V. Banothu, P.K. Dubey, K. Mukkanti, S. Pal, Synthesis, biological evaluation and docking study of a series of 1,4-disubstituted 1,2,3-triazole derivatives with an indole-triazole-peptide conjugate, J. Heterocycl. Chem. 57 (2020) 3126–3141 DOI: 10.1002/jhet.4020.
- [7] S. Srinivas, P. Neeraja, K. Naveen, V. Banothu, P.K. Dubey, K. Mukkanti, S. Pal, Synthesis, chemotherapeutic screening and docking studies of NSAID inserted peptide-triazole hybrid molecules, Chem. Select 5 (2020) 6786–6791 doi.org/10.1002/slct.202000492.

- [8] S. Srinivas, P. Neeraja, V. Banothu, P.K. Dubey, K. Mukkanti, S Pal, Synthesis, biological evaluation and *in silico* studies of compounds based on tryptophan-naproxen-triazole hybrids, Chem. Select 5 (2020) 14741–14746 doi.org/10.1002/slct.202003786.
- [9] K.S.S. Praveena, S. Durgadas, N.S. Babu, S. Akkenapally, C.G. Kumar, G.S. Deora, N.Y.S. Murthy, K. Mukkanti, S. Pal, Synthesis of 2,2,4-trimethyl-1,2-dihydroquinolinyl substituted 1,2,3-triazole derivatives: their evaluation as potential PDE 4B inhibitors possessing cytotoxic properties against cancer cells, Bioorg. Chem. 53 (2014) 8–14.
- [10] P. Goldhoff, N.M. Warrington, D.D. Limbrick, J.A. Hope, B.M. Woerner, E. Jackson, A. Perry, D. Piwnica-Worms, J.B. Rubin, Targeted inhibition of cyclic AMP phosphodiesterase-4 promotesbrain tumor regression, Clin. Cancer Res. 14 (2008) 7717–7725.
- [11] L. Yang, E. Jackson, B.M. Woerner, A. Perry, D. Piwnica-Worms, J.B. Rubin, Blocking CXCR4-mediated cyclic AMP suppression inhibits brain tumor growth *in vivo*, Cancer Res. 67 (2007) 651–658.
- [12] S.S. Pullamsetti, G.A. Banat, M. Szibor, D. Pomagruk, J. Hänze, E. Kolosionek, J. Wilhelm, T. Braun, F. Grimminger, W. Seeger, R.T. Schermuly, R. Savai, Phosphodiesterase-4 promotes proliferation and angiogenesis of lung cancer by crosstalk with HIF, Oncogene 32 (2013) 1121–1134.
- [13] F. Mentz, M.D. Mossalayi, F. Ouaaz, S. Baudet, F. Issaly, S. Ktorza, M. Semichon, J.L. Binet, H. Merle-Beral, Theophylline synergizes with chlorambucil in inducing apoptosis of B- chronic lymphocytic leukemia cells, Blood 88 (1996) 2172–2182.
- [14] P. Csizmadia, MarvinSketch and MarvinView: molecule applets for the World Wide Web, in: Proceedings of ECSOC-3, the Third International Electronic Conference on Synthetic Organic Chemistry, 1999, pp. 367–369. September 1-30.
- [15] R. Huey, G.M. Morris, AutoDock Tools, The Scripps Research Institute, La Jolla, CA, USA, 2003.
- [16] O. Trott, A.J. Olson, AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading, J. Comput. Chem. 31 (2010) 455–461.
- [17] H.C. Kolb, K.B. Sharpless, The growing impact of click chemistry on drug discovery, Drug Discov. Today 8 (2003) 1128–1137.
- [18] S. Pericherla, J. Mareddy, D.P.G. Rani, P.V. Gollapudi, S. Pal, Chemical modifications of nimesulide, J. Braz. Chem. Soc. 18 (2007) 384–390.
- [19] K. Kankanala, V.R. Reddy, Y.P. Devi, L.N. Mangamoori, K. Mukkanti, S. Pal, Nimesulide based novel glycolamide esters: their design, synthesis, and pharmacological evaluation, J. Chem. (2013) 8 Article ID 816769pages.
- [20] P. Wang, J.G. Myers, P. Wu, B. Cheewatrakoolpong, R.W. Egan, M.M. Billah, Expression, purification, and characterization of human cAMP-specific phosphodiesterase (PDE4) subtypes A, B, C, and D, Biochem. Biophys. Res. Commun. 234 (1997) 320–324.
- [21] A. Daina, O. Michielin, V. Zoete, SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules, Sci. Rep. 7 (2017) Article number: 42717. https://doi.org/10.1038/srep42717.