

View Article Online View Journal

ChemComm

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: L. Sanchez, J. S. Valera, J. Calbo, R. Gómez and E. Orti, *Chem. Commun.*, 2015, DOI: 10.1039/C5CC03616J.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/chemcomm

Published on 13 May 2015. Downloaded by Freie Universitaet Berlin on 13/05/2015 16:49:35.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxx

Blue-emitting pyrene-based aggregates

Jorge S. Valera,^{a‡} Joaquín Calbo,^{b‡} Rafael Gómez,^a Enrique Ortí^{*b} and Luis Sánchez*^a

Received (in XXX, XXX) Xth XXXXXXXX 200X, Accepted Xth XXXXXXXX 200X DOI: 10.1039/b000000x

⁵ The supramolecular polymerization of pyreneimidazoles 1 and 2, governed by H-bonding and C-H $\cdots \pi$ interactions, yields aggregates showing the characteristic blueish emission pattern of pyrene-based monomers.

Since its discovery in 1837 by Laurent,¹ pyrene has become 10 one of the most studied polycyclic aromatic hydrocarbons due to its attractive electronic and photophysical properties, and it has been widely tested as an active component in organic electronics.^{2,3} Pyrene is a blue-emitting material with limited applicability in optoelectronic devices due to its strong tendency 15 to form excimers both in solution and in the solid state. The formation of excimers red-shifts the emission of pyrene and reduces its intensity.^{2,3} To improve the emissive properties, big efforts have been devoted to the structural modification of pyrene to avoid aggregation and to suppress the quenching effect of the $_{20}$ π - π interactions. The optimization of the functional unit of pyrene and the control of its supramolecular organization are essential requirements to achieve high-performance in organic electronics devices. Therefore, a challenging task in pyrene chemistry is the design of derivatives in which supramolecular order and 25 appropriate blue-emitting features can coexist.

Herein, we describe the synthesis and the supramolecular polymerization of the pyreneimidazoles **1** and **2** endowed with peripheral achiral and chiral alkyl chains, respectively (Fig. 1). Using a synergistic experimental and theoretical investigation, a ³⁰ detailed insight into the supramolecular polymerization mechanism is presented, what is unprecedented for pyrene-based molecules despite the number of reports that refer to the self-

- assembly of pyrene derivatives.⁴ 1 and 2 have been designed with a variety of supramolecular interaction motifs that guide their self-assembly by H-bonding interactions between amide groups and imidazole moieties together with a number of C-H… π dispersion interactions. The organized supramolecular structures
- are able to immobilize apolar solvents like cyclohexane forming gels.⁵ Notably, the aggregates formed by **1** and **2** show the ⁴⁰ characteristic blue-emitting pattern of pyrene-based monomers.



Fig. 1 Chemical structure of pyreneimidazoles 1 and 2

The preparation of **1** and **2** was accomplished by the methodology reported by Kumar et al.,⁶ starting from pyrene, 4-45 carboxybenzaldehyde and the corresponding amine (Scheme S1, ESI). Prior to the synthesis of the final derivatives, two *tert*-butyl groups were incorporated at positions 2 and 7 of the pyrene moiety to increase the solubility of the target compounds. The chemical structure of **1** and **2** was confirmed by NMR, FT-IR and 50 HRMS analysis (see the ESI[†]).

To unravel the supramolecular polymerization mechanism of **1** and **2**, temperature-dependent UV-Vis spectroscopy measurements were performed in methylcyclohexane (MCH) as solvent (Fig. S1 and S2, ESI[†]). The bands observed at around 355 ⁵⁵ and 400 nm in the UV-Vis spectra of both **1** and **2** experience a slight bathochromic shift and a broadening when decreasing the temperature, what is diagnostic of the aggregation process. Plotting the variation of the absorbance at 404 nm with the temperature results in cooling curves with an apparent sigmoidal ⁶⁰ shape, characteristic of an isodesmic supramolecular polymerization mechanism (Fig. S1b and S2b, ESI[†]).⁷

To obtain a complete set of thermodynamic parameters, the UV-Vis cooling curves derived for 1 and 2 were fitted to the model recently reported by ten Eikelder et al. (EQ model), in ⁶⁵ which the equilibria between monomers and supramolecular polymers are considered, together with a modified version of the van't Hoff analysis.⁸ The thermodynamic parameters obtained for 1 and 2 demonstrate that the branched nature of the peripheral alkyl chain in 2 plays a relevant role, showing significantly ⁷⁰ different values of the enthalpy of elongation (ΔH_e), the entropy of elongation (ΔS) and the nucleation penalty (ΔH_n) (Table 1). **Table 1.** Thermodynamic parameters for 1 and 2 (MCH).

	1	2		
$\Delta H_{\rm e} [{\rm kJ/mol}]$	-72±1	-52±1		
ΔS [J/Kmol]	-122±3	-58±2		
$\Delta H_{\rm n} [{\rm kJ/mol}]^a$	-7.1±0.2	-11.5±0.2		
$\Delta G [kJ/mol]^a$	-35.88	-34.73		
$\sigma[-]^b$	5.5×10^{-2}	9.3×10^{-3}		
$K_{\rm e} \left[{\rm L/mol} \right]^b$	$2.04 imes 10^6$	1.5×10^6		
$K_{\rm n} [\rm L/mol]^b$	1.13×10^{5}	1.21×10^{4}		

^a The nucleation penalty ΔH_n is negative, which implies that the enthalpy gain is smaller for nucleation compared to elongation. ^b The equilibrium constants 75 for elongation (K_e) and dimerization (K_n), and the cooperativity factor ($\sigma = K_n / K_e$) are calculated at 298 K.

In contrast, the total free energy (ΔG) is very similar in both pyrene derivatives (~35 kJ/mol). Especially interesting is the calculated value of the degree of cooperativity σ , which so corresponds to the quotient between the nucleation (K_n) and the elongation (K_e) binding constant values. Despite the sigmoidal shape of the cooling curves, σ presents values of 5.5 × 10⁻² and 9.3 × 10⁻³ for compounds 1 and 2, respectively, which denote a certain degree of cooperativity in the supramolecular polymerization of these pyrene derivatives (Table 1).

⁵ Interestingly, the self-assembling properties of compounds **1** and **2** are similar and both readily form gels in cyclohexane as solvent at a concentration of *ca*. 1 wt%. The organogels exhibit blue emission upon irradiation at 364 nm (Fig. 2).

The emission properties of 1 and 2 were investigated in CHCl₃ ¹⁰ and MCH. The emission spectra in MCH at 1×10^{-5} M, conditions under which 1 and 2 are aggregated, show the typical pattern of monomeric pyrene with maxima at 400, 423 and 447 nm (Fig. 2 and Fig. S3, ESI[†]). In contrast, the emission spectra in CHCl₃, a good solvent that facilitates the solvation of isolated 15 molecules, at 1×10^{-5} M feature broad and structureless bands centred at 448 nm, evidencing the formation of pyrene excimers (Figure 2 and S3).^{9,10} Compounds 1 and 2 show a remarkable fluorescence quantum yield (ϕ_f) in the aggregated state (MCH, 1 $\times 10^{-5}$ M) of 0.43. A lower ϕ_f value is measured for the excimer 20 formed in CHCl₃ (0.38 and 0.22 for 1 and 2, respectively). To the best of our knowledge, this unusual emissive behaviour has only been described for a pyrene derivative endowed with a chiral oligo(glutamic acid) side chain,¹¹ and it was ascribed to a helical supramolecular organization.



Published on 13 May 2015. Downloaded by Freie Universitaet Berlin on 13/05/2015 16:49:35.

Fig. 2 a) Emission spectra of 1 in MCH and CHCl₃ (298 K, 1×10^{-5} M, $\lambda_{exc} = 334$ nm). The inset shows a picture of the cyclohexane gels obtained from compounds 1 and 2 with illumination at 364 nm.

The presence of the sterogenic center on the side chain of 2 ³⁰ was utilized to put in evidence the possible helical organization of the aggregates formed by 1 and 2 in an attempt to justify the observed emission properties. Compound 2 shows a weak dichroic response with a poorly defined bisignated Cotton effect centered at around 250 nm (Fig. S5a, ESI†). In addition, ³⁵ sergeants-and-soldiers (SaS) experiments were performed by mixing achiral 1 and chiral 2 to investigate the amplification of chirality.¹² The SaS experiment shows a linear increase of the dichroic response upon increasing the amount of chiral sergeant 2 (Fig. S5b, ESI†). The dichroic data reveal that compounds 1 and ⁴⁰ 2 do not self-assemble forming helical structures and, therefore,

this supramolecular organization cannot be invoked to justify the observed emission features.

The extracted experimental data arise two important questions: 1) which are the non-covalent forces participating in the self-45 assembly of the investigated pyreneimidazoles?, and 2) what kind of supramolecular aggregate is formed upon their self-assembly

capable to exhibit the above-mentioned emissive features? To give an appropriate answer to these questions, a synergistic experimental and theoretical study was performed. First, FTIR ⁵⁰ and ¹H NMR spectroscopies were used. In the FTIR spectra of **1** and 2, the Amide I and Amide II bands at around 1635 and 1560 cm⁻¹, diagnostic of the formation of H-bonds between the amide, are visible (Fig. S6, ESI[†]).¹³ In the concentration-dependent ¹H NMR experiments, the triplet at $\delta \sim 6.3$, corresponding to the 55 amide groups, and the resonance corresponding to the imidazole N-H group ($\delta \sim 13$) show a downfield shift with increasing concentrations that is indicative of the formation of an array of Hbonds (Fig. S7, ESI⁺). On the other hand, the aromatic resonances, used to envisage the intermolecular interactions ⁶⁰ between the aromatic fragments, show an opposite shifting of the δ values upon increasing the concentration. Thus, whilst the resonance of pyrene at $\delta \sim 9$ experiences a shielding effect, one of the anisochronous protons at the *para* position of the benzamide moiety ($\delta \sim 8.2$) shifts to lower δ values. A rotating-frame 65 Overhauser effect spectroscopy (ROESY) NMR experiment in a concentrated solution of 1 (45 \times 10⁻³ M, CDCl₃, 298 K) was therefore conducted to extract more information about the supramolecular organization of this pyrene derivative.¹⁴



⁷⁰ Fig. 3 ROESY NMR spectrum (CDCl₃, 300 MHz, 45 \times 10⁻³ M, 293 K) of **1**. The dotted blue, green and black lines highlight intra-, intermolecular or both through-space coupling signals, respectively. The inset shows the chemical structure of **1** depicting the anisochronous protons by coloured circles.

The ROESY experiment clearly shows a number of intramolecular through-space coupling signals between the protons spatially close within the molecule, like for instance both *para* protons of the benzamide moiety (see the dotted blue circles in Fig. 3). In addition, ROESY signals corresponding to ⁸⁰ intermolecular contacts are also observed (dotted green lines in Fig. 3). Especially interesting are the contacts observed between the methylene units of the alkyl side chain and most of the aromatic protons of the pyrene unit. These contacts suggest an alternated distribution of the pyrene units in the aggregated state ⁸⁵ that could justify the observed monomeric emission properties.

Theoretical calculations were performed on compound 1 within the density functional theory (DFT) framework to shed light on the structural and energetic aspects of the

Published on 13 May 2015. Downloaded by Freie Universitaet Berlin on 13/05/2015 16:49:35.

pyreneimidazole system upon aggregation (see the ESI[†] for computational details). The isolated molecule shows a planar minimum-energy structure with an out-of-plane disposition of the amide group (Fig. S8, ESI[†]) that facilitates H-bonding s interactions between neighbouring molecules. To unravel the

possible contribution of the different parts of the molecule in the operation of non-covalent forces, the molecular electrostatic potential (MEP) calculated for **1** (Fig. S9, ESI[†]) was analysed. The MEP surface shows that the largest potential energy points ¹⁰ are concentrated in the amide and the imidazole groups. Negative potential energies of -55.7 and -54.6 kcal/mol are calculated in the oxygen and nitrogen vicinity, respectively, whereas positive potentials of +101.2 and +121.0 kcal/mol are located in the N–H region of the amide and imidazole, respectively. Strong ¹⁵ electrostatic interactions are then expected between monomers by means of H-bonding interactions involving the amide and imidazole units, as well as weak π - π and/or C-H \cdots π dispersion interactions between the aromatic units and the aliphatic chains.

A number of dimers, as the minimal supramolecular unit, can ²⁰ be modelled for **1** owing to the large variety of supramolecular interaction motifs present in the monomer: (i) the π -extended aromatic structure, (ii) the polar amide and imidazole groups and (iii) the nonpolar long aliphatic chain. In a first approach, two monomers may accommodate either in a parallel or antiparallel ²⁵ disposition by the π -stacking of the pyreneimidazole cores with themselves or with the benzamide group, respectively (dimers A and B in Fig. 4). In dimer A, an amide-amide N-H...O bond is formed due to the rotation of $\sim 30^{\circ}$ of the upper monomer around the perpendicular axis. This structure would grow up in a helical 30 column upon self-assembly, but this possibility has been discarded by CD measurements. Similar to A, dimer C is arranged in a parallel mode but with the amide groups pointing in opposite directions and forming a non-linear H-bond (Fig. 4). A weak H-bonding N-H...N interaction between the imidazole 35 rings is also present in antiparallel dimer B.



Fig. 4 Minimum-energy geometries calculated for the different dimers postulated for 1 at the B97D3/6-31G** level of theory.

Besides the π - π recognition motif that gives rise to structures ⁴⁰ A–C, the polar imizadole and amide groups constitute high potential energy regions for the supramolecular stabilization of the dimers. Thereby, dimers D–F, in which the formation of strong H-bonds is the recognition driving force, have been modelled (Fig. 4). Whereas dimers D and E are only stabilized by ⁴⁵ the H-bond interaction between the imidazole or the amide groups, respectively, dimer F forms two H-bonds between the amide and the imidazole groups together with an extended CH… π stabilization due to the interaction between the aliphatic chains and the pyreneimidazole cores.

Table 2 summarizes the interaction energy (E_{int}) and the main intermolecular forces governing the self-assembly of dimers A–F calculated at the B97D3/6-31G** level. A topological analysis of the electron density was also carried out by means of the NCIPLOT software to obtain the non-covalent interaction (NCI) ⁵⁵ surfaces, which provide additional information on the origin of the supramolecular aggregation (Fig. S10 and S11, ESI⁺).

Table 2 Interaction energy (E_{int}) , number of H-bond contacts (n_{HB}) , distance of the closest H-bond contact (d_{HB}) and dominant dispersion interactions $(\pi$ - π , CH \cdots π and CH \cdots CH) that ⁶⁰ characterize dimers A–F.

	Eint (kcal/mol)	n _{HB}	$d_{\mathrm{HB}}(\mathrm{\AA})$	π-π	$CH{\cdots}\pi$	СН⋯СН
А	-54.6	1	1.918	~	-	~
В	-51.0	1	2.064	~	_	_
С	-53.7	1	2.051	~	-	-
D	-32.0	1	1.826	_	~	-
Е	-7.7	2	1.822	-	-	-
F	-61.7	2	1.839	-	~	-

Dimers A, B and C are computed very close in energy owing to the similar number and nature of the interactions stabilizing the aggregate. Dimers B and C (E_{int} of -51.0 and -53.7 kcal/mol, respectively) display the typical green surface indicative of dispersion interactions along the π - π skeleton between monomers (Fig. S10, ESI[†]). Stabilizing CH···CH interactions between the aliphatic chains are additionally found for A, which is calculated slightly more stable ($E_{int} = -54.6$ kcal/mol) than B and C. In these three dimers, weak H-bonds are formed as indicated by the 70 closest NH···N/O contacts in the range of 1.92–2.06 Å (Table 2).

The situation is different for dimers D, E and F. Dimer E is stabilized by two H-bonding interactions between the two amides in a *cis* disposition. The occurrence of dimer E is however very unlikely because of: (i) the high barrier (15 kcal/mol) that has to 75 be overcome to convert a trans amide into a cis amide (Fig. S12, ESI [†]), and (ii) the small interaction energy (-7.7 kcal/mol) resulting from the lower stability of the cis conformation and the additional steric interactions that it introduces. The cross shape of dimer D enables a strong H-bond interaction between the ⁸⁰ imidazole groups (N···H-N distance of 1.83 Å) and incorporates C-H··· π interactions between the *tert*-butyl groups of one monomer and the aromatic pyrene core of the other (Fig. S10, ESI[†]). Nonetheless, the E_{int} computed for this dimer (-32.0 kcal/mol) cannot compete with the stabilization obtained for the 85π - π patterned dimers A–C. Finally, dimer F combines H-bonding interactions between the imidazole and amide groups and C-H $\cdots\pi$ stabilizing interactions between the alkyl chains and the pyrene cores (Fig. 4 and Fig. S10, ESI⁺). The additive effect of these interactions leads to the highest stability ($E_{int} = -61.7$ kcal/mol) 90 among all the computed dimers. The structure of dimer F allows to identify all the ROESY signals observed experimentally and, in particular, the short contacts between the alkyl chains and the pyrene cores (Fig. S13, ESI[†]). In addition, the absence of π - π interactions between the pyrene cores in dimer F explains the

emission properties similar to the monomer. Dimer F is therefore proposed as the most plausible associate for compounds 1 and 2.

- To rationalize how dimer F can grow up from such nonplanar, distorted disposition of the monomers, the tetramer shown in Fig 5 5 was built up through an imidazole-amide-imidazole H-bond sequence. The minimum-energy geometry computed at the B97D3/6-31G** level for the tetramer clearly reveals that the Hbonding motifs are preserved along the different monomeric pairs. The average NH(amide)-N(imizadole) contact is computed ¹⁰ at 2.18 Å, whereas the CO(amide)–HN(imizadole) interaction is
- calculated at 1.82 Å. The total interaction energy in the tetramer reaches -182.1 kcal/mol, which corresponds with an $E_{int} = -60.7$ kcal/mol per monomer pair. This value is very close to the E_{int} computed for dimer F (-61.7 kcal/mol), suggesting that: (i) the 15 oligomer can easily grow keeping the strong H-bond pattern and the stabilizing C-H $\cdots\pi$ forces, and (ii) the degree of cooperativity

a) b)

Fig. 5. Top (a) and side (b) views of the B97D3/6-31G**-optimized 20 geometry calculated for the tetramer of 1. Dotted lines emphasise the Hbonding array established between the imidazole and amide units. Only hydrogen atoms involved in H-bonding interactions are shown.

In summary, we have synthesised pyreneimidazoles 1 and 2 that prevents the formation of excimers upon aggregation. The 25 number and type of non-covalent interactions operating in the aggregation of these pyrene derivatives results in an amazing scenario of possibilities for 1 and 2 to self-assemble. Both experimental and theoretical studies demonstrate that the

- formation of the aggregates is mainly governed by the H-bonding 30 interaction between the amide functionality and the imidazole moiety, together with a number of C-H $\cdots\pi$ interactions. The absence of π - π interactions between the pyrene cores determines that the aggregate preserves the blue-emitting properties of the monomer. The results presented herein demonstrate the complex
- 35 pathway followed by 1 and 2 to finally yield blue-emitting supramolecular aggregates and contribute to shed light into the relationship between functionality and supramolecular polymerization.

Financial support by the MINECO of Spain (CTQ2011-22581 40 and CTQ2012-31914), European Feder funds (CTQ2012-31914), UCM (UCM-SCH-PR34/07-15826), Generalitat Valenciana

(PROMETEO/2012/053) and Comunidad de Madrid (NanoBIOSOMA, S2013/MIT-2807) is acknowledged. JC thanks the MECD for a doctoral FPU grant.

45 Notes and references

^aDepartamento de Química Orgánica, Facultad de Ciencias Químicas, Universidad Complutense de Madrid, 28040 Madrid (Spain); E-mail: lusamar@guim.ucm.es

- ^bInstituto de Ciencia Molecular, Universidad de Valencia, 46980 Paterna 50 (Spain); E-mail: enrique.orti@uv.es
 - † Electronic Supplementary Information (ESI) available: Figures S1-S12, theoretical computational details and experimental section. See DOI: 10 1039/b00000x/
- [‡] These authors contributed equally
- A. Laurent, Ann. Chim. Phys. 1837, 66, 136. 55 1
- 2 C. Wang, H. Dong, W. Hu, Y. Liu and D. Zhu, Chem Rev. 2012, 112, 2208
- 3 T. M. Figueira-Duarte and K. Müllen, Chem. Rev. 2011, 111, 7260.
- 4 (a) N. Tchebotareva, X. M. Yin, M. D. Watson, P. Samori, J. P. Rabe and K. Müllen, J. Am. Chem. Soc. 2003, 125, 9734; (b) D. Venkataraman, S.; Lee, J. S. Zhang and J. S.; Moore, Nature 1994, 371. 591.
- 5 (a) A. Ajayaghosh, V. K. Praveen and C. Vijayakumar, Chem. Soc. Rev., 2008, 37, 109; (c) J. W. Steed, Chem. Soc. Rev., 2010, 46, 3686.
- D. Kumar, K. R. J. Thomas, C.-C. Lin and J.-H. Jou, Chem. 65 6 Asian J. 2013, 8, 2111.
- 7 T. F. A. De Greef, M. M. J. Smulders, M. Wolffs, A. P. H. J. Schenning, R. P. Sijbesma and E. W. Meijer, Chem. Rev. 2009, 109, 5687
- H. M. M. ten Eikelder, A. J. Markvoort, T. F. A. de Greef and P. A. J. 70 8 Hilbers, J. Phys. Chem. B 2012, 116, 5291.
- 9 F. M. Winnik, Chem. Rev. 1993, 93, 587.
- The emision corresponding to the pyrene monomer is only reached in 10 CHCl₃ by using highly diluted (7×10^{-9} M), deoxygenated solutions (Fig. S4, ESI[†]).
- 11 Y. Kamikawa and T. Kato, Langmuir, 2007, 23, 274.
- 12 (a) M. M. J. Smulders, A. P. H. J. Schenning and E. W. Meijer, J. Am. Chem. Soc., 2008, 130, 606; (b) F. García, P. A. Korevaar, A. Verlee, E. W. Meijer, A. R. A. Palmans and L. Sánchez, Chem. Commun. 2013. 49, 8674.
- 13 (a) Y. Matsunaga, N. Miyajima, Y. Nakayasu, S. Sakai and M. Yonenaga, Bull. Chem. Soc. Jpn. 1988, 61, 207; (b) F. García, P. M. Viruela, E. Matesanz, E. Ortí and L. Sánchez, Chem. Eur. J. 2011, 17, 7755
- 85 14 Prior to this ROESY experiment, the aggregation of 1 was assessed by diffusion-ordered spectroscopy (DOSY) NMR, which reveals a larger value of the diffusion coefficient when compared to a more diluted solution (Fig. S6b, ESI[†]). A. Wong, R. Ida, L. Spindler and G. Wu, J. Am. Chem. Soc. 2005, 127, 6990.

4 | Journal Name, [year], [vol], oo-oo



Published on 13 May 2015. Downloaded by Freie Universitaet Berlin on 13/05/2015 16:49:35.

15

Published on 13 May 2015. Downloaded by Freie Universitaet Berlin on 13/05/2015 16:49:35.

Blue-emitting pyrene-based aggregates

J. S. Valera, ^a J. Calbo, ^b R. Gómez, ^a E. Ortí, ^{b*} and L. $_{\rm 5}$ Sánchez^{a*}

The self-assembling features and gel formation of two pyreneimidazoles is presented. The supramolecular aggregation of these synthesised molecules results in an unusual blue-monomeric emission, which is rationalized by a ¹⁰ combined experimental and theoretical investigation.

