

# Phospha-Fischer Carbenes: Synthesis, Structure, Bonding, and Reactions of Pd(0)– and Pt(0)–Phosphenium Complexes

Christine A. Caputo,<sup>†</sup> Michael C. Jennings,<sup>†</sup> Heikki M. Tuononen,<sup>‡</sup> and Nathan D. Jones<sup>\*,†</sup>

Department of Chemistry, The University of Western Ontario, London, Ontario, Canada N6A 5B7, and  
Department of Chemistry, University of Jyväskylä, P.O. Box 35, FI-40014 Jyväskylä, Finland

Received October 9, 2008

The analogy between cationic group 10 metal–phosphenium complexes and Fischer carbenes has been formalized through structural and reactivity studies and by energy decomposition analysis (EDA) of the M–P bond. The studied compounds were the three-coordinate, 16-electron species [(NHP<sup>Mes</sup>)M(PPh<sub>3</sub>)<sub>2</sub>]OTf (M = Pt (**1**) and Pd (**2**); [NHP<sup>Mes</sup>]<sup>+</sup> is the N-heterocyclic phosphenium (NHP) cation, [PN(2,4,6-Me<sub>3</sub>-C<sub>6</sub>H<sub>2</sub>)CH<sub>2</sub>CH<sub>2</sub>N(2,4,6-Me<sub>3</sub>-C<sub>6</sub>H<sub>2</sub>)]<sup>+</sup>, OTf = trifluoromethanesulfonate); these were made by reaction of [NHP<sup>Mes</sup>]OTf with M(PPh<sub>3</sub>)<sub>4</sub>. The metal–phosphenium bond in both compounds was dominated by metal-to-ligand  $\pi$ -donation. This differed from the M–C bonds in the analogous N-heterocyclic carbene (NHC) complexes, (NHC<sup>Mes</sup>)M(PPh<sub>3</sub>)<sub>2</sub> (M = Pt (**6**), Pd (**7**)), which were instead predominantly  $\sigma$ -type. Structural determination of **1** by X-ray crystallography revealed the shortest yet reported Pt–P bond of 2.107(3) Å, consistent with significant double-bond character, and trigonal planar geometries at both the P-atom within the [NHP<sup>Mes</sup>]<sup>+</sup> ligand ( $\Sigma(\text{angles}) = 359.99^\circ$ ) and at the Pt-atom ( $\Sigma(\text{angles}) = 360.00^\circ$ ), which indicated that **1** was better described as a Pt(0)–phosphenium rather than as a Pt(II)–phosphide. Reactions of **1** and **2** with excess PMe<sub>3</sub> cleanly gave the four-coordinate species [(NHP<sup>Mes</sup>)M(PMe<sub>3</sub>)<sub>3</sub>]OTf (M = Pt (**3**) and Pd (**4**)), while reaction of **1** with bis(diphenylphosphino)ethane (dppe) gave [(NHP<sup>Mes</sup>)Pt(dppe- $\kappa^2$ P)(dppe- $\kappa$ P)]OTf (**5**). Hydrolysis of these complexes resulted in metal hydrides and oxidation of the NHP to phosphine oxide via initial nucleophilic attack of water at the P-atom in the coordinated [NHP<sup>Mes</sup>]<sup>+</sup> ligand, which was calculated to bear a significantly positive charge in **1**.

## Introduction

The unfurling of new molecular metal chemistry relies to a great extent on the discovery and development of novel ancillary and reactive ligand types with unusual electronic and steric demands. These initially strange ligands often permit the generation of unfamiliar metal coordination geometries and the stabilization of unusual oxidation states. They also impart unique reactivity on the resulting complexes, which may in turn lead to significant applications. There are few better examples than that of metal–carbenes, which have revolutionized organometallic chemistry and catalysis. The pioneering work in this area by Fischer,<sup>1,2</sup> Schrock,<sup>3,4</sup> Grubbs,<sup>5</sup> Arduengo,<sup>6</sup> and Bertrand<sup>7,8</sup> has been extended to include analogous complexes of the main

group elements, e.g., boron (boryls, :BR<sub>2</sub><sup>−</sup>),<sup>9,10</sup> silicon (silylenes, :SiR<sub>2</sub>),<sup>11,12</sup> and phosphorus (phospheniums, :PR<sub>2</sub><sup>+</sup>).<sup>13–17</sup>

N-Heterocyclic phosphenium cations (NHPs) are the “carbon copies” of the now ubiquitous Arduengo-type carbenes (NHCs).<sup>18</sup> Both species may have saturated or unsaturated “backbones”, but this report deals only with the saturated variety, specifically the mesityl-substituted compound [NHP<sup>Mes</sup>]OTf (Chart 1). The N-heterocyclic carbenes and phospheniums are isostructural and isovalent, but electronically inverse. NHCs are strong  $\sigma$ -donors and weak  $\pi$ -acceptors, whereas NHPs are weak  $\sigma$ -donors and good  $\pi$ -acceptors due to the formal positive charge and isotropic s- (as opposed to directional sp<sup>2</sup>-) character of the “lone pair” orbital on phosphorus.<sup>19</sup> The two families should therefore show reciprocal reactivity in transition metal chemistry.<sup>20</sup>

\* Corresponding author. Fax: 1-519-661-3022. E-mail: njones26@uwo.ca.

<sup>†</sup> The University of Western Ontario.

<sup>‡</sup> University of Jyväskylä.

(1) Fischer, E. O.; Maasböl, A. *Angew. Chem., Int. Ed. Engl.* **1964**, *3*, 580.

(2) Dötz, K. H.; Fischer, H.; Hofmann, P.; Kreissl, F. R.; Schubert, U.; Weiss, K. *Transition Metal Carbene Complexes*; VCH: Weinheim, 1983.

(3) Schrock, R. R. *J. Am. Chem. Soc.* **1974**, *96*, 6796.

(4) Schrock, R. R. *Acc. Chem. Res.* **1979**, *12*, 98.

(5) Nguyen, S. T.; Johnson, L. K.; Grubbs, R. H. *J. Am. Chem. Soc.* **1992**, *114*, 3974.

(6) Arduengo, A. J., III; Harlow, R. L.; Kline, M. *J. Am. Chem. Soc.* **1991**, *113*, 361.

(7) Lavallo, V.; Canac, Y.; Präsang, C.; Donnadiou, B.; Bertrand, G. *Angew. Chem., Int. Ed.* **2005**, *44*, 5705.

(8) Dyker, C. A.; Lavallo, V.; Donnadiou, B.; Bertrand, G. *Angew. Chem., Int. Ed.* **2008**, *47*, 3206.

(9) Aldridge, S.; Coombs, D. L. *Coord. Chem. Rev.* **2004**, *248*, 535.

(10) Irvine, G. J.; Lesley, M. J. G.; Marder, T. B.; Norman, N. C.; Rice, C. R.; Robins, E. G.; Roper, W. R.; Whittell, G. R.; Wright, L. J. *Chem. Rev.* **1998**, *98*, 2685.

(11) Okazaki, M.; Tobita, H.; Ogino, H. *Dalton Trans.* **2003**, 493.

(12) Waterman, R.; Hayes, P. G.; Tilley, T. D. *Acc. Chem. Res.* **2007**, *40*, 712.

(13) Gudat, D. *Coord. Chem. Rev.* **1997**, *163*, 71.

(14) Nakazawa, H. *J. Organomet. Chem.* **2000**, *611*, 349.


(15) Nakazawa, H. In *Advances in Organometallic Chemistry*; West, R. H., A. F., Ed.; Academic Press: Amsterdam, The Netherlands, 2004; Vol. 50, p 107.

(16) Cowley, A. H.; Kemp, R. A. *Chem. Rev.* **1985**, *85*, 367.

(17) Sanchez, M.; Mazières, M.-R.; Lamandé, L.; Wolf, R. *Phosphorus Compounds with Coordination Number 2: Phosphenium Cations*; Thieme: Stuttgart, Germany, 1990.

(18) Dillon, K. B.; Mathey, F.; Nixon, J. F. *Phosphorus: The Carbon Copy: From Organophosphorus to Phospha-organic Chemistry*; John Wiley: Chichester, England, 1998.

**Chart 1. Comparison between N-Heterocyclic Carbenes (NHCs) and Phospheniums (NHPs) and the Specific Ligand under Study, [NHP<sup>Mes</sup>]OTf (Mes = 2,4,6-trimethylphenyl, TfO<sup>-</sup> = trifluoromethanesulfonate)**



Often considered “phosphine replacement” ligands,<sup>21–23</sup> NHCs have triggered an explosion of research in the field of homogeneous catalysis. By comparison, NHPs have garnered much less attention as ancillary ligands in catalysis. However, reinvigorated investigation of their metal chemistry over the last several years has begun to uncover their remarkable properties.<sup>13,24</sup>

The use of diamminophospheniums as ligands dates to Parry’s 1978 synthesis of  $[(\text{CO})_4\text{Fe}(\text{P}\{\text{NR}_2\}_2)]\text{PF}_6$ .<sup>25</sup> Since then, the synthesis, structure, bonding, and reactivity of early and mid transition metal complexes (e.g., of Cr,<sup>26,27</sup> Mo,<sup>28,29</sup> W,<sup>26,27</sup> Fe,<sup>30</sup> and Ru<sup>31</sup>) have been explored, principally by Nakazawa.<sup>14</sup> To date, however, there are exceedingly few reported examples of late (groups 9 and 10) transition metal–NHP complexes. The groups of Baker<sup>32</sup> and Richeson<sup>20</sup> have both made NHP-containing analogues of Wilkinson’s catalyst (**I–III**; Chart 2), and Gudat has disclosed a Co complex (**IV**).<sup>33</sup> To the best of our knowledge, there are currently only four reported NHP complexes of the group 10 metals. The first of these was  $[(\text{NHP}^{\text{Mes}})\text{Ni}(\text{CO})_3][\text{BH}^+\text{Bu}_3]$  (**V**) by Parry.<sup>34</sup> Niecke has described a Ni complex with a zwitterionic phosphenium-containing ligand (**VI**),<sup>35</sup> while Baker has shown that  $[\text{NHP}^{\text{Mes}}]\text{OTf}$  reacts directly with  $\text{Pt}(\text{PPh}_3)_3$  to give  $[(\text{NHP}^{\text{Mes}})\text{Pt}(\text{PPh}_3)_2]\text{OTf}$  (**1**).<sup>36</sup> One of the  $\text{PPh}_3$  ligands in this complex can be replaced by an Enders-type NHC to give  $[(\text{NHP}^{\text{Mes}})\text{Pt}(\text{PPh}_3)(\text{NHC})]\text{OTf}$

(**VII**); the same compound results from reaction between  $\text{Pt}(\text{PPh}_3)_3$  and the NHC→NHP adduct.

On the basis of their strong  $\pi$ -acidity and reasonably straightforward modular synthesis, NHPs have been described as tunable CO equivalents.<sup>32</sup> However, it may be better to think of them as tunable  $\text{NO}^+$  surrogates instead, not only because they are cationic but also because, in contrast to terminal CO ligands that bind only linearly, they may in principle adopt either planar or pyramidal (at P) coordination modes.<sup>37</sup> These are analogous to the linear and “bent” modes, respectively, of metal nitrosyls and have identical implications for oxidation state formalisms (Chart 3). The planar ligand is properly considered a coordinated phosphenium, while the pyramidal one is a metal phosphide. Both of these modes have been observed structurally by Paine in  $\text{CpMo}(\text{CO})_2(\text{NHP}^{\text{Me}})$  (planar)<sup>38</sup> and  $\text{Cp}^*\text{Fe}(\text{CO})_2(\text{NHP}^{\text{Me}})$  (pyramidal).<sup>39</sup> The analogy between NHP and  $\text{NO}^+$  (and Fischer carbenes, *vide infra*) was first drawn in passing in Paine’s pioneering work and also later in a review by Gudat,<sup>13</sup> but to the best of our knowledge the relationship has not been formalized via definitive structure, bonding, and reactivity relationships.

Just as there is inconsistency in the ways that metal–NHC bonds are drawn in the literature, representation of the metal–NHP interaction has proven problematic (see Chart 2 for the ways that group 9 and 10 metal complexes have been drawn, and Chart 4 for various generic representations). Although a range of descriptions is warranted on the basis of differences in structure and reactivity, and therefore a universally applicable drawing is impossible, historical representation of the metal–NHP interaction appears to be complicated by the ligand’s carbene-like nature, formal positive charge, dual bonding modes, and capacity for strong  $\pi$ -back-bonding. The representations in Chart 4 may be considered to be on the same “resonance continuum”, sometimes by deconvolution of the dative bond,  $\text{D} \rightarrow \text{A}$ , to its charge-separated, valence bond representation,  $\text{D}^+ - \text{A}^-$ ,<sup>40</sup> but these diagrams have different chemical meanings and should have significant implications in interpreting and predicting the structures and reactivity of the NHP complexes they represent. For example, it should be possible to substitute the phosphenium ligand in **A**, and **B** should contain a trigonal pyramidal phosphorus center with a lone pair (as in Chart 3), but it has been used in the literature to represent coordinated planar phospheniums.

We recently reported an improved synthesis of a family of free saturated NHP ligands.<sup>41</sup> In this article, we present a series of cationic Pt(0)– and Pd(0)–NHP complexes (Chart 5, including **1** via a new synthesis) followed by their reactions with both monodentate and chelating tertiary phosphines, and also with  $\text{H}_2\text{O}$ .

The solid state structure of **1** presented herein, together with the results of electronic structure calculations and reactivity patterns, affords insight into the nature of the bonding in late metal–NHP complexes and allows the analogy with carbenes and linear nitrosyls to be put on a solid footing. The Pt and Pd compounds are best described as phosphorus analogues of

(19) Tuononen, H. M.; Roesler, R.; Dutton, J. L.; Ragogna, P. J. *Inorg. Chem.* **2007**, *46*, 10693.

(20) Spinney, H. A.; Yap, G. P. A.; Korobkov, I.; DiLabio, G.; Richeson, D. S. *Organometallics* **2006**, *25*, 3541.

(21) Arduengo, A. J., III. *Acc. Chem. Res.* **1999**, *32*, 913.

(22) Bourissou, D.; Guerret, O.; Gabbai, F. P.; Bertrand, G. *Chem. Rev.* **2000**, *100*, 39.

(23) Herrmann, W. A.; Köcher, C. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 2162.

(24) Gudat, D.; Haghverdi, A.; Hupfer, H.; Nieger, M. *Chem.–Eur. J.* **2000**, *6*, 3414.

(25) Montemayor, R. G.; Sauer, D. T.; Fleming, S.; Bennett, D. W.; Thomas, M. G.; Parry, R. W. *J. Am. Chem. Soc.* **1978**, *100*, 2231.

(26) Nakazawa, H.; Yamaguchi, Y.; Miyoshi, K. *J. Organomet. Chem.* **1994**, *465*, 193.

(27) Yamaguchi, Y.; Nakazawa, H.; Itoh, T.; Miyoshi, K. *Bull. Chem. Soc. Jpn.* **1996**, *69*, 983.

(28) Gudat, D.; Haghverdi, A.; Nieger, M. *J. Organomet. Chem.* **2001**, *383*, 617.

(29) Nakazawa, H.; Miyoshi, Y.; Katayama, T.; Mizuta, T.; Miyoshi, K.; Tsuchida, N.; Ono, A.; Takano, K. *Organometallics* **2006**, *25*, 5913.

(30) Nakazawa, H.; Kishishita, M.; Nakamoto, T.; Nakamura, N.; Ishiyama, T.; Miyoshi, K. *Chem. Lett.* **2000**, 230.

(31) Kawamura, K.; Nakazawa, H.; Miyoshi, K. *Organometallics* **1999**, *18*, 4785.

(32) Abrams, M. B.; Scott, B. L.; Baker, R. T. *Organometallics* **2000**, *19*, 4944.

(33) Burck, S.; Daniels, J.; Gans-Eichler, T.; Gudat, D.; Näntinen, K.; Nieger, M. *Z. Anorg. Allg. Chem.* **2005**, *631*, 1403.

(34) Snow, S. S.; Jiang, D. X.; Parry, R. W. *Inorg. Chem.* **1987**, *26*, 1629.

(35) Oberdörfer, R.; Nieger, M.; Niecke, E. *Chem. Ber.* **1994**, *127*, 2397.

(36) Hardman, N. J.; Abrams, M. B.; Pribisko, M. A.; Gilbert, T. M.; Martin, R. L.; Kubas, G. J.; Baker, R. T. *Angew. Chem., Int. Ed. Engl.* **2004**, *43*, 1955.


(37) Collman, J. P.; Hegedus, L. S.; Norton, J. R.; Finke, R. G. *Principles and Applications of Organotransition Metal Chemistry*; University Science Books: Mill Valley, CA, 1987.

(38) Hutchins, L. D.; Paine, R. T.; Campana, C. F. *J. Am. Chem. Soc.* **1980**, *102*, 4521.

(39) Hutchins, L. D.; Duesler, E. N.; Paine, R. T. *Organometallics* **1982**, *1*, 1254.

(40) Parkin, G. *Organometallics* **2006**, *25*, 4744.

(41) Caputo, C. A.; Price, J. T.; Jennings, M. C.; McDonald, R.; Jones, N. D. *Dalton Trans.* **2008**, 3461.

Chart 2. Summary of Reported Group 9 and 10 Metal–NHP Complexes<sup>a</sup>

<sup>a</sup> With the exception of **1** and **VII**, for which graphics were not given in the original reports, the cations are drawn exactly as they have been represented in the literature. In all Pt and Rh compounds, the anion is TfO<sup>−</sup>; for Ni, it is [HB(<sup>t</sup>Bu)<sub>3</sub>]<sup>−</sup>.

Chart 3. Structural Analogy between Metal–Nitrosyls and –Phospheniums






Chart 4. Selection of Generic Representations of Metal–NHP Complexes Found in the Literature

Chart 5. NHP-Containing Compounds Made in This Study (all anions are TfO<sup>−</sup>)

Fischer carbenes with P=M double bond character and localization of positive charge at P (as in **D**, Chart 4). To demonstrate this analogy with Fischer carbenes, we show that the coordinated phosphenium ligand is inert to substitution by monodentate and chelating phosphines. We show also that the complexes undergo attack by H<sub>2</sub>O at the NHP P-atom to generate metal–hydrides in a mechanism akin to that proposed for the hydrolysis of Fischer carbenes.<sup>42–44</sup>

## Experimental Section

**General Considerations.** Reagents were obtained from commercial sources and used as supplied, unless otherwise indicated. These were of American Chemical Society (ACS) grade or finer. Solvents were dried and deoxygenated either by N<sub>2</sub> purge followed by passage through alumina columns (Innovative Technology or MBraun solvent purification systems) or by distillation under N<sub>2</sub> from the appropriate drying agent. All reactions were carried out under N<sub>2</sub> atmosphere using standard Schlenk techniques unless stated otherwise. Pt(PPh<sub>3</sub>)<sub>4</sub> and Pd(PPh<sub>3</sub>)<sub>4</sub> were made according to literature procedures.<sup>45,46</sup>

<sup>1</sup>H, <sup>13</sup>C{<sup>1</sup>H}, <sup>19</sup>F{<sup>1</sup>H}, and <sup>31</sup>P{<sup>1</sup>H} NMR data were recorded on a 400 MHz Varian Mercury (400.085 MHz for <sup>1</sup>H, 100.602 MHz for <sup>13</sup>C, 376.458 MHz for <sup>19</sup>F) or 400 MHz Varian Inova spectrometer (399.762 MHz for <sup>1</sup>H, 100.520 MHz for <sup>13</sup>C, 161.825 MHz for <sup>31</sup>P). Unless otherwise indicated, spectra were recorded at rt in CDCl<sub>3</sub> solution using residual solvent proton (relative to external SiMe<sub>4</sub>, δ 0.00) or solvent carbon (relative to external SiMe<sub>4</sub>, δ 0.00) as an internal reference. All <sup>31</sup>P{<sup>1</sup>H} NMR spectra were recorded relative to an external standard (85% H<sub>3</sub>PO<sub>4</sub>, δ 0.00). Downfield shifts were taken as positive, and all coupling constants are given in hertz.

High-resolution mass spectrometry data were recorded using a Finnigan MAT 8200 instrument. Elemental analyses (C, H, N) were performed by Guelph Chemical Laboratories Inc., Guelph, ON, Canada.

**Synthesis of Pt and Pd Metal Complexes.** [(NHP<sup>Mes</sup>)Pt-(PPh<sub>3</sub>)<sub>2</sub>OTf (**1**). The synthesis of this compound (previously reported by Baker and co-workers)<sup>36</sup> was modified to use Pt(PPh<sub>3</sub>)<sub>4</sub> instead of Pt(PPh<sub>3</sub>)<sub>3</sub>. In a small vial, Pt(PPh<sub>3</sub>)<sub>4</sub> (0.53 g, 0.42 mmol) was dissolved in C<sub>6</sub>H<sub>6</sub> (10 mL). A solution of [NHP<sup>Mes</sup>]OTf (0.20 g, 0.42 mmol) in C<sub>6</sub>H<sub>6</sub> (5 mL) was added

(42) Bernasconi, C. F.; Leyes, A. E. *J. Chem. Soc., Perkin Trans. 2* **1997**, 1641.

(43) Zoloff Michoff, M. E.; de Rossi, R. H.; Granados, A. M. *J. Org. Chem.* **2006**, *71*, 2395.

(44) Bernasconi, C. F.; Flores, F. X.; Kittredge, K. W. *J. Am. Chem. Soc.* **1997**, *119*, 2103.

(45) Ugo, R.; Cariati, F.; La Monica, G. In *Reagents for Transition Metal Complex and Organometallic Syntheses*; Angelici, R. J., Ed.; John Wiley & Sons Inc.: New York, 1990; Vol. 28, p 123.

(46) Coulson, D. R. In *Reagents for Transition Metal Complex and Organometallic Syntheses*; Angelici, R. J., Ed.; John Wiley & Sons Inc.: New York, 1990; Vol. 28, p 107.

dropwise to give a dark yellow solution. The solvent was frozen and removed *in vacuo* to give a yellow solid, which was washed with Et<sub>2</sub>O (2 × 10 mL) to remove free PPh<sub>3</sub>, then dried *in vacuo* to give the title compound as a yellow solid. Yield: 0.26 g (49%). X-ray quality crystals of **1** · 1/2C<sub>6</sub>H<sub>6</sub> were grown from concentrated C<sub>6</sub>H<sub>6</sub> solution by slow evaporation. The <sup>1</sup>H NMR spectroscopic data were identical to those reported by Baker.<sup>36</sup> However, chemical shift differences due to solvent were observed in the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum. <sup>31</sup>P{<sup>1</sup>H} NMR (CDCl<sub>3</sub>): δ 44.3 (d, <sup>1</sup>J<sub>Pt</sub> = 4243, <sup>2</sup>J<sub>PP</sub> = 228), 290.0 (t, <sup>1</sup>J<sub>Pt</sub> = 6446, <sup>2</sup>J<sub>P-P</sub> = 228). In addition, the following additional characterization data were collected. <sup>19</sup>F{<sup>1</sup>H} NMR (CDCl<sub>3</sub>): δ -78.1 (s). ESI-MS: C<sub>56</sub>H<sub>56</sub>N<sub>2</sub>P<sub>3</sub>Pt<sup>+</sup> calcd (found) 1044.3299 (1044.3367, M<sup>+</sup>). Anal. Calcd for C<sub>57</sub>H<sub>58</sub>F<sub>3</sub>N<sub>2</sub>O<sub>3</sub>P<sub>3</sub>PtS: C, 57.2; H, 4.9; N, 2.3. Found: C, 57.2; H, 5.0; N, 2.2.

[(NHP<sup>Mes</sup>)Pd(PPh<sub>3</sub>)<sub>2</sub>OTf (**2**). This compound was made in the same way as **1**. Thus, reaction of Pd(PPh<sub>3</sub>)<sub>4</sub> (0.59 g, 0.52 mmol) and [NHP<sup>Mes</sup>]OTf (0.27 g, 0.57 mmol) gave 0.30 g (52%) of a yellow solid. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>): δ 2.05 (s, 6H, *p*-CH<sub>3</sub>), 2.08 (s, 12H, *o*-CH<sub>3</sub>), 4.10 (d, 4H, CH<sub>2</sub>, <sup>3</sup>J<sub>HP</sub> = 4.0), 6.54 (s, 4H, Ar), 7.01 (pt, 18H, Ar), 7.13 (pq, 12H, Ar). <sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>, 263 K): δ 17.8, 21.3, 52.5, 128.8, 130.0, 130.3, 131.8, 133.3, 135.0, 136.0, 138.8. <sup>31</sup>P{<sup>1</sup>H} NMR (CDCl<sub>3</sub>): δ 25.1 (d, <sup>2</sup>J<sub>PP</sub> = 149), 260.0 (t, <sup>2</sup>J<sub>PP</sub> = 149). <sup>19</sup>F NMR (CDCl<sub>3</sub>): δ -78.2 (s). ESI-MS: C<sub>56</sub>H<sub>56</sub>N<sub>2</sub>P<sub>3</sub>Pd<sup>+</sup> calcd (found) 955.3 (955.3, M<sup>+</sup>). Anal. Calcd for C<sub>57</sub>H<sub>56</sub>F<sub>3</sub>N<sub>2</sub>O<sub>3</sub>P<sub>3</sub>PdS: C, 61.9; H, 5.1; N, 2.5. Found: C, 62.0; H, 5.1; N, 2.5.

[(NHP<sup>Mes</sup>)Pt(PMe<sub>3</sub>)<sub>3</sub>] (**3**). Neat PMe<sub>3</sub> (0.25 mL, 2.4 mmol) was added dropwise to a yellow CH<sub>2</sub>Cl<sub>2</sub> (2 mL) solution containing **1** (0.33 g, 0.28 mmol) in a small vial. The reaction mixture was stirred for 10 min at rt, during which it turned red. The solvent was removed *in vacuo*, and the orange residue was washed with Et<sub>2</sub>O (2 × 10 mL) to remove free PPh<sub>3</sub>, then dried under vacuum. Yield: 0.22 g (88%). <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.17 (m (br m, 27H, CH<sub>3</sub>), 2.25 (s, 6H, *p*-CH<sub>3</sub>), 2.34 (s, 12H, *o*-CH<sub>3</sub>), 3.79 (d, 4H, CH<sub>2</sub>, <sup>3</sup>J<sub>HP</sub> = 2.0), 7.28 (br s, 4H, *m*-CH). <sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>, 273 K): δ 18.7, 21.0, 24.6, 50.7, 129.9, 134.3, 136.6, 138.6. <sup>31</sup>P{<sup>1</sup>H} NMR (CDCl<sub>3</sub>): δ -47.1 (d, <sup>1</sup>J<sub>Pt</sub> = 3283, <sup>2</sup>J<sub>PP</sub> = 125), 207.6 (q, <sup>1</sup>J<sub>Pt</sub> = 6162, <sup>2</sup>J<sub>PP</sub> = 125). <sup>19</sup>F NMR (CDCl<sub>3</sub>): δ -78.1 (s). ESI-MS: C<sub>29</sub>H<sub>53</sub>N<sub>2</sub>P<sub>4</sub>Pt<sup>+</sup> calcd (found) 748.2807 (748.2743, M<sup>+</sup>). Anal. Calcd for C<sub>30</sub>H<sub>53</sub>F<sub>3</sub>N<sub>2</sub>O<sub>3</sub>P<sub>4</sub>PtS: C, 40.1; H, 5.95; N, 3.1. Found: C, 39.65; H, 6.1; N, 2.6.

[(NHP<sup>Mes</sup>)Pd(PMe<sub>3</sub>)<sub>3</sub>]OTf (**4**). This compound was made in the same way as **3**. Thus reaction of **2** (0.30 g, 0.27 mmol) and PMe<sub>3</sub> (0.112 mL, 1.09 mmol) gave 0.13 g (58%) of an orange solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.03 (br s, 27H, CH<sub>3</sub>), 2.24 (s, 6H, *p*-CH<sub>3</sub>), 2.34 (s, 12H, *o*-CH<sub>3</sub>), 3.86 (d, 4H, CH<sub>2</sub>, <sup>3</sup>J<sub>HP</sub> = 4.4), 6.95 (s, 4H, Ar). <sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>): δ 18.7, 21.2, 21.4, 52.1 (d, <sup>2</sup>J<sub>PC</sub> = 6.1), 128.6 (d, <sup>2</sup>J<sub>PC</sub> = 7.6), 130.3, 133.8 (d, <sup>2</sup>J<sub>PC</sub> = 17.7), 134.3 (d, <sup>2</sup>J<sub>CP</sub> = 9.2), 136.3 (d, <sup>2</sup>J<sub>CP</sub> = 3.1), 139.0. <sup>31</sup>P{<sup>1</sup>H} NMR (CDCl<sub>3</sub>, 253 K): δ -29.6 (d, <sup>2</sup>J<sub>PP</sub> = 73), 242.8 (d, <sup>2</sup>J<sub>PP</sub> = 73). <sup>19</sup>F NMR (CDCl<sub>3</sub>): δ -74.2 (s). ESI-MS: C<sub>29</sub>H<sub>53</sub>N<sub>2</sub>P<sub>4</sub>Pd<sup>+</sup> calcd (found) 585.2 (583.1, M<sup>+</sup> - PMe<sub>3</sub>), (507.1, M<sup>+</sup> - 2 PMe<sub>3</sub>). Anal. Calcd for C<sub>30</sub>H<sub>53</sub>F<sub>3</sub>N<sub>2</sub>O<sub>3</sub>P<sub>4</sub>PdS: C, 44.53; H, 6.60; N, 3.46. Found: C, 44.25; H, 6.52; N, 3.16.

[(NHP<sup>Mes</sup>)Pt(dppe-κ<sup>2</sup>P)(dppe-κP)]OTf (**5**). This compound was prepared in the same way as **3**. Thus, reaction of **1** (102 mg, 8.50 mmol) and dppe (68.0 mg, 17.0 mmol) gave 94.7 mg (76%) of a dark orange solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.26 (br s, 4H, PCH<sub>2</sub>), 1.58 (br s, 4H, PCH<sub>2</sub>), 2.10 (s, 6H, *p*-CH<sub>3</sub>), 2.26 (s, 12H, *o*-CH<sub>3</sub>), 3.94 (br s, 4H, NCH<sub>2</sub>), 6.59 (br s, 4H, Ar), 6.93–7.25 (br m, 40H, Ar). <sup>31</sup>P{<sup>1</sup>H} NMR (CDCl<sub>3</sub>, 263 K): δ -14.2 (d, <sup>3</sup>J<sub>PP</sub> = 28.5), 9.7 (m, <sup>1</sup>J<sub>Pt</sub> = 3502), 26.3 (m, <sup>1</sup>J<sub>Pt</sub> = 2994, <sup>2</sup>J<sub>PP</sub> = 105), 217.5 (m, <sup>1</sup>J<sub>Pt</sub> = 6914). <sup>19</sup>F NMR (CDCl<sub>3</sub>): δ -78.0 (s). ESI-MS: C<sub>72</sub>H<sub>77</sub>N<sub>2</sub>P<sub>5</sub>Pt<sup>+</sup> calcd (found) 1319.4 (1320.4, M<sup>+</sup> + 1), (921.9, M<sup>+</sup> - dppe). The inevitable appearance of [PtH(dppe)<sub>2</sub>]OTf hydrolysis product in all samples precluded satisfactory elemental analysis.

**Computational Details.** Molecular geometries of all studied compounds were optimized with DFT using the GGA PBE/PBE exchange–correlation functional<sup>47–49</sup> with the def-TZVP basis sets;<sup>50,51</sup> ECP basis sets of similar valence quality were used for the transition metal nuclei.<sup>52</sup> Although the use of the GGA functional leads to a slight overestimation of all bond lengths, it was preferred over the hybrid counterpart PBE1PBE for reasons of computational efficiency, i.e., the ability to use multipole accelerated RI approximation and to run calculations in parallel. All optimizations were carried out with the Turbomole 5.10 program package.<sup>53,54</sup> Energy decomposition analyses for optimized structures were performed with the ADF 2007.01 program.<sup>55</sup> The analysis followed the Morokuma–Rauk–Ziegler partition scheme<sup>56–58</sup> and utilized the PBE/PBE GGA functional<sup>47–49</sup> in combination with STO-type all-electron basis sets of TZP quality.<sup>59,60</sup> Scalar relativistic effects were taken into account in all EDA calculations by employing the ZORA Hamiltonian.<sup>61–63</sup> Representations of the molecular orbitals of **2** and **7** given in Figure 2 were constructed using the gOpenMol program.<sup>64,65</sup>

## Results and Discussion

**Synthesis and Structure.** The phosphonium triflate [NHP<sup>Mes</sup>]OTf reacts quickly and cleanly with equimolar M(PPh<sub>3</sub>)<sub>4</sub> precursors at rt to generate [(NHP<sup>Mes</sup>)M(PPh<sub>3</sub>)<sub>2</sub>]OTf (**1**, M = Pt; **2**, M = Pd) with elimination of 2 equiv of PPh<sub>3</sub> (Scheme 1). As mentioned in the Introduction, **1** has previously been made (although not structurally characterized) by reaction of [NHP<sup>Mes</sup>]OTf with Pt(PPh<sub>3</sub>)<sub>3</sub>; **2** represents the first well-defined Pd–NHP complex, although related species have been assumed to form *in situ* in catalytic processes.<sup>66,67</sup>

The geometries of mixed phosphine–phosphonium complexes of the type [(NHP<sup>Mes</sup>)M(PR<sub>3</sub>)<sub>n</sub>]OTf are easily elucidated from <sup>31</sup>P{<sup>1</sup>H} NMR data. The spectrum of **1** in CDCl<sub>3</sub> exhibits an upfield doublet (with Pt satellites) for the PPh<sub>3</sub> ligands (δ 44.3,

(47) Perdew, J. P.; Burke, K.; Ernzerhof, M. *Phys. Rev. Lett.* **1996**, *77*, 3865.

(48) Perdew, J. P.; Burke, K.; Ernzerhof, M. *Phys. Rev. Lett.* **1997**, *78*, 1396.

(49) Perdew, J. P.; Ernzerhof, M.; Burke, K. *J. Chem. Phys.* **1996**, *105*, 9982.

(50) Schäfer, A.; Huber, C.; Ahlrichs, R. *J. Chem. Phys.* **1994**, *100*, 5829.

(51) Eichkorn, K.; Weigend, F.; Treutler, O.; Ahlrichs, R. *Theor. Chem. Acc.* **1997**, *97*, 119.

(52) Andrae, D.; Haeussermann, U.; Dolg, M.; Stoll, H.; Preuss, H. *Theor. Chem. Acc.* **1990**, *77*, 123.

(53) TURBOMOLE, Program Package for *ab initio* Electronic Structure Calculations; Version 5.10 ed.; Theoretical Chemistry Group, University of Karlsruhe: Karlsruhe, Germany, 2008.

(54) Ahlrichs, R.; Bär, M.; Häser, M.; Horn, H.; Kölmel, C. *Chem. Phys. Lett.* **1989**, *162*, 165.

(55) ADF2007.01 Theoretical Chemistry Program; SCM, V. U.: Amsterdam, The Netherlands, 2007.

(56) Morokuma, K. *J. Chem. Phys.* **1971**, *55*, 1236.

(57) Kitaura, K.; Morokuma, K. *Int. J. Quantum Chem.* **1976**, *10*, 325.

(58) Ziegler, T.; Rauk, A. *Theor. Chim. Acta* **1977**, *46*, 1.

(59) van Lenthe, E.; Baerends, E. J. *J. Comput. Chem.* **2003**, *24*, 1142.

(60) Chong, D. P.; van Lenthe, E.; van Gisbergen, S.; Baerends, E. J. *J. Comput. Chem.* **2004**, *25*, 1030.

(61) van Lenthe, E.; Baerends, E. J.; Snijders, J. G. *J. Chem. Phys.* **1993**, *99*, 4597.

(62) van Lenthe, E.; Baerends, E. J.; Snijders, J. G. *J. Chem. Phys.* **1994**, *101*, 9783.


(63) van Lenthe, E.; Ehlers, A. E.; Baerends, E. J. *J. Chem. Phys.* **1999**, *110*, 8943.

(64) Laaksonen, L. *J. Mol. Graph.* **1992**, *10*, 33.


(65) Bergman, D. L.; Laaksonen, L.; Laaksonen, A. *J. Mol. Graph. Model.* **1997**, *15*, 301.

(66) Ackermann, L.; Born, R. *Angew. Chem., Int. Ed.* **2005**, *44*, 2444.

(67) Sakakibara, K.; Yamashita, M.; Nozaki, K. *Tetrahedron Lett.* **2005**, *46*, 959.




**Figure 1.** Stacked  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra for  $[(\text{NHP}^{\text{Mes}})\text{Pd}(\text{PPh}_3)_2]\text{OTf}$  (**2**, bottom) and  $[(\text{NHP}^{\text{Mes}})\text{Pd}(\text{PMe}_3)_3]\text{OTf}$  (**4**).



**Figure 2.** ORTEP representation of the molecular structure of the cation of **1**. Thermal ellipsoids are drawn at 30% probability. H-atoms and all but the *ipso*-C-atoms of the  $\text{PPh}_3$  ligands have been omitted for clarity. Selected experimental [and calculated] bond distances ( $\text{\AA}$ ) and angles (deg): Pt–P1, 2.1073(9) [2.185]; P1–N2, 1.640(3) [1.695]; P1–N5, 1.628(3) [1.695]; Pt–P2, 2.3178(9) [2.383]; Pt–P3, 2.3113(8) [2.383]; N5–P1–N2, 93.68(16) [92.0]; N5–P1–Pt, 134.15(12) [134.0]; N2–P1–Pt, 132.16(12) [134.0]; P1–Pt–P3, 121.41(3) [122.0]; P1–Pt–P2, 121.70(3) [122.0]; P2–Pt–P3, 116.89(3) [115.9]; P2–Pt–P1–N5,  $-61.66(19)$  [ $-69.1$ ]; P3–Pt–P1–N5, 119.02(19) [110.9].

**Scheme 1. Synthesis of Pd(0)– and Pt(0)–NHP Metal Complexes (anion is  $\text{TfO}^-$ )**



$^1J_{\text{PtP}} = 4143$  Hz,  $^2J_{\text{PP}} = 228$  Hz) and a dramatically downfield triplet (with Pt satellites) for the coordinated  $[\text{NHP}^{\text{Mes}}]^+$  ligand ( $\delta$  290.0,  $^1J_{\text{PtP}} = 6446$  Hz,  $^2J_{\text{PP}} = 228$  Hz) with relative 2:1 peak intensity. These data are similar to those reported by Baker for **1** in  $\text{CD}_2\text{Cl}_2$  solution.<sup>36</sup> They unequivocally confirm the presence of two equivalent phosphine and one phosphonium ligand in a trigonal-planar coordination environment. The shift


on coordination ( $\Delta\delta$ ) of  $[\text{NHP}^{\text{Mes}}]^+$  to Pt is 86.9 ppm downfield, which implies a dramatic deshielding of the phosphorus center. The Pt–P coupling constant is very much larger than those obtained by Glueck and co-workers<sup>68</sup> for a range of Pt(II)–phosphine complexes (ca. 900–1000 Hz), which implies that the bound NHP in **1** is probably not pyramidal in solution. Similar NMR data are obtained for **2**: an upfield doublet ( $\delta$  25.1,  $^2J_{\text{PP}} = 149$  Hz) for the  $\text{PPh}_3$  ligands and a downfield triplet ( $\delta$  260.0,  $^2J_{\text{PP}} = 149$  Hz;  $\Delta\delta = 56.9$ ) for the coordinated NHP in a 2:1 ratio show that **1** and **2** are isostructural (Figure 1).

Crystals of  $\mathbf{1} \cdot 1/2\text{C}_6\text{H}_6$  suitable for X-ray crystallographic analysis were grown at rt by slow evaporation of  $\text{C}_6\text{H}_6$  solution; the solid state structure is given in Figure 2. As expected from the NMR data, the structure is trigonal planar at Pt ( $\Sigma(\text{angles}) = 360.00^\circ$ ). The geometry at the phosphonium P-atom is also trigonal planar ( $\Sigma(\text{angles}) = 359.99^\circ$ ). The N–P–N plane of the  $[\text{NHP}^{\text{Mes}}]^+$  ligand makes a torsion angle of approximately  $62^\circ$  with the coordination plane of the metal. The Pt–NHP bond (Pt–P1: 2.107(3)  $\text{\AA}$ ) is significantly shorter than the Pt– $\text{PPh}_3$  bonds (Pt–P2: 2.3178(9)  $\text{\AA}$  and Pt–P3: 2.3113(8)  $\text{\AA}$ ) by ca. 0.21  $\text{\AA}$ , consistent with the large Pt–P coupling constant, and indicates a strong Pt–P bond and greater back-bonding to the phosphonium than to the  $\text{PPh}_3$  ligands. To the best of our knowledge, this Pt–P bond is the shortest reported to date; the previous record of 2.116(3)  $\text{\AA}$  was held by Baker’s complex, **VII** (Chart 2);<sup>36</sup> formal Pt=P double bonds in phosphinidenes lie in the range 2.2–2.3  $\text{\AA}$ ,<sup>69</sup> which contains the average Pt– $\text{PPh}_3$  single bond of 2.28  $\text{\AA}$ .<sup>70</sup> These data imply that by metrical considerations alone the Pt–NHP interaction in **1** may be thought of as a double bond. No short contacts between either the NHP P-atom or the Pt-atom and the  $\text{TfO}^-$  anion are observed in the solid state; the closest distances are 3.282 and 3.370  $\text{\AA}$  between a “backbone” C-atom and the  $\text{OTf}^-$  O-atoms.

(68) Scriban, C.; Glueck, D. S.; DiPasquale, A. G.; Rheingold, A. L. *Organometallics* **2006**, 25, 5435.

(69) Kourkine, I. V.; Chapman, M. B.; Glueck, D. S.; Eichele, K.; Wasylishen, R. E.; Yap, G. P. A.; Liable-Sands, L. M.; Rheingold, A. L. *Inorg. Chem.* **1996**, 35, 1478.

(70) Orpen, A. G.; Brammer, L.; Allen, F. H.; Kennard, O.; Watson, D. G.; Taylor, R. *J. Chem. Soc., Dalton Trans.* **1989**, S1.

Scheme 2. Phosphine Substitution Reactions of **1** and **2** with  $\text{PMe}_3$  (all anions are  $\text{TfO}^-$ )


The question arises whether to consider **1** as a  $\text{Pt(0)}$ –NHP complex or a  $\text{Pt(II)}$ –phosphide (Chart 3). In **1**, and all other structurally characterized group 9 and 10 metal–NHP complexes,<sup>20,32,35,36</sup> the geometry at the NHP P-atom is trigonal planar, which clearly rules out the  $\text{Pt(II)}$ –phosphide description (analogous to “bent” metal–nitrosyls) and thereby discounts formal two-electron oxidation of the metal by the phosphonium. Hence, structural characteristics alone imply that **1** and **2** are best described as 16-electron metal(0)–NHP complexes.

Although **1** should be considered as a compound of  $\text{Pt(0)}$ , it reacts differently from homoleptic  $\text{Pt(0)}$ –phosphine complexes: it does not dissociate phosphine in solution (within the limits of  $^{31}\text{P}$  NMR detection); it is inert to oxidative addition by  $\text{MeI}$  and  $\text{MeOTf}$ ; and it does not bind the electron-poor alkene dimethyl maleate. These findings indicate that **1** is significantly less electron rich than other group 10 metal(0) phosphine compounds, probably due to the formal positive charge borne by the NHP ligand.

Chart 4 shows a selection of generic representations of metal–NHP complexes found in the literature. Clearly, **B** is not a good description of either **1** or **2** because it implies a trigonal-pyramidal geometry at the NHP P-atom. Representation of the  $\text{M}$ –NHP interaction in **1** and **2** as **D** (equivalent to **E**) should mean (i) that the NHP is not easily substituted (the opposite of what would be predicted by **A**) and (ii) that the ligand should suffer attack by good nucleophiles; representation instead as **C** would predict attack by nucleophiles at the metal rather than the phosphorus center. In fact, both predictions (i) and (ii) are borne out by experiment.

To test the first hypothesis, we surveyed the reaction of **1** and **2** with a variety of monodentate and chelating phosphines, and in no case was the NHP ligand displaced from the inner coordination sphere of the metal. This finding is consistent with Baker’s observation that reaction of **1** with an NHC leads to the mixed NHP/NHC compound **VII**, in which the phosphonium ligand is retained.<sup>36</sup> In our study, reaction of **1** and **2** with an excess of the small, basic phosphine  $\text{PMe}_3$  quickly and cleanly gives  $[(\text{NHP}^{\text{Mes}})\text{M}(\text{PMe}_3)_3]\text{OTf}$  (**3**,  $\text{M} = \text{Pt}$ ; **4**,  $\text{M} = \text{Pd}$ ). The substitution of  $\text{PPh}_3$  by  $\text{PMe}_3$  is cooperative, so that when only a single equivalent of  $\text{PMe}_3$  is added, **3** and **4** are generated, but two-thirds of the starting material is present in the reaction mixture at completion (Scheme 2). In no case is a mixed  $\text{PMe}_3/\text{PPh}_3$  complex detected, and the NHP ligand is always retained.

The formation of four- rather than the three-coordinate products could be explained by the smaller steric demand of  $\text{PMe}_3$  compared to  $\text{PPh}_3$  (Tolman cone angles:  $118^\circ$  and  $143^\circ$ , respectively)<sup>71</sup> and also to the higher Lewis basicity of the

Scheme 3. Substitution Reactions of **1** with  $\text{dppe}$  (anion is  $\text{TfO}^-$ )

incoming ligand. The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopic data of **3** and **4** are similar to those of the parent compounds, except that in these cases the downfield peaks due to the coordinated NHP are quartets, rather than triplets, due to coupling of the NHP to three equivalent phosphines in the four-coordinate products (Figure 1). These complexes are best described as 18-electron metal(0)–NHP complexes. Putative metal(II) compounds bearing three phosphine ligands should be square planar and therefore have two chemically distinct  $\text{PR}_3$  environments, one *trans* to the NHP and the other *cis*. However, in **3** and **4** only a single  $\text{PMe}_3$  environment is evident by NMR spectroscopy, which indicates the presence of the three equivalent phosphine ligands in a tetrahedral complex and thereby reinforces the idea that these compounds (together with **1** and **2**) are in fact metal(0) species.

Reaction of **1** with the chelating phosphine, bis(diphenylphosphino)ethane ( $\text{dppe}$ ), gives **5** with substitution of both  $\text{PPh}_3$  ligands (Scheme 3). This compound has a distinctive  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum that unequivocally describes the structure. The spectrum of **5** displays a low-field multiplet (with Pt “satellites”) at 215.6, an upfield multiplet (26.4, showing Pt “satellites”), yet another multiplet in this region (9.73, showing Pt “satellites”), and a doublet even further upfield ( $-14.2$ , no coupling to Pt). The chemical shifts, multiplicities, and relative integrations of these four chemically inequivalent  $^{31}\text{P}$  environments are consistent with the presence of coordinated  $[\text{NHP}^{\text{Mes}}]^+$  and one chelating and one monodentate  $\text{dppe}$  ligand, i.e., the compound  $[(\text{NHP}^{\text{Mes}})\text{Pt}(\text{dppe-}\kappa^2\text{P})(\text{dppe-}\kappa\text{P})]\text{OTf}$  (Scheme 3). This once again is clear evidence of the substitutionally inert nature of the coordinated NHP and reinforces the notion that **1** is properly considered as a phospha-Fischer carbene.

**Bonding.** The qualitative bonding scheme describing cyclic aminophosphonium ions as  $\sigma$ -donors and  $\pi$ -acceptors emerged concurrently with the publication of the first X-ray crystal structures of NHP–metal complexes.<sup>38</sup> By contrast, quantitative treatments of the  $\text{M}$ – $\text{P}$  bond in these systems are rare and vastly outnumbered by analyses of the  $\text{M}$ – $\text{C}$  interaction in metal–NHC complexes, which have been intensely scrutinized using a

(71) Tolman, C. A. *Chem. Rev.* **1977**, *77*, 313.

myriad of computational approaches.<sup>72–78</sup> The most authoritative investigations into M–NHP bonding have been published by Gudat and co-workers.<sup>24,33,79</sup> Together with others in the field,<sup>80</sup> they have analyzed the M–P bond primarily with orbital-based approaches, such as Mulliken and natural population analyses. To the best of our knowledge, there are no reports of an energy decomposition analysis (EDA) of M–NHP bonding within the framework of a fragment molecular orbital (FMO) approach.<sup>56–58</sup> This combination of methods is aptly suited for the analysis of donor–acceptor interactions because it allows a simple quantification of the relative importance of  $\sigma$ - and  $\pi$ -contributions to bonding.

The EDA approach has been extensively described in the literature,<sup>81</sup> and only a brief overview is presented herein. The energy associated with bond formation between two (or more) fragments—typically the metal and the ligand(s)—both possessing the geometry they take in the optimized complex, is referred to as the total bonding energy  $\Delta E_{\text{tot}}$ .<sup>82</sup> According to EDA,  $\Delta E_{\text{tot}}$  can be expressed as a sum of steric,  $\Delta E^0$ , and orbital,  $\Delta E_{\text{orbint}}$ , interactions, i.e.,  $\Delta E_{\text{tot}} = \Delta E^0 + \Delta E_{\text{orbint}}$ . The steric interaction consists of a purely electrostatic term,  $\Delta E_{\text{elstat}}$ , and a Pauli repulsion term,  $\Delta E^{\text{Pauli}}$ , between occupied orbitals on both fragments. The former stabilizes the bond, while the latter weakens it. It is the relative importance of the two terms that determines the effect of  $\Delta E^0$  on the total bonding energy. The orbital interaction energy has a stabilizing contribution and is calculated as the energy gain resulting from the relaxation of the FMOs to their optimal form in the complex. This term has an instructive interpretation as a sum of contributions from the various irreducible representations of the molecular point group. Therefore, by assessing the symmetry of the system, the  $\sigma$ - and  $\pi$ -characteristics of the interaction can be determined in a straightforward manner.

We have examined the bonding in complexes **1–4** using the EDA method. Because the M–P bond is the interaction of interest, the complexes were divided into two parts: the  $[\text{NHP}^{\text{Mes}}]^+$  ligand and the formally 14- and 16-electron fragments  $\text{M}(\text{PPh}_3)_2$  and  $\text{M}(\text{PMe}_3)_3$ , respectively ( $\text{M} = \text{Pd}, \text{Pt}$ ). Analogous valence isoelectronic complexes involving the neutral NHC<sup>Mes</sup> ligand (**6–9**, Chart 6) were included in the analysis for comparative purposes; the results can also be contrasted with those published earlier for a variety of Fischer carbenes.<sup>75,83,84</sup>

(72) Hu, X.; Tang, Y.; Gantzel, P.; Meyer, K. *Organometallics* **2003**, *22*, 612.

(73) Hu, X.; Castro-Rodriguez, I.; Olsen, K.; Meyer, K. *Organometallics* **2004**, *23*, 755.

(74) Nemcsok, D.; Wichmann, K.; Frenking, G. *Organometallics* **2004**, *23*, 3640.

(75) Frenking, G.; Solà, M.; Vyboishchikov, S. F. *J. Organomet. Chem.* **2005**, *690*, 6178.

(76) Jacobsen, H.; Correa, A.; Costabile, C.; Cavallo, L. *J. Organomet. Chem.* **2006**, *691*, 4350.

(77) Boehme, C.; Frenking, G. *Organometallics* **1998**, *17*, 5801.

(78) Jacobsen, H.; Correa, A.; Poater, A.; Costabile, C.; Cavallo, L. *Coord. Chem. Rev.* **2008**, doi:10.1016/j.ccr.2008.06.006.

(79) Gudat, D. *Eur. J. Inorg. Chem.* **1998**, 1087.

(80) Takano, K.; Tsumura, H.; Nakazawa, H.; Kurakata, M.; Hirano, T. *Organometallics* **2000**, *19*, 3323.


(81) Bickelhaupt, F. M.; Baerends, E. J. *Reviews in Computational Chemistry*; Wiley-VCH: New York, 2000; Vol. 15.

(82) The total bonding energy calculated in this manner is, however, not the negative of bond dissociation enthalpy, as the former quantity does not take into account the fragment relaxation and reorganization processes inherent in real bonding schemes.

(83) Lein, M.; Szabó, A.; Kovács, A.; Frenking, G. *Faraday Discuss.* **2003**, *124*, 365.

(84) Cases, M.; Frenking, G.; Duran, M.; Solà, M. *Organometallics* **2002**, *21*, 4182.

**Chart 6. Theoretical Pt and Pd NHC Complexes, with M=C Bonds Drawn Analogously to the M=P Bonds of 1–4**



**Table 1. Results of Energy Decomposition Analyses (kJ mol<sup>-1</sup>) of 1–4 and 6–9**

	$\Delta E_{\text{tot}}$	$\Delta E^0$	$\Delta E_{\text{elstat}}$	$\Delta E^{\text{Pauli}}$	$\Delta E_{\text{orbint}}$	
					<i>a</i>	<i>b</i>
<b>1</b>	-318.5	195.5	-641.9	837.3	-191.0 (37%)	-323.0 (63%)
<b>2</b>	-265.1	138.8	-496.5	635.2	-133.2 (33%)	-270.7 (67%)
<b>3</b>	-359.0	175.9	-644.3	820.2	-534.9	
<b>4</b>	-293.2	136.7	-506.6	643.2	-429.9	
<b>6</b>	-226.3	121.5	-803.8	925.3	-230.2 (66%)	-117.5 (34%)
<b>7</b>	-157.4	67.2	-555.7	622.8	-140.4 (63%)	-84.2 (37%)
<b>8</b>	-191.1	114.9	-718.0	832.8	-305.9	
<b>9</b>	-126.9	68.2	-495.1	563.3	-195.1	

The geometries of all systems were fully optimized prior to EDA calculations. Selected calculated metrical parameters for **1** are given in the legend for Figure 2, and complete structural data are deposited as Supporting Information (SI). The calculated geometries are in good overall agreement with the experimental data and display all the expected trends; therefore an in-depth analysis of geometrical parameters is not warranted. However, special attention should be drawn to three points: (i) The M–P bond lengths in **1–4** are not significantly affected by the identity of the metal fragment and indicate multiple bonding in all compounds (the calculated bond lengths lie in the range 2.179–2.199 Å). In contrast, the calculated M–C bond lengths in the analogous NHC complexes **6–9** vary more widely (from 2.050 to 2.145 Å) and are slightly longer than typical measured NHC–Pt(0) and NHC–Pd(0) single bonds in two coordinate complexes (2.02–2.09<sup>85</sup> and 1.99–2.11 Å, respectively).<sup>86–90</sup> (ii) The geometry at the P- and C-atoms of **1–4** and **6–9**, respectively, is invariably trigonal planar, with the sum of bond angles being very close to 360°. (iii) The optimized geometries of **3** and **4** are tetrahedral at the metal center, in agreement with the NMR data.

Results of the EDA calculations are summarized in Table 1. They show that changing the metal from Pd to Pt strengthens the M–P bond by ca. 60 kJ mol<sup>-1</sup>; a change in the composition of the metal fragment from  $\text{M}(\text{PPh}_3)_2$  to  $\text{M}(\text{PMe}_3)_3$  further strengthens the bond by 40 kJ mol<sup>-1</sup>. The overall effect of steric interaction in **1–4** is destabilizing, unlike the case of Fischer carbenes,<sup>75,83,84</sup> because of the increased Pauli repulsion. However, the total orbital interaction, which is the more informative of the two contributors to  $\Delta E_{\text{tot}}$ , outweighs the steric

(85) Gelloz-Berthon, G.; Buisine, O.; Brière, J.-F.; Michaud, G.; Stérin, S.; Mignani, G.; Tinant, B.; Declercq, J.-P.; Chapon, D.; Markó, I. E. *J. Organomet. Chem.* **2005**, *690*, 6156.

(86) Arnold, P. L.; Cloke, F. G. N.; Geldbach, T.; Hitchcock, P. B. *Organometallics* **1999**, *18*, 3228.

(87) Caddick, S.; Cloke, F. G. N.; Hitchcock, P. B.; Leonard, J.; Lewis, A. K. d. K.; McKerrecher, D.; Titcomb, L. R. *Organometallics* **2002**, *21*, 4318.

(88) Jackstell, R.; Harkal, S.; Jiao, H.; Spannenberg, A.; Borgmann, C.; Röttger, D.; Nierlich, F.; Elliot, M.; Niven, S.; Cavell, K. J.; Navarro, O.; Viciu, M. S.; Nolan, S. P.; Beller, M. *Chem.–Eur. J.* **2004**, *10*, 3891.


(89) Clement, N. D.; Cavell, K. J.; Ooi, L. *Organometallics* **2006**, *25*, 4155.

(90) Fantasia, S.; Nolan, S. P. *Chem.–Eur. J.* **2008**, *14*, 6987.

term for all systems. Furthermore, the  $C_2$ -symmetry of **1** and **2** allows us to describe the M–P bond by clean deconvolution of  $\Delta E_{\text{orbint}}$  in these complexes into  $\sigma$ - and  $\pi$ -MOs, which belong to the irreducible representations  $a$  and  $b$  of the  $C_2$  point group, respectively<sup>91</sup> (Table 1). This treatment reveals that the bond has approximately 2/3  $\pi$ - and 1/3  $\sigma$ -character. Although the same analysis cannot be conducted for the tetrahedral compounds **3** and **4**, we note that their similarity to **1** and **2** in terms of other bonding characteristics (Table 1), together with the FMO analysis (*vide infra*), imply predominant  $\pi$ -bonding also in these complexes.

These findings are in stark contrast to those of NHC complexes **6–9**. Although carbenes have long been considered pure  $\sigma$ -donors, recent computational treatments have demonstrated a small to moderate (10–30%)  $\pi$ -bonding character, the significance of which increased with the increasing formal d-electron count.<sup>75,76,92</sup> Energy decomposition analysis for **6–9** yielded similar  $\pi$ -bonding contributions of 34% and 37% for **6** and **7**, respectively. For Fischer carbenes, the percentage of  $\Delta E_{\text{orbint}}$  attributable to  $\pi$ -bonding has been reported to lie between 25% and 50%.<sup>75,83,84</sup> Therefore, the bonding in **1–4** is clearly of Fischer type also at the quantitative level, and as anticipated,<sup>79</sup> the M–NHP bond is dominated by  $\pi$ -back-bonding.

The bonding in Fischer carbenes and M–NHC complexes can be explained qualitatively by dative and retrodative components, whose relative contributions lead to formal double and single M–C bonds in the former and latter species, respectively.<sup>93,94</sup> That is, in both cases there generally exists one  $\sigma$ -type MO that is a linear combination of occupied ligand-based orbitals (typically the highest occupied FMO) and unoccupied metal-based d-orbitals, and one  $\pi$ -type MO that is a linear combination of unoccupied ligand-based orbitals (typically the lowest unoccupied FMO) and occupied metal-based d-orbitals. The MOs of **1** and **2** differ significantly from the above classical description, however, and reflect the dominant nature of the  $\pi$ -interaction. Figure 3 illustrates this difference by comparing the important bonding MOs of **2** and **7**. In the NHP complexes **1** and **2**, the unoccupied metal-based d-orbitals make only a negligible to small contribution to the HOMOs of the resultant complexes (6.8% and 0.0%), whereas the  $\sigma$ -bonding MOs of the NHC complexes **6** and **7** contain 12.8% and 6.0% contributions, respectively, from the unoccupied metal-based d-orbitals. By contrast, there is a much greater mixing between the occupied metal-based d-orbitals and the lowest unoccupied FMO of the [NHP<sup>Mes</sup>]<sup>+</sup> fragment: this ligand-based FMO contributes 18.8% and 15.3% to the HOMOs of **1** and **2**, respectively. The analogous contributions to the  $\pi$ -bonding MOs of complexes **6** and **7** are only 8.3% and 5.2%, respectively. There is also a third bonding contribution in **1** and **2** that is absent from the analogous NHC complexes: due to the cationic charge borne by the [NHP<sup>Mes</sup>]<sup>+</sup> fragment, the  $\pi$ -symmetric LUMO+1 of the ligand is also sufficiently low in energy to accept electron density from the occupied metal-based d-orbitals. This results in a second back-bonding contribution that is, however, of lesser importance than the primary



**Figure 3.** Important bonding MOs in [(NHP<sup>Mes</sup>)Pd(PPh<sub>3</sub>)<sub>2</sub>]<sup>+</sup> (**2**) and (NHC<sup>Mes</sup>)Pd(PPh<sub>3</sub>)<sub>2</sub> (**7**). In all cases the coordination plane of the metal is oriented parallel to the page.

$\pi$ -interaction; the percentage contributions of LUMO+1 to the bonding MOs of the metal complex are 6.0% and 5.1% in **1** and **2**, respectively.

The qualitative bonding description of **3** and **4** (as well that of **8** and **9**) closely follows the above discussion. The most notable difference is the increased importance of the  $\sigma$ -type dative interaction: the unoccupied metal-based d-orbitals have 8.0% and 4.4% contributions to the occupied MOs of the resulting complexes **3** and **4**, respectively. Even so, the retrodative  $\pi$ -interaction dominates the bonding in both compounds. It can therefore be concluded that the net flow of electrons in **1–4** takes place from metal to ligand, but as evidenced by the calculated data, the transfer of electrons involves significantly less than an electron pair. Thus, the NHP complexes can be considered as containing the metal center in a formal zero oxidation state.

**Hydrolysis.** Serendipitous discovery of the reaction between **1** and H<sub>2</sub>O confirmed the second of the two hypotheses given above, i.e., that these phospha-Fischer carbenes should be susceptible to attack by nucleophiles. Indeed, **1** is highly sensitive to moisture and needs to be handled under a rigorously inert atmosphere. Despite repeated drying of solvents, the presence of adventitious water eventually leads to the decomposition of these complexes. Overnight incubation of reaction mixtures of [NHP<sup>Mes</sup>]OTf and Pt(PPh<sub>3</sub>)<sub>4</sub> in wet CH<sub>2</sub>Cl<sub>2</sub> or CDCl<sub>3</sub> solution invariably gives a mixture of the well-known Pt(II)–hydride [PtH(PPh<sub>3</sub>)<sub>3</sub>]OTf (**10**)<sup>95–98</sup> and the phosphine oxide **11**,

(91) We note that this is only a rough division, as proper definition of symmetry implies the presence of a mirror plane and not a  $C_2$  axis. Orbitals that should be classified antisymmetric and symmetric transfer as the  $a$  and  $b$  representations of the  $C_2$  point group, respectively. Consequently, the reported contributions to bonding are slightly under- and overestimated. However, the error is expected to be only minor, as the metal–phosphorus bond lies on the  $C_2$  axis.

(92) Xile, H.; Yongjun, T.; Gantzel, P.; Meyer, K. *Organometallics* **2003**, *22*, 612.

(93) Dewar, M. J. S. *Bull. Soc. Chim. Fr.* **1951**, *18*, C79.

(94) Chatt, J.; Duncanson, L. A. *J. Chem. Soc.* **1953**, 2939.



(95) Thomas, K.; Dumler, J. T.; Renoe, B. W.; Nyman, C. J.; Roundhill, D. M. *Inorg. Chem.* **1972**, *11*, 1795.

(96) Chinakov, V. D.; Il'inich, G. N.; Zudin, V. N.; Likhobolov, V. A.; Nekipelov, V. M. *J. Organomet. Chem.* **1989**, *366*, 421.

(97) Goel, R. G.; Srivastava, R. C. *J. Organomet. Chem.* **1983**, *244*, 303.

(98) Caputo, R. E.; Mak, D. K.; Willett, R. D.; Roundhill, S. G. N.; Roundhill, D. M. *Acta Crystallogr.* **1977**, *B33*, 215.



Scheme 4. Complete *in Situ* Hydrolysis of **1** in Wet CH<sub>2</sub>Cl<sub>2</sub> Solution (anion is TfO<sup>-</sup>)Scheme 5. Proposed Mechanisms for the Hydrolysis of **1** (anion is TfO<sup>-</sup>)


which has been independently characterized by Ackermann and Born (Scheme 4).<sup>66</sup>

Hydrolysis of **1** may reasonably occur via one of two routes, both of which formally involve hydration of the Pt=P double bond, but are differentiated by the regiochemistry of addition: (Scheme 5a) initial attack of H<sub>2</sub>O at the NHP, followed by protonation of the Pt center and subsequent release and tautomerization of the modified NHP ligand to give **11**; coordination of excess PPh<sub>3</sub> in solution either prior to, or after this release, would generate the hydride **10**; (Scheme 5b) nucleophilic attack of H<sub>2</sub>O at the Pt center and subsequent deprotonation of coordinated water by NHP-turned-phosphide. The resultant Pt(II)-hydroxide may then decompose by either of two reductive elimination pathways (both analogous to that determined by Alper<sup>99</sup> for the oxidation of phosphines by Pd(OH<sup>-</sup>) either to generate O=PPh<sub>3</sub> and an analogue of **10** that incorporates a secondary phosphine derived from the NHP and/or to generate **10** and **11**. Although the formation of O=PPh<sub>3</sub> is favored statistically by a 2:1 margin in route (b), we never observe it, nor do we see by NMR spectroscopy any metallic species other than **10** that incorporate phosphine and hydride.

The calculated natural atomic orbital charge distribution<sup>100</sup> in **1** (+1.09 for P, +0.03 for Pt) also implicates the more cationic NHP phosphorus atom as the likely site for nucleophilic attack by water. In addition, the total energies calculated for the P-OH/Pt-H and Pt-OH/P-H bonded intermediates in Scheme 5 indicate strong energetic preference for pathway (a): the initial attack of H<sub>2</sub>O at P leads to a structure that is approximately 90

kJ mol<sup>-1</sup> more stable than the alternative platinum hydroxide. We therefore conclude that route (a) is the more reasonable pathway. This mechanism is further supported by the calculated LUMO of **1**, which is localized predominantly on the NHP P-atom, with smaller contributions from the metal and N-atoms (Figure 4), and by the analogous reactivity of group 6 metal-NHP complexes, which undergo nucleophilic attack by Me<sup>-</sup> and EtO<sup>-</sup> at the NHP, although dissociation of the resultant phosphine is not observed in these cases.<sup>101</sup>



In both routes (a) and (b), an additional molar equivalent of PPh<sub>3</sub> is necessary for formation of the observed products. In crude reaction mixtures toward the preparation of **1** (Scheme 4), we never observe deposition of Pt black, presumably owing



**Figure 4.** Calculated representation of the LUMO of **1** showing localization predominantly at the NHP P-atom. The metal coordination plane is approximately parallel to the page, and that of the NHP ligand is perpendicular to it.

(99) Grushin, V. V.; Alper, H. *Organometallics* **1993**, *12*, 1890.

(100) Reed, A. E.; Weinstock, R. B.; Weinhold, F. *J. Chem. Phys.* **1985**, *83*, 735.

Scheme 6. Hydrolysis of Isolated **1**Scheme 7. Overall Hydrolysis Reactions of **1–5** in  $\text{CH}_2\text{Cl}_2$  or  $\text{CDCl}_3$  Solution at rt (all anions not shown are  $\text{TfO}^-$ )

to the presence of excess  $\text{PPh}_3$  from the  $\text{Pt}(\text{PPh}_3)_4$  starting material in these systems. However, deliberate addition of  $\text{H}_2\text{O}$  to concentrated  $\text{CH}_2\text{Cl}_2$  solutions containing pure, isolated samples of **1** gives the expected hydrolysis products **10** and **11**, and Pt black because of the stoichiometric deficiency of  $\text{PPh}_3$  in these reactions (Scheme 6). No intermediates en route to the bulk metal are detected.

The phosphine-containing hydrolysis product depends on the  $\text{M-NHP}$  starting material. Thus, the expected hydride  $[\text{HPt}(\text{PMe}_3)_3]\text{OTf}^{102}$  does not form as the final product when **3** is hydrolyzed; instead, we observe the known homoleptic complex  $[\text{Pt}(\text{PMe}_3)_4]\text{OTf}_2^{103,104}$  (Scheme 7). Presumably, the predicted cation  $[\text{HPt}(\text{PMe}_3)_3]^+$  is formed initially, but due to the greater Lewis basicity of  $\text{PMe}_3$  relative to  $\text{PPh}_3$ , it is much more hydridic than **10** and very easily protonated by adventitious  $\text{H}_2\text{O}$ ; the resultant coordinatively unsaturated  $[\text{Pt}(\text{PMe}_3)_3]^{2+}$  is then trapped by  $\text{PMe}_3$  to give the final product.

Hydrolysis of **5** gives the five-coordinate platinum hydride  $[\text{PtH}(\text{dppe-}\kappa^2\text{P})_2]\text{OTf}$ . A characteristic pentet (with Pt satellites) is observed in the high-field region of the  $^1\text{H}$  NMR spectrum of this species ( $\delta -10.6$ ,  $^1J_{\text{HPt}} = 645$  Hz,  $^2J_{\text{HP}} = 32$  Hz), which unambiguously indicates the presence of a hydride coupled to four equivalent phosphines and platinum. The  $^{31}\text{P}\{^1\text{H}\}$  spectrum

( $\delta 22.2$ ,  $^2J_{\text{HP}} = 22$  Hz,  $^1J_{\text{PPT}} = 2344$  Hz) $^{105}$  is very similar to that of the known compound  $[\text{PtH}(\text{dppe-}\kappa^2\text{P})_2]\text{PF}_6^{106}$  and analogous to that of the Pd complex, which has been made previously from  $[\text{Pd}(\text{dppe-}\kappa^2\text{P})_2]^{2+}$ . $^{107}$

## Conclusions

Taken together, metrical parameters, computational results, and spectroscopic and reactivity data collectively support the interpretation that the most appropriate representation of **1** (and **2**) is that which shows an  $\text{M}=\text{P}$  double bond and a formal positive charge residing on the NHP phosphorus, i.e., the phospho-Fischer carbene **D**, which is formally equivalent to **E** (Chart 4), albeit with diminished and augmented  $\sigma$ - and  $\pi$ -interactions, respectively, when compared to traditional carbon-based systems. This representation of the group 10 metal(0) compounds is warranted also on the grounds of direct analogy to the characteristics generally attributed to Fischer carbenes $^{108}$  and highlights the “diagonal relationship” $^{18}$  between C and P: Fischer carbenes are derived from singlet  $:\text{CR}_2$ , contain an electrophilic carbene center, have  $\pi$ -donor groups adjacent to the carbene center, are typically associated with late metals in low oxidation states, and are found in conjunction with  $\pi$ -accepting ancillary ligands. We note also that Fischer carbenes are recognized as two-electron

(101) Nakazawa, H.; Ohta, M.; Miyoshi, K.; Yoneda, H. *Organometallics* **1989**, *8*, 638.

(102) Packett, D. L.; Syed, A.; Troglor, W. C. *Organometallics* **1988**, *7*, 159.

(103) Hahn, F. E.; Langenhahn, V.; Lügger, T.; Pape, T.; Le Van, D. *Angew. Chem., Int. Ed.* **2005**, *44*, 3759.

(104) Kozelka, J.; Lüthi, H.-P.; Dubler, E.; Kunz, R. W. *Inorg. Chim. Acta* **1984**, *86*, 155.

(105) The  $^1\text{H}$  decoupler was on for this  $^{31}\text{P}$  experiment; however it was unable to fully decouple the hydride proton. As a result, a doublet was still visible, but the  $^1J_{\text{HP}}$  value was slightly smaller than that observed in the  $^1\text{H}$  NMR spectrum.

(106) Berning, D. E.; Noll, B. C.; DuBois, D. L. *J. Am. Chem. Soc.* **1999**, *121*, 1432.

(107) Aresta, M.; Quaranta, E. *J. Organomet. Chem.* **2002**, *662*, 112.

(108) Crabtree, R. H. *The Organometallic Chemistry of the Transition Metals*; John Wiley & Sons, Inc.: New Haven, CT, 2001.

donors in both the covalent and ionic counting schemes and do not change the formal oxidation state of the metal. We attribute these same characteristics to NHP complexes of Pt(0) and Pd(0).

Despite advances in synthesis, the onward reactivity of late metal–NHP complexes is all but unknown, and our initial investigations into substitution and hydrolysis reactions described herein will be followed by catalytic studies. The unique sterically encumbered and  $\pi$ -acidic character of NHP ligands holds great potential for rich and exciting new chemistry, as does the “operational unsaturation”<sup>109,110</sup> that lies in the possibility for trigonal-planar to pyramidal isomerism.

**Acknowledgment.** We acknowledge financial support from the Natural Sciences and Engineering Research Council

of Canada (NSERC), a Petro-Canada Young Innovator’s Award (to N.D.J.), the Academy of Finland (to H.M.T.), and an Ontario Graduate Scholarship (to C.A.C.).

**Supporting Information Available:** Calculated coordinates for **1–4** and **6–9** and X-ray crystallographic data for **1** (CIF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

OM800973V

---

(109) Johnson, T. J.; Huffman, J. C.; Caulton, K. G. *J. Am. Chem. Soc.* **1992**, *114*, 2725.

(110) Johnson, T. J.; Coan, P. S.; Caulton, K. G. *Inorg. Chem.* **1993**, *32*, 4594.