

PII: S0040-4039(96)01786-8

## Application of 1,2-Bis(2-bromo-3,5-di-t-butylphenyl)ethane to Preparation of Compounds Having Two Diphosphene Units

Masaaki Yoshifuji,\* Naoyuki Shinohara, and Kozo Toyota

Department of Chemistry, Faculty of Science, Tohoku University, Aoba, Sendai 980-77, Japan

Abstract: A sterically crowded bromobenzene, 1,2-bis(2-bromo-3,5-di-t-butylphenyl)ethane, was prepared and converted to the corresponding phosphonous dichloride, which reacted with lithium (2,4,6-tri-t-butylphenyl)phosphinde to give 1,2-bis[3,5-di-t-butyl-2-{(2,4,6-tri-t-butylphenyl)phosphinidenephosphino}phenyl]ethane; the photolysis of the compound gave *E*-bis(2,4,6-tri-t-butylphenyl)diphosphene and an internal Z-diphosphene; the tungsten complex of the latter was analyzed by X-ray crystallography. Copyright © 1996 Elsevier Science Ltd

Compounds with low coordinated heavier main group elements such as phosphorus are of current interest.<sup>1</sup> The 2,4,6-tri-*t*-butylphenyl group (hereafter abbreviated to Ar) is one of the most powerful protecting groups and by utilizing this substituent we have been successful in the first preparation of diphosphenes.<sup>2,3</sup> We report here on the utilization of 1,2-bis(2-bromo-3,5-di-*t*-butylphenyl)ethane (3)<sup>4</sup> for preparation of compounds containing two diphosphene units within one molecule.

2-Bromo-1-(bromomethyl)-3,5-di-*t*-butylbenzene (2) was prepared according to the method reported previously.<sup>5</sup> Reaction of 2 with phenyllithium afforded a homo-coupling product 3. To a solution of 2 (2.991 g, 8.26 mmol) in ether (40 mL) was added 4.24 mmol of phenyllithium (1.01 M in cyclohexane-ether) at room temperature and the resulting solution was stirred for 30 min. The reaction mixture was worked up as usual and recrystallization from hexane afforded 1.487 g of 3 in 64% yield.<sup>6</sup>



Lithiation of 3 in THF with butyllithium at -78 °C led to the formation of dilithio derivative 4, which was quenched with methanol to give 1 (abbreviated to ArraH<sub>2</sub>)<sup>7</sup> in 67% yield. When 4 was allowed to react with phosphorus trichloride, 1,2-bis[3,5-di-t-butyl-2-(dichlorophosphino)phenyl]ethane (5) was obtained. To a solution of 3 (236.5 mg, 0.419 mmol) in THF (10 mL) was added 0.924 mmol of butyllithium (1.68 M in hexane) at -78 °C, then 2.12 mmol of phosphorus trichloride was added, and the resulting mixture was warmed to room temperature to give 5 as a pentane-insoluble material (182.3 mg, 72%).<sup>6</sup> Attempts to prepare an internal diphosphene 6 from 5 with either magnesium metal, t-butyllithium, or lithium naphthalenide as a coupling reagent,<sup>1</sup> were not successful. But preparation of a compound containing two diphosphene units was successful. To a solution of ArPH<sub>2</sub> (100.0 mg, 0.36 mmol) in THF (10 mL) was added 0.36 mmol of



butyllithium (1.64 M in hexane) at -78 °C and the resulting solution of lithium (2,4,6-tri-*t*-butylphenyl)phosphide (ArPHLi)<sup>8</sup> was added dropwise to a THF (100 mL) solution of 5 (153.1 mg, 0.25 mmol) over a 15-min period at -78 °C. Then the mixture was warmed to room temperature and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU; 80  $\mu$ L, 0.53 mmol) was added to the solution at 0 °C. After usual workup, 1,2-bis[3,5-di-*t*-butyl-2-{(2,4,6-tri-*t*-butylphenyl)phosphinidenephosphino}phenyl]ethane (*E,E-*7; 37.3 mg)<sup>6</sup> was obtained in 20% yield based on ArPH<sub>2</sub>.

On the other hand, we and others have reported on generation of the symmetrical diphosphene 8 during the photolysis of some unsymmetrical diphosphenes such as ArP=PMes,<sup>9</sup> 2,4,6-(CF<sub>3</sub>)<sub>3</sub>C<sub>6</sub>H<sub>2</sub>P=PAr,<sup>10</sup> or  $(Me_5C_5)P=PAr$ .<sup>11</sup> When a benzene-d<sub>6</sub> solution of the diphosphene E,E-7 was irradiated with a 500-W Xelamp through a Toshiba Y-49 filter at room temperature for 30 min, a mixture consisting of E,Z-7,<sup>6</sup> E-8 ( $\delta_P$  = 494),<sup>1</sup> Z-8 ( $\delta_P = 370$ ),<sup>12,13</sup> and an internal Z-diphosphene Z-6 ( $\delta_P = 394$ ) together with E,E-7 was obtained in a molar ratio of 48:10:1:10:17, respectively.<sup>14</sup> Upon 18-h irradiation of *E,E-7* (86.8 mg, 0.0851 mmol), however, E-8 (almost quantitative yield) and a cyclotetraphosphane 9 (24.3 mg, 0.026 mmol)<sup>6</sup> were isolated. In a separate experiment, when a THF solution of an excess amount of W(CO)<sub>5</sub>(THF) was added to a photolysis mixture of E,E-7 (46.1 mg, 0.0452 mmol) after 30-min irradiation, [Z-6][W(CO)5]2 (10)<sup>6</sup> was isolated as red crystals (3.4 mg, 7% yield) together with  $[E-8][W(CO)_5]$  (11) as the major product (ca. 20%) yield).<sup>6,15</sup> The structure of the complex 10 was unambiguously confirmed by X-ray crystallography.<sup>16</sup> Figure 1 shows a molecular structure drawing of 10 and the structure is similar to that of [(Z)-Mes-P=P-Ar]Cr(CO)5<sup>17</sup> or, more likely, to that of [(Z)-Mes-P=P-Mes][Mo(CO)5]2.<sup>18</sup> The P-P bond length is 2.041(4) Å and is similar to 2.034(2) Å for E-8, 2.039(3) Å for [(Z)-Mes-P=P-Ar]Cr(CO)5, and 2.026(2) Å for [(Z)-Mes-P=P-Mes][Mo(CO)<sub>5</sub>]<sub>2</sub>. The dihedral angle  $\angle C(1)$ -P(1)-P(2)-C(7) is -2.0(5)°, indicating that the Zdiphosphene system [C(1),P(1),P(2),C(7)] is coplanar within 0.03(1) Å, while the atoms [W(1),P(1),P(2),W(2)] form another plane within 0.030(3) Å, making a dihedral angle of 15.8° to each other. Two phenyl rings are planar within 0.02 Å, respectively, faced at a dihedral angle of 51.1°. The atoms [C(6), C(21), C(30), C(12)] are coplanar within 0.03 Å, indicating the ethylene bridge takes an eclipsed conformation. Some important bond lengths and angles are shown in the caption to Fig. 1.

While Niecke *et al.* has reported an amino-stabilized Z-diphosphene,<sup>19</sup> during the photolysis of *E,E-7* an olefin-metathesis type of reaction<sup>20</sup> occurred to give an internal Z-diphosphene 6 as well as 8. Furthermore,



Figure 1. Molecular structure for 10 showing atom labeling scheme. The methyl groups C27–C29 are disordered but only the structure with major occupancy factor is displayed. Hydrogen atoms are omitted for clarity. Some important bond lengths and angles: W(1)–P(1), 2.491(3); P(1)–P(2), 2.041(4); P(1)–C(1), 1.835(10); W(2)–P(2), 2.484(3); W(1)–P(1), 2.491(3); P(2)–C(7), 1.83(1); W(1)–P(1)–P(2), 131.2(1); W(1)–P(1)–C(1), 120.9(4); P(2)–P(1)–C(1), 105.7(4); W(2)–P(2)–P(1), 129.3(1); W(2)–P(2)–C(7), 119.3(4); P(1)–P(2)–C(7), 109.9(4); C(6)–C(21)–C(30), 116.8(9); C(12)–C(30)–C(21), 114.5(9); W(1)–P(1)–P(2)–W(2), -4.7(3); W(1)–P(1)–P(2)–C(7), 160.9(4); W(2)–P(2)–P(1)–C(1), -167.6(4); C(6)–C(21)–C(30)–C(12), -3(1).

E/Z photoisomerization<sup>12,13</sup> was observed for both diphosphenes 7 and 8. Upon prolonged irradiation, compound 6 appeared to be dimerized to a cyclotetraphosphane 9. Although we have not been successful in isolation of Z-6 during the photolysis of E, E-7, the Z-diphosphene was trapped by a complex formation with tungsten carbonyl.

The authors thank Dr. Shigeru Sasaki at Tohoku University for his help in the X-ray crystallographic analysis of **10**. This work was supported in part by Grants-in-Aid for Scientific Research (Nos. 07216210 and 08454193) from the Ministry of Education, Science, Sports and Culture, Japanese Government.

## **References and Notes**

- 1. Regitz, M.; Scherer, O. J. Eds. Multiple Bonds and Low Coordination in Phosphorus Chemistry; Georg Thieme Verlag: Stuttgart, 1990.
- Yoshifuji, M.; Shima, I.; Inamoto, N.; Hirotsu, K.; Higuchi, T. J. Am. Chem. Soc. 1981, 103, 4587– 4589; 1982, 104, 6167.
- 3. Yoshifuji, M.; Shibayama, K.; Inamoto, N.; Matsushita, T.; Nishimoto, K. J. Am. Chem. Soc. 1983, 105, 2495-2497.
- A part of this work was presented at the 21st Symposium on Heteroatom Chemistry, Toyama (December 1993); Yoshifuji, M.; Toyota, K.; An, D.-L.; Otoguro, A.; Kamijo, K.; Sangu, S.; Nakamoto, R. Abstr. No. 44.
- 5. Yoshifuji, M.; Kamijo, K.; Toyota, K. Bull. Chem. Soc. Jpn. 1993, 66, 3440-3443.
- 6. 3: Colorless needles, mp 207 208 °C (hexane); <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.22 (18H, s, t-Bu), 1.57 (18H, s, t-Bu), 3.17 (4H, s, CH<sub>2</sub>), 6.93 (2H, d, <sup>4</sup>J = 2.5 Hz, arom.), and 7.31 (2H, d, <sup>4</sup>J = 2.5 Hz, arom.)

Hz, arom.); MS (70 eV) m/z (rel intensity) 566 (M++4; 3), 564 (M++2; 6), 562 (M+; 3), 549 (M+-Me+2; 2), 281 (M+/2; 5), and 57 (t-Bu+; 100). Found: C, 63.74; H, 7.93; Br, 28.08%. Calcd for C<sub>30</sub>H<sub>44</sub>Br<sub>2</sub>: C, 63.83; H, 7.86; Br, 28.31%. 5: Colorless grains, mp 234-235 °C (decomp); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  = 1.34 (18H, s, *p*-*t*-Bu), 1.61 (18H, d, <sup>5</sup>*J*<sub>PH</sub> = 1.2 Hz, *o*-*t*-Bu), 3.78 (4H, s, CH<sub>2</sub>), 7.36 (2H, dd,  ${}^{4}J_{PH}$  = 6.8 Hz and  ${}^{4}J_{HH}$  = 1.9 Hz, arom.), and 7.75 (2H, d,  ${}^{4}J_{HH}$  = 1.9 Hz, arom.);  ${}^{31}P{}^{1}H$  NMR (81 MHz, CDCl<sub>3</sub>)  $\delta$  = 166.9; MS (70 eV) m/z (rel intensity) 608 (M<sup>+</sup>+2; 5), 606 (M<sup>+</sup>; 4), 571 (M<sup>+</sup>-Cl; 9), 535 (M<sup>+</sup>-Cl<sub>2</sub>-1; 99), 504 (M<sup>+</sup>-PCl<sub>2</sub>-1; 60), and 57 (t-Bu<sup>+</sup>; 100). Found: m/z 606.1674. Calcd for  $C_{30}H_{44}Cl_4P_2$ : M, 606.1672. E,E-7: Orange solid, mp 186 – 187 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta =$ 1.32 (18H, s, p-t-Bu, Arra), 1.33 (18H, s, p-t-Bu, Ar), 1.35 (36H, s, o-t-Bu, Ar), 1.45 (18H, s, o-t-Bu, Arra), 3.18 (4H, s, CH<sub>2</sub>), 7.14 (2H, d,  ${}^{4}J_{HH} = 1.8$  Hz, arom., Arra), 7.39 (4H, s, *m*-Ar), and 7.44 (2H, d,  ${}^{4}J_{HH} = 1.8$  Hz, arom., Arra);  ${}^{31}P{}^{1}H{}$  NMR (81 MHz, CDCl<sub>3</sub>)  $\delta$  = ABq, 476.7 and 515.9,  ${}^{1}J_{PP}$  = 582.8 Hz; UV (hexane) 280 (log  $\varepsilon$  4.26), 327 (3.86), and 468 nm (2.68); MS (70 eV) m/z (rel intensity) 1017 (M+-1; <1), 742 (M+-ArP; 1), 710 (M+-ArP2; 4), 685 (M+-ArP-Bui; 33), 496 (M+-Ar<sub>2</sub>P<sub>2</sub>; 95), and 277 (ArP++1; 100). E,Z-7: ( $\delta_P$  = ABq, 519 and 487,  ${}^{1}J_{PP}$  = 580 Hz;  $\delta_P$  = ABq, 402 and 389,  ${}^{1}J_{PP} = 555$  Hz). 9: Colorless, mp >300 °C;  $\delta_{P} = -62$ ,  $\delta_{H} = 1.35$  (36H, s, t-Bu), 1.38 (36H, s, t-Bu), 3.00 (4H, m, CHH-CHH), 5.60 (4H, m, CHH-CHH), 7.17 (4H, m, arom), 7.31 (4H, m, arom); MS (FAB) 931 (M<sup>+</sup>-1). 10: Red crystals, mp 120 °C (decomp);  $\delta_P = 332$ ,  $J_{PW} = 145$  and 116 Hz;  $\delta_{\rm H}$  = 1.14 (18H, s, t-Bu), 1.45 (18H, s, t-Bu), 3.14 (2H, m, CHH-CHH), 3.50 (2H, m, CHH-CHH), 6.74 (2H, m, arom), 7.22 (2H, m, arom).  $11:^{15} \delta_{\rm P} = ABq$ , 486 and 375 ( $J_{\rm PW} = 247$  Hz),  ${}^{1}J_{\rm PP}$ = 561.5 Hz.

- 7. Aurich, H. G.; Eidel, J.; Schmidt, M. Chem. Ber. 1986, 119, 18-35.
- Issleib, K.; Schmidt, H.; Wirkner, Chr. Z. Anorg. Allg. Chem. 1982, 488, 75-79; Yoshifuji, M.; Toyota, K.; Shibayama, K.; Inamoto, N. Chem. Lett. 1983, 1653-1656.
- 9. Yoshifuji, M.; Sato, T.; Inamoto, N. Bull. Chem. Soc. Jpn. 1989, 62, 2394-2395.
- Yoshifuji, M.; Abe, M.; Toyota, K.; Goto, K.; Inamoto, N. Bull. Chem. Soc. Jpn. 1993, 66, 1572– 1575.
- 11. Jutzi, P.; Meyer, U. J. Organomet. Chem. 1987, 333, C6-C20.
- 12. Diphosphene Z-8 is not stable at room temperature and so results isomerization to E-8.
- a) Caminade, A.-M.; Verrier, M.; Ades, C.; Paillous, N.; Koenig, M. J. Chem. Soc., Chem. Commun. 1984, 875–877; b) Yoshifuji, M.; Sato, T.; Inamoto, N. Chem. Lett. 1988, 1735–1738.
- 14. The molar ratio was estimated by the peak heights in the <sup>31</sup>P NMR. In addition to these peaks, an unknown peak at  $\delta_P = 435$  was observed, which could be assignable to *E*-6; however, no evidence for the structure was obtained.
- Yoshifuji, M.; Toyota, K.; Sato, T.; Inamoto, N. Science Reports, Tohoku Univ., Ser. 1 1990, 73, 1– 7.
- 16. Crystal data for 10: Recrystallized from benzene, P21/n (#14), C40H44O10P2W2, M<sub>r</sub> = 1114.43. Monoclinic, a = 16.597(5) Å, b = 11.175(7) Å, c = 23.411(3) Å, β = 98.38(2)°, V = 4295(2) Å<sup>3</sup>, Z = 4, T = 223 K; 8004 unique reflections with 2θ ≤ 50.0° were recorded on a four-circle diffractometer (MoKα radiation, graphite monochrometer). Of these, 5512 with I > 3σ(I) was judged as observed. The structure was solved with SHELXS86.<sup>21</sup> The methyl groups of one of the *p*-t-butyl groups (C27–C29) are disordered and the predominant occupancy factor for C27A–C29A is 0.532 while that for C27B–C29B is 0.468. The non-hydrogen atoms were refined anisotropically. Hydrogen atoms were included at calculated positions. R = 0.046, R<sub>w</sub> = 0.048. Further details of the crystal structure investigation are available on request from the Director of the Cambridge Crystallographic Data Centre, 12 Union Road, GB-Cambridge CB2 1EZ (UK).
- 17. Yoshifuji, M.; Hashida, T.; Inamoto, N.; Hirotsu, K.; Horiuchi, T.; Higuchi, T.; Ito, K.; Nagase, S. Angew. Chem., Int. Ed. Engl. 1985, 24, 211-212.
- 18. Lang, H.; Orama, O.; Huttner, G. J. Organomet. Chem. 1985, 291, 293-309.
- 19. Niecke, E.; Kramer, B.; Nieger, M. Angew. Chem., Int. Ed. Engl. 1989, 28, 215-217.
- 20. Dillon, K. B.; Gibson, V. C.; Sequeira, L. J. J. Chem. Soc., Chem. Commun. 1995, 2429-2430.
- a) Sheldrick, G. M. SHELXS86: Programs for the Automatic Solution of Crystal Structures, University of Göttingen, Germany (1986); b) Johnson, C. K. ORTEP-II, Oak Ridge National Laboratory Report, ORNL-TM-5138, Oak Ridge, TN (1976).

(Received in Japan 17 August 1996; revised 4 September 1996; accepted 10 September 1996)