

Asymmetric Synthesis of Simplactones A and B


by **Peddikotla Prabhakar, Dasari Ramesh, Singanaboina Rajaram, Dorigondla Kumar Reddy,**
and **Yenamandra Venkateswarlu***

Natural Products Laboratory, Organic Chemistry Division – I, Indian Institute of Chemical Technology,
Hyderabad – 500 007, India

(phone: +91-40-27193167; fax: +91-40-27160512; e-mail: luchem@iict.res.in)


A new, simple, and short route for the synthesis of simplactones A (**1**) and B (**2**) was achieved from a synthetically prepared chiral auxiliary, *i.e.*, the *Oppolzer* camphor-derived sultam **4**, and (4-methoxybenzyl)-protected 3-hydroxypropanal, in 52 and 48% overall yield, respectively, and with high diastereoselectivity (*Schemes 2 and 3*).

Introduction. – Simplactones are pharmacologically active marine secondary metabolites isolated from the Caribbean sponge *Plakortis simplex* [1]. Simplactones show *in vitro* cytotoxic activity against WEHI 164, murine fibrosarcoma cells. Simplactone A (**1**) and simplactone B (**2**) were first isolated by *Fattorusso* and co-workers in 1999 [1]. The configuration of the structures was revised by *Ogasawara* and co-workers by the synthesis from enantiomerically pure 4-(cumyloxy)cyclopent-2-en-1-one [2], and the structures were asymmetrically synthesized by *Osorio-Lozada* and *Olivo* through a double-diastereoselective acetate aldol reaction [3]. Recently, *Kamal* and co-workers [4] and *Rama Rao* and co-workers [5] reported a synthesis of simplactone A [4] and simplactone B [5], respectively. Structurally, this type of lactones show very good cytotoxic activity against WEHI 164, and many biologically active compounds like mevinolin, massiolactone [6], compactin, pironetin, phomalactone, and asperlin [7] contained this type of lactone moiety.



The biological potential of these compounds has stimulated us to synthesize **1** and **2** with *Oppolzer*'s camphor-derived sultam, which can be simply prepared compared with other chiral auxiliaries. The retrosynthetic analysis for the synthesis of **1** and **2** is shown in *Scheme 1* starting from (4-methoxybenzyl)-protected 3-hydroxypropanal and *N*-butanoylbornane-10,2-sultam (**3**). In this *Oppolzer* aldol addition, the 'syn' aldol reaction is giving a higher diastereomer excess (94% *de*) when compared to the 'anti'

Scheme 1. Retrosynthetic Analysis




aldol reaction with a *de* of 86%. Hence, we followed this approach for the synthesis of simpliclactones A and B.

Results and Discussion. – The stereoselective synthesis of simpliclactones A (**1**) and B (**2**) was carried out as shown in *Schemes 2* and *3*. Thus, the (2*S*)-*N*-butanoylbornane-10,2-sultam (**3**) [8] was subjected to asymmetric aldol reactions under two different conditions. When the reaction was carried out with 3.0 equiv. of (4-methoxybenzyl)-protected 3-hydroxypropanal, 3.0 equiv. of TiCl₄, and 2.2 equiv of ⁱPr₂NEt, the resulting major product was the ‘*anti*’ product **4** [9] (*de* 86% by chiral HPLC; 84% yield) (*Scheme 2*). However, when the reaction was carried out with 1.0 equiv. of (4-methoxybenzyl)-protected 3-hydroxypropanal, 1.0 equiv. of TiCl₄, and 1.2 equiv. of ⁱPr₂NEt, the resulting major product was the ‘*syn*’ product **10** [8–10] (*de* 94% by chiral HPLC; 92% yield) (*Scheme 3*). Hence we are reporting the same route for the synthesis of both simpliclactone isomers.

The ‘*anti*’ aldol compound **4** was further treated with LiAlH₄ [10] in dry Et₂O at 0° for 4 h, to give ‘*anti*’ diol **5** in 95% yield (*Scheme 2*), and in the same way, the ‘*syn*’ diol **11** was obtained in 95% yield (*Scheme 3*). The two OH groups in both isomers **5** and **11** were protected with ^tBuMe₂SiOTf in the presence of 2,6-lutidine [11] in dry CH₂Cl₂ to give **6** and **12**, respectively, in 95% yield. The (4-methoxybenzyl)-protecting group in **6** and **12** was removed with DDQ [12] to give **7** in 89% yield and **13** in 90% yield, respectively, which were oxidized with iodobenzene diacetate [13] and TEMPO (cat.) in dry CH₂Cl₂ to give **8** in 90% yield and **14** in 88% yield. The removal of the ^tBuMe₂Si groups in **8** and **14** was achieved by treatment with Bu₄NF [14] in dry THF furnishing lactols **9** and **15** in 90% yield. Finally, compounds **9** and **15** were oxidized with iodobenzene diacetate and TEMPO (cat.) in dry CH₂Cl₂ at 0° to give the required compounds **1** and **2** in good yield. The structures of **1** and **2** were established by their IR, ¹H- and ¹³C-NMR, and mass spectra, and their optical rotations were identical with those of the natural products reported by *Osorio–Lozada* and *Olivo* [3].

Scheme 2



a) (4-Methoxybenzyl)-protected 3-hydroxypropanal, TiCl₄, ⁱPr₂NEt, dry CH₂Cl₂, 3 h; 84%. b) LiAlH₄, dry Et₂O, 4 h; 95%. c) (*tert*-Butyl)dimethylsilyl trifluoromethanesulfonate (^tBuMe₂SiOTf = TBSOTf), 2,6-lutidine (=2,6-dimethylpyridine), dry CH₂Cl₂, 1 h; 95%. d) DDQ (4,5-dichloro-3,6-dioxocyclohexa-1,4-diene-1,2-dicarbonitrile), CH₂Cl₂, H₂O, 2 h; 89%. e) Iodobenzene diacetate, (=bis(acetato- κ O)phenyliodine), cat. 2,2,6,6-tetramethylpiperidin-1-yloxy (TEMPO), dry CH₂Cl₂, 3 h; 90%. f) Bu₄NF, THF, 2 h; 90%. g) Iodobenzene diacetate, cat. TEMPO, dry CH₂Cl₂, 3 h; 89%.


In conclusion, we have achieved a simple, short, and efficient total synthesis of simplactones A (**1**) and B (**2**) from (4-methoxybenzyl)-protected 3-hydroxypropanal and (2*S*)-*N*-butanoylbornane-10,2-sultam (**3**) in an overall yield of 52 and 48%, respectively. The advantage of this synthesis compared with the previous report [3] is the use of only one chiral auxiliary for the ‘*anti*’ and ‘*syn*’ isomers, the difference in the reaction conditions being the amount of TiCl₄ and aldehyde with respect to the chiral auxiliary **3**. In the synthesis of simplactones A and B reported by *Osorio-Lozada* and *Olivio* [3], two different chiral auxiliaries were used in a first step, and a further chiral auxiliary was present in a subsequent step.

The authors are thankful to the *Council of Scientific and Industrial Research (CSIR)*, New Delhi, India, for the financial support, and to Dr. *J. S. Yadav*, Director, Indian Institute of Chemical Technology (IICT), for his encouragement.

Experimental Part

General. Solvents were dried over standard drying agents and freshly distilled prior to use. The reagents were purchased from *Aldrich* and *Acros* and were used without further purification unless otherwise stated. All moisture-sensitive reactions were carried out under N₂. Org. solns. were concentrated *in vacuo* below 40°. Column chromatography (CC): silica gel (*Acme's* 60–120 mesh). Optical rotations: *Horiba* high sensitive polarimeter *SEPA-300* at 25°. IR Spectra: *Perkin-Elmer-IR-683*

Scheme 3



a) (4-Methoxybenzyl)-protected 3-hydroxypropanal, TiCl₄, ⁱPr₂NEt, dry CH₂Cl₂, 3 h; 92%. b) LiAlH₄, dry Et₂O, 4 h; 95%. c) ^tBuMe₂SiOTf, 2,6-lutidine, dry CH₂Cl₂, 1 h; 95%. d) DDQ, CH₂Cl₂, H₂O, 2 h; 90%. e) Iodobenzene diacetate, cat. TEMPO, dry CH₂Cl₂, 3 h; 88%. f) Bu₄NF, THF, 2 h; 90%. g) Iodobenzene diacetate, cat. TEMPO, dry CH₂Cl₂, 3 h; 88%.

spectrophotometer with NaCl optics; $\tilde{\nu}$ in cm⁻¹. ¹H- (300 MHz) and ¹³C-NMR (75 MHz) Spectra: Bruker-Avance-300 instrument; in CDCl₃; δ in ppm rel. to Me₄Si as internal standard, *J* in Hz. MS: Agilent-Technologies-1100 instrument (Agilent Chemistation software); in *m/z*.

(2S)-N-Butanoylbornane-10,2-sultam (=1-[(3aR,6R,7aS)-Tetrahydro-8,8-dimethyl-2,2-dioxido-3H-3a,b-methano-2,1-benzisothiazol-1(4H)-yl]butan-1-one; **3**). To a stirred soln. of (+)-camphor-derived sultam (5 g, 23.25 mmol) and butanoyl chloride (4.88 ml, 46.51 mmol) in dry benzene (50 ml) under N₂ was added anh. CuCl₂ (0.62 g, 4.65 mmol), and the mixture was refluxed for 12 h. The hot mixture was filtered, the filter, washed with CH₂Cl₂ (20 ml), the combined filtrate concentrated and, the residue purified by CC: **3** (6.0 g, 92%). Colorless crystals. $[\alpha]_D^{25} = +91.9$ (*c* = 1.0, CHCl₃). ¹H-NMR: 3.82 (*t*, *J* = 6.2, 1 H); 3.39 (*q*, *J* = 13.5, 2 H); 2.63 (*q*, *J* = 8.3, 2 H); 2.01–2.11 (*m*, 2 H); 1.81–1.91 (*m*, 3 H); 1.68 (*q*, *J* = 7.2, 2 H); 1.32–1.45 (*m*, 2 H); 1.14 (*s*, 3 H); 0.96 (*s*, 3 H); 0.96 (*t*, *J* = 7.2, 3 H). ESI-MS: 286 (*[M + 1]*⁺).

(2R,3R)-2-Ethyl-3-hydroxy-5-[(4-methoxyphenyl)methoxy]-1-[(3aR,6R,7aS)-tetrahydro-8,8-dimethyl-2,2-dioxido-3H-3a,6-methano-2,1-benzisothiazol-1(4H)-yl]-pentan-1-one (**4**). To a cooled (–78°) soln. of **3** (1.0 g, 3.5 mmol) in dry CH₂Cl₂ (20 ml) and TiCl₄ (1.15 ml, 3.0 equiv.) was added slowly ⁱPr₂NEt (1.85 ml, 2.2 equiv.), and the mixture was stirred for 90 min, followed by addition of (4-methoxybenzyl)-protected 3-hydroxypropanal (2.0 g, 10.3 mmol) at –78°, and further stirring for 90 min. After completion of the reaction (TLC), the reaction was quenched with sat. NH₄Cl soln. (10 ml), the mixture extracted with CH₂Cl₂ (3 × 20 ml), the extract dried (Na₂SO₄) and concentrated, and the residue purified by CC (hexane/AcOEt 4:1): **4** (1.42 g, 84%). Colorless liquid. $[\alpha]_D^{25} = +40$ (*c* = 0.8, CHCl₃). IR (neat): 3479, 2959, 2929, 1688, 1513, 1244. ¹H-NMR: 7.20 (*d*, *J* = 8.3, 2 H); 6.80 (*d*, *J* = 8.3, 2 H); 4.41 (*s*, 2 H); 3.98–4.09 (*m*, 1 H); 3.87 (*t*, *J* = 6.0, 1 H); 3.78 (*s*, 3 H); 3.52–3.67 (*m*, 2 H); 3.42 (*q*, *J* = 13.5, 2 H); 3.18 (br. *s*, 1 H); 3.07 (*q*, *J* = 5.2, 1 H); 2.07 (*d*, *J* = 6.0, 2 H); 1.77–1.94 (*m*, 6 H); 1.33–1.46 (*m*, 2 H); 1.26 (*s*, 1 H); 1.16 (*s*, 3 H); 0.98 (*s*, 3 H); 0.92 (*t*, *J* = 7.5, 3 H). ¹³C-NMR: 171.1; 159.58; 129.35; 128.65 (2 C);

113.54 (2 C); 72.86; 72.10; 67.47; 65.40; 55.26; 53.28; 52.51; 44.58 (2 C); 40.93; 34.86; 32.90; 26.40; 21.68; 20.69; 19.94 (2 C); 11.62. ESI-MS: 480 ($[M+1]^+$).

(2S,3R)-2-Ethyl-5-[(4-methoxyphenyl)methoxy]pentane-1,3-diol (**5**). To a cooled (0°) stirred suspension of LiAlH_4 (168 mg, 4.42 mmol) in dry Et_2O (20 ml) was added **4** (1.0 g, 2.96 mmol), and the mixture was stirred for 3 h at 0°. After completion of the reaction (TLC), the reaction was quenched with sat. NH_4Cl soln., the mixture extracted with AcOEt (5×10 ml), the extract dried (Na_2SO_4) and concentrated, and the residue purified by CC (hexane/ AcOEt 7:3): pure **5** (760 mg, 95%). Colorless liquid. $[\alpha]_D^{25} = -4.54$ ($c = 0.6$, CHCl_3). IR (neat): 3419, 2926, 2863, 1612, 1512, 1247, 1086, 1033. $^1\text{H-NMR}$: 7.19 (d , $J = 8.3$, 2 H); 6.83 (d , $J = 9.0$, 2 H); 4.44 (s , 2 H); 3.81 (dt , $J = 2.2$, 5.2, 2 H); 3.79 (s , 3 H); 3.67–3.75 (m , 1 H); 3.54–3.67 (m , 2 H); 1.67–1.77 (m , 1 H); 1.51–1.55 (m , 1 H); 1.23–1.46 (m , 3 H); 0.94 (t , $J = 7.55$, 3 H). $^{13}\text{C-NMR}$: 159.5; 129.45; 129.36 (2 C); 113.87 (2 C); 76.46; 73.15; 69.64; 63.89; 55.26; 46.63; 34.64; 21.37; 11.77. ESI-MS: 291 ($[M+\text{Na}]^+$).

(5R,6S)-6-Ethyl-5-[2-[(4-methoxyphenyl)methoxy]ethyl]-2,2,3,3,9,9,10,10-octamethyl-4,8-dioxo-3,9-disilaundecane (**6**). To a cooled (0°) soln. of **5** (400 mg, 1.49 mmol) and 2,6-lutidine (0.672 ml, 6.28 mmol, 2 equiv.) in dry CH_2Cl_2 (10 ml) was added $\text{BuMe}_2\text{SiOTf}$ (0.696 ml, 2.86 mmol), and the mixture was stirred at r.t. for 45 min. After completion of the reaction (TLC), the reaction was quenched with sat. NH_4Cl soln. (5 ml), the mixture extracted with CH_2Cl_2 (3×10 ml), the extract dried (Na_2SO_4) and concentrated, and the residue purified by CC (hexane/ AcOEt 9:1): pure **6** (700 mg, 95%). Pale yellow liquid. $[\alpha]_D^{25} = -4.11$ ($c = 0.8$, CHCl_3). IR (neat): 2929, 2856, 1639, 1250, 1092. $^1\text{H-NMR}$: 7.25 (d , $J = 9.06$, 2 H); 6.87 (d , $J = 9.06$, 2 H); 4.41 (d , $J = 2.26$, 2 H); 3.92–4.04 (s , 1 H); 3.80 (s , 3 H); 3.42–3.60 (m , 4 H); 1.60–1.85 (m , 1 H); 1.23–1.60 (m , 3 H); 1.04–1.21 (m , 1 H); 0.90 (t , $J = 7.55$, 3 H); 0.88 (s , 9 H); 0.87 (s , 9 H); 0.04 (s , 6 H); 0.02 (s , 6 H). $^{13}\text{C-NMR}$: 159.5; 129.34; 129.25 (2 C); 113.86 (2 C); 73.13; 72.81; 69.67; 63.90; 55.20; 48.43; 43.3; 34.58; 25.85 (3 C); 25.59 (3 C); 19.1 (2 C); 10.38; –3.6 (2 C); –5.7 (2 C). ESI-MS: 519 ($[M+\text{Na}]^+$).

(3R,4S)-3-[[tert-Butyl]dimethylsilyl]oxy]-4-[[[tert-butyl]dimethylsilyl]oxy]methyl]hexan-1-ol (**7**). To a soln. of **6** (600 mg, 1.2 mmol) in $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$ 9:1 (20 ml) was added DDO (329 mg, 1.44 mmol), and the mixture was stirred at r.t. for 2 h. After completion of the reaction (TLC), the mixture was neutralized with sat. NaHCO_3 soln. (5 ml) and extracted with CH_2Cl_2 (3×10 ml), the combined org. phase washed with brine (10 ml), dried (Na_2SO_4), and concentrated, and the residue purified by CC (hexane/ AcOEt 7:3): pure **7** (400 mg, 89%). Pale yellow liquid. $[\alpha]_D^{25} = +2.72$ ($c = 0.6$, CHCl_3). IR (neat): 3462, 2955, 1635, 1219. $^1\text{H-NMR}$: 4.03–4.11 (m , 1 H); 3.67 (q , $J = 6.0$, 2 H); 3.43–3.64 (m , 2 H); 1.56–1.75 (m , 2 H); 1.23–1.37 (m , 2 H); 1.02–1.17 (m , 1 H); 0.92 (t , $J = 7.55$, 3 H); 0.89 (s , 18 H); 0.08 (s , 3 H); 0.07 (s , 3 H); 0.04 (s , 3 H); 0.03 (s , 3 H). $^{13}\text{C-NMR}$: 71.52; 61.68; 60.88; 47.90; 35.37; 29.70; 25.87 (6 C); 25.63; 20.34; 12.35; –4.46 (2 C); –5.49 (2 C). ESI-MS: 377 ($[M+1]^+$).

(3R,4S)-3-[[tert-Butyl]dimethylsilyl]oxy]-4-[[[tert-butyl]dimethylsilyl]oxy]methyl]hexanal (**8**). To a cooled (0°) soln. of **7** (300 mg, 0.79 mmol) in dry CH_2Cl_2 (15 ml) were added iodobenzene diacetate (384 mg, 1.19 mmol) and TEMPO (24.9 mg, 0.15 mmol), and the mixture was stirred at r.t. for 4 h. After completion of the reaction (TLC), the reaction was quenched with sat. NH_4Cl soln. (5 ml), the mixture extracted with CH_2Cl_2 (3×10 ml), the combined org. layer washed with brine (10 ml), dried (Na_2SO_4), and concentrated, and the residue purified by CC (hexane/ AcOEt 9:1): pure **8** (270 mg, 90%). Pale yellow liquid. $[\alpha]_D^{25} = +6.0$ ($c = 1.0$, CHCl_3). IR (neat): 2956, 2857, 1728, 1253, 1089. $^1\text{H-NMR}$: 9.80–9.76 (m , 1 H); 4.43–4.50 (m , 1 H); 3.57–3.68 (m , 1 H); 3.48 (dd , $J = 7.5$, 10.5, 1 H); 2.44–2.64 (m , 1 H); 1.51–1.69 (m , 1 H); 1.05–1.20 (m , 2 H); 0.92 (t , $J = 7.5$, 3 H); 0.89 (s , 9 H); 0.87 (s , 9 H); 0.07 (s , 3 H); 0.04 (s , 9 H). $^{13}\text{C-NMR}$: 202.85; 68.36; 62.34; 48.64; 47.30; 25.86 (6 C); 18.5 (2 C); 18.1; 12.4; –4.48 (2 C); –5.46 (2 C). ESI-MS: 397 ($[M+\text{Na}]^+$).

(4R,5S)-5-Ethyltetrahydro-2H-pyran-2,4-diol (**9**). To a cooled (0°) soln. of **8** (200 mg, 0.53 mmol) in dry THF (10 ml) was added 1.0M Bu_4NF in THF (0.53 ml, 0.53 mmol), and the mixture was stirred for 2 h at 0°. After completion of the reaction (TLC), the reaction was quenched with sat. NH_4Cl soln. (5 ml), the mixture extracted with AcOEt (3×10 ml), the combined org. phase washed with brine (10 ml), dried (Na_2SO_4), and concentrated, and the residual liquid purified by CC (hexane/ AcOEt 7:3): pure **9** (70 mg, 90%). Pale yellow liquid. $[\alpha]_D^{25} = +16.5$ ($c = 3.0$, CHCl_3). IR (neat): 3453, 2924, 1639, 1381, 1048. $^1\text{H-NMR}$: 5.31 (t , $J = 3.0$, 1 H); 4.72 ($br. s$, 1 H); 4.63 (dd , $J = 3.0$, 6.0, 1 H); 4.17–4.10 (m , 1 H); 3.69 (dd , $J = 3.7$, 10.3, 1 H); 3.53 (dd , $J = 6.7$, 11.3, 1 H); 1.98 ($br. s$, 1 H); 1.77–1.86 (m , 1 H); 1.50–1.69 (m , 2 H);

1.30–1.47 (*m*, 1 H); 0.96 (*t*, $J = 6.7$, 3 H). ^{13}C -NMR: 92.83; 67.73; 61.52; 43.76; 37.03; 21.08; 11.28. ESI-MS: 147 ($[M + 1]^+$).

(4*R*,5*S*)-5-Ethyltetrahydro-4-hydroxy-2H-pyran-2-one (**1**). To a cooled (0°) soln. of **9** (40 mg, 0.27 mmol) in dry CH_2Cl_2 (10 ml) were added iodobenzene diacetate (88 mg, 0.27 mmol) and TEMPO (cat.), and the mixture was stirred at r.t. for 4 h. After completion of the reaction (TLC), the reaction was quenched with sat. NH_4Cl soln. (5 ml), the mixture extracted with CH_2Cl_2 (3×10 ml), the combined org. layer washed with brine (10 ml), dried (Na_2SO_4), and concentrated, and the residual liquid purified by CC (hexane/ AcOEt 4:1): pure **1** (35 mg, 89%). Colourless liquid. $[\alpha]_{\text{D}}^{25} = +23.0$ ($c = 1.0$, CHCl_3) [3]. IR (neat): 3420, 2945, 1721, 1415, 1205. ^1H -NMR: 4.38 (*dd*, $J = 10.4$, 11.2, 1 H); 4.21 (*dd*, $J = 5.1$, 11.2, 1 H); 4.10 (*br. s*, 1 H); 2.90 (*br. s*, 1 H); 2.72 (*dd*, $J = 3.6$, 17.8, 1 H); 2.61 (*dd*, $J = 3.6$, 17.8, 1 H); 1.80–1.98 (*m*, 1 H); 1.40–1.50 (*m*, 1 H); 1.25–1.38 (*m*, 1 H); 0.98 (*t*, $J = 7.5$, 3 H). ^{13}C -NMR: 170.6; 69.2; 64.4; 39.2; 39.1; 19.2; 11.1. ESI-MS: 145 ($[M + 1]^+$).

(2*S*,3*R*)-2-Ethyl-3-hydroxy-5-[(4-methoxyphenyl)methoxy]-1-[(3*aR*,6*R*,7*aS*)-tetrahydro-8,8-dimethyl-2,2-dioxido-3*H*-3*a*,6-methano-2,1-benzisothiazol-1(4*H*)-yl]pentan-1-one (**10**). As described for **4**, with **3** (1.0 g, 3.5 mmol), TiCl_4 (0.38 ml, 1.0 equiv.), CH_2Cl_2 (20 ml), $^i\text{Pr}_2\text{NEt}$ (0.73 ml, 1.2 equiv.), and (4-methoxybenzyl)-protected 3-hydroxypropanal (0.68 g, 3.5 mmol): pure **10** (1.55 g, 92%). White solid. $[\alpha]_{\text{D}}^{25} = +33.07$ ($c = 0.7$, CHCl_3). IR (KBr): 3509, 2956, 2878, 1691, 1613, 1515, 1022. ^1H -NMR: 7.20 (*d*, $J = 8.3$, 2 H); 6.81 (*d*, $J = 8.3$, 2 H); 4.41 (*s*, 2 H); 3.99–4.11 (*m*, 1 H); 3.87 (*t*, $J = 6.0$, 1 H); 3.78 (*s*, 3 H); 3.53–3.67 (*m*, 2 H); 3.43 (*q*, $J = 13.5$, 2 H); 3.19 (*d*, $J = 1.51$, 1 H); 3.08 (*q*, $J = 5.2$, 1 H); 2.07 (*d*, $J = 6.79$, 2 H); 1.65–1.96 (*m*, 7 H); 1.33–1.51 (*m*, 2 H); 1.16 (*s*, 3 H); 0.97 (*s*, 3 H); 0.92 (*t*, $J = 7.5$, 3 H). ^{13}C -NMR: 175.1; 159.0; 130.1; 129.22 (2 C); 113.68 (2 C); 72.66; 70.05; 68.33; 65.25; 55.18; 53.20; 52.49; 48.3; 47.6; 44.56; 38.57; 33.89; 32.89; 26.32; 21.67; 20.71; 19.83; 11.02. ESI-MS: 502 ($[M + \text{Na}]^+$).

(2*R*,3*R*)-2-Ethyl-5-[(4-methoxyphenyl)methoxy]pentane-1,3-diol (**11**). As described for **5**, with LiAlH_4 (166 mg, 4.36 mmol), Et_2O (20 ml), and **10** (1.4 g, 2.92 mmol) (quenching with sat. Na_2SO_4 soln.): pure **11** (740 mg, 95%). Colorless liquid. $[\alpha]_{\text{D}}^{25} = -16.0$ ($c = 0.7$, CHCl_3). IR (neat): 3419, 2926, 2863, 1612, 1086, 1033. ^1H -NMR: 7.19 (*d*, $J = 8.3$, 2 H); 6.83 (*d*, $J = 9.0$, 2 H); 4.44 (*s*, 2 H); 4.03 (*td*, $J = 3.0$, 9.8, 1 H); 3.79 (*s*, 3 H); 3.55–3.75 (*m*, 4 H); 1.84–1.99 (*m*, 1 H); 1.51–1.65 (*m*, 2 H); 1.22–1.39 (*m*, 2 H); 0.95 (*t*, $J = 7.5$, 3 H). ^{13}C -NMR: 159.67; 129.43; 129.35 (2 C); 113.86 (2 C); 75.97; 73.12; 69.89; 64.20; 55.25; 46.09; 31.79; 19.03; 12.28. ESI-MS: 291 ($[M + \text{Na}]^+$).

(5*R*,6*R*)-6-Ethyl-5-[(2-[(4-methoxyphenyl)methoxy]ethyl)-2,2,3,3,9,9,10,10-octamethyl-4,8-dioxo-3,9-disilaundecane (**12**). As described for **6**, with **11** (400 mg, 1.49 mmol), CH_2Cl_2 (20 ml), 2,6-lutidine (0.672 ml, 6.28 mmol), and $^t\text{BuMe}_2\text{SiOTf}$ (0.696 ml, 2.86 mmol) (quenching with sat. NH_4Cl soln. (10 ml)): **12** (700 mg, 95%). Pale yellow liquid. $[\alpha]_{\text{D}}^{25} = +3.75$ ($c = 0.8$, CHCl_3). IR (neat): 2931, 2858, 1513, 1464, 1250, 1094. ^1H -NMR: 7.25 (*d*, $J = 8.3$, 2 H); 6.87 (*d*, $J = 8.3$, 2 H); 4.41 (*s*, 2 H); 3.91–4.00 (*m*, 1 H); 3.80 (*s*, 3 H); 3.6–3.68 (*dd*, $J = 5.2$, 9.8, 1 H); 3.41–3.58 (*m*, 3 H); 1.54–1.86 (*m*, 2 H); 1.23–1.48 (*m*, 3 H); 0.91 (*t*, $J = 7.5$, 3 H); 0.88 (*s*, 9 H); 0.87 (*s*, 9 H); 0.04 (*s*, 6 H); 0.02 (*s*, 6 H). ^{13}C -NMR: 159.72; 129.34; 129.22 (2 C); 113.83 (2 C); 75.90; 73.10; 69.85; 67.40; 64.14; 55.21; 48.17; 46.40; 25.86 (4 C); 25.60 (2 C); 18.99 (2 C); 12.23; –3.63 (2 C); –4.62 (2 C). ESI-MS: 519 ($[M + \text{Na}]^+$).

(3*R*,4*R*)-3-[[[(tert-Butyl)dimethylsilyl]oxy]-4-[[[(tert-butyl)dimethylsilyl]oxy]methyl]hexan-1-ol (**13**). As described for **7**, with **12**, $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$ 9:1, and DDQ. After completion of the reaction, the mixture was diluted with H_2O (10 ml) and extracted as described for **7**: pure **13** (410 mg, 90%). Pale yellow liquid. $[\alpha]_{\text{D}}^{25} = -2.7$ ($c = 0.5$, CHCl_3). IR (neat): 3453, 2929, 2858, 1638, 1253, 1090. ^1H -NMR: 3.94–4.04 (*m*, 1 H); 3.61–3.73 (*m*, 3 H); 3.51–3.59 (*m*, 1 H); 1.59–1.79 (*m*, 2 H); 1.32–1.46 (*m*, 1 H); 1.23–1.32 (*m*, 2 H); 0.93 (*t*, $J = 7.5$, 3 H); 0.89 (*s*, 9 H); 0.88 (*s*, 9 H); 0.09 (*s*, 3 H); 0.06 (*s*, 3 H); 0.03 (*s*, 6 H). ^{13}C -NMR: 71.52; 61.68; 60.89; 47.90; 35.37; 25.87 (6 C); 20.34; 18.72 (2 C); 12.35; –4.54 (2 C); –5.45 (2 C). ESI-MS: 377 ($[M + 1]^+$).

(3*R*,4*R*)-3-[[[(tert-Butyl)dimethylsilyl]oxy]-4-[[[(tert-butyl)dimethylsilyl]oxy]methyl]hexanal (**14**). As described for **8**, with **13**, CH_2Cl_2 , iodobenzene diacetate, and TEMPO: pure **14** (260 mg, 88%). Pale yellow liquid. $[\alpha]_{\text{D}}^{25} = -4.0$ ($c = 1.5$, CHCl_3). IR (neat): 2927, 2856, 1635, 1219. ^1H -NMR: 9.78–9.74 (*m*, 1 H); 4.31–4.37 (*m*, 1 H); 3.57–3.68 (*m*, 2 H); 2.43–2.63 (*m*, 2 H); 1.69–1.92 (*m*, 1 H); 1.18–1.66 (*m*, 1 H); 1.02–1.17 (*m*, 1 H); 0.92 (*t*, $J = 7.5$, 3 H); 0.89 (*s*, 9 H); 0.86 (*s*, 9 H); 0.06 (*s*, 3 H); 0.04 (*s*, 3 H); 0.04 (*s*, 6 H). ^{13}C -NMR: 202.56; 68.25; 60.88; 48.52; 48.38; 25.86 (4 C); 25.77 (2 C); 20.36; 18.18; 18.03; 12.21; –4.56 (2 C); –4.97 (2 C). ESI-MS: 397 ($[M + \text{Na}]^+$).

(4R,5R)-5-Ethyltetrahydro-2H-pyran-2,4-diol (**15**). As described for **9**, with **14**, THF, and 1.0M Bu₄NF in THF: pure **15** (70 mg, 90%). Pale yellow liquid. $[\alpha]_D^{25} = -32.3$ ($c = 1.0$, CHCl₃). IR (neat): 3450, 2923, 2854, 1639, 1460, 1220. ¹H-NMR: 5.35 (*t*, $J = 3.0$, 1 H); 4.95 (br. s, 1 H); 3.63–3.90 (*m*, 2 H); 3.27 (*dd*, $J = 6.0, 12.0$, 1 H); 2.15 (br. s, 1 H); 1.70–1.86 (*m*, 1 H); 1.55–1.69 (*m*, 2 H); 1.15–1.52 (*m*, 2 H); 0.94 (*t*, $J = 6.7$, 3 H). ¹³C-NMR: 93.67; 69.66; 62.41; 45.19; 38.97; 21.37; 11.70. ESI-MS: 147 ($[M + 1]^+$).

(4R,5R)-5-Ethyltetrahydro-4-hydroxy-2H-pyran-2-one (**2**). As described for **1**, with **15**, CH₂Cl₂, iodobenzene diacetate, and TEMPO: pure **2** (34 mg, 88%). Colorless liquid. $[\alpha]_D^{25} = -24.0$ ($c = 1.0$, CHCl₃) [3]. IR (neat): 3475, 2989, 2930, 1721, 1265, 1014. ¹H-NMR: 4.44 (*dd*, $J = 5.2, 10.4$, 1 H); 3.90–4.05 (*m*, 1 H); 2.78 (*dd*, $J = 5.8, 11.6$, 1 H); 2.52 (*dd*, $J = 5.8, 11.6$, 1 H); 1.50–1.88 (*m*, 2 H); 1.20–1.45 (*m*, 1 H); 1.01 (*t*, $J = 7.55$, 3 H). ¹³C-NMR: 171.0; 69.1; 67.8; 42.5; 38.4; 20.8; 11.1. ESI-MS: m/z 145 ($[M + 1]^+$).

REFERENCES

- [1] F. Cafieri, E. Fattorusso, O. Taglialatela-Scafati, M. Di Rosa, A. Ianaro, *Tetrahedron* **1999**, *55*, 13831.
- [2] M. Sato, H. Nakashima, K. Hanada, M. Hayashi, M. Honzumi, T. Taniguchi, K. Ogasawara, *Tetrahedron Lett.* **2001**, *42*, 2833.
- [3] A. Osorio-Lozada, H. F. Olivo, *J. Org. Chem.* **2009**, *74*, 1360.
- [4] A. Kamal, P. Venkat Reddy, S. Prabhakar, P. Suresh, *Tetrahedron: Asymmetry* **2009**, *20*, 1798.
- [5] M. Somi Reddy, M. Narender, K. Rama Rao, *Tetrahedron* **2007**, *63*, 11011.
- [6] G. W. K. Cavill, D. V. Clark, F. B. Whitefield, *Aust. J. Chem.* **1968**, *21*, 2819.
- [7] T. Honda, T. Kametani, K. Kanai, Y. Tatsuzaki, M. Tsubuki, *J. Chem. Soc., Perkin Trans. 1* **1990**, 1733; A. D. Argoudelis, J. F. Zieserl, *Tetrahedron Lett.* **1966**, *18*, 1969.
- [8] W. Oppolzer, J. Blagg, I. Rodriguez, E. Walther, *J. Am. Chem. Soc.* **1990**, *112*, 2767.
- [9] B. H. Fraser, D. M. Gelman, P. Perlmutter, F. Vountatos, *Tetrahedron: Asymmetry* **2006**, *17*, 1152; W. Oppolzer, C. Starkemann, I. Rodriguez, G. Bernardinelli, *Tetrahedron Lett.* **1991**, *32*, 61; P. Prabhakar, S. Rajaram, Y. Venkateswarlu, *Tetrahedron: Asymmetry* **2009**, *20*, 1806.
- [10] G. Kumaraswamy, M. Padmaja, B. Markondaiah, N. Jena, B. Sridhar, M. Udaya Kiran, *J. Org. Chem.* **2006**, *71*, 337.
- [11] P. V. Murphy, C. McDonnell, L. Hämig, D. E. Paterson, R. J. K. Taylor, *Tetrahedron: Asymmetry* **2003**, *14*, 79.
- [12] H. P. Acharya, Y. Kobayashi, *Tetrahedron Lett.* **2005**, *46*, 8435.
- [13] C. P. Narasimhulu, J. Iqbal, K. Mukkanti, P. Das, *Tetrahedron Lett.* **2008**, *49*, 3185.
- [14] I. Safir, J. I. Candela Lena, L. Finet, N. Birlirakis, S. Arseniyadis, *Tetrahedron: Asymmetry* **2005**, *16*, 3436.

Received January 13, 2011