



## Discovery of a synthetic dual inhibitor of HIV and HCV infection based on a tetrabutoxy-calix[4]arene scaffold

Lun K. Tsou<sup>a</sup>, Ginger E. Dutschman<sup>b</sup>, Elizabeth A. Gullen<sup>b</sup>, Maria Telpoukhovskaia<sup>a</sup>, Yung-Chi Cheng<sup>b,\*</sup>, Andrew D. Hamilton<sup>a,\*</sup>

<sup>a</sup> Department of Chemistry, Yale University, New Haven, CT 06520, United States

<sup>b</sup> Department of Pharmacology, School of Medicine, Yale University, New Haven, CT 06520, United States

### ARTICLE INFO

#### Article history:

Received 8 January 2010

Accepted 10 February 2010

Available online 13 February 2010

#### Keywords:

Protein surface recognition

Calixarenes

HIV–HCV coinfection inhibitors

### ABSTRACT

A potential anti-HIV and HCV drug candidate is highly desirable as coinfection has become a worldwide public health challenge. A potent compound based on a tetrabutoxy-calix[4]arene scaffold that possesses dual inhibition for both HIV and HCV is described. Structural activity relationship studies demonstrate the effects of lower-rim alkylation in maintaining cone conformation and upper-rim interacting head groups on the calix[4]arene play key roles for its potent dual antiviral activities.

© 2010 Elsevier Ltd. All rights reserved.

Coinfection with both human immunodeficiency virus 1 (HIV) and hepatitis C virus (HCV) has become a public health challenge, afflicting more than 10 million people worldwide.<sup>1,2</sup> Complicating the challenge is the realization that with the advancement in highly active antiretroviral therapy (HAART), liver disease has emerged as a leading cause of death among HIV-infected patients.<sup>3</sup> Treatment of coinfecting patients with HAART has resulted in more liver toxicity,<sup>4</sup> creating a major health problem as there is currently no specific anti-HCV drug available. Therefore, the discovery of a dual compound as a potential drug candidate that can block HIV and HCV infection would be desirable. Herein, we report the first synthetic dual inhibitor of HIV and HCV infection in vitro based on a new class of compounds derived from a tetrabutoxy-calix[4]arene scaffold.

The secretion of vascular endothelial growth factor (VEGF) is elevated in HIV infected T-cells and is a major factor in the development of Kaposi's sarcoma.<sup>5</sup> In addition, platelet derived growth factor receptor (PDGFR) is expressed by uninfected T-cell lines, in which secretion of PDGF is also observed.<sup>6</sup> Although the impact of these processes is known for the development of Kaposi's sarcoma, the effect of these growth factors and receptors on HIV replication itself is still under investigation. The discovery of calix[4]arene derivatives that block VEGF and PDGF with their respective receptors<sup>7–10</sup> provided an opportunity to investigate the effect of this compound series on the replication of HIV. More-

over, a number of studies have shown that certain calixarene derivatives have interesting activities against viruses, enveloped viruses, bacteria, fungi, and cancerous cells.<sup>11</sup> In these reports,<sup>12,13</sup> the macrocyclic scaffolds are based on pyrogallol calixarenes, calixarenes and resorcurenes of different ring sizes (4, 6, and 8 units) without functionality at the lower rim and with sulfite, phosphite or simple carboxylic acid groups on the top rim. These studies suggested that the antiviral activity was derived from the compounds binding to the viral envelope and inhibiting its interaction with cells.

In this study, we utilized the series of calix[4]arene derivatives that were previously shown to inhibit the binding of VEGF, PDGF to their respective receptors.<sup>9</sup> The three initial compounds tested for antiviral activity and cytotoxicity **1–3** contained a lower ring *n*-butyl group and an upper ring carboxamide-linked isophthalic group with a benzyl ester, free carboxylic acid and cyclohexylamide arm, respectively, on one of the carboxylic acids. Compound **2** showed the most promising anti-HIV activity (Table 1) with an EC<sub>50</sub> value of 0.3 μM and an ID<sub>50</sub> value of 90 μM in an MT-2 cellular assay. Compounds **1** and **3** showed little to no antiviral activity. Notably, these results showed no correlation between antiviral activity and the ability of the compounds to inhibit the interaction of VEGF and PDGF with their receptors. Compound **2** was equally potent against three different laboratory HIV strains: HIV-IIIB, NL-43, and LaiM184 V. Moreover, antiviral activity was determined in different cell lines including MT-2, which are human T-cells with HTLV-1 virus, CEMx174, which is a human T cell line, and TZM-bl, which is a HeLa cell line with a firefly luciferase, that is, controlled by an HIV promoter. These cell lines are different in their acquired viruses, extracellular proteins, and cytoplasm contents.


\* Corresponding authors. Tel./fax: +1 203 214 1089.

E-mail addresses: [ketsou@rockefeller.edu](mailto:ketsou@rockefeller.edu) (L.K. Tsou), [yccheng@yale.edu](mailto:yccheng@yale.edu) (Y.-C. Cheng), [andrew.hamilton@yale.edu](mailto:andrew.hamilton@yale.edu) (A.D. Hamilton).

**Figure 1.** Synthetic routes for: (A) upper rim modifications, mono-*t*-butyl-5-aminoisophthalate. (B) Lower rim *n*-butyl modifications on compounds **1**, **2** and **4–7**. (C) Lower rim benzylated compound **10** and tetrahydroxy calix[4]arene **8**. D. Compounds **9** and **11**.

**Table 2**

HIV and HCV antiviral activities and cytotoxicity assay results for compounds 1–11



Compds	HIV inhibition IC <sub>50</sub> <sup>a</sup> (μM)	Cytotoxicity IC <sub>50</sub> <sup>a</sup> (μM)	HCV inhibition IC <sub>50</sub> <sup>a</sup> (μM)	CEM IC <sub>50</sub> <sup>a</sup> (μM)
<b>1</b>	>50	>50	25.6	12.5
<b>2</b>	0.36 (±0.12)	>50	1.8 (±0.3)	50
<b>3</b>	>50	>50	4.1	>50
<b>4</b>	2.9 (±1.4)	13.0 (±1.4)	30.1	12.5
<b>5</b>	3.0 (±2.8)	>50	1.5 (±0.3)	82
<b>6</b>	2.4 (±1.4)	>50	4.1 (±1.8)	100
<b>7</b>	<16	16	5.7	<12.5
<b>8</b>	>50	>50	>50	>50
<b>9</b>	6.7 (±2.9)	>50	>50	ND
<b>10</b>	37.1 (±6.1)	>50	>50	50
<b>11</b>	0.3	18.0	1.4	0.46

<sup>a</sup> 50 μM is the cutoff for assays. Standard deviation is given in parentheses.

conformation through intramolecular hydrogen bonding. However, the cone conformation is in equilibrium with the inverted cone conformation through a partial cone intermediate.<sup>14</sup> Preorganization of the scaffold into a cone conformation for projection of the recognition groups appears to be important for both anti-HIV and anti-HCV activities.

While alkylation at the bottom rim was essential for activity, the importance of projected diacid groups on the top rim was also clear. Among the seven compounds with a butyl chain at the bottom rim, compound **2** exhibited the most potent anti-HIV activity. When the isophthalic derivative was replaced with aspartic acid in compound **5**, anti-HIV activity decreases about 10-fold but anti-HCV activity slightly improved, while cytotoxicity remained the same. This suggested that aromatic substitutions are superior to aliphatic ones at the upper rim for HIV inhibition, while anti-HCV activity appears not to be as sensitive to this change. Compounds **1** and **3** contained benzyl ester and cyclohexylamide derivatives of the isophthalate linkers, respectively, suggesting that substituting both acid groups are required for potent antiviral activity. Overall, these observations suggest that four charges on the top rim are important for anti-HIV and anti-HCV activities. The lack of antiviral activity in positively charged **7**, confirms the importance of the negative charges on the projected periphery of the compounds.

In conclusion, the results demonstrate remarkable anti-HIV and anti-HCV activities for a series of compounds based on the tetrabutyl-calix[4]arene scaffold. We have shown that maintaining the cone conformation of the scaffold is important for antiviral activity. In addition, aromatic isophthalate spacers at the upper rim are essential for anti-HIV activities and the diacid groups are also necessary for the observed anti-HCV effects. Furthermore, we have identified a potent compound that possesses dual inhibition for both HIV and HCV in vitro. Moreover, it retains potency against different HIV strains in different cell lines while maintaining low

cytotoxicity. The molecular targets and mechanisms for anti-HIV and anti-HCV activities with these calix[4]arene compounds are under investigation and will provide valuable insight for future attempts to improve potency.

### Acknowledgment

We thank the National Institutes of Health (GM 35208 to A.D.H) and (AI 38204 to Y.-C.C.) for financial support of this work.

### References and notes

- Kim, A. Y.; Chung, R. T. *Gastroenterology* **2009**, *137*, 795.
- Singal, A. K.; Anand, B. S. *World J. Gastroenterol.* **2009**, *15*, 3713.
- Weber, R.; Sabin, C. A.; Friis-Møller, N.; Reiss, P.; El-Sadr, W. M.; Kirk, O.; Dabis, F.; Law, M. G.; Pradier, C.; De Wit, S.; Akerlund, B.; Calvo, G.; Monforte, A.; Rickenbach, M.; Ledergerber, B.; Phillips, A. N.; Lundgren, J. D. *Arch. Intern. Med.* **2006**, *166*, 1632.
- Velasco, M.; Guijarro, C. *JAMA* **2000**, *283*, 2526.
- Ascherl, G.; Hohenadl, C.; Schatz, O.; Shumay, E.; Bogner, J.; Eckhart, L.; Tschachler, E.; Monini, P.; Ensoli, B.; Sturzl, M. *Blood* **1999**, *93*, 4232.
- Chi, K. D.; McPhee, R. A.; Wagner, A. S.; Dietz, J. J.; Pantazis, P.; Goustin, A. S. *Oncogene* **1997**, *15*, 1051.
- Blaskovich, M. A.; Lin, Q.; Delarue, F. L.; Sun, J.; Park, H. S.; Coppola, D.; Hamilton, A. D.; Sebt, S. M. *Nat. Biotechnol.* **2000**, *18*, 1065.
- Sun, J.; Blaskovich, M. A.; Jain, R. K.; Delarue, F.; Paris, D.; Brem, S.; Wotoczek-Obadia, M.; Lin, Q.; Coppola, D.; Choi, K.; Mullan, M.; Hamilton, A. D.; Sebt, S. M. *Cancer Res.* **2004**, *64*, 3586.
- Sun, J.; Wang, D. A.; Jain, R. K.; Carie, A.; Paquette, S.; Ennis, E.; Blaskovich, M. A.; Baldini, L.; Coppola, D.; Hamilton, A. D.; Sebt, S. M. *Oncogene* **2005**, *24*, 4701.
- Zhou, H.; Wang, D. A.; Baldini, L.; Ennis, E.; Jain, R.; Carie, A.; Sebt, S. M.; Hamilton, A. D. *Org. Biomol. Chem.* **2006**, *4*, 2376.
- Perret, F.; Lazar, A. N.; Coleman, A. W. *Chem. Commun. (Cambridge: England)* **2006**, 2425.
- Hwang, K.M.Q.; Mao, Y.; Liu, S.Y.; Choy, W.; Chen, J. Genelabs Technologies, I., USA, 1994.
- Coveney, D. C. Benjamin (AIDS Care Pharma Limited, I.), 2005.
- Veggel, F. C. J. M. V. In *Calixarenes in Action*; Cook, H., Ed.; Imperial College Press: London, 2000.