

Accepted Manuscript

Synthesis, cytotoxicity, antimicrobial and anti-biofilm activities of novel pyrazolo[3,4-b]pyridine and pyrimidine functionalized 1,2,3-triazole derivatives

P. Nagender, G. Malla Reddy, R. Naresh Kumar, Y. Poornachandra, C. Ganesh Kumar, B. Narsaiah

PII: S0960-894X(14)00441-7

DOI: <http://dx.doi.org/10.1016/j.bmcl.2014.04.084>


Reference: BMCL 21577

To appear in: *Bioorganic & Medicinal Chemistry Letters*

Received Date: 5 March 2014

Revised Date: 8 April 2014

Accepted Date: 22 April 2014



Please cite this article as: Nagender, P., Malla Reddy, G., Naresh Kumar, R., Poornachandra, Y., Ganesh Kumar, C., Narsaiah, B., Synthesis, cytotoxicity, antimicrobial and anti-biofilm activities of novel pyrazolo[3,4-b]pyridine and pyrimidine functionalized 1,2,3-triazole derivatives, *Bioorganic & Medicinal Chemistry Letters* (2014), doi: <http://dx.doi.org/10.1016/j.bmcl.2014.04.084>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.


Graphical Abstract

To create your abstract, type over the instructions in the template box below
Fonts or abstract dimensions should not be changed or altered.

Leave this area blank for abstract info.

Synthesis, cytotoxicity, antimicrobial and anti-biofilm activities of novel pyrazolo[3,4-b]pyridine and pyrimidine functionalized 1,2,3-triazole derivatives

P. Nagender, G. Malla Reddy, R. Naresh Kumar, Y. Poornachandra, C. Ganesh Kumar, B. Narsaiah*





Synthesis, cytotoxicity, antimicrobial and anti-biofilm activities of novel pyrazolo[3,4-b]pyridine and pyrimidine functionalized 1,2,3-triazole derivatives

P. Nagender^a, G. Malla Reddy^a, R. Naresh Kumar^a, Y. Poornachandra^b, C. Ganesh Kumar^b, B. Narsaiah^{a,*}

^aFluoroorganic Division, CSIR-Indian Institute of Chemical Technology, Tarnaka, Hyderabad-500007, India.

^bMedicinal Chemistry and Pharmacology Division, CSIR-Indian Institute of Chemical Technology, Tarnaka, Hyderabad-500007, India.

ARTICLE INFO

Article history:

Received

Revised

Accepted

Available online

Key words:

Pyrazolo[3,4-b]pyridine

3H-pyrimidin-4-one

Triazole

cytotoxicity

antimicrobial

anti-biofilm

ABSTRACT

A series of novel pyrazolo[3,4-b]pyridine and pyrimidine functionalized 1,2,3-triazole derivatives **8a-g** and **9a-g** were prepared starting from 6-trifluoromethylpyridine-2(1H)one **2** via selective *O*-alkylation, followed by cyclisation using hydrazine hydrate to obtain 6-(trifluoromethyl)-1*H*-pyrazolo[3,4-b]pyridin-3-amine **4**. Compound **4** was diazotized followed by reaction with sodium azide, resulted in 3-azido-6-(trifluoromethyl)-1*H*-pyrazolo[3,4-b]pyridine **5**. Compound **5** was further cyclized with *N/O*- propargylated pyrimidine derivatives under Sharpless conditions and obtained compounds **6** and **7**, respectively. Each set of compounds **6** and **7** were alkylated with different alkyl halides and obtained respective products **8** and **9**. All the products were screened for cytotoxicity against four human cancer cell lines such as A549 - Lung (CCL-185), MCF7 - Breast (HTB-22), DU145 - Prostate (HTB-81) and HeLa - Cervical (CCL-2), compounds **9d**, **9e** and **9f** which showed promising activity have been identified. The products were also screened for antimicrobial, anti bio-film and MBC activities. Promising compounds in each case have been identified.

Cancer is a multifaceted disease that represents one of the leading causes of mortality in developed countries. Worldwide, one in eight deaths is due to cancer and it is the second most common cause of death in the US, exceeded only by heart disease.¹ Chemotherapy is the mainstay for cancer treatment, the use of available chemotherapeutics is often limited due to undesirable side effects. It is important to identify new agents and new targets for the treatment of cancer.

Nitrogen heterocycles have received a great deal of attention in the literature due to their role as active pharmacophores of historical significance. Among these heterocyclic systems, especially those containing pyridine rings are associated with diverse pharmacological properties such as anticancer,^{2,3} antimicrobial,⁴⁻⁶ anticonvulsant,⁷ antiviral,⁸ anti-HIV,⁹ antifungal and antimycobacterial activities.¹⁰ The pyrazolo[3,4-b]pyridine framework is a key structural fragment of many heterocyclic compounds showing a broad spectrum of biological activities.¹¹⁻¹³

In the last decade, some heterocycles of this class have been found to regulate the cardiovascular system and possess antiviral,^{14,15} antileishmanial,¹⁶ GSK-3 inhibitor (Fig 1)¹⁷ and antimicrobial properties.¹⁸ Recently, it was found that the fluorine¹⁹ or trifluoromethyl^{20,21} group at a strategic position of an organic molecule dramatically alters the properties of molecule in

terms of lipid solubility, oxidative thermal stability, permeability and oral bioavailability thereby showed an enhancement of transport mechanism.

During the last two decades, several pyrimidine derivatives have been developed as chemotherapeutic agents and found wide clinical applications.²²⁻²⁴ Some of the promising pyrimidine derivatives such as Flufenimer,²⁵ Primsulfuron-methyl,²⁶ Diflufenimer²⁷ and Fluropyrimidol²⁸ were identified as insecticides, herbicides, fungicides, and plant growth regulators, respectively. The 5-fluorouracil²⁹ (Fig 1) and their congeners belong to a family of the most potent anticancer and antiviral drugs^{30,31}.





Fig 1. Bioactive compounds based on pyrimidine and pyridine derivatives.


* Corresponding author. Tel.: +91-40-27193185; fax: +91-40-27160387 ; e-mail: narsaiah@iict.res.in

Based on the importance of the two scaffolds i.e., the pyrazolo[3,4-b]pyridine and pyrimidine, it was proposed to conceive both the scaffolds in a single molecule to have promising activity. The two scaffolds were connected through the click reaction under Sharpless conditions^{32,33} and obtained pyrazolo[3,4-b]pyridine and pyrimidine functionalized 1,2,3-triazoles. The products were evaluated for cytotoxic activity against four human cancer cell lines such as A549 - Lung (CCL-185), MCF7 - Breast (HTB-22), DU145 - Prostate (HTB-81) and HeLa - Cervical (CCL-2). The products also screened for antimicrobial, anti biofilm and MBC activities. The compounds which showed promising activity in each case have been identified.



Scheme 1. Synthesis of 3-azido-6-(trifluoromethyl)-1H-pyrazolo[3,4-b]pyridine

Reagents and conditions: i) Pyridine, DCM, r.t., 4 h ii) Cyanoacetamide, Na/EtOH, reflux, 6 h iii) Ethyl 2-bromoacetate, K₂CO₃, acetone, reflux, 3 h, iv) Hydrazine hydrate, 100°C, 4 h v) NaNO₂, Conc. HCl, H₂O vi) NaN₃, H₂O.



Scheme 2. Synthesis of pyrazolo[3,4-b]pyridine and pyrimidine functionalized 1,2,3-triazole derivatives **8a-g** and **9a-g**

Reagents and conditions: i) CuI, THF, r.t., 14 h ii) K₂CO₃, Acetone, R-X, r.t., 4h.

The 3-cyano-6-(trifluoromethyl)-2(1H) pyridone **2**³⁴ was prepared by acylation of vinyl ethyl ether in pyridine at room temperature, followed by reaction with cyanoacetamide in presence of sodium ethoxide in ethanol. Compound **2** was selectively *O*-alkylated³⁵ with α -bromoethylacetate in acetone reflux using potassium carbonate as a base, followed by reaction with hydrazine hydrate by an efficient method to obtain 6-(trifluoromethyl)-1H-pyrazolo[3,4-b]pyridin-3-amine **4** in high yield. Compound **4**

was diazotized with sodium azide at room temperature as a result 3-azido-6-(trifluoromethyl)-1H-pyrazolo [3,4-b]pyridine **5** was formed. The synthetic sequence is depicted in Scheme 1.

The compounds 2-amino-3-prop-2-ynyl-6-trifluoromethyl-3H-pyrimidin-4-one (**A**) and 4-prop-2-ynoxy-6-trifluoromethyl-pyrimidin-2-ylamine (**B**) reported³⁶ by us, was further reacted independently with compound **5** in THF using Cu(I) as catalyst under Sharpless conditions and obtained exclusively 1,4-disubstituted pyrazolo[3,4-b]pyridine and pyrimidine functionalized 1,2,3-triazole derivatives **6** and **7** respectively. Compounds **6**, **7** were further alkylated with different alkyl halides and obtained exclusively *N*-alkylated pyrazolo[3,4-b]pyridine and pyrimidine functionalized 1,2,3-triazole derivatives **8a-g** and **9a-g** respectively as outlined in Scheme 2. All the products formed have been tabulated in Table 1.

Compd	R	Yield (%) ^a	M.P (°C)
6	H	71	221-223
8a	CH ₃ CH ₂ -	91	254-256
8b	CH ₃ CH ₂ OC(O)CH ₂ -	89	192-194
8c	CH ₂ =CH-CH ₂ -	93	240-242
8d	HC≡C-CH ₂ -	87	217-219
8e	CH ₃ (CH ₂) ₆ CH ₂ -	94	140-142
8f	CH ₃ (CH ₂) ₈ CH ₂ -	81	129-131
8g	CF ₃ (CF ₂) ₇ CH ₂ CH ₂ -	79	111-113
7	H	73	164-166
9a	CH ₃ CH ₂ -	93	181-183
9b	CH ₃ CH ₂ OC(O)CH ₂ -	87	145-147
9c	CH ₂ =CH-CH ₂ -	91	136-138
9d	HC≡C-CH ₂ -	83	198-200
9e	CH ₃ (CH ₂) ₆ CH ₂ -	93	102-104
9f	CH ₃ (CH ₂) ₈ CH ₂ -	85	115-117
9g	CF ₃ (CF ₂) ₇ CH ₂ CH ₂ -	89	161-163

Table 1. Physical properties of compounds **6**, **7**, **8a-g** and **9a-g**

^a isolated yields.

The cytotoxicity of all the synthesized compounds were screened against four human cancer cell lines namely, A549 - Lung (CCL-185), MCF7 - Breast (HTB-22), DU145 - Prostate (HTB-81) and HeLa - Cervical (CCL-2) using MTT assay.³⁷ 5-Fluorouracil was used as a positive control. IC₅₀ values of the test compounds for 24 h on each cell line was calculated and presented in Table 2. Most of the compounds were found to show significant cytotoxicity on all the cell lines. More specifically, compounds **6**, **8b**, **8d**, **8e**, **8f** and **9b** exhibited promising activity against all the four cell lines with IC₅₀ value of ≤ 21 μ M, whereas compound **8a**, **8c** and **9c** showed moderate activity against MCF7 and HeLa cell lines. Similarly, compound **7**, **9a**, **9d** and **9g** also showed specific activity against A549 and DU145 with IC₅₀ for compounds **7** and **9a** in the range of 22.2 – 28.9 μ M, while compound **9d** showed IC₅₀ values of 4.1 and 4.7 μ M, respectively. Among all the compounds, *O*-alkylated derivatives **9d**, **9e** and **9f** exhibited promising activity against A549 and DU145 cell lines with IC₅₀ value of ≤ 6.3 μ M. The remaining compounds did not show cytotoxicity up to the concentration of 100 μ g/mL.

The structure-activity relationship studies revealed that the alkyl (decyl and octyl) chain on the nitrogen promotes cytotoxicity as observed in compounds **8e**, **8f**, **9e** and **9f** against A549 and DU145 cell lines. Whereas the perfluoroalkyl chain on nitrogen did not show any activity against MCF7, HeLa cell lines and decreased activity against A549 and DU145 cell lines compared to the hydrogen substituent on nitrogen in compounds **6** and **7**. There was an increase in activity, by increasing the n-alkyl chain length (ethyl-octyl-decyl) (**9b**, **9e** and **9f**). From the data, *O*-alkylated triazole derivatives were more potent than *N*-alkylated triazole derivatives on A549 and DU145 cancer cell lines. Alternatively, *N*-alkylated triazole derivatives were more active than *O*-alkylated triazole derivatives on MCF7 and HeLa cancer cell lines. Among all the tested compounds, the compounds **9d**, **9e** and **9f** were considered as promising candidates.

Compounds **6**, **8a-g**, **7** and **9a-g** were also tested for antimicrobial activity³⁸ against several bacterial organisms: *Micrococcus luteus* MTCC 2470, *Staphylococcus aureus* MTCC 96, *Staphylococcus aureus* MLS-16 MTCC 2940, *Bacillus subtilis* MTCC 121, *Escherichia coli* MTCC 739, *Pseudomonas aeruginosa* MTCC 2453, *Klebsiella planticola* MTCC 530 and *Candida albicans* MTCC 3017 and the results are tabulated in Table 3. *O*-alkylated triazole derivatives **7**, **9b** and **9f** showed promising activity ($\leq 15.6 \mu\text{g/mL}$) against all the bacterial organisms except *Candida albicans* MTCC 3017. These conjugates were further studied for the bio-film inhibition and minimum bactericidal assays^{39, 40} and the results were tabulated in Tables 4 and 5. Compounds **7**, **9b** and **9f** exhibited good bio-film inhibitory activity against *Micrococcus luteus* MTCC 2470, *Staphylococcus aureus* MTCC 96, *Staphylococcus aureus* MLS-16 MTCC 2940, *Pseudomonas aeruginosa* MTCC 2453 and *Klebsiella planticola* MTCC 530 bacterial strains with IC_{50} values up to $7.8 \pm 0.11 \mu\text{g/mL}$ (except **9b** which showed IC_{50} value of $15.6 \pm 0.31 \mu\text{g/mL}$). The compound **9f** was found to be more promising with IC_{50} value of $3.9 \pm 0.14 \mu\text{g/mL}$ against *Staphylococcus aureus* MTCC 96, *Staphylococcus aureus* MLS-16 MTCC 2940, *Pseudomonas aeruginosa* MTCC 2453 and *Klebsiella planticola* MTCC 530 bacterial strains. Minimum bactericidal concentration (MBC) results showed that the compound **7** exhibited significant activity against *Escherichia coli* MTCC 739 ($7.8 \mu\text{g/mL}$), while compound **9f** showed promising activity ($7.8 \mu\text{g/mL}$) against *Staphylococcus aureus* MLS-16 MTCC 2940, and *Klebsiella planticola* (MTCC 530) organisms. Compound **9f** was found to be most potent among all the tested compounds.

Table 2. *In vitro* cytotoxicity of pyrazolo[3,4-b] pyridine and pyrimidine functionalized 1,2,3-triazole derivatives

IC_{50} = compound concentration required to inhibit tumor cell proliferation by 50%.

Test compds	IC_{50} values (in μM)			
	A549	MCF7	DU145	HeLa
6	9.5 \pm 0.19	14.4 \pm 0.26	10.5 \pm 0.22	16.1 \pm 0.32
8a	-	95.2 \pm 0.58	-	62.3 \pm 0.46
8b	16.2 \pm 0.22	12.2 \pm 0.46	18.3 \pm 0.16	11.4 \pm 0.28
8c	-	20.2 \pm 0.34	-	18.2 \pm 0.31
8d	21.2 \pm 0.32	12.4 \pm 0.18	23.4 \pm 0.32	10.2 \pm 0.30
8e	10.1 \pm 0.18	11.3 \pm 0.22	12.2 \pm 0.20	9.4 \pm 0.26
8f	14.3 \pm 0.21	15.7 \pm 0.18	17.1 \pm 0.35	13.4 \pm 0.19
8g	68.2 \pm 0.54	-	88.8 \pm 0.52	-
7	22.8 \pm 0.14	-	28.9 \pm 0.48	-
9a	25.4 \pm 0.22	-	27.7 \pm 0.26	-
9b	16.3 \pm 0.16	12.4 \pm 0.22	18.2 \pm 0.23	9.8 \pm 0.26
9c	-	19.4 \pm 0.24	-	18.4 \pm 0.23
9d	4.1 \pm 0.12	-	4.7 \pm 0.18	-
9e	5.7 \pm 0.22	24.7 \pm 0.16	6.3 \pm 0.21	22.7 \pm 0.11
9f	4.2 \pm 0.31	37.2 \pm 0.31	5.8 \pm 0.14	34.3 \pm 0.32
9g	33.7 \pm 0.25	-	37.7 \pm 0.26	-
4	22.4 \pm 0.22	58.4 \pm 0.42	29.1 \pm 0.33	60.2 \pm 0.29
5-FU	1.3 \pm 0.11	1.4 \pm 0.09	1.5 \pm 0.12	1.3 \pm 0.14

- activity >100 μM , A549 – Lung cancer (ATCC No. CCL-185), MCF7– Breast cancer (ATCC No. HTB-22), DU145 – Prostate cancer (ATCC No. HTB-81), HeLa - Cervical cancer (ATCC No. CCL-2)

Table 3. MIC values of the compounds **7** and **9a-g** against several standard strains

S. No	Test Compds	Minimum inhibitory concentration ($\mu\text{g/mL}$)							
		<i>M.l</i> ^a	<i>S.a</i> ^b	<i>S.a</i> ^c	<i>B.s</i> ^d	<i>B.s</i> ^d	<i>E.c</i> ^e	<i>P.a</i> ^f	<i>K.p</i> ^g
1	7	7.8	15.6	15.6	15.6	7.8	7.8	15.6	- ⁱ
2	9a	>250	>250	>250	>250	>250	>250	>250	- ⁱ
3	9b	>250	15.6	7.8	15.6	15.6	15.6	7.8	- ⁱ
4	9c	7.8	>250	>250	>250	>250	>250	>250	- ⁱ
5	9d	>250	>250	>250	>250	>250	>250	>250	- ⁱ
6	9e	>250	>250	>250	>250	>250	>250	>250	- ⁱ
7	9f	15.6	7.8	7.8	15.6	7.8	7.8	7.8	- ⁱ
8	9g	>250	>250	>250	>250	>250	>250	>250	- ⁱ
	Ciprofloxacin	0.9	0.9	0.9	0.9	0.9	0.9	0.9	- ⁱ
	Miconazole	-	-	-	-	-	-	-	7.8

^a *Micrococcus luteus* MTCC 2470, ^b *Staphylococcus aureus* MTCC 96, ^c *Staphylococcus aureus* MLS-16 MTCC 2940, ^d *Bacillus subtilis* MTCC 121,

^e *Escherichia coli* MTCC 739, ^f *Pseudomonas aeruginosa* MTCC 2453, ^g *Klebsiella planticola* MTCC 530, ^h *Candida albicans* MTCC 3017, ⁱ No activity

Table 4. Anti bio-film activity of compounds **7**, **9b** and **9f** against five different bacterial strains

S. No.	Test Compounds	IC ₅₀ values in (µg/ml)				
		<i>Micrococcus luteus</i> MTCC 2470	<i>Staphylococcus aureus</i> MTCC 96	<i>Staphylococcus aureus</i> MLS16 MTCC 2940	<i>Klebsiella planticola</i> MTCC 530	<i>Pseudomonas aeruginosa</i> MTCC 2453
1	7	7.8 ± 0.11	7.8 ± 0.22	7.8 ± 0.21	7.8 ± 0.34	3.9 ± 0.26
2	9b	7.8 ± 0.22	3.9 ± 0.19	7.8 ± 0.32	3.9 ± 0.15	15.6 ± 0.31
3	9f	3.9 ± 0.14	7.8 ± 0.36	3.9 ± 0.18	3.9 ± 0.24	3.9 ± 0.21
Erythromycin (Standard)		0.24 ± 0.31	0.32 ± 0.14	0.32 ± 0.14	0.21 ± 0.12	0.26 ± 0.11

Table 5. MBC values of the compounds **7**, **9b** and **9f** against several bacterial strains

S. No.	Test Compounds	Minimum bactericidal concentration (µg/ml)						
		<i>Micrococcus luteus</i> MTCC 2470	<i>Staphylococcus aureus</i> MTCC 96	<i>Staphylococcus aureus</i> MLS-16 MTCC 2940	<i>Bacillus subtilis</i> MTCC 121	<i>Escherichia coli</i> MTCC 739	<i>Pseudomonas aeruginosa</i> MTCC 2453	<i>Klebsiella planticola</i> MTCC 530
1	7	15.6	31.2	31.2	31.2	7.8	15.6	31.2
2	9b	15.6	15.6	15.6	15.6	31.2	15.6	15.6
3	9f	15.6	15.6	7.8	31.2	15.6	15.6	7.8
Ciprofloxacin (Standard)		0.58	0.58	1.17	1.17	0.58	0.58	1.17

A series of novel pyrazolo[3,4-b]pyridine and pyrimidine functionalized 1,2,3-triazole derivatives **6,7**, **8a-g** and **9a-g** were prepared and screened for their *in vitro* cytotoxicity against four human cancer cell lines as well as their antimicrobial, anti-bio-film and MBC activities. The compounds **9d**, **9e** and **9f** showed very good activity among all the compounds against A549 and DU145 cell lines. *O*-alkylated triazole derivatives were more active than *N*-alkylated triazole derivatives on A549 and DU145 cancer cell lines, whereas *N*-alkylated triazole derivatives were more active than *O*-alkylated triazole derivatives on MCF7 and HeLa cancer cell lines. From the antimicrobial, anti bio-film and MBC data, the compound **9f** was found to be the most potent amongst all the tested compounds.

Acknowledgement

Authors are thankful to the Council of Scientific and Industrial Research (CSIR), New Delhi, India for providing funding through XIIth five year plan project, DITSF, code:CSC-0204. Authors (P. Nagender, G. Malla Reddy, R. Naresh Kumar, Y. Poornachandra) are also thankful to CSIR for financial assistance in the form of senior research fellowship and contingency grant.

References and notes

- Jemal, A.; Siegel, R.; Ward, E.; Murray, T.; Xu, J. *CA Cancer J. Clin.* **2007**, *57*, 51.
- Srivastava, A.; Pandeya, S. N. *Int. J. Curr. Pharm. Rev. Res.* **2011**, *74*, 5.
- Mosaad Sayed Mohamed; Yara Essam El-Deen Awad1.; Salwa M. El-Hallouty.; Moustafa El-Araby. *Open Journal of Medicinal Chemistry*. **2012**, *2*, 78.
- Patel, N. B.; Agravat, S. N.; Shaikh, F.M. *Med. Chem. Res.* **2011**, *20*, 1033.
- Patel, N. B.; Agravat, S. N. *Chem. Heterocycl. Compd.* **2009**, *45*, 1343.
- El-Emary, T. I.; Shawkat A. A. *Molecules* **2012**, *17*, 14464.
- Paronikyan, E. G.; Noravyan, A. S.; Dzhagatspany, I. A.; Nazaryan, I. M.; Paronikyan, R.G. *Pharm. Chem. J.* **2002**, *36*, 465.
- Bernardino, A. M. R.; De Azevedo, A. R.; Pinheiro, L. C. D.; Borges, J. C.; Carvalho, V. L.; Miranda, M. D.; De Meneses, M. D. F.; Nascimento, M.; Ferreira, D.; Rebello, M. A. *Med.Chem. Res.* **2007**, *16*, 352.
- Tucker, T. J.; Sisko, J. T.; Tynebor, R. M.; Williams, T. M.; Felock, P. J.; Flynn, J. A.; Lai, M.; Liang, Y.; McGaughey, G.; Liu, M.; *J. Med. Chem.* **2008**, *51*, 6503.
- Mamolo, M. G.; Zampieri, D.; Falagiani, V.; Vio, L.; Fermeglia, M.; Ferrone, M.; Priol, S.; Banfi, E.; Scialino, G.; *Arkivoc.* **2004**, *5*, 231.
- Kurumurthy, C.; Veeraswamy, B.; Sambasiva Rao, P.; Santhosh Kumar, G.; Shanthan Rao, P.; Loka Reddy, V.; Venkateswara Rao, J.; Narsaiah, B. *Bioorg. Med. Chem. Lett.* **2014**, *24*, 746.
- Hao, Y.; Xiao-Ping, X.; Chen, T.; LiZhao, L.; JunJi, S. *Org. Biomol. Chem.* **2012**, *10*, 724.
- Golubev, A. S.; Starostin, G. S.; Ghunikhin, K. S.; Peregudov, A. S.; Rodygin, K. C.; Rubtsova, S. A.; Slepukhin, P. A.; Kuchin, A. V.; Chkamonkov, N. D. *Russ. Chem. Bull.* **2011**, *60*, 733.
- Molecules. **2012**, *17*, 10915.
- De Clercq, E. *Curr. Opin. Microbiol.* **2005**, *8*, 552.
- Eizuru, Y. *Antivir. Chem. Chemother.* **2003**, *14*, 299.
- Witherington, J.; Bordas, V.; Garland, S. L.; Hickey, M. B.; Ife R. J.; John, L.; Saunders, M.; Smith, D. G.; Ward, R. W. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 1577.
- Mellol, H.; Echevarria, A.; Bernardino, A. M. R.; Canto-Cavalheiro, M.; Leon, L. L. *J. Med. Chem.* **2004**, *47*, 5427.
- Tarik, E. S. A. *Eur. J. Med. Chem.* **2009**, *44*, 4385.
- Hertel, L. W.; Kroin, J. S.; Misner, J. W.; Tustin, J. M. *J. Org. Chem.* **1988**, *53*, 2406.
- Filler, R.; *Organofluorine Compounds in Medicinal Chemistry and Biomedical Applications, Studies in Organic Chemistry.* **1993**, *48*, 362.
- Chae, J.; Konno, T.; Ishihara, T.; Yamanaka, H. *Chem. Lett.* **2004**, *33*, 314.
- Jain, K. S.; Chitre, T. S.; Miniya, P. B.; Kathiravan, M. K.; Bendre, V. S.; Veer, V. S.; Shahane, S. R.; Shishoo, C. J. *Current science.* **2006**, *90*, 6.
- Horton, T. M. *Clin. Cancer Res.* **2005**, *11*, 1884.
- Ambrus, J. L.; Stadler, I.; Kulaylat, M.; Koreschi, A.; Akhtar, S. J. *Med. Chem.* **1996**, *27*, 21.
- Fujii, K.; Nakamoto, Y.; Jpn. Kokai Tokkyo Koho JP 1149759. 1999; *Chem. Abstr.* **1999**, *130*, 209720 g.
- Meyer, W.; Forey, W. *Eur. Pat. Appl* EP 72347 (Ciba-Geigy. Basle; Feb. 1983); *Chem. Abstr.* **1983**, *98*, 215616q.
- Fujii, K.; Fukuda, Y.; Yamanaka Y.; Jpn. Kokai Tokyo koho JP 04235976. 1992; *Chem. Abstr.* **1993**, *118*, 38944d.
- Benefiel, R. L.; Belg. 815245 (Eli Lilly; 20 Nov. 1974; applic. 20 May 1974). *Chem. Abstr.* **1975**, *8*, 97354t.
- Cox, R. A. *Quart. Rev.* **1968**, *22*, 934.
- S. Raic-Malic, D. Sverduzic, T. Gazivoda, A. Marunovic, A. Hergold-Brundic, A. Nagl, J. Balzarini, E. De Clercq, M. Mintas, *J. Med. Chem.* **2000**, *43*, 4806.
- Douglas, K.T. *Med. Res. Rev.* **1987**, *7*, 441.
- Rostovtsev, V. V.; Green, L. G.; Fokin, V. V.; Sharpless, K. B. *Angew. Chem.* **2002**, *41*, 2596.
- Tornoe, C. W.; Christensen, C.; Meldal, M. *J. Org. Chem.* **2002**, *67*, 3057.

34. Langa, R.W.; Wenk, P.F. *Helv. Chim. Acta.* **1988**, 71.
35. Reddy, A. C. S.; Narsaiah, B.; Venkataratnam, R. V.; *J. Fluorine Chem.* **1996**, 78, 21.
36. Nagender, P.; Malla Reddy, G.; Naresh Kumar, R.; Chandrashekar Reddy, A.; Lokka Reddy, V.; Rajesh, P.; Venkateswara Rao, J.; Narsaiah, B. *Lett. Drug Design Discovery* **2013**, 10, 865.
37. Mosmann, T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods* **1983**, 65, 55.
38. Amsterdam, D. Susceptibility testing of antimicrobials in liquid media. In: Loman, V. (Ed.) *Antibiotics in Laboratory Medicine*, 4th Edition, Williams and Wilkins, Baltimore, MD, **1996**, 52–111.
39. Furlani, R. E.; Yeagley, A. A.; Melander, C. *Eur. J. Med. Chem.* **2013**, 62: 59.
40. National Committee for Clinical Laboratory Standards, NCCLS. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard fifth edition, NCCLS: Wayne, PA, **2000**.