

# Preparation and Properties of Sterically Protected Diphosphene and Fluorenylidene phosphine Bearing the 2,6-Di-*tert*-butyl-4-methoxyphenyl Group


Kozo Toyota, Subaru Kawasaki, Akitake Nakamura, and Masaaki Yoshifuji\*

Department of Chemistry, Graduate School of Science, Tohoku University, Aoba, Sendai 980-8578

(Received January 30, 2003; CL-030091)


A new bulky bromobenzene, 2-bromo-1,3-di-*tert*-butyl-5-methoxybenzene, was prepared and utilized to preparations of the corresponding diphosphene and fluorenylidene phosphine including low-coordinate phosphorus atom(s). An electronic perturbation of the *p*-methoxy group in the system was indicated by UV-vis spectra and  $^{31}\text{P}$  NMR chemical shifts.

Intramolecular interactions between a reactive site and the neighboring functional group are of current interest.<sup>1</sup> Sterically protected diphosphenes include an essentially reactive  $\text{P}=\text{P}$  bond in the molecule and they are kinetically stabilized by bulky groups such as the 2,4,6-tri-*tert*-butylphenyl group.<sup>2</sup> We have studied modification of the bulky aryl substituents at the *o*-position(s) and prepared 1,2-bis[2,4-di-*tert*-butyl-6-(dimethylamino)phenyl]diphosphene (**1b**) as an *o*-amino analogue of **1a**.<sup>3</sup> It should be noted that an attempted preparation of 1,2-bis(2,4-di-*tert*-butyl-6-methoxyphenyl)diphosphene (**1c**) has been unsuccessful.<sup>4</sup> In contrast, studies concerning the effect of *p*-substituent(s) in diaryldiphosphenes have been limited.<sup>5</sup> Although we have reported preparation of diphosphenes lacking the *p*-*tert*-butyl groups, i.e., 1,2-bis(2,6-di-*tert*-butylphenyl)diphosphene (**1d**),<sup>6</sup> the effect of introduction of an electron-donating or an electron-withdrawing substituent at the *p*-position of the 2,6-di-*tert*-butylphenyl group has remained unclear. We report here preparation and properties of *p*-methoxy substituted diphosphene **1e**, as well as fluorenylidene phosphines **2d** and **2e**.



**1a:**  $\text{R}^1=\text{R}^2=\text{t-Bu}$   
**b:**  $\text{R}^1=\text{t-Bu}$ ,  $\text{R}^2=\text{NMe}_2$   
**c:**  $\text{R}^1=\text{t-Bu}$ ,  $\text{R}^2=\text{OMe}$   
**d:**  $\text{R}^1=\text{H}$ ,  $\text{R}^2=\text{t-Bu}$   
**e:**  $\text{R}^1=\text{OMe}$ ,  $\text{R}^2=\text{t-Bu}$


**2a:**  $\text{R}^1=\text{R}^2=\text{t-Bu}$   
**d:**  $\text{R}^1=\text{H}$ ,  $\text{R}^2=\text{t-Bu}$   
**e:**  $\text{R}^1=\text{OMe}$ ,  $\text{R}^2=\text{t-Bu}$



First, we tried to prepare a bulky bromobenzene **3d**. Although the compound **3d** was first prepared by Rundel, starting from 2,4,6-tri-*tert*-butylaniline,<sup>7</sup> the reported yield was not good.<sup>6,7</sup> Thus, we have sought an alternative synthetic route and found that the Wolff-Kishner (the Huang-Minlon method)

reduction<sup>8</sup> of **4d**<sup>9</sup> afforded **3d** in good yield (70%).


Then we applied this methodology to the preparation of **3e** as follows: 2-bromo-1,3-bis(bromomethyl)-5-methoxybenzene<sup>10</sup> was allowed to react with KCN in acetonitrile (containing ca. 10% of water) in the presence of 18-crown-6 at room temperature for 24 h to give **5** in 66% yield.<sup>11</sup> Methylation of **5** to **6** (quant.) followed by treatment with diisobutylaluminum hydride gave **4e** (96% yield), which was then reduced to **3e** (48% yield)<sup>11</sup> by the Huang-Minlon method. Lithiation of **3e** with butyllithium followed by reaction with  $\text{PCl}_3$  gave **7** [ $^{31}\text{P}$  NMR ( $\text{THF}-\text{C}_6\text{D}_6$ )  $\delta_{\text{P}} = 153.3$ ], which was then converted to **1e** in 12% isolated yield.<sup>12</sup> Reaction of **7** with 9-(trimethylsilyl)-9-fluorenyllithium followed by treatment with  $n\text{-Bu}_4\text{N}^+\text{F}^-$  gave **2e**<sup>12</sup> (38% yield). Compound **2d** was prepared from **3d** by a method similar to that for **2e** (86% yield).<sup>12</sup>



18-c-6 = 18-crown-6; TmsFluLi = 9-(trimethylsilyl)-9-fluorenyllithium

Table 1 shows  $^{31}\text{P}$  NMR chemical shifts and UV-vis data of **1a,d,e** and **2a,d,e**. As for the  $^{31}\text{P}$  NMR chemical shift, electron-donating substituent in **1a,e** and **2a,e** causes a shift to a lower field, compared to that of **1d** and **2d**, respectively. The order of the  $\delta_{\text{P}}$  values is  $\text{d} < \text{a} < \text{e}$ , for both diphosphene and fluorenylidene phosphine series. In the case of UV-vis spectra of **1a,d,e**, the *p*-substitution causes red shifts of absorption of the longest  $\lambda_{\text{max}}$  attributable to  $n \rightarrow \pi^*$  transition (again, the order is  $\text{d} < \text{a} < \text{e}$ ). In contrast, *p*-substitution effect on the second  $\lambda_{\text{max}}$  ( $\pi \rightarrow \pi^*$  transition) is small. These facts may indicate that the *p*-methoxy group affects the electronic states of the phosphorus lone pair, although the effect on the  $\text{P}=\text{P}$   $\pi$  system is small, probably because the  $\pi$ -conjugation between the aromatic rings and the phosphorus double bonds is restricted, due to steric hindrance that prevents coplanar conformation.<sup>2,13</sup>

Structure of **1e** was unambiguously determined by X-ray crystallography.<sup>16</sup> Figure 1 shows a molecular structure of **1e**. Although the  $\text{P}-\text{C}_{\text{ipso}}$  bond lengths for **1e** [1.860(4) Å for  $\text{P}(1)-\text{C}(1)$  and 1.869(4) Å for  $\text{P}(2)-\text{C}(7)$ ] are very close to those



**Figure 1.** Molecular structure of **1e**, showing the atomic labeling scheme with thermal ellipsoids (50% probability). Hydrogen atoms are omitted for clarity. Some selected bond lengths (Å) and angles (°): P(1)–P(2), 2.043(1); P(1)–C(1), 1.860(4); P(2)–C(7), 1.869(4); O(1)–C(4), 1.372(4); O(2)–C(10), 1.370(4); P(1)–P(2)–C(7), 98.7(1); P(2)–P(1)–C(1), 100.5(1).

**Table 1.**  $^{31}\text{P}$  NMR and UV-vis data of compounds **1** and **2**

| Compd.    | $\delta_{\text{P}}^{\text{a}}$ | $\lambda/\text{nm}$ (log $\epsilon$ )                    |
|-----------|--------------------------------|----------------------------------------------------------|
| <b>1a</b> | 490.0 <sup>b</sup>             | 284 (4.19) <sup>c,d</sup> 340 (3.89) 460 (3.13)          |
| <b>1d</b> | 488.7 <sup>c</sup>             | 290 (4.03) <sup>c,e</sup> 340 (3.85) 450 (3.00)          |
| <b>1e</b> | 495.9                          | 291 (4.21) <sup>c</sup> 346 (3.75) 477 (3.12)            |
| <b>2a</b> | 256.5 <sup>f</sup>             | 238 (4.48) <sup>g</sup> 266 (4.30) 275 (4.32) 365 (4.05) |
| <b>2d</b> | 254.8                          | 237 (4.55) <sup>g</sup> 266 (4.40) 275 (4.45) 362 (4.25) |
| <b>2e</b> | 258.0                          | 239 (4.61) <sup>g</sup> 266 (4.42) 275 (4.46) 369 (4.30) |

<sup>a</sup>Measured in  $\text{CDCl}_3$ . <sup>b</sup>Data taken from Ref. 14. <sup>c</sup>Measured in  $\text{CH}_2\text{Cl}_2$ . <sup>d</sup>Data taken from Ref. 2a. <sup>e</sup>Data taken from Ref. 6. <sup>f</sup>Data taken from Ref. 1b, see also, Ref. 15. <sup>g</sup>Measured in hexane.

for **1a** [1.862(2) Å],<sup>2a</sup> the P=P bond length for **1e** [2.043(1) Å] is slightly longer than that for **1a** [2.034(2) Å]. The interplanar angles between the average plane [P(1), P(2), C(1), C(7)] and the aromatic rings [C(1)–C(6) and C(7)–C(12)] are 70.18(10) and 108.08(10)°, respectively, whereas the corresponding angle is 63.9° in **1a**. The interplanar angle between the two aromatic rings of **1e** is 70.7(1)°. The dihedral angle C(1)–P(1)–P(2)–C(7) for **1e** [165.0(2)°] is smaller than that for **1a** [172.2(1)°], which means that the deviation from the planarity of the –P=P– moiety is larger in **1e** than in **1a** in the crystal.

In summary, we have developed a new bulky substituent bearing a *p*-methoxy group. Some low-coordinated phosphorus compounds bearing the substituent were prepared and the effects of the *p*-substituent were evaluated. The method described here is promising for preparations of various *p*-functionalized bulky bromobenzenes. Studies on the reactivities of **1e** and **2e** are now in progress.

This work was supported in part by the Grants-in-Aid for Scientific Research (Nos. 13304049, 14044012, and 13640522) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

#### References and Notes

- a) J. G. Verkade, *Acc. Chem. Res.*, **26**, 483 (1993). b) K. Toyota, S. Kawasaki, and M. Yoshifuji, *Tetrahedron Lett.*, **43**, 7953 (2002).
- a) M. Yoshifuji, I. Shima, N. Inamoto, K. Hirotsu, and T. Higuchi, *J.*

- Am. Chem. Soc.*, **103**, 4587 (1981); M. Yoshifuji, I. Shima, N. Inamoto, K. Hirotsu, and T. Higuchi, *J. Am. Chem. Soc.*, **104**, 6167 (1982). b) M. Yoshifuji, *J. Organomet. Chem.*, **611**, 210 (2000).
- M. Yoshifuji, M. Hirano, and K. Toyota, *Tetrahedron Lett.*, **34**, 1043 (1993).
- M. Yoshifuji, D.-L. An, K. Toyota, and M. Yasunami, *Chem. Lett.*, **1993**, 2069.
- p*-Phenylene-bridged bis(diphosphene) derivative has recently been reported. See, S. Shah, T. Concolino, A. L. Rheingold, and J. D. Protasiewicz, *Inorg. Chem.*, **39**, 3860 (2000).
- M. Yoshifuji, T. Niitsu, D. Shiomi, and N. Inamoto, *Tetrahedron Lett.*, **30**, 5433 (1989).
- W. Rundel, *Chem. Ber.*, **101**, 2956 (1968).
- K. Toyota, Y. Matsushita, N. Shinohara, and M. Yoshifuji, *Heteroat. Chem.*, **12**, 418 (2001).
- K. Toyota, A. Nakamura, and M. Yoshifuji, *Chem. Commun.*, **2002**, 3012.
- L. A. van de Kuil, H. Luitjes, D. M. Grove, J. W. Zwikker, J. G. M. van der Linden, A. M. Roelofs, L. W. Jenneskens, W. Drenth, and G. van Koten, *Organometallics*, **13**, 468 (1994).
- 3e**: mp 59–61 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  = 1.63 (18H, s, *t*-Bu), 3.85 (3H, s, OMe), 7.02 (2H, s);  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  = 31.2 ( $\text{CMe}_3$ ), 38.7 ( $\text{CMe}_3$ ), 55.5 (OMe), 112.7, 115.9, 151.0, 157.9; MS (70 eV)  $m/z$  300 ( $\text{M}^+$ +2; 100), 298 ( $\text{M}^+$ ; 100), 285 ( $\text{M}^+$ –Me+2; 33), 283 ( $\text{M}^+$ –Me; 33). Found:  $m/z$  298.0934. Calcd for  $\text{C}_{15}\text{H}_{23}\text{BrO}$ : M, 298.0932. **4e**: mp 98–100 °C;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 1.54 (12H, s,  $\text{CMe}_2$ ), 3.88 (3H, s, OMe), 6.99 (2H, s), 9.78 (2H, s, CHO);  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 24.3 ( $\text{CMe}_2$ ), 53.2 ( $\text{CMe}_2$ ), 55.9 (OMe), 114.5, 115.7, 145.9, 159.6, 203.0 (CHO); MS  $m/z$  328 ( $\text{M}^+$ +2; 6), 326 ( $\text{M}^+$ ; 6), 247 ( $\text{M}^+$ –Br; 100). Found:  $m/z$  326.0521. Calcd for  $\text{C}_{15}\text{H}_{19}\text{BrO}_3$ : M, 326.0518. **5**: mp 155–157 °C;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 3.86 (4H, s,  $\text{CH}_2\text{CN}$ ), 3.88 (3H, s, OMe), 7.10 (2H, s);  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 26.1 ( $\text{CH}_2$ ), 56.2 (OMe), 115.2, 115.9, 116.9, 132.7, 159.8; MS  $m/z$  266 ( $\text{M}^+$ +2; 98), 264 ( $\text{M}^+$ ; 100). Found:  $m/z$  263.9905. Calcd for  $\text{C}_{11}\text{H}_9\text{BrN}_2\text{O}$ : M, 263.9898. **6**: mp 149–151 °C;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 1.95 (12H, s,  $\text{CMe}_2\text{CN}$ ), 3.87 (3H, s, OMe), 7.05 (2H, s);  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 28.7 ( $\text{CMe}_2$ ), 38.6 ( $\text{CMe}_2$ ), 56.0 (OMe), 113.6, 114.8, 123.7, 142.2, 159.2; MS  $m/z$  322 ( $\text{M}^+$ +2; 99), 320 ( $\text{M}^+$ ; 100). Found:  $m/z$  320.0520. Calcd for  $\text{C}_{15}\text{H}_{17}\text{BrN}_2\text{O}$ : M, 320.0524.
- 1e**: Orange prisms, mp 165–169 °C (dec.);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  = 1.41 (36H, s, *t*-Bu), 3.56 (6H, s, OMe), and 7.14 (4H, s);  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  = 55.4 (OMe);  $^{31}\text{P}\{^1\text{H}\}$  NMR (162 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  = 498.8; MS (70 eV)  $m/z$  500 ( $\text{M}^+$ ; 16) and 249 ( $\text{ArP}^+$ –1; 100). Found:  $m/z$  500.3000. Calcd for  $\text{C}_{30}\text{H}_{46}\text{O}_2\text{P}_2$ : M, 500.2973. **2d**: Yellow needles, mp 162–164 °C;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 1.46 (18H, s, *t*-Bu), 5.29 (1H), 6.78 (1H), 7.17 (1H), 7.29–7.37 (2H), 7.45–7.59 (4H), 7.64 (1H), and 8.28 (1H);  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 170.3 (d,  $^1J_{\text{PC}}$  = 42.2 Hz, P=C);  $^{31}\text{P}\{^1\text{H}\}$  NMR ( $\text{C}_6\text{D}_6$ )  $\delta$  = 253.5; MS  $m/z$  384 ( $\text{M}^+$ ; 27) and 219 ( $\text{ArP}^+$ –1; 100). Found:  $m/z$  384.2007. Calcd for  $\text{C}_{27}\text{H}_{29}\text{OP}$ : M, 384.2007. **2e**: Yellow prisms, mp 205–207 °C (dec.);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 1.44 (18H, s, *t*-Bu), 3.96 (3H, s, OMe), 5.56 (1H), 6.84 (1H), 7.14 (2H), 7.18 (1H), 7.28–7.37 (2H), 7.57 (1H), 7.64 (1H), and 8.27 (1H);  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 171.2 (d,  $^1J_{\text{PC}}$  = 43.2 Hz, P=C);  $^{31}\text{P}\{^1\text{H}\}$  NMR ( $\text{C}_6\text{D}_6$ )  $\delta$  = 257.2; MS  $m/z$  414 ( $\text{M}^+$ ; 18), 249 ( $\text{ArP}^+$ –1; 100), and 165 ( $\text{Flu}^+$ +1; 33). Found:  $m/z$  414.2104. Calcd for  $\text{C}_{28}\text{H}_{31}\text{OP}$ : M, 414.2113.
- R. Gleiter, G. Friedrich, M. Yoshifuji, K. Shibayama, and N. Inamoto, *Chem. Lett.*, **1984**, 313.
- D.-L. An, K. Toyota, M. Yasunami, and M. Yoshifuji, *J. Organomet. Chem.*, **508**, 7 (1996).
- V. D. Romanenko, A. V. Ruban, M. I. Povolotskii, L. K. Polyachenko, and L. N. Markovskii, *Zh. Obshch. Khim.*, **56**, 1186 (1986).
- Recrystallized from  $\text{CH}_2\text{Cl}_2$ –hexane, measured at 153 K.  $\text{C}_{30}\text{H}_{46}\text{O}_2\text{P}_2$ ,  $M$  = 500.64. Monoclinic, space group  $P2_1/n$  (#14),  $a$  = 10.181(7),  $b$  = 19.299(4),  $c$  = 15.413(3) Å,  $\beta$  = 104.75(3)°,  $V$  = 2928(2) Å<sup>3</sup>,  $Z$  = 4,  $D_c$  = 1.135 g cm<sup>–3</sup>,  $\mu$  (Mo K $\alpha$ ) = 1.72 cm<sup>–1</sup>. 3955 Unique reflections with  $2\theta \leq 50.0^\circ$ . Of these, 3683 with  $I > 2.0 \sigma(I)$  were used for  $R_1$  calculation. The structure was solved by direct methods. Non-hydrogen atoms were refined anisotropically. Hydrogen atoms were included but not refined.  $R_1$  = 0.073,  $R_w$  = 0.178. Crystallographic data have been deposited at the Cambridge Crystallographic Data Centre (No. CCDC-202408).