



1-Bromo-2-(diphenylphosphinoyl)ethyne and 1-bromo-2-(*p*-tolylsulfinyl)ethyne: versatile reagents eventually leading to benzocyclotrimers

Pierluigi Padovan^a, Stefano Tartaggia^a, Silvia Lorenzon^a, Enrico Rosso^a, Cristiano Zonta^b, Ottorino De Lucchi^a, Fabrizio Fabris^{a,*}

^a Dipartimento di Chimica, Università Ca' Foscari di Venezia, Dorsoduro 2137, I-30123 Venezia, Italy

^b Dipartimento di Scienze Chimiche, Università degli Studi di Padova, via Marzolo 1, I-35131 Padova, Italy

ARTICLE INFO

Article history:

Received 13 January 2009

Revised 6 February 2009

Accepted 10 February 2009

Available online 14 February 2009

Keywords:

Dienophile

Alkyne

Sulfoxide

Phosphine oxide

Cyclotrimerization

Synthetic equivalent


ABSTRACT

The title compounds form Diels–Alder cycloadducts with a number of dienes, which can be transformed into *vic*-bromo(trimethylstannyl)olefins, ultimate precursors for the synthesis of benzocyclotrimers via copper-mediated cyclotrimerization. The overall result is a formal condensation of three diatomic carbons with three dienes.

© 2009 Elsevier Ltd. All rights reserved.

Benzocyclotrimers are useful scaffolds in supramolecular applications,¹ such as complexation of fullerene,² chiral recognition of ammonium salts³ and gas hosting in molecular capsule.⁴ Furthermore, one of these cup-shaped compounds is the crucial precursor for the synthesis of the smallest, non planar fragment of fullerene, that is, sumanene.⁵ These and further applications call for general and effective synthetic methodologies. In principle, the Diels–Alder cycloaddition is the most valuable reaction for the synthesis of suitable substrates for the cyclotrimerization provided that the functional groups needed to activate the dienophile are convertible into halogen (bromine) and metal (trimethyltin) moieties which are needed to promote cyclotrimerization.^{1–3,5–7} In fact, copper(I) 2-thiophenecarboxylate (CuTC)-mediated cyclotrimerization of *vic*-bromo(trimethyltin)olefins is still the most reliable method to prepare benzocyclotrimers with a wide range of shapes and functional groups.^{6,8} In an alternative strategy, we have proved that these two functionalities can be introduced in the diene rather than in the dienophile (Scheme 1).^{8,9}

Both sulfoxide and phosphine oxide groups can efficiently activate triple bonds for Diels–Alder reactions.^{10,11} In addition, when treated with organometallic reagents they can act as leaving groups in olefins bearing vicinal halogens.^{12,13} For these reasons, 1-bromo-2-(diphenylphosphinoyl)ethyne and 1-bromo-2-(*p*-tolyl-



Scheme 1. Possible strategies for the synthesis of precursors of benzocyclotrimers.


sulfinyl)ethyne can be regarded as synthetic equivalents of 1-bromo-2-(trimethyltin)ethyne.

The synthesis of 1-bromo-2-(*p*-tolylsulfinyl)ethyne **1** turned out to be low yielding and troublesome,¹⁴ such as the cycloaddition reaction was tested with 1,3-cyclopentadiene only providing the expected cycloadducts in good yield as a separable 1:1 mixture of diastereomers.¹⁵ The latter underwent facile substitution at sulfur with phenylmagnesiumbromide. The resulting organomagnesium derivative was treated with trimethyltinchloride to afford 2-bromo-3-(trimethyltin)bicyclo[2.2.1]hepta-2,5-diene **3**^{7b,16} in good yields (Scheme 2).


In order to overcome the poor stability of **1**, 1-bromo-2-(diphenylphosphinoyl)ethyne was synthesized. Commercially available trimethylsilyl ethyne or sodium acetylide was converted into intermediate diphenylphosphinoethyne (DPE) in good yields.¹⁷ The

* Corresponding author. Tel.: +39 41 2348908; fax: +39 41 2348517.

E-mail address: fabrisfa@unive.it (F. Fabris).



Scheme 2. Synthesis of norbornadiene benzocyclotrimers, via cycloaddition of bromo sulfinyl alkyne **1** and sulfinyl replacement.



Scheme 3. Synthesis of dienophile **7**.


Table 1
Cycloaddition of **7** with 1,3-dienes

Entry	Diene	Cycloadduct	Yield ^a (%)
1			27 ^b
2			48 ^b
3			80
4			53
5			27
6			22 ^c
7			94
8			34
9			72


^a Isolated yields.

^b Aromatization occurred.


^c Aromatization occurred via retro Diels–Alder.



Scheme 4. High-pressure Diels–Alder reaction of (–)-nopadiene with **7**.



Scheme 5. Possible paths of organometallic intermediate generated by **10**.



Scheme 6. Alternative disconnections for benzocyclotrimers: tribenzynes cycloaddition (left), versus cyclo-condensation of diatomic carbon (right).

bromination of DPE was achieved with *N*-bromosuccinimide in the presence of catalytic amounts of silver(I) nitrate,¹⁸ to afford **7** in good yield and reasonable purity after a simple filtration over basic alumina (Scheme 3).¹⁹

The cycloaddition of **7** was tested with a number of dienes under standard conditions at 120 °C, affording the expected products in variable yields (Table 1).²⁰ Nevertheless in the case of cyclopentadiene, when milder conditions (80 °C) were applied for a prolonged time (24 h), a nearly quantitative amount of cycloadduct was obtained.

Other 1,3-dienes (1,3,5,7-cyclooctatetraene, α -phellandrene, naphthalene, 1*H*-pyrrol-*N*-(*t*-butyl)carbamate, 1-tosyl-1*H*-pyrrole and nopadiene) failed to afford the expected cycloadducts, because of concomitant decomposition. In order to avoid the isomerization, nopadiene **17**²¹ was reacted with the dienophile at rt under high pressure conditions (12 kbar)²² for 48 hours, to obtain cycloadduct **18** in very good yield as a single stereo- and regio-isomer, as demonstrated by NOESY experiments (Scheme 4).²³

The cycloadduct **10** was treated with various Grignard reagents²⁴ in THF, followed by quenching with trimethyltin chloride: when methyl magnesium chloride was used the desired precursor **3** was obtained, although in modest yield (Scheme 5).

The poor yield can be imputed to the long reaction time required for the complete consumption of the starting phosphine oxide: during this time the resulting *vic*-bromo(chloromagnesium)olefin can decompose with elimination of the dihalomagnesium salt, leading to the formation of the very reactive strained alkyne **19**.²⁵ Furthermore, when the reaction was carried out in the presence of a catalytic amount of nickel(II) dichloro[1,2-bis(diphenylphosphino)ethane], a small amount of *syn*- and *anti*-cyclooligomers was observed in the reaction mixture. Under these

conditions nickel can either mediate the coupling between organo-magnesium species with halides²⁶ or stabilize the strained alkyne by complexation.²⁷

In conclusion, in this Letter we describe the use of two different dienophiles for the synthesis of cycloadducts that can be converted to precursors of functionalized cyclotrimers. The overall path is such that the dienophile can be regarded as a synthetic equivalent of diatomic carbon, or it can resemble a cycloaddition of cyclohexatriene with three dienes (Scheme 6).

Acknowledgement

This work was co-funded by MIUR (Rome) within the national PRIN framework.

References and notes

1. Rieth, S.; Yan, Z.; Xia, S.; Gardlik, M.; Chow, A.; Fraenkel, G.; Hadad, C. M.; Badjić, J. D. *J. Org. Chem.* **2008**, *73*, 5100–5109; Yan, Z.; McCracken, T.; Xia, S.; Maslak, V.; Gallucci, J.; Hadad, C. M.; Badjić, J. D. *J. Org. Chem.* **2008**, *73*, 355–363; Yan, Z.; Xia, S.; Gardlik, M.; Seo, W.; Maslak, V.; Gallucci, J.; Hadad, C. M.; Badjić, J. D. *Org. Lett.* **2007**, *9*, 2301–2304; Maslak, V.; Yan, Z.; Xia, S.; Gallucci, J.; Hadad, C. M.; Badjić, J. D. *J. Am. Chem. Soc.* **2006**, *128*, 5887–5894.
2. Zonta, C.; Cossu, S.; De Lucchi, O. *Eur. J. Org. Chem.* **2000**, 1965–1971.
3. Fabris, F.; Pellizzaro, L.; Zonta, C.; De Lucchi, O. *Eur. J. Org. Chem.* **2007**, 283–291.
4. Scarso, A.; Pellizzaro, L.; De Lucchi, O.; Linden, A.; Fabris, F. *Angew. Chem., Int. Ed.* **2007**, *46*, 4972–4975.
5. Higashibayashi, S.; Sakurai, H. *J. Am. Chem. Soc.* **2008**, *130*, 8592–8593; Amaya, T.; Sakane, H.; Hirao, T. *Angew. Chem., Int. Ed.* **2007**, *46*, 8376–8379; Higashibayashi, S.; Sakurai, H. *Chem. Lett.* **2007**, *36*, 18–19.
6. Borsato, G.; Brussolo, S.; Crisma, M.; De Lucchi, O.; Lucchini, V.; Zambon, A. *Synlett* **2005**, 1125–1128; Borsato, G.; De Lucchi, O.; Fabris, F.; Groppo, L.; Lucchini, V.; Zambon, A. *J. Org. Chem.* **2002**, *67*, 7894–7897; Borsato, G.; De Lucchi, O.; Fabris, F.; Lucchini, V.; Pasqualotti, M.; Zambon, A. *Tetrahedron Lett.* **2003**, *44*, 561–563; Fabris, F.; Bellotto, L.; De Lucchi, O. *Tetrahedron Lett.* **2003**, *44*, 1211–1213; Dastan, A.; Fabris, F.; De Lucchi, O.; Güney, M.; Balci, M. *Helv. Chim. Acta* **2003**, *86*, 3411–3416; De Lucchi, O.; Dastan, A.; Altundag, A.; Fabris, F.; Balci, M. *Helv. Chim. Acta* **2004**, *87*, 2364–2368.
7. (a) Zonta, C.; Cossu, S.; Peluso, P.; De Lucchi, O. *Tetrahedron Lett.* **1999**, *40*, 8185–8188; (b) Durr, R.; Cossu, S.; Lucchini, V.; De Lucchi, O. *Angew. Chem., Int. Ed.* **1997**, *36*, 2805–2807.
8. Zonta, C.; Fabris, F.; De Lucchi, O. *Org. Lett.* **2005**, *7*, 1003–1006.
9. Dalkılıç, E.; Güney, M.; Dastan, A.; Saracoglu, N.; De Lucchi, O.; Fabris, F. *Tetrahedron Lett.* **2009**, doi:10.1016/j.tetlet.2009.02.070.
10. CC-SOAr: Lee, A. W. M.; Chan, W. H.; Wong, M. S. J. *Chem. Soc., Chem. Commun.* **1989**, 1585–1586; Maignan, C.; Belkasmoui, F. *Tetrahedron Lett.* **1988**, *29*, 2823–2824.
11. CC-POAr: Huang, Y. D.; Yu, H. T.; Meng, J. B.; Matsuura, T.; Wang, Y. M. *J. Mol. Struct.* **2002**, *610*, 247–252; Huang, Y. D.; Meng, J. B.; Yu, H. T.; Matsuura, T.; Wang, Y. M. *J. Mol. Struct.* **2002**, *604*, 119–124; Rheingold, A. L.; Liable-Sands, L. M.; Trofimenko, S. *Angew. Chem., Int. Ed.* **2000**, *39*, 3321–3324; Arnaud-Neu, F.; Browne, J. K.; Byrne, D.; Marrs, D. J.; McKervey, M. A.; O'Hagan, P.; Schwing-Weill, M. J.; Walker, A. *Chem. Eur. J.* **1999**, *1*, 175–186; Fu, T. Y.; Liu, Z.; Olovsson, G.; Scheffer, J. R.; Trotter, J. *Acta Crystallogr., Sect. B* **1997**, *53*, 293–299; Palacios, F.; Ochoa de Retana, A. M.; Pagalday, J. *Heterocycles* **1994**, *38*, 95–102.
12. Capozzi, M. A. M.; Cardellicchio, C.; Naso, F. *Eur. J. Org. Chem.* **2004**, 1855–1863.
13. Cardellicchio, C.; Fianadese, V.; Naso, F.; Pacifico, S.; Koprowski, M.; Pietrusiewicz, K. M. *Tetrahedron Lett.* **1994**, *35*, 6343–6346; Cardellicchio, C.; Fianadese, V.; Naso, F.; Pietrusiewicz, K. M.; Wisniewski, W. *Tetrahedron Lett.* **1993**, *34*, 3135–3138.
14. Obtained according to the reported methodology for 1-bromo-2-(phenylsulfanyl)ethyne: Hori, I.; Oishi, T. *Tetrahedron Lett.* **1979**, *42*, 4087–4090. An analytical sample of 1-bromo-2-(p-tolylsulfanyl)ethyne (**1**) was obtained by FC on silica-gel (eluant Et₂O/hexanes 3:7): pale yellow oil stable for several months in CDCl₃ solution in refrigerator, which rapidly polymerizes when concentrated. ¹H NMR (CDCl₃, 200 MHz, TMS) δ 7.68 (2H, d, J = 7.0 Hz), 7.36 (2H, d, J = 7.0 Hz), 2.44 (3H, s); ¹³C NMR (CDCl₃, 50 MHz, TMS) δ 142.9, 140.2, 130.2, 125.1, 78.2, 66.8, 21.4; IR (film) ν 3056, 2925, 2130, 1089, 1059, 809, 537; GC–MS (EI, 70 eV): m/z 194–196 (13), 155 (54), 139 (100), 91 (31), 65 (40).
15. Diastereomers of 2-bromo-3-(phenylsulfanyl)-bicyclo[2.2.1]hepta-2,5-diene **2a** and **2b** were separated by flash-chromatography (eluant 6:4 diethylether/hexanes). First eluant: colourless oil (44% yield). ¹H NMR (CDCl₃, 300 MHz) δ (ppm): 7.55 (2H, d, J = 8.3 Hz), 7.34 (2H, d, J = 8.3 Hz), 6.92 (1H, dd, J = 5.0 and 3.0 Hz), 6.84 (1H, dd, J = 5.0 and 3.0 Hz), 3.70 (1H, br s), 3.65 (1H, br s), 2.44 (3H, s), 2.04 (1H, dt, J = 6.6 and 1.8 Hz), 1.99 (1H, dd, J = 6.6 and 1.4 Hz); ¹³C NMR (CDCl₃, 75 MHz) δ (ppm): 153.0, 143.4, 142.9, 141.09, 140.1, 129.9, 129.7, 124.0, 72.0, 60.1, 50.1, 21.4; MS (70 eV): m/z 260–262 (M⁺–SO, 24), 229 (50), 181 (56), 166 (55), 89 (91), 77 (100%). Second eluate: colourless oil, (39% yield). ¹H NMR (CDCl₃, 300 MHz) δ (ppm): 7.35 (2H, d, J = 8.2 Hz), 7.28 (2H, d, J = 8.2 Hz), 6.50 (1H, dd, J = 3.2 and 4.7 Hz), 6.01 (1H, dd, J = 3.2 and 4.7 Hz), 3.82 (1H, br s), 3.71 (1H, br s), 2.41 (3H, s), 2.34 (1H, dt, J = 6.7 and 1.5 Hz), 2.07 (1H, dd, J = 6.7 and 1.7 Hz); ¹³C NMR (CDCl₃, 75 MHz) δ (ppm): 153.0, 142.4, 141.0, 140.9, 137.8, 136.3, 129.7, 124.0, 70.8, 60.1, 48.3; MS (70 eV): m/z: 260–262 (M⁺–SO, 25), 229 (51), 181 (57), 166 (56), 89 (89), 77 (100%).
16. Cossu, S.; Cimenti, C.; Peluso, P.; Paulon, A.; De Lucchi, O. *Angew. Chem., Int. Ed.* **2001**, *40*, 4086–4089. Noteworthy, compound **3** and the related syn-cyclotrimer are key intermediates for the synthesis of sumanene: Ref. 5.
17. Pellizzaro, L.; Tatibouët, A.; Fabris, F.; Rollin, P.; De Lucchi, O. *Tetrahedron Lett.* **2009**, *50*, 101–103.
18. Bellina, F.; Carpita, A.; Mannocci, L.; Rossi, R. *Eur. J. Org. Chem.* **2004**, 2610–2619; Zhang, C.; Ballay, C. J., II; Trudell, M. L. *J. Chem. Soc., Perkin Trans. 1* **1999**, 675–676; Leroy, J. *Org. Synth.* **1998**, Coll. Vol. 9, 129–133.
19. An analytical sample of 1-bromo-2-(diphenylphosphinoyl)ethyne (**7**) was obtained by FC on silica-gel (eluant AcOEt/hexanes 6:4): mp 67 °C. ¹H NMR (CDCl₃, 200 MHz, TMS) δ 7.91–7.75 (4H, Ar, m), 7.36–7.42 (6H, Ar, m); ¹³C NMR (CDCl₃, 50 MHz, TMS) δ 132.2 (Ar, J_{CPiso} = 122.8 Hz), 132.4 (Ar, J_{CPpara} = 3.0 Hz), 130.9 (Ar, J_{CPmeta} = 11.4 Hz), 128.7 (Ar, J_{CPortho} = 13.7 Hz), 76.4 (C≡C, J_{CP} = 161.0 Hz), 69.2 (C≡C, J_{CP} = 26.2 Hz); ³¹P NMR (CDCl₃, 80 MHz, H₃PO₄) δ 9.17; IR (KBr) ν 3058, 2141, 1200. GC–MS (EI, 70 eV) m/z 306–304 (8), 225 (100), 178 (24), 77 (23%).
20. General cycloaddition procedure. In a screw-capped Pyrex test tube, a solution of **7** (915 mg, 3.0 mmol), diene (1 equiv for entry 4; 1.2 equiv for entry 7; 1.5 equiv for entries 5, 6, 9; 2.0 equiv for entries 2, 3, 8; 4.0 equiv for entry 1) and hydroquinone (5 mg) in toluene (1 mL) was purged with argon, sealed and heated at 120 °C for 18 h. The mixture was purified by flash-chromatography (eluant 1:1 AcOEt/hexanes). 1-Bromo-2-(diphenylphosphinoyl)cyclohexa-1,4-diene (**8**). Waxy solid. ¹H NMR (200 MHz, CDCl₃, TMS) δ (ppm): 7.85–7.63 (4H, m), 7.60–7.30 (6 H, m), 5.74–5.47 (2H, m), 3.39–3.23 (2H, m), 2.82–2.66 (2H, m); ³¹P NMR (81 MHz, CHCl₃, H₃PO₄) δ (ppm): 29.7; ¹³C NMR (50 MHz, CHCl₃, TMS) δ (ppm): 135.9 (d, J_{C-P} = 10.6 Hz), 134.5 (d, J_{C-P} = 24.8 Hz), 133.9 (d, J_{C-P} = 104.0 Hz), 133.4 (d, J_{C-P} = 2.4 Hz), 131.9 (d, J_{C-P} = 2.8 Hz), 131.7 (d, J_{C-P} = 10.0 Hz), 128.6 (d, J_{C-P} = 12.4 Hz), 123.0 (d, J_{C-P} = 8.8 Hz), 122.7 (d, J_{C-P} = 1.3 Hz), 39.6 (d, J_{C-P} = 7.7 Hz), 33.0 (d, J_{C-P} = 10.2 Hz); IR (KBr) ν 1187 (PO) cm⁻¹; m/z (EI, 70 eV): 358–360 (M⁺, 20), 359–357 (M⁺–H, 55), 357–355 (80), 277 (80), 199 (90), 152 (97), 77 (100%). 1-Bromo-2-(diphenylphosphinoyl)-4,5-dimethylcyclohexa-1,4-diene (**9**). Waxy solid. ¹H NMR (200 MHz, CDCl₃, TMS) δ (ppm): 7.84–7.67 (4H, m), 7.60–7.38 (6 H, m), 3.28–3.13 (2H, m), 2.81–2.64 (2H, m), 1.60 (3H, s), 1.52 (3H, s); ³¹P NMR (81 MHz, CHCl₃, H₃PO₄) δ (ppm): 29.1; ¹³C NMR (50 MHz, CHCl₃, TMS) δ (ppm): 133.2 (d, J_{C-P} = 20.5 Hz), 131.8 (d, J_{C-P} = 2.8 Hz), 131.7 (d, J_{C-P} = 10.0 Hz), 131.0 (d, J_{C-P} = 9.6 Hz), 129.9 (d, J_{C-P} = 93.6 Hz), 128.5 (d, J_{C-P} = 12.4 Hz), 122.5 (d, J_{C-P} = 6.4 Hz), 122.4 (d, J_{C-P} = 0.8 Hz), 46.0 (d, J_{C-P} = 8.0 Hz), 39.0 (d, J_{C-P} = 9.2 Hz), 17.7, 17.5; IR (KBr) ν 1199 (P=O) cm⁻¹; m/z (EI, 70 eV): 388–386 (M⁺, 25), 385–388 (M⁺–H, 85), 305–307 (80), 227 (80), 201 (100), 77 (97%). 2-Bromo-3-(diphenylphosphinoyl)bicyclo[2.2.1]hepta-2,5-diene (**10**). Oil. ¹H NMR (200 MHz, CDCl₃, TMS) δ (ppm): 7.76–7.37 (10 H, series of m), 6.86 (1H, dd, J = 4.9 and 3.0 Hz), 6.76 (1H, dd, J = 4.9 and 3.0 Hz), 3.95 (1H, br s), 3.75 (1H, br s), 2.42 (1H, dt, J = 6.7 and 1.8 Hz), 2.07 (1H, br d, J = 6.7 Hz); ³¹P NMR (81 MHz, CHCl₃, H₃PO₄) δ (ppm): 24.1; ¹³C NMR (50 MHz, CHCl₃, TMS) δ (ppm): 151.1 (d, J_{C-P} = 4.4 Hz), 142.9 (d, J_{C-P} = 0.8 Hz), 141.5 (d, J_{C-P} = 112.8 Hz), 140.0 (d, J_{C-P} = 2.0 Hz), 132.4 (d, J_{C-P} = 38.9 Hz), 132.0 (d, J_{C-P} = 2.8 Hz), 131.9 (d, J_{C-P} = 2.8 Hz), 131.6 (d, J_{C-P} = 10.4 Hz, two overlapping C), 130.2 (d, J_{C-P} = 38.5 Hz), 128.5 (d, J_{C-P} = 12.4 Hz), 128.4 (d, J_{C-P} = 12.4 Hz), 72.3 (d, J_{C-P} = 4.4 Hz), 63.0 (d, J_{C-P} = 9.3 Hz), 55.1 (d, J_{C-P} = 8.8 Hz); IR (film) ν 1182 (P=O) cm⁻¹; m/z (EI, 70 eV): 370–372 (M⁺, 3), 369–371 (M⁺–H, 4), 355–307 (8), 291 (100), 201 (50), 77 (70%). 2-Bromo-3-(diphenylphosphinoyl)-7-isopropylidenebicyclo[2.2.1]hepta-2,5-diene (**11**). Colourless crystals mp = 102–103 °C (dec); ¹H NMR (200 MHz, CDCl₃, TMS) δ (ppm): 7.27–7.32 (10H, series of m), 6.89 (1H, ddd, J = 4.3, 3.0 and 1.2 Hz), 6.79 (1H, dd, J = 4.3 and 3.0 Hz), 4.30–4.21 (1H, m), 4.18–4.11 (1H, m), 1.45 (3H, s), 1.35 (3H, s); ³¹P NMR (81 MHz, CHCl₃, H₃PO₄) δ (ppm): 24.1; ¹³C NMR (50 MHz, CHCl₃, TMS) δ (ppm): 160.2 (d, J_{C-P} = 5.2 Hz), 149.5 (d, J_{C-P} = 4.8 Hz), 142.3 (d, J_{C-P} = 1.2 Hz), 141.3 (d, J_{C-P} = 110.8 Hz), 140.0 (d, J_{C-P} = 2.0 Hz), 132.3 (d, J_{C-P} = 9.6 Hz), 131.91 (d, J_{C-P} = 2.8 Hz), 131.89 (d, J_{C-P} = 2.8 Hz), 131.6 (d, J_{C-P} = 10.4 Hz), 131.4 (d, J_{C-P} = 10.4 Hz), 130.2 (d, J_{C-P} = 9.2 Hz), 128.42 (d, J_{C-P} = 12.4 Hz), 128.38 (d, J_{C-P} = 12.8 Hz), 99.2, 62.5 (d, J_{C-P} = 7.6 Hz), 55.3 (d, J_{C-P} = 9.2 Hz), 18.4, 18.1; IR (KBr) ν 1198 (P=O) cm⁻¹; m/z (EI, 70 eV): 410–412 (M⁺, 7), 409–411 (M⁺–H, 15), 201 (25), 141 (75), 57 (100%). 2-Bromo-3-(diphenylphosphinoyl)-7-oxabicyclo[2.2.1]hepta-2,5-diene (**12**). Colourless crystals mp = 125–127 °C (dec); ¹H NMR (200 MHz, CDCl₃, TMS) δ (ppm): 7.78–7.39 (10 H, series of m), 7.16 (1H, dd, J = 5.4 and 1.8 Hz), 7.06 (1H, dd, J = 5.4 and 1.8 Hz), 5.69 (1H, br q, J = 1.5 Hz), 5.37–5.34 (1H, m); ³¹P NMR (81 MHz, CHCl₃, H₃PO₄) δ (ppm): 21.9; ¹³C NMR (50 MHz, CHCl₃, TMS) δ (ppm): 151.6 (d, J_{C-P} = 4.0 Hz), 144.6 (d, J_{C-P} = 0.8 Hz), 142.8 (d, J_{C-P} = 113.2 Hz), 140.5 (d, J_{C-P} = 1.6 Hz), 132.4 (d, J_{C-P} = 3.2 Hz, two overlapping C), 131.7 (d, J_{C-P} = 10.4 Hz), 131.5 (d, J_{C-P} = 10.4 Hz), 130.7 (d, J_{C-P} = 110.8 Hz), 130.2 (d, J_{C-P} = 110.8 Hz), 128.8 (d, J_{C-P} = 12.8 Hz), 128.7 (d, J_{C-P} = 12.8 Hz), 90.5 (d, J_{C-P} = 9.2 Hz), 87.2 (d, J_{C-P} = 11.6 Hz); IR (KBr) ν 1186 (P=O) cm⁻¹; m/z (EI, 70 eV): 355–357 (M⁺–O, 10), 280–282 (15), 133 (60), 95 (65), 76 (100%). 2-Bromo-3-(diphenylphosphinoyl)bicyclo[2.2.2]octa-2,5-diene (**13**). Colourless crystals mp = 130–4 °C (dec); ¹H NMR (200 MHz, CDCl₃, TMS) δ (ppm): 7.78–7.30 (10 H, series of m), 6.40–6.30 (2H, m), 4.16–4.00 (1H, m), 3.96–3.82 (1H, m), 1.50–1.20 (4H, m); ³¹P NMR (81 MHz, CHCl₃, H₃PO₄) δ (ppm): 26.4; ¹³C NMR (50 MHz, CHCl₃, TMS) δ (ppm): 138.8 (d, J_{C-P} = 2.8 Hz), 135.5 (d, J_{C-P} = 106.0 Hz), 133.8 (d, J_{C-P} = 3.6 Hz), 132.8 (d, J_{C-P} = 2.0 Hz), 131.8 (d, J_{C-P} = 108.0 Hz), 131.7 (d, J_{C-P} = 3.2 Hz), 131.6 (d, J_{C-P} = 2.8 Hz), 131.5 (d, J_{C-P} = 107.2 Hz), 131.5 (d, J_{C-P} = 10.4 Hz), 131.4 (d,

J_{C-P} = 10.4 Hz), 128.3 (d, J_{C-P} = 12.8 Hz, two overlapping C), 51.2 (d, J_{C-P} = 7.2 Hz), 41.7 (d, J_{C-P} = 8.4 Hz), 24.9 (d, J_{C-P} = 1.6 Hz), 24.6 (d, J_{C-P} = 2.0 Hz); IR (KBr) ν 1190 (P=O) cm^{-1} ; m/z (EI, 70 eV): 358–356 (M^+ -C₂H₄, 55), 357–355 (70), 277 (60), 199 (85), 183 (70), 152 (100), 77 (97%). 8-Bromo-9-(diphenylphosphinoyl)-4,4-dimethyl-3,5-dioxatricyclo[5.2.2.0^{2,6}]undeca-8,10-diene (**14**). Colourless crystals mp = 153–155 °C (dec.) (Et₂O). ¹H NMR (200 MHz, CDCl₃, TMS) δ (ppm): 7.77–7.40 (10 H, series of m), 6.43 (1H, *pseudo-t*, J = 6.3 Hz), 6.33 (1H, *pseudo-t*, J = 6.3 Hz), 4.51 (1H, dd, J = 6.9 and 3.5 Hz), 4.31 (1H, dd, J = 6.9 and 3.5 Hz), 4.20–4.07 (2H, m), 1.29 (3H, s), 1.25 (3H, s); ³¹P NMR (81 MHz, CHCl₃, H₃PO₄) δ (ppm): 26.3; ¹³C NMR (50 MHz, CHCl₃, TMS) δ (ppm): 139.2 (d, J_{C-P} = 3.2 Hz), 134.6 (d, J_{C-P} = 104.2 Hz), 131.3 (d, J_{C-P} = 108.4 Hz), 131.2 (d, J_{C-P} = 108.0 Hz), 132.19 (d, J_{C-P} = 2.9 Hz), 132.17 (d, J_{C-P} = 2.8 Hz), 131.7 (d, J_{C-P} = 11.5 Hz), 131.5 (d, J_{C-P} = 11.1 Hz), 131.4 (d, J_{C-P} = 3.4 Hz), 131.0 (d, J_{C-P} = 1.9 Hz), 128.8 (d, J_{C-P} = 12.8 Hz), 128.7 (d, J_{C-P} = 12.5 Hz), 113.6, 78.0 (d, J_{C-P} = 3.0 Hz), 77.5 (d, J_{C-P} = 2.9 Hz), 56.6 (d, J_{C-P} = 7.1 Hz), 46.6 (d, J_{C-P} = 8.0 Hz), 25.7, 25.6; IR (KBr) ν 1189 (P=O) cm^{-1} ; m/z (EI, 70 eV): 358–356 (M^+ -C₅H₈O₂, 27), 357–355 (M^+ -C₅H₉O₂, 70), 277 (30), 199 (60), 152 (70), 77 (100%). 8-Bromo-9-(diphenylphosphinoyl)tetracyclo[4.3.0.2.4^{0,3,7}]non-8-ene (**15**). Colourless crystals mp = 130–3 °C. ¹H NMR (200 MHz, CDCl₃, TMS) δ (ppm): 7.80–7.62 (4H, m), 7.60–7.37 (6 H, m), 2.97–2.88 (2H, m), 2.37 (1H, br s), 1.84–1.76 (1H, m), 1.70–1.62 (1H, m), 1.57 (2H, br s), 1.49–1.41 (1H, m); ³¹P NMR (81 MHz, CHCl₃, H₃PO₄) δ (ppm): 23.4; ¹³C NMR (50 MHz, CHCl₃, TMS) δ (ppm): 140.9 (d, J_{C-P} = 2.4 Hz), 136.7 (d, J_{C-P} = 110.0 Hz), 134.0 (d, J_{C-P} = 65.8 Hz), 132.0 (d, J_{C-P} = 75.5 Hz), 131.8 (d, J_{C-P} = 3.2 Hz, two overlapping C), 131.6 (d, J_{C-P} = 10.4 Hz), 131.5 (d, J_{C-P} = 10.4 Hz), 131.2 (d, J_{C-P} = 62.6 Hz), 128.4 (d, J_{C-P} = 13.6 Hz, two overlapping C), 61.0 (d, J_{C-P} = 9.6 Hz), 56.8 (d, J_{C-P} = 4.8 Hz), 53.8 (d, J_{C-P} = 8.8 Hz), 32.4, 26.7, 23.8 (d, J_{C-P} = 0.8 Hz), 23.4 (d, J_{C-P} = 2.8 Hz); IR (KBr) ν 1194 (P=O) cm^{-1} ; m/z (EI, 70 eV): 398–396 (M^+ , 7), (M^+ -H, 17), 317 (28), 201 (100), 115 (60), 77 (50%). 11-Bromo-12-(diphenylphosphinoyl)-9,10-dihydro-9,10-ethenylantracene (**16**). Colourless crystals mp = 130–132 °C (dec); ¹H NMR (200 MHz, CDCl₃, TMS) δ (ppm): 7.60–7.32 (12H, m), 7.14–6.94 (8 H, m), 5.34–5.25 (2H, m); ³¹P NMR (81 MHz, CHCl₃, H₃PO₄) δ (ppm): 28.0; ¹³C NMR (50 MHz, CHCl₃, TMS) δ (ppm): 146.2 (d, J_{C-P} = 2.8 Hz), 143.2 (d, J_{C-P} = 2.8 Hz), 143.1 (d, J_{C-P} = 2.0 Hz), 138.8 (d, J_{C-P} = 103.6 Hz), 132.1 (d, J_{C-P} = 2.8 Hz), 131.8 (d, J_{C-P} = 10.4 Hz), 131.1 (d, J_{C-P} = 108.8 Hz), 128.5 (d, J_{C-P} = 12.4 Hz), 125.7, 125.2, 123.6, 123.4, 64.3 (d,

J_{C-P} = 7.2 Hz), 55.0 (d, J_{C-P} = 8.8 Hz); IR (KBr) ν 1184 (P=O) cm^{-1} ; m/z (EI, 70 eV): 484–482 (M^+ , 1), 403 (M^+ -Br, 7), 201 (30), 178 (100%).

21. Paquette, L. A.; Gugelcuck, M.; McLaughlin, M. L. *J. Org. Chem.* **1987**, 52, 4732–4740.
22. Minuti, L.; Taticchi, A.; Marrocchi, A.; Broggi, A. *Tetrahedron: Asymmetry* **2004**, 15, 1187–1119.
23. A solution of **7** (1.04 g, 3.4 mmol) and nopadiene (666 mg, 4.5 mmol) in DCM was placed in a 3 mL screw-capped Teflon capsule that was submitted to 12 Kbar pressure for 48 hours at r.t. The volatile materials were removed, and the residue was recrystallized from AcOEt to afford 1.39 g (89% yield) of 5-bromo-6-(diphenylphosphinoyl)-9,9-dimethyl-1,2,3,4,4a,7-hexahydro-1,3-methanonaphthalene (**18**) as colourless crystals, mp = 184–185 °C. $[\alpha]_D^{22}$ –12 (c 1.4, CHCl₃); ¹H NMR (300 MHz, CDCl₃, TMS) δ (ppm): 7.68–7.76 (4H, series of m), 7.41–7.56 (6H, series of m), 5.31 (1H, m), 3.75 (1H, m), 2.48–2.82 (4H, series of m), 2.08–2.38 (3H, series of m), 1.27 (3H, s), 0.97 (3H, s), 0.90 (1H, d, J = 8.9 Hz); ³¹P NMR (122 MHz, CDCl₃, H₃PO₄) δ (ppm): 30.6; ¹³C NMR (75 MHz, CDCl₃, TMS) δ (ppm): 143.7 (d, J = 1.5 Hz), 143.3, 132.6 (d, J = 106.2 Hz), 132.3 (d, J = 105.0 Hz), 132.1 (d, J = 3.4 Hz), 132.0 (d, J = 3.4 Hz), 131.7 (d, J = 9.7 Hz), 131.6 (d, J = 9.9 Hz), 128.6 (d, J = 101.2 Hz), 128.5 (d, J = 12.3 Hz), 128.4 (d, J = 12.3 Hz), 115.4 (d, J = 9.0 Hz), 51.1, 42.3, 40.2 (d, J = 7.9 Hz), 39.5, 35.7, 34.0, 33.8 (d, J = 10.6 Hz), 27.0, 23.2; IR (KBr) ν 1187 (P=O) cm^{-1} .
24. PhMgBr furnished no products, presumably because of steric hindrance. Me₂Mg/MgI₂ in THF furnished only 10–15% yields of **3**.
25. Laird, D. W.; Gilbert, J. C. *J. Am. Chem. Soc.* **2001**, 123, 6704–6705; Kitamura, T.; Kotani, M.; Yokoyama, T.; Fujiwara, Y. *J. Org. Chem.* **1999**, 64, 680–681; Kennedoff, J.; Polborn, K.; Szeimies, G. *J. Am. Chem. Soc.* **1990**, 112, 6117–6118; Harold, H.; Shamouil, S.; Yoshikazu, T. *J. Org. Chem.* **1981**, 46, 4427–4432; Brewer, J. P. N.; Heaney, H.; Marples, B. A. *Chem. Commun. (London)* **1967**, 27–28.
26. Busacca, C. A.; Eriksson, M. C.; Fiaschi, R. *Tetrahedron Lett.* **1999**, 40, 3101–3104; Kottirsch, G.; Szeimies, G. *Chem. Ber.* **1990**, 123, 1495–1505; Walker, J. A.; Bitler, S. A.; Wudl, F. *J. Org. Chem.* **1984**, 49, 4733–4734.
27. Bennet, M. A. *Pure Appl. Chem.* **1989**, 61, 1700–1965; Pericàs, M. A.; Riera, A.; Rossell, O.; Serratos, F.; Seco, M. *J. Chem. Soc., Chem. Commun.* **1988**, 942–943; Iyoda, M.; Kuwatani, Y.; Yamauchi, T.; Oda, M. *J. Chem. Soc., Chem. Commun.* **1988**, 65–66.