

### Supramolecular Allosteric Cofacial Porphyrin Complexes

Christopher G. Oliveri,<sup>†</sup> Nathan C. Gianneschi,<sup>†</sup> SonBinh T. Nguyen,<sup>\*,†</sup>  
Chad A. Mirkin,<sup>\*,†</sup> Charlotte L. Stern,<sup>†</sup> Zdzislaw Wawrzak,<sup>‡</sup> and Maren Pink<sup>§</sup>

*Contribution from the Department of Chemistry and the International Institute for Nanotechnology, Northwestern University, 2145 Sheridan Road, Evanston, Illinois, 60208-3113, Department of Biochemistry, Molecular Biology and Cell Biology, Northwestern University, 2205 Tech Drive, Evanston, Illinois 60208-3500, and Department of Chemistry, Indiana University, Bloomington, Indiana 47405*

Received August 22, 2006; E-mail: stn@northwestern.edu; chadnano@northwestern.edu

**Abstract:** Nature routinely uses cooperative interactions to regulate cellular activity. For years, chemists have designed synthetic systems that aim toward harnessing the reactivity common to natural biological systems. By learning how to control these interactions *in situ*, one begins to allow for the preparation of man-made biomimetic systems that can efficiently mimic the interactions found in Nature. To this end, we have designed a synthetic protocol for the preparation of flexible metal-directed supramolecular cofacial porphyrin complexes which are readily obtained in greater than 90% yield through the use of new hemilabile porphyrin ligands with bifunctional ether–phosphine or thioether–phosphine substituents at the 5 and 15 positions on the porphyrin ring. The resulting architectures contain two hemilabile ligand–metal domains (Rh<sup>I</sup> or Cu<sup>I</sup> sites) and two cofacially aligned porphyrins (Zn<sup>II</sup> sites), offering orthogonal functionalities and allowing these multimetallic complexes to exist in two states, “condensed” or “open”. Combining the ether–phosphine ligand with the appropriate Rh<sup>I</sup> or Cu<sup>I</sup> transition-metal precursors results in “open” macrocyclic products. In contrast, reacting the thioether–phosphine ligand with Rh<sup>I</sup> or Cu<sup>I</sup> precursors yields condensed structures that can be converted into their “open” macrocyclic forms via introduction of additional ancillary ligands. The change in cavity size that occurs allows these structures to function as allosteric catalysts for the acyl transfer reaction between *X*-pyridylcarbinol (where *X* = 2, 3, or 4) and 1-acetylimidazole. For 3- and 4-pyridylcarbinol, the “open” macrocycle accelerates the acyl transfer reaction more than the condensed analogue and significantly more than the porphyrin monomer. In contrast, an allosteric effect was not observed for 2-pyridylcarbinol, which is expected to be a weaker binder and is unfavorably constrained inside the macrocyclic cavity.

#### Introduction

Over the past two decades, chemists have been designing synthetic systems that incorporate cofacial porphyrin entities.<sup>1–10</sup> These systems are of particular interest owing to their unique

photophysical properties,<sup>11–15</sup> their ability to catalyze small molecule transformations,<sup>16–20</sup> and their use in molecular recognition.<sup>21,22</sup> Among these, cofacial porphyrin compounds exhibit unique properties attributed to their spatial arrangement with respect to each other, and therefore, the ability to control that arrangement via incorporation of multiple porphyrins in supramolecular arrays offers a potentially facile route to


<sup>†</sup> Department of Chemistry and the International Institute for Nanotechnology, Northwestern University.

<sup>‡</sup> Department of Biochemistry, Molecular Biology and Cell Biology, Northwestern University.

<sup>§</sup> Indiana University.

- (1) Cowan, J. A.; Sanders, J. K. M. *J. Chem. Soc., Chem. Commun.* **1985**, 1213–14.
- (2) Fletcher, J. T.; Therien, M. J. *J. Am. Chem. Soc.* **2000**, *122*, 12393–12394.
- (3) Collman, J. P.; Bencosme, C. S.; Durand, R. R., Jr.; Kreh, R. P.; Anson, F. C. *J. Am. Chem. Soc.* **1983**, *105*, 2699–703.
- (4) Feiters, M. C.; Fyfe, M. C. T.; Martinez-Diaz, M. V.; Menzer, S.; Nolte, R. J. M.; Stoddart, J. F.; van Kan, P. J. M.; Williams, D. J. *J. Am. Chem. Soc.* **1997**, *119*, 8119–8120.
- (5) Yagi, S.; Yonekura, I.; Awakura, M.; Ezoe, M.; Takagishi, T. *Chem. Commun.* **2001**, 557–558.
- (6) Chang, C. J.; Yeh, C.-Y.; Nocera, D. G. *J. Org. Chem.* **2002**, *67*, 1403–1406.
- (7) Jokic, D.; Asfari, Z.; Weiss, J. *Org. Lett.* **2002**, *4*, 2129–2132.
- (8) Chng, L. L.; Chang, C. J.; Nocera, D. G. *J. Org. Chem.* **2003**, *68*, 4075–4078.
- (9) Collman, J. P.; Tyvoll, D. A.; Chng, L. L.; Fish, H. T. *J. Org. Chem.* **1995**, *60*, 1926–31.
- (10) Chen, W.-H.; Yan, J.-M.; Tagashira, Y.; Yamaguchi, M.; Fujita, K. *Tetrahedron Lett.* **1999**, *40*, 891–894.

- (11) Faure, S.; Stern, C.; Guillard, R.; Harvey, P. D. *J. Am. Chem. Soc.* **2004**, *126*, 1253–1261.
- (12) Splan, K. E.; Stern, C. L.; Hupp, J. T. *Inorg. Chim. Acta.* **2004**, *357*, 4005–4014.
- (13) Hajjaj, F.; Yoon, Z. S.; Yoon, M. C.; Park, J.; Satake, A.; Kim, D.; Kobuke, Y. *J. Am. Chem. Soc.* **2006**, *128*, 4612–4623.
- (14) Fletcher, J. T.; Therien, M. J. *J. Am. Chem. Soc.* **2002**, *124*, 4298–4311.
- (15) Kobuke, Y. *Struct. Bonding* **2006**, *121*, 145–165.
- (16) Collman, J. P.; Fu, L.; Herrmann, P. C.; Zhang, X. *Science* **1997**, *275*, 949–951.
- (17) Chang, C. J.; Loh, Z.-H.; Shi, C.; Anson, F. C.; Nocera, D. G. *J. Am. Chem. Soc.* **2004**, *126*, 10013–10020.
- (18) Rosenthal, J.; Pistorio, B. J.; Chng, L. L.; Nocera, D. G. *J. Org. Chem.* **2005**, *70*, 1885–1888.
- (19) Collman, J. P.; Wagenknecht, P. S.; Hutchison, J. E. *Angew. Chem., Int. Ed. Engl.* **1994**, *106*, 1620–39.
- (20) Rosenthal, J.; Luckett, T. D.; Hodgkiss, J. M.; Nocera, D. G. *J. Am. Chem. Soc.* **2006**, *128*, 6546–6547.
- (21) Brettar, J.; Gisselbrecht, J.-P.; Gross, M.; Solladie, N. *Chem. Commun.* **2001**, 733–734.
- (22) Jokic, D.; Boudon, C.; Pognon, G.; Bonin, M.; Schenk, K. J.; Gross, M.; Weiss, J. *Chem.–Eur. J.* **2005**, *11*, 4199–4209.

**Scheme 1.** Design of Allosteric Porphyrin-Based Supramolecules Whose Cavity Sizes Can Be Modified by the Binding of a Ligand L<sup>a</sup>

<sup>a</sup> **I**: Condensed Macrocycle, **II**: Open Macrocycle. PPh<sub>2</sub> = diphenylphosphine and MES = 1,3,5-trimethylbenzene.

biomimetic structures with novel physicochemical properties.<sup>23,24</sup> Taking inspiration from nature, a primary goal of our research has been to prepare supramolecular structures that exhibit allosteric behavior in the context of catalysis and small-molecule/ion sensing analogous to that displayed ubiquitously in natural enzymatic regulation.<sup>25–28</sup> In this context, we reasoned that the ability to manipulate the distance between porphyrins by altering the shape and size of supramolecular cavities in which they reside, via external stimuli, may provide a means for designing abiotic allosteric mimics of biological systems. Indeed, a clear inspiration for this kind of allosteric control over the function of multi-porphyrin assemblies comes from perhaps the most famous example of allosteric control in biology, the cooperative homoallosteric protein hemoglobin, in which O<sub>2</sub> binding at one subunit initiates conformational changes that facilitate subsequent O<sub>2</sub> binding at other porphyrin domains.<sup>29</sup>

To design synthetic allosteric porphyrin-based structures, an attractive strategy entails the preparation of structures that provide facile control over both the orientation and distance of both porphyrins in the context of a small molecule-mediated reaction. To date, most approaches for the synthesis of cofacial porphyrins rely on the use of covalently attached organic molecules as spacers between the porphyrin units.<sup>6,9,21,30–32</sup> As cofacial porphyrins have been prepared predominantly through synthetic organic approaches, only a few examples exist in which transition metals have been incorporated to facilitate the formation of cofacial porphyrins.<sup>24,33–36</sup> Additionally, the backbones used to couple these cofacial porphyrins are often based upon rigid linker molecules, which restrict the overall flexibility of the molecule and make the incorporation of

different substrates with varying size and shape challenging. To address this issue, several cofacial porphyrin structures have been designed which incorporate flexible linkages.<sup>22,37–40</sup> However, few are able to specifically access a particular conformation via the introduction or removal of small-molecule effectors at a distal site (i.e., in an allosteric fashion). To this end, we hypothesized that coordination complexes of type **I** and **II** (Scheme 1), whose cavity size can be modified *in situ* with small molecules to form **II** would be very useful for designing stimulant-responsive porphyrin-based biomimetic systems.

Herein, we present a general, high-yielding synthetic methodology for the synthesis of cofacial porphyrin complexes that utilizes flexible porphyrin-based hemilabile ligands and M<sup>I</sup> precursors (where M<sup>I</sup> = Rh<sup>I</sup> or Cu<sup>I</sup>) to form macrocyclic products that can be chemically stimulated to change shape. Our strategy, based upon the Weak-Link Approach (WLA),<sup>41–43</sup> provides rapid and convergent access to unique cofacial porphyrin systems because it relies on the use of metal-heteroatom interactions to facilitate the formation of the desired macrocyclic structures, thus allowing for the controlled and selective modification of the resulting structures *in situ* via introduction of specific external stimuli. Such stimulation can be used to tune the efficiency of a distance-dependent bimolecular acyl transfer reaction and discriminate geometric isomers of substituted pyridylcarbinol substrates.

## General Methods and Instrument Details

All reactions were carried out under an inert atmosphere of nitrogen using standard Schlenk techniques or an inert atmosphere glovebox unless otherwise noted. Tetrahydrofuran (THF), diethyl ether (Et<sub>2</sub>O), dichloromethane (CH<sub>2</sub>Cl<sub>2</sub>), acetonitrile (CH<sub>3</sub>CN), and hexanes were purified according to published methods.<sup>44</sup> All solvents were deoxygenated with nitrogen or argon prior to use. 1-Chloro-2-diphenylphosphinoethane (Organometallics Inc.), deuterated solvents (Cambridge Isotope Laboratories Inc.), [Rh(NBD)Cl]<sub>2</sub> (NBD = norbornadiene, Strem Chemicals), and 4-bromothiophenol (Alfa Aesar) were obtained from commercial sources and used as received. 4-(2-Chloroethoxy)-

- (23) Merlau, M.; Grande, W. J.; Nguyen, S. T.; Hupp, J. T. *J. Mol. Cat. A: Chem.* **2000**, *156*, 79–84.
- (24) Merlau, M. L.; del Pilar Mejia, M.; Nguyen, S. T.; Hupp, J. T. *Angew. Chem., Int. Ed.* **2001**, *40*, 4239–4242.
- (25) Heo, J.; Mirkin, C. A. *Angew. Chem., Int. Ed.* **2006**, *45*, 941–944.
- (26) Gianneschi, N. C.; Bertin, P. A.; Nguyen, S. T.; Mirkin, C. A.; Zakharov, L. N.; Rheingold, A. L. *J. Am. Chem. Soc.* **2003**, *125*, 10508–10509.
- (27) Gianneschi, N. C.; Cho, S.-H.; Nguyen, S. T.; Mirkin, C. A. *Angew. Chem., Int. Ed.* **2004**, *43*, 5503–5507.
- (28) Gianneschi, N. C.; Nguyen, S. T.; Mirkin, C. A. *J. Am. Chem. Soc.* **2005**, *127*, 1644–1645.
- (29) Stryer, L. *Biochemistry*, 4th ed.; W.H. Freeman and Company: New York, 1995.
- (30) Brothers, P. J.; Collman, J. P. *Acc. Chem. Res.* **1986**, *19*, 209–15.
- (31) Chng, L. L.; Chang, C. J.; Nocera, D. G. *J. Org. Chem.* **2003**, *68*, 4075–4078.
- (32) Guillard, R.; Burdet, F.; Barbe, J.-M.; Gros, C. P.; Espinosa, E.; Shao, J.; Ou, Z.; Zhan, R.; Kadish, K. M. *Inorg. Chem.* **2005**, *44*, 3972–3983.
- (33) Slone, R. V.; Hupp, J. T. *Inorg. Chem.* **1997**, *36*, 5422–5423.
- (34) Dinolfo, P. H.; Hupp, J. T. *Chem. Mater.* **2001**, *13*, 3113–3125.
- (35) Dinolfo, P. H.; Lee, S. J.; Coropceanu, V.; Bredas, J. L.; Hupp, J. T. *Inorg. Chem.* **2005**, *44*, 5789–5797.
- (36) Benkstein, K. D.; Stern, C. L.; Splan, K. E.; Johnson, R. C.; Walters, K. A.; Vanhelmont, F. W. M.; Hupp, J. T. *Eur. J. Inorg. Chem.* **2002**, 2818–2822.

- (37) Ohsaki, K.; Konishi, K.; Aida, T. *Chem. Commun.* **2002**, 1690–1691.
- (38) Tomohiro, Y.; Satake, A.; Kobuke, Y. *J. Org. Chem.* **2001**, *66*, 8442–8446.
- (39) Sanders, J. K. M. *Chem. Eur. J.* **1998**, *4*, 1378–1383.
- (40) Pognon, G.; Boudon, C.; Schenk, K. J.; Bonin, M.; Bach, B.; Weiss, J. J. *Am. Chem. Soc.* **2006**, *128*, 3488–3489.
- (41) Gianneschi, N. C.; Masar III, M. S.; Mirkin, C. A. *Acc. Chem. Res.* **2005**, *38*, 825–837.
- (42) Farrell, J. R.; Mirkin, C. A.; Guzei, I. A.; Liable-Sands, L. M.; Rheingold, A. L. *Angew. Chem., Int. Ed.* **1998**, *37*, 465–467.
- (43) Holliday, B. J.; Mirkin, C. A. *Angew. Chem., Int. Ed.* **2001**, *40*, 2022–2043.
- (44) Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J. *Organometallics*. **1996**, *15*, 1518–20.

benzaldehyde,<sup>45</sup> 2-(4-bromophenylsulfanyl)ethyldiphenylphosphane,<sup>46</sup> 5-mesityldipyrromethane,<sup>47</sup> and (2-(mesitylthio)ethyl)diphenylphosphine<sup>48</sup> were prepared according to literature procedures. The synthesis of [5,10,15,20-tetraphenylporphyrinato]zinc<sup>II</sup> (Zn(TPP))<sup>49</sup> was adapted from a literature synthetic procedure. All other chemicals were used as received from Aldrich Chemical Co. <sup>1</sup>H NMR (300.22 MHz) and <sup>13</sup>C{<sup>1</sup>H} NMR (75.50 MHz) spectra were recorded on a Varian Mercury 300 MHz FT-NMR spectrometer and referenced relative residual proton resonances. <sup>31</sup>P{<sup>1</sup>H} NMR (121.53 MHz) spectra were recorded on a Varian Mercury 300 MHz FT-NMR spectrometer and referenced relative to an external 85% H<sub>3</sub>PO<sub>4</sub> standard. All chemical shifts are reported in ppm. Electrospray ionization mass spectra (ESIMS) were recorded on a Micromass Quattro II triple quadrupole mass spectrometer or a Micromass Q-ToF Ultima mass spectrometer. Electron impact mass spectra (EIMS) were recorded on a Fisons VG 70-250 SE mass spectrometer. Elemental analyses were performed by Quantitative Technologies Inc., Whitehouse, NJ. Gas chromatography (GC) analyses of reaction mixtures were carried out on a computer-interfaced Agilent Technologies 6890 Network instrument equipped with a flame ionization detector (FID). The column used was a 30-m HP-5 capillary column with a 0.32-mm inner diameter and a 0.25-μm film thickness. GC yields were determined through integration of the product peak against biphenyl (internal standard) using pre-established response factors. GC retention times of products were confirmed with analytically pure samples.

**2-[4-(2-Chloroethoxy)-phenyl]-[1,3]dithiane (1).** Under ambient conditions, 4-(2-chloroethoxy)benzaldehyde (3.00 g, 16.3 mmol) and 1,3-propanedithiol (1.95 mL, 19.2 mmol) were combined with CH<sub>3</sub>CN (100 mL) in a 250-mL round-bottom flask and allowed to stir at room temperature for 5 min at which point Y(OTf)<sub>3</sub> (437 mg, 5 mol %) was added. The resulting solution was stirred for another 30 min when it gradually became turbid due to product precipitation. The mixture was dried on a rotary evaporator to give an oily residue which was then dissolved in CH<sub>2</sub>Cl<sub>2</sub> (100 mL), washed with H<sub>2</sub>O (2 × 25 mL), dried over MgSO<sub>4</sub>, and concentrated *in vacuo*. The crude product was purified via column chromatography (1:1 v/v, CH<sub>2</sub>Cl<sub>2</sub>/hexanes as an eluent) to yield **1** as a white solid (3.38 g, 75% yield). <sup>1</sup>H NMR (CD<sub>2</sub>Cl<sub>2</sub>): δ 1.80 (m, 1H, CH<sub>2</sub>–CH<sub>A</sub>H<sub>B</sub>–CH<sub>2</sub>), 2.14 (m, 1H, CH<sub>2</sub>–CH<sub>A</sub>H<sub>B</sub>–CH<sub>2</sub>), 2.86 (m, 2H, SCH<sub>2</sub>), 2.92 (m, 2H, SCH<sub>2</sub>), 3.81 (t, 2H, CH<sub>2</sub>Cl), 4.21 (t, 2H, OCH<sub>2</sub>), 5.15 (s, 1H, S–CH–S), 6.87 (d, 2H, J<sub>H–H</sub> = 8.4 Hz, ArH), 7.37 (d, 2H, J<sub>H–H</sub> = 8.7 Hz, ArH). <sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>): δ 25.2 (CH<sub>2</sub>–CH<sub>2</sub>–CH<sub>2</sub>), 32.4 (CH<sub>2</sub>–CH<sub>2</sub>–CH<sub>2</sub>), 42.0 (ClCH<sub>2</sub>), 50.8 (S–CH–S), 68.2 (OCH<sub>2</sub>), 115.0 (ArC), 129.3 (ArC), 132.3 (CHC<sub>ipso</sub>), 158.3 (OC<sub>ipso</sub>). EIMS (*m/z*): Calcd. 274.03 [M]<sup>+</sup>. Found: 274.02. Elemental analysis for C<sub>12</sub>H<sub>15</sub>ClOS<sub>2</sub>: Calcd. C, 52.44; H, 5.50. Found: C, 52.35; H, 5.44.

**[2-(4-[1,3]Dithian-2-yl-phenoxy)ethyl]diphenylphosphane (2).** In a 100-mL Schlenk round-bottom flask, **1** (1.00 g, 3.64 mmol) was dissolved in THF (50 mL). To this solution, KPPH<sub>2</sub> (7.28 mL of a 0.5 M solution in THF, 3.64 mmol) was added over 10 min and allowed to stir for an additional 30 min. The solvent was removed, and the residue was extracted with degassed CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O. The solvent was removed from the organic fraction, yielding **2** as an off-white solid, which was recrystallized from CH<sub>2</sub>Cl<sub>2</sub>/hexanes (1.37 g, 86% yield). <sup>1</sup>H NMR (CD<sub>2</sub>Cl<sub>2</sub>): δ 1.83 (m, 1H, CH<sub>2</sub>–CH<sub>A</sub>H<sub>B</sub>–CH<sub>2</sub>), 2.11 (m, 1H, CH<sub>2</sub>–CH<sub>A</sub>H<sub>B</sub>–CH<sub>2</sub>), 2.54 (t, 2H, CH<sub>2</sub>P), 2.84 (m, 2H, SCH<sub>2</sub>), 3.01 (m, 2H, SCH<sub>2</sub>), 4.07 (q, 2H, OCH<sub>2</sub>), 5.12 (s, 1H, S–CH–S), 6.73 (d, 2H, J<sub>H–H</sub> = 8.7 Hz, ArH), 7.31–7.50 (br m, 12H, P(Ar–H)). <sup>13</sup>C{<sup>1</sup>H} NMR (CD<sub>2</sub>Cl<sub>2</sub>): δ 25.3 (CH<sub>2</sub>–CH<sub>2</sub>–CH<sub>2</sub>), 28.3 (CH<sub>2</sub>–CH<sub>2</sub>–CH<sub>2</sub>), 28.5 (CH<sub>2</sub>–CH<sub>2</sub>–CH<sub>2</sub>), 32.3 (CH<sub>2</sub>P), 50.8 (S–CH–S), 65.5, 65.8, 114.8 (ArC), 128.7 (ArC), 128.8 (ArC), 129.0 (ArC), 129.0 (ArC), 131.9 (ArC), 132.7 (ArC), 133.0 (ArC), 138.3 (ArC), 138.4 (CHC<sub>ipso</sub>), 158.7 (OC<sub>ipso</sub>). <sup>31</sup>P{<sup>1</sup>H} NMR (CD<sub>2</sub>Cl<sub>2</sub>): δ –21.6 (s). EIMS (*m/z*): Calcd. 421.11 [M]<sup>+</sup>. Found: 421.10. Elemental analysis for C<sub>24</sub>H<sub>25</sub>OPS<sub>2</sub>: Calcd. C, 67.90; H, 5.94. Found: C, 67.02; H, 5.86.

**2-[4-[2-(Diphenylphosphinothioyl)ethoxy]phenyl]-(1,3)dithiane (3).** In a 100-mL Schlenk round-bottom flask, **2** (1.18 g, 2.76 mmol) and elemental sulfur (88.5 mg, 2.76 mmol) were stirred in THF (50 mL) for 4 h at which point the solvent was removed *in vacuo* and recrystallized from CH<sub>2</sub>Cl<sub>2</sub>/hexanes to yield **3** as a light-yellow microcrystalline solid (1.23 g, 97% yield). <sup>1</sup>H NMR (CD<sub>2</sub>Cl<sub>2</sub>): δ 1.83 (m, 1H, CH<sub>2</sub>–CH<sub>A</sub>H<sub>B</sub>–CH<sub>2</sub>), 2.11 (m, 1H, CH<sub>2</sub>–CH<sub>A</sub>H<sub>B</sub>–CH<sub>2</sub>), 2.84 (m, 6H, (SCH<sub>2</sub>)<sub>2</sub> and CH<sub>2</sub>P), 4.35 (m, 2H, OCH<sub>2</sub>), 5.11 (s, 1H, S–CH–S), 6.62 (d, 2H, J<sub>H–H</sub> = 8.7 Hz, ArH), 7.27 (d, 2H, J<sub>H–H</sub> = 8.4 Hz, ArH), 7.46–7.91 (br m, 10H, P(ArH)). <sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>): δ 25.2 (CH<sub>2</sub>–CH<sub>2</sub>–CH<sub>2</sub>), 32.4 (CH<sub>2</sub>–CH<sub>2</sub>–CH<sub>2</sub>), 33.0 (CH<sub>2</sub>–CH<sub>2</sub>–CH<sub>2</sub>), 50.9 (CH<sub>2</sub>P=S), 53.6 (S–CH–S), 62.5 (OCH<sub>2</sub>), 114.8 (ArC), 128.8 (ArC), 129.0 (ArC), 129.1 (ArC), 131.1 (ArC), 131.3 (ArC), 131.8 (ArC), 131.9 (ArC), 132.2 (ArC), 133.3 (ArC), 158.1 (OC<sub>ipso</sub>). <sup>31</sup>P{<sup>1</sup>H} NMR (CD<sub>2</sub>Cl<sub>2</sub>): δ 39.3 (s). EIMS (*m/z*): Calcd. 456.08 [M]<sup>+</sup>. Found: 456.07. Elemental analysis for C<sub>24</sub>H<sub>25</sub>OPS<sub>3</sub>: Calcd. C, 68.84; H, 5.23. Found: C, 67.43; H, 4.90.

**4-[2-(Diphenylphosphinothioyl)ethoxy]benzaldehyde (4).** Under ambient conditions, NaNO<sub>2</sub> (226 mg, 3.27 mmol) and acetyl chloride (0.233 mL, 3.27 mmol) were stirred in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) a 100-mL round-bottom flask at 0 °C for 10 min. A solution of **3** (500 mg, 1.09 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was added and stirred for an additional 5 min at 0 °C. At this point, H<sub>2</sub>O (5 mL) was added and the reaction was brought to room temperature and allowed to stir for an hour. The reaction was neutralized with a saturated aqueous solution of NaHCO<sub>3</sub> and was extracted with CH<sub>2</sub>Cl<sub>2</sub> (50 mL). The organic layer was washed with H<sub>2</sub>O (2 × 10 mL) and dried over MgSO<sub>4</sub>. After removing the solvent *in vacuo*, the crude product was purified via column chromatography (1:1 v/v, ethyl acetate/hexanes as eluent) yielding **4** as a light-yellow microcrystalline solid (352 mg, 88% yield). <sup>1</sup>H NMR (CD<sub>2</sub>Cl<sub>2</sub>): δ 2.98 (m, 2H, CH<sub>2</sub>P), 4.46 (m, 2H, OCH<sub>2</sub>), 6.78 (d, 2H, J<sub>H–H</sub> = 9 Hz, ArH), 7.48–7.91 (br m, 12H, P(ArH)), 9.84 (s, 1H, CHO). <sup>13</sup>C{<sup>1</sup>H} NMR (CD<sub>2</sub>Cl<sub>2</sub>): δ 32.1 (d, J<sub>C–P</sub> = 56 Hz, CH<sub>2</sub>P=S), 63.0 (OCH<sub>2</sub>), 114.8 (ArC), 128.8 (C<sub>ipso</sub>CHO), 128.9 (ArC), 130.4 (ArC), 131.1 (ArC), 131.2 (ArC), 131.9 (ArC), 132.4 (ArC), 133.5 (ArC), 163.2 (C<sub>ipso</sub>O), 190.7 (CHO). <sup>31</sup>P{<sup>1</sup>H} NMR (CD<sub>2</sub>Cl<sub>2</sub>): δ 39.2 (s). EIMS (*m/z*): Calcd. 366.08 [M]<sup>+</sup>. Found: 366.08. Elemental analysis for C<sub>21</sub>H<sub>19</sub>O<sub>2</sub>PS: Calcd. C, 63.13; H, 5.52. Found: C, 62.68; H, 5.22.

**5,15-Bis-[4-[2-(diphenylphosphinothioyl)ethoxy]phenyl]-10,20-bis-(mesityl)porphyrin (5).** In an aluminum-foil-wrapped 1000-mL Schlenk round-bottom flask, **4** (1.47 g, 4 mmol), 5-mesityldipyrromethane (1.06 g, 4 mmol), and activated molecular sieves (4 Å) were stirred in CHCl<sub>3</sub> (600 mL) and degassed under a stream of N<sub>2</sub> for 15 min. BF<sub>3</sub>•OEt<sub>2</sub> (0.420 mL) was added dropwise to this solution, and the resulting mixture was allowed to stir for 3 h under N<sub>2</sub>. DDQ (1.09 g, 4.8 mmol) was then added as a solid under a stream of N<sub>2</sub>, and the reaction was allowed to stir for an additional 30 min at which point NEt<sub>3</sub> (4 mL) was added. The reaction mixture was stirred for 1 min before being filtered through a pad of Celite to remove the sieves. The solution was concentrated *in vacuo* and the resulting residue was dissolved in a minimum amount CH<sub>2</sub>Cl<sub>2</sub> and poured on top of a silica gel column and purified via flash chromatography (eluent CH<sub>2</sub>Cl<sub>2</sub>) to yield **5** as a purple microcrystalline solid (1.01 g, 41% yield). <sup>1</sup>H NMR (CD<sub>2</sub>Cl<sub>2</sub>): δ –2.69 (s, 2H, NH), 1.82 (s, 12H, CH<sub>3</sub>), 2.66 (s, 6H, CH<sub>3</sub>), 3.16 (m, 4H, CH<sub>2</sub>P=S), 4.66 (m, 4H, OCH<sub>2</sub>), 7.07 (d, 4H, J<sub>H–H</sub> = 7.8 Hz, ArH), 7.30 (s, 4H), 7.59 (br s, 12H, ArH), 7.96 (m, 12H, ArH), 8.65 (d, 4H, J<sub>H–H</sub> = 5.1 Hz, ArH), 8.79 (d, 4H, J<sub>H–H</sub> = 4.5 Hz, ArH). <sup>31</sup>P{<sup>1</sup>H} NMR (CD<sub>2</sub>Cl<sub>2</sub>): δ 39.5 (s). ESIMS (*m/z*): Calcd. 1219.4 [M]<sup>+</sup>.

(45) Mao, P. C.-M.; Mouscadet, J.-F.; Leh, H.; Auclair, C.; Hsu, L.-Y. *Chem. Pharm. Bull.* **2002**, *50*, 1634–1637.

(46) Brown, A. M.; Ovchinnikov, M. V.; Stern, C. L.; Mirkin, C. A. *J. Am. Chem. Soc.* **2004**, *126*, 14316–14317.

(47) Laha, J. K.; Dhanalekshmi, S.; Taniguchi, M.; Ambrose, A.; Lindsey, J. S. *Org. Process Res. Dev.* **2003**, *7*, 799–812.

(48) Brown, A. M.; Ovchinnikov, M. V.; Mirkin, C. A. *Angew. Chem., Int. Ed.* **2005**, *44*, 4207–4209.

(49) Tomizaki, K.-y.; Yu, L.; Wei, L.; Bocian, D. F.; Lindsey, J. S. *J. Org. Chem.* **2003**, *68*, 8199–8207.



Found: 1219.3. Elemental analysis for  $C_{78}H_{68}N_4O_2P_2S_2$ : Calcd. C, 76.82; H, 5.62; N, 4.59. Found: C, 76.04; H, 5.18; N, 4.65.

**[5,15-Bis-{4-[2-(diphenylphosphinothioyl)ethoxy]phenyl}-10,20-bis-(mesityl)porphyrinato]zinc (II) (6).** Under ambient conditions, **5** (500 mg, 0.399 mmol) and  $Zn(OAc)_2 \cdot 2H_2O$  (700 mg, 3.19 mmol) were combined in a 500-mL round-bottom flask and stirred under reflux for 4 h in a  $CHCl_3/CH_3OH$  (4:1 v/v, 350 mL) solution. The solution was then washed with  $H_2O$  (100 mL) and extracted with  $CHCl_3$  ( $2 \times 100$  mL). The organic layer was further washed with  $H_2O$  (100 mL), dried over  $Na_2SO_4$ , and concentrated to give **6** as a purple microcrystalline solid (507 mg, 96% yield).  $^1H$  NMR (THF- $d_8$ ):  $\delta$  1.83 (s, 12H,  $CH_3$ ), 2.60 (m, 6H,  $CH_3$ ), 2.79 (m, 4H,  $CH_2P=S$ ), 4.36 (m, 4H,  $OCH_2$ ), 7.16 (br m, 28H, ArH), 8.09 (d, 4H,  $J_{H-H} = 8.4$  Hz, ArH), 8.61 (d, 4H,  $J_{H-H} = 4.5$  Hz, ArH), 8.77 (d, 4H,  $J_{H-H} = 4.5$  Hz, ArH).  $^{31}P\{^1H\}$  NMR (THF- $d_8$ ):  $\delta$  39.2 (s). ESIMS ( $m/z$ ): Calcd. 1282.8 [ $M^+$ ]. Found: 1282.2. Elemental analysis for  $C_{78}H_{66}N_4O_2P_2S_2Zn$ : Calcd. C, 73.03; H, 5.19; N, 4.37. Found: C, 72.89; H, 5.23; N, 4.17.

**[5,15-Bis-{4-(2-diphenylphosphanylethoxy)phenyl}-10,20-bis-(mesityl)porphyrinato]zinc(II) (7).** In a 50-mL Schlenk round-bottom flask, **6** (300 mg, 0.234 mmol) and  $Cp_2ZrHCl$  (392 mg, 1.52 mmol) were stirred in THF under  $N_2$  (40 mL) at 60 °C for 4 h. The solvent was removed and the reaction was purified via column chromatography (silica gel, THF) in a glove box under an atmosphere of  $N_2$ . The solvent was removed *in vacuo* to yield **7** as a purple microcrystalline solid (254 mg, 89% yield).  $^1H$  NMR (THF- $d_8$ ):  $\delta$  1.83 (s, 12H,  $CH_3$ ), 2.60 (s, 6H,  $CH_3$ ), 2.77 (m, 4H,  $CH_2P$ ), 4.36 (m, 4H,  $CH_2O$ ), 7.17 (d, 4H,  $J_{H-H} = 6.6$  Hz, ArH), 7.30–7.61 (bm, 24H, ArH), 8.01 (d, 4H,  $J_{H-H} = 8.7$  Hz, ArH), 8.64 (d, 4H,  $J_{H-H} = 4.5$  Hz, ArH), 8.78 (d, 4H,  $J_{H-H} = 4.5$  Hz, ArH).  $^{31}P\{^1H\}$  NMR (THF- $d_8$ ):  $\delta$  -21.2 (s). ESIMS ( $m/z$ ): Calcd. 1218.74 [ $M^+$ ]. Found: 1217.4. Elemental analysis for  $C_{78}H_{66}N_4O_2P_2Zn$ : Calcd. C, 76.87; H, 5.46; N, 4.60. Found: C, 76.37; H, 5.18; N, 4.45.

**[(7)RhCl(CO)]<sub>2</sub> Macrocycle (8a).** A small vial was charged with  $[Rh(CO)_2(Cl)]_2$  (4.80 mg, 0.0123 mmol) and  $CH_2Cl_2$  (2 mL). The resulting solution was stirred for 1 min, at which point a solution of **7** (30.0 mg, 0.0246 mmol) in THF (5 mL) was added dropwise over 1 min. The resulting solution was stirred for 3 h. The solvent was removed and the product was recrystallized from  $CH_2Cl_2$ /pentane (32.1 mg, 94% yield).  $^1H$  NMR ( $CD_2Cl_2$ ):  $\delta$  1.54 (s, 24H,  $CH_3$ ), 2.56 (s, 12H,  $CH_3$ ), 3.34 (br m, 8H,  $CH_2P$ ), 4.81 (br m, 8H,  $CH_2O$ ), 7.10 (br m, 22H, ArH), 7.47 (m, 22H, ArH), 7.90 (br m, 20H, ArH), 8.79 (br m, 16H, ArH).  $^{31}P\{^1H\}$  NMR ( $CD_2Cl_2$ ):  $\delta$  21.6 (d,  $J_{Rh-P} = 124$  Hz). ESIMS ( $m/z$ ) for  $[C_{156}H_{132}N_8O_4P_4Rh_2Zn_2(CO)_2]^{2+}$ : Calcd. 1349.6. Found: 1349.3. Elemental analysis for  $C_{158}H_{132}Cl_2N_8O_6P_4Rh_2Zn_2$ : Calcd. C, 68.51; H, 4.80; N, 4.04. Found: C, 68.38; H, 4.91; N, 3.52.

**[(7)Cu(CH<sub>3</sub>CN)<sub>2</sub>(PF<sub>6</sub>)<sub>2</sub> Macrocycle (8b).** A 50-mL Schlenk flask was charged with  $[Cu(CH_3CN)_4]PF_6$  (29.8 mg, 0.0799 mmol) and  $CH_2Cl_2$  (5 mL). A THF solution of **7** (100 mg, 0.0799 mmol, 20 mL) was added to the “Cu” solution dropwise over 5 min at room temperature to give a red/purple solution, which was then allowed to stir for 3 h. The solvent was removed to yield a purple microcrystalline solid which was recrystallized from  $CH_2Cl_2$ /pentane (107 mg, 92% yield).  $^1H$  NMR ( $CD_2Cl_2$ ):  $\delta$  1.66 (s, 24H,  $CH_3$ ), 2.07 (s, 12H,  $CH_3CN$ ), 2.43 (s, 12H,  $CH_3$ ), 2.95 (br m, 8H,  $CH_2P$ ), 4.46 (br m, 8H,  $CH_2O$ ), 7.10 (s, 22H, ArH), 7.50 (s, 22H, ArH), 7.62 (12H, ArH), 8.00 (d, 8H,  $J_{H-H} = 7.8$  Hz, ArH), 8.64 (d, 8H,  $J_{H-H} = 4.2$  Hz, ArH), 8.78 (d, 8H,  $J_{H-H} = 4.2$  Hz, ArH).  $^{31}P\{^1H\}$  NMR ( $CD_2Cl_2$ ):  $\delta$  -11.5 (s). ESIMS ( $m/z$ ) for  $[C_{156}H_{132}N_8S_4P_4Cu_2Zn_2]^{2+}$ : Calcd. 1282.2. Found: 1282.4. Elemental analysis for  $[C_{164}H_{144}N_{12}O_4P_4F_{12}Zn_2Cu_2]$ : Calcd. C, 65.25; H, 4.81; N, 5.57. Found: C, 65.32; H, 4.56; N, 6.35.

**1-Bromo-4-[2-(diphenylphosphinothioyl)ethylsulfanyl]-benzene (9).** In a 100-mL Schlenk round-bottom flask, 2-(4-bromo-phenylsulfanyl)-ethylidiphenylphosphane (3.00 g, 7.48 mmol), and elemental sulfur (264 mg, 8.22 mmol) were stirred in THF (150 mL) under  $N_2$  for 3 h. The reaction mixture was concentrated *in vacuo*, and the product was purified via column chromatography (1:1 v/v,  $CH_2Cl_2$ /hexanes).

Compound **9** was isolated as an off-white microcrystalline solid (3.02 g, 92% yield).  $^1H$  NMR ( $CD_2Cl_2$ ):  $\delta$  2.68 (m, 2H,  $CH_2P=S$ ), 3.10 (m, 2H,  $SCH_2$ ), 7.12 (d, 2H,  $J_{H-H} = 8.7$  Hz, ArH), 7.39 (d, 2H,  $J_{H-H} = 8.7$  Hz, ArH), 7.48–7.79 (m, 10H, P(ArH)).  $^{13}C\{^1H\}$  NMR ( $CDCl_3$ ):  $\delta$  26.9 ( $SCH_2$ ), 32.3 (d,  $CH_2P=S$ ,  $J_{C-P} = 51.3$  Hz), 128.9 (ArC), 129.1 (ArC), 131.0 (ArC), 131.1 (ArC), 131.2 (ArC), 131.6 (ArC), 132.0 (ArC), 132.3 (ArC).  $^{31}P\{^1H\}$  NMR ( $CD_2Cl_2$ ):  $\delta$  41.4 (s). EIMS ( $m/z$ ): Calcd. 431.97 [ $M^+$ ]. Found: 431.97. Elemental analysis for  $C_{20}H_{18}BrP_2S_2$ : Calcd. C, 55.43; H, 4.19. Found: C, 55.59; H, 4.07.

**4-[2-(Diphenylphosphinothioyl)ethylsulfanyl]benzaldehyde (10).** Compound **9** (3.00 g, 6.90 mmol) was dissolved in THF (60 mL) in a 100-mL Schlenk round-bottom flask and was cooled to -78 °C. *n*-BuLi (2.76 mL, 6.90 mmol, 2.5 M in hexanes) was added dropwise to the solution over 5 min and the mixture was allowed to stir for 30 min, before DMF (0.802 mL, 10.35 mmol) was added to the flask. This solution was cooled to -78 °C and allowed to stir for an additional 30 min before the temperature was allowed to rise back to room temperature. The mixture was quenched with  $H_2O$  followed by extraction with  $CH_2Cl_2$ . The organic layer was dried over  $MgSO_4$ , filtered, and concentrated *in vacuo* to give a yellow crude product which was then recrystallized ( $CH_2Cl_2$ /pentane) to yield **10** as a light-yellow microcrystalline solid (2.35 g, 88% yield).  $^1H$  NMR ( $CD_2Cl_2$ ):  $\delta$  2.76 (m, 2H,  $CH_2P=S$ ), 3.23 (m, 2H,  $SCH_2$ ), 7.29 (d, 2H,  $J_{H-H} = 6.6$  Hz, ArH), 7.47 (m, 6H, P(ArH)), 7.73 (d, 2H,  $J_{H-H} = 8.7$  Hz, ArH), 7.79 (m, 4H, P(ArH)), 9.92 (s, 1H, CHO).  $^{13}C\{^1H\}$  NMR ( $CDCl_3$ ):  $\delta$  25.0 ( $SCH_2$ ), 32.0 (d,  $CH_2P=S$ ,  $J_{C-P} = 51.3$  Hz), 126.8 (ArC), 129.0 (ArC), 129.1 (ArC), 130.4 (ArC), 131.1 (ArC), 131.3 (ArC), 132.1 (ArC), 132.5 (ArC), 191.4 (CHO).  $^{31}P\{^1H\}$  NMR ( $CD_2Cl_2$ ):  $\delta$  41.5 (s). EIMS ( $m/z$ ): Calcd. 382.01 [ $M^+$ ]. Found: 382.06. Elemental analysis for  $C_{21}H_{19}OPS_2$ : Calcd. C, 65.94; H, 5.01. Found: C, 65.23; H, 4.71.

**[5,15-Bis-{4-(2-diphenylphosphinothioylethylsulfanyl)phenyl}-10,20-bis-(mesityl)porphyrin (11).** In an aluminum-foil-wrapped 1000-mL Schlenk flask, compound **10** (1.23 g, 3.20 mmol), 5-mesityldipyrromethane (848 mg, 3.20 mmol), and activated molecular sieves (4 Å) were stirred in  $CHCl_3$  (600 mL) and degassed under a stream of  $N_2$  for 15 min.  $BF_3 \cdot OEt_2$  (0.350 mL) was added dropwise to this solution, and the resulting mixture was allowed to stir for 3 h under  $N_2$ . DDQ (862 mg, 3.8 mmol) was then added as a solid under a stream of  $N_2$ , and the reaction was allowed to stir for an additional 30 min at which point  $NEt_3$  (4 mL) was added. The reaction mixture was stirred for 1 min before being filtered through a pad of Celite to remove the sieves. The solution was concentrated *in vacuo* and the resulting residue was dissolved in  $CH_2Cl_2$  and poured on top of a silica gel column (eluent  $CH_2Cl_2$ ). Subsequent chromatography yielded **11** as a purple microcrystalline solid (879 mg, 44% yield).  $^1H$  NMR ( $CD_2Cl_2$ ):  $\delta$  -2.67 (s, 2H, NH), 1.84 (s, 12H,  $CH_3$ ), 2.63 (s, 6H,  $CH_3$ ), 3.03 (m, 4H,  $CH_2P=S$ ), 3.42 (m, 4H,  $SCH_2$ ), 7.31 (s, 4H, mesityl H), 7.55 (m, 16H, ArH), 7.89 (m, 8H, ArH), 8.13 (d, 4H,  $J_{H-H} = 8.1$  Hz, ArH), 8.69 (d, 4H,  $J_{H-H} = 5.1$  Hz, ArH), 8.82 (d, 4H,  $J_{H-H} = 4.8$  Hz, ArH).  $^{31}P\{^1H\}$  NMR ( $CD_2Cl_2$ ):  $\delta$  41.6 (s). ESIMS ( $m/z$ ): Calcd. 1251.6 [ $M^+$ ]. Found: 1251.3. Elemental analysis for  $C_{78}H_{68}N_4P_2S_4$ : Calcd. C, 74.85; H, 5.48; N, 4.48. Found: C, 74.35; H, 4.98; N, 4.43.

**[5,15-Bis-{4-(2-diphenylphosphinothioylethylsulfanyl)phenyl}-10,20-bis(mesityl)porphyrinato] zinc(II) (12).** In a 250-mL round-bottom flask, **11** (400 mg, 0.319 mmol) and  $Zn(OAc)_2 \cdot 2H_2O$  (700 mg, 3.19 mmol) were refluxed for 3 h in a  $CHCl_3/CH_3OH$  (4:1 v/v, 200 mL) solution. The solution was washed with  $H_2O$  (100 mL) and extracted with  $CHCl_3$  ( $2 \times 50$  mL). The organic layer was washed again with  $H_2O$  (50 mL), dried over  $Na_2SO_4$ , and concentrated to give **12** as a purple microcrystalline solid (412 mg, 98% yield).  $^1H$  NMR ( $CD_2Cl_2$ ):  $\delta$  1.82 (s, 12H,  $CH_3$ ), 2.63 (s, 6H,  $CH_3$ ), 3.03 (m, 4H,  $CH_2P=S$ ), 3.41 (m, 4H,  $SCH_2$ ), 7.31 (s, 4H, ArH), 7.54 (m, 12H, ArH), 7.89 (m, 12H, ArH), 8.12 (d, 4H,  $J_{H-H} = 8.4$  Hz, ArH), 8.76 (d, 4H,  $J_{H-H} = 5.1$  Hz, ArH), 8.89 (d, 4H,  $J_{H-H} = 4.8$  Hz, ArH).  $^{31}P\{^1H\}$  NMR ( $CD_2Cl_2$ ):  $\delta$  41.6 (s). ESIMS ( $m/z$ ): Calcd. 1314.9 [ $M^+$ ]. Found: 1314.1.

Elemental analysis for  $C_{78}H_{66}N_4P_2S_4Zn$ : Calcd. C, 71.24; H, 5.06; N, 4.26. Found: C, 70.69; H, 4.97; N, 3.94.

**[5,15-Bis-[4-(2-diphenylphosphanylethylsulfanyl)phenyl]-10,20-bis(mesityl)porphyrinato]zinc(II) (13).** In a 50-mL Schlenk flask, **12** (300 mg, 0.228 mmol) and  $Cp_2ZrHCl$  (382 mg, 1.48 mmol) were stirred in THF (40 mL) at 60 °C for 3 h. The solvent was removed, and the product was purified via column chromatography (THF, silica gel) in a glove box under an atmosphere of  $N_2$ . The solvent was removed *in vacuo* to yield **13** as a purple microcrystalline solid (251 mg, 88% yield).  $^1H$  NMR (THF- $d_8$ ):  $\delta$  1.84 (s, 12H,  $CH_3$ ), 2.47 (m, 4H,  $CH_2P$ ), 2.61 (s, 6H,  $CH_3$ ), 3.23 (m, 4H,  $SCH_2$ ), 7.29 (br m, 28H,  $ArH$ ), 8.09 (d, 4H,  $J_{H-H} = 7.8$  Hz,  $ArH$ ), 8.64 (d, 4H,  $J_{H-H} = 4.8$  Hz,  $ArH$ ), 8.79 (d, 4H,  $J_{H-H} = 4.2$  Hz,  $ArH$ ).  $^{31}P\{^1H\}$  NMR (THF- $d_8$ ):  $\delta$  -15.2 (s). ESIMS ( $m/z$ ): Calcd. 1250.8 [ $M^+$ ]. Found: 1249.4. Elemental analysis for  $C_{78}H_{66}N_4P_2S_4Zn$ : Calcd. C, 74.90; H, 5.32; N, 4.48. Found: C, 73.15; H, 4.94; N, 4.16.

**[(13)Rh(BF<sub>4</sub>)<sub>2</sub>] Condensed Macrocycle (14a).** A small vial was charged with  $[Rh(NBD)Cl]_2$  (31.0 mg, 0.068 mmol),  $AgBF_4$  (26.0 mg, 0.135 mmol), and  $CH_2Cl_2$  (4 mL). This solution was stirred for 1 h and then filtered dropwise through Celite into a Schlenk flask. The red solution was then diluted with  $CH_2Cl_2$  (20 mL) to give a clear yellow/orange solution. A solution of ligand **13** (170 mg, 0.135 mmol) in THF (20 mL) was added to the “Rh” solution dropwise over 5 min at room temperature to give a dark-purple solution, which was then stirred for an additional 3 h. The solvent was removed to yield **14a** as a purple microcrystalline solid which was recrystallized from  $CH_2Cl_2$ /pentane (176 mg, 90% yield).  $^1H$  NMR ( $CD_2Cl_2$ ):  $\delta$  1.48 (br s, 24H,  $CH_3$ ), 2.44 (br s, 12H,  $CH_3$ ), 2.76 (br m, 8H,  $CH_2P$ ), 3.20 (br m, 8H,  $CH_2S$ ), 6.98 (br m, 8H,  $ArH$ ), 7.37 – 7.59 (br m, 40H,  $ArH$ ), 8.02 (br m, 16H,  $ArH$ ), 8.50 (br m, 16H,  $ArH$ ).  $^{31}P\{^1H\}$  NMR ( $CD_2Cl_2$ ):  $\delta$  64.1 (d,  $J_{Rh-P} = 162$  Hz). ESIMS ( $m/z$ ) for  $[C_{156}H_{132}N_8S_4P_4Rh_2Zn_2]^{2+}$ : Calcd. 1353.7. Found: 1354.1. Elemental analysis for  $[C_{156}H_{132}N_8S_4P_4Rh_2Zn_2]$ : Calcd. C, 65.03; H, 4.62; N, 3.89. Found: C, 65.56; H, 4.11; N, 3.46.

**[(13)Cu(PF<sub>6</sub>)<sub>2</sub>] Condensed Macrocycle (14b).** A 50-mL Schlenk flask was charged with  $[Cu(CH_3CN)_4]PF_6$  (29.8 g, 0.0799 mmol). The  $Cu^I$  precursor was dissolved in  $CH_2Cl_2$  (5 mL), and a THF solution of ligand **13** (100 mg, 0.0799 mmol, 20 mL) was added to the “Cu” solution dropwise over 5 min at room temperature to give a red/purple solution. The solution was then allowed to stir for 3 h. The solvent was removed to yield **14b** as a purple microcrystalline solid which was recrystallized from  $CH_2Cl_2$ /pentane (105 mg, 90% yield).  $^1H$  NMR ( $CD_2Cl_2$ ):  $\delta$  1.75 (br s, 24H,  $CH_3$ ), 2.55 (br s, 12H,  $CH_3$ ), 3.00 (br m, 8H,  $CH_2P$ ), 3.65 (br m, 8H,  $CH_2S$ ), 7.24 – 7.82 (br m, 56H,  $ArH$ ), 8.16 (br m, 8H,  $ArH$ ), 8.76 (br s, 16H,  $ArH$ ).  $^{31}P\{^1H\}$  NMR ( $CD_2Cl_2$ ):  $\delta$  1.0 (s). ESIMS ( $m/z$ ) for  $[C_{156}H_{132}N_8S_4P_4Cu_2Zn_2]^{2+}$ : Calcd. 1314.4. Found: 1314.5. Elemental analysis for  $[C_{156}H_{132}N_8S_4P_4F_{12}Cu_2Zn_2]CH_2Cl_2$ : Calcd. C, 62.78; H, 4.50; N, 3.73. Found: C, 61.95; H, 4.18; N, 3.31.

**[(13)RhCl(CO)]<sub>2</sub> Macrocycle (15a).** Compound **14a** (20.0 mg, 0.00694 mmol) was dissolved in  $CD_2Cl_2$  (1 g) and placed in an air-free NMR tube. PPNCl (Bis(triphenylphosphoranylidene) ammonium chloride, 2 equiv) was added and the NMR tube was then pressurized with CO (1 atm) for 30 s. Compound **14a** was quantitatively converted to **15a** as determined by  $^{31}P$  NMR spectroscopy.  $^1H$  NMR ( $CD_2Cl_2$ ):  $\delta$  1.77 (br s, 24H,  $CH_3$ ), 2.54 (br s, 12H,  $CH_3$ ), 3.14 (br m, 8H,  $CH_2P$ ), 3.48 (br m, 8H,  $CH_2S$ ), 7.26 – 8.01 (br m, 124H,  $(C_6H_5)_3PPN$ ,  $P(C_6H_5)_2$ ,  $ArH$ ), 8.65 – 8.80 (br m, 16H,  $ArH$ ).  $^{31}P\{^1H\}$  NMR ( $CD_2Cl_2$ ):  $\delta$  22.1 (s, PPNCl),  $\delta$  25.1 (d,  $J_{Rh-P} = 123$  Hz). ESIMS ( $m/z$ ) for  $[C_{156}H_{132}N_8S_4P_4Rh_2Zn_2]^{2+}$  ( $M - 2(Cl^-/CO)$ ): Calcd. 1353.7. Found: 1353.7.

**[(13)Cu(pyridine- $d_5$ )<sub>2</sub>(PF<sub>6</sub>)<sub>2</sub>] Macrocycle (15b).** Compound **14b** (20.0 mg, 0.00640 mmol) was dissolved in  $CD_2Cl_2$  (1 g) and placed in an air-free NMR tube.  $C_5D_5N$  (4 equiv.) was added to this solution and the conversion of **15b** was quantitative as determined by  $^1H$  NMR and  $^{31}P$  NMR spectroscopy.  $^1H$  NMR ( $CD_2Cl_2$ ):  $\delta$  1.72 (br s, 24H,  $CH_3$ ), 2.45 (br s, 12H,  $CH_3$ ), 2.82 (br m, 8H,  $CH_2P$ ), 3.38 (br m, 8H,

$CH_2S$ ), 7.17 – 7.55 (br m, 56H,  $ArH$ ), 8.04 (d, 8H,  $J_{H-H} = 7.8$  Hz,  $ArH$ ), 8.64 (d, 8H,  $J_{H-H} = 4.5$  Hz,  $ArH$ ), 8.70 (d, 8H,  $J_{H-H} = 4.8$  Hz,  $ArH$ ).  $^{31}P\{^1H\}$  NMR ( $CD_2Cl_2$ ):  $\delta$  -8.0 (s). ESIMS ( $m/z$ ) for  $[C_{156}H_{132}N_8S_4P_4Cu_2Zn_2]^{2+}$ : Calcd. 1314.4. Found: 1314.9.


**[(Ph<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>S-C<sub>6</sub>H<sub>2</sub>(CH<sub>3</sub>)<sub>3</sub>)<sub>2</sub>Rh]BF<sub>4</sub> Condensed Tweezer-Type Complex (16a).** A 20-mL vial was charged with  $[Rh(NBD)Cl]_2$  (63.2 mg, 0.137 mmol),  $AgBF_4$  (53.4 mg, 0.274 mmol), and  $CH_2Cl_2$  (4 mL). This solution was stirred for 1 h and then filtered dropwise through Celite into a Schlenk flask. The orange solution was then diluted with  $CH_2Cl_2$  (20 mL) to give a clear yellow solution. A solution of (2-(mesitylthio)ethyl)diphenylphosphine (200 mg, 0.549 mmol) in  $CH_2Cl_2$  (20 mL) was added to the “Rh” solution dropwise over 5 min at room temperature to give a clear yellow solution, which was then stirred for an additional 3 h. The solvent was removed to yield **16a** as a yellow microcrystalline solid which was recrystallized from  $CH_2Cl_2$ /pentane (113 mg, 90% yield).  $^1H$  NMR ( $CD_2Cl_2$ ):  $\delta$  2.22 (d, 12H,  $CH_3$ ), 2.51 (br s, 10H,  $CH_3$  and  $SCH_2CH_2P$ ), 2.83 (br m, 4H,  $CH_2P$ ), 6.71 (s, 4H,  $ArH$ ), 7.23 – 7.34 (br m,  $P(Ar-H)_2$ , 20H).  $^{31}P\{^1H\}$  NMR ( $CD_2Cl_2$ ):  $\delta$  63.0 (d,  $J_{Rh-P} = 163$  Hz). ESIMS ( $m/z$ ) for  $[C_{46}H_{50}N_8S_2P_2Rh]^+$ : Calcd. 831.8. Found: 831.6.

**(Ph<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>S-C<sub>6</sub>H<sub>2</sub>(CH<sub>3</sub>)<sub>3</sub>)<sub>2</sub>RhCl(CO) Open Tweezer-Type Complex (16b).** Compound **16a** (15.0 mg, 0.0163 mmol) was dissolved in  $CD_2Cl_2$  (1 g) and placed in an air-free NMR tube. PPNCl (bis(triphenylphosphoranylidene)ammonium chloride, 9.38 mg, 0.0163 mmol) was added as a solid and the NMR tube was then pressurized with CO (1 atm) for 30 s. Compound **16a** was quantitatively converted to **16b** as determined by  $^{31}P\{^1H\}$  NMR spectroscopy.  $^1H$  NMR ( $CD_2Cl_2$ ):  $\delta$  2.25 (s, 6H,  $CH_3$ ), 2.37 (s, 12H,  $CH_3$ ), 2.67 (br m, 4H,  $CH_2P$ ), 2.82 (br m, 4H,  $CH_2S$ ), 6.90 (s, 4H,  $ArH$ ), 7.34 – 7.65 (br m, 50H,  $(C_6H_5)_3PPN$ ,  $P(Ar-H)_2$ ).  $^{31}P\{^1H\}$  NMR ( $CD_2Cl_2$ ):  $\delta$  22.1 (s, PPNCl),  $\delta$  23.5 (d,  $J_{Rh-P} = 123$  Hz). ESIMS ( $m/z$ ) for  $[C_{46}H_{50}N_8S_2P_2Rh]^+$  ( $M - (Cl^-/CO)$ ): Calcd. 831.8. Found: 831.2.

**General Procedure for Catalysis Experiments.** The formation of 2-, 3-, and 4-acetoxymethylpyridine were monitored by GC relative to an internal standard (biphenyl) and quantified using a previously established calibration curve. Experiments containing **14a**, **15a**, or a mixture of an analogous monomeric  $Rh^I$  complex (**16a** or **16b**) and **Zn(TPP)** (as the control), respectively, were run concurrently in separate vials at room temperature in  $CH_2Cl_2$  under an atmosphere of  $N_2$  inside a glove box.

**GC Experiments for the Closed Macrocycle (14a).** Inside a glove box,  $CH_2Cl_2$  stock solutions of complex **14a** (0.58 mL of a 2.6 mM solution), biphenyl (0.5 mL of a 25 mM solution), and pyridylcarbinol (0.5 mL of a 90 mM solution) were added to a 20-mL vial. Fresh  $CH_2Cl_2$  was added to the vial bringing the total reaction volume up to 4.5 mL. After stirring for 5 min, 1-acetylimidazole (0.5 mL of a 60 mM solution) was added to the vial ( $t = 0$ ). At various times, an aliquot (100  $\mu$ L) was taken from the solution and added to diethyl ether (2 mL). This was then passed down a plug of Celite (3 cm  $\times$  0.5 cm) to remove the catalyst. The plug was further treated with fresh diethyl ether (5 mL). The combined organics were used for GC analysis.

**GC Experiments for the Open Macrocycle (15a).** Inside a glove box, complex **14a** (0.58 mL of a 2.6 mM solution) and benzyltriethylammonium chloride (0.5 mL of a 6.4 mM solution) were added to a 10-mL Schlenk flask. The flask was removed from the glove box and placed on a Schlenk line where CO (1 atm) was bubbled through the solution for 30 s resulting in the formation of **15a**. The flask was brought back into the glove box and the contents were transferred into a vial. Fresh  $CH_2Cl_2$  was added to the flask and transferred into the vial bringing the total reaction volume up to approximately 4 mL. Pyridylcarbinol (0.5 mL of a 90 mM solution) and biphenyl (0.5 mL of a 25 mM solution) were added to the vial and after stirring for 5 min, 1-acetylimidazole (0.5 mL of a 60 mM solution) was added to the flask ( $t = 0$ ). At various times, an aliquot (100  $\mu$ L) was taken from the solution and added to diethyl ether (2 mL). This was then passed down a plug of Celite (3 cm  $\times$  0.5 cm) to remove the catalyst. The

**Scheme 2.** Synthesis of Ether-Based Ligand **7** and Macrocycles **8a** and **8b**<sup>a</sup>

plug was further treated with fresh diethyl ether (5 mL). The resulting samples were used for GC analysis.

**GC Control Experiments for Rh<sup>I</sup>-Monomer (**16a**) and Porphyrin Monomer (**Zn(TPP)**).** Inside a glove box, CH<sub>2</sub>Cl<sub>2</sub> stock solutions of complex **16a** (0.892 mL of a 8.0 mM solution), biphenyl (0.5 mL of a 25 mM solution), Zn(TPP) (0.20 mL of a 15 mM solution), and pyridylcarbinol (0.5 mL of a 90 mM solution) were added to a 20-mL vial. Fresh CH<sub>2</sub>Cl<sub>2</sub> was added to the vial bringing the total reaction volume up to 4.5 mL. After stirring for 5 min, 1-acetylimidazole (0.5 mL of a 60 mM solution) was added to the vial (*t* = 0). At various times, an aliquot (100 μL) was taken from the solution and added to diethyl ether (2 mL). This was then passed down a plug of Celite (3 cm × 0.5 cm) to remove the catalyst. The plug was further treated with fresh diethyl ether (5 mL). The combined organics were used for GC analysis.

**GC Control Experiments for Rh<sup>I</sup>-Monomer (**16b**) and Porphyrin Monomer.** Inside a glove box, complex **16a** (0.892 mL of a 8.0 mM solution), Zn(TPP) (0.20 mL of a 15 mM solution), and benzyltriethylammonium chloride (1.0 mL of a 6.4 mM solution) were added to a 10-mL Schlenk flask. The flask was removed from the glove box and placed on a Schlenk line where CO (1 atm) was bubbled through the solution for 30 s resulting in the formation of **16b**. The flask was brought back into the glove box and the contents were transferred into a vial. Fresh CH<sub>2</sub>Cl<sub>2</sub> was added to the flask and transferred into the

vial bringing the total reaction volume up to approximately 4 mL. Pyridylcarbinol (0.5 mL of a 90 mM solution) and biphenyl (0.5 mL of a 25 mM solution) were added to the vial and after stirring for 5 min, 1-acetylimidazole (0.5 mL of a 60 mM solution) was added to the flask (*t* = 0). At various times, an aliquot (100 μL) was taken from the solution and added to diethyl ether (2 mL). This was then passed down a plug of Celite (3 cm × 0.5 cm) to remove the catalyst. The plug was further treated with fresh diethyl ether (5 mL). The combined organics were used for GC analysis.

## Results and Discussion

**The Design of Ether–Phosphine Hemilabile Ligand **7**.** To design porphyrin-based supramolecular systems capable of *in situ* allosteric activity, flexibility must be engineered into the backbone of the framework as well as “weak” and “strong” binding structural domains. To this end, we synthesized two new hemilabile porphyrin ligands which incorporate “weak” ether or thioether functionalities in addition to “strong” phosphine binding sites. Ether-based ligand **7** was obtained in eight steps from 4-hydroxybenzaldehyde (Scheme 2). 4-Hydroxybenzaldehyde was alkylated with excess 1-bromo-2-chloroethane to yield 4-(2-chloroethoxy)benzaldehyde, which was transformed into **3** after a dithiane protection of the aldehyde,




conversion of the chloride to the corresponding phosphine, and reaction with elemental sulfur. Deprotection of the dithiane moiety in **3**, followed by condensation of the resulting aldehyde with 5-mesityldipyrromethane affords the freebase porphyrin **5** in gram quantities after column chromatography. Metalation with  $\text{Zn}(\text{OAc})_2 \cdot 2\text{H}_2\text{O}$  gave compound **6**, and desulfurization with Schwartz's reagent gave ligand **7** in 12% overall yield. The use of a sulfide as the protecting group for the phosphine moiety not only allowed for a convenient nonaqueous protection/deprotection sequence, but also significantly increased the scalability and the yield of the porphyrin synthesis while simplifying the complicated purification and isolation process that often accompanies porphyrin syntheses. Indeed, if one uses an oxide protecting group as opposed to the sulfide, the ligand adheres to the silica column used for chromatography and makes the isolation of the pure product difficult.

**Reactivity of Ligand 7 with  $\text{Rh}^{\text{I}}$  and  $\text{Cu}^{\text{I}}$ .** Initially, our attention was focused on the design, synthesis, and isolation of  $\text{Rh}^{\text{I}}$  and  $\text{Cu}^{\text{I}}$  condensed intermediates of type **I** using ligand **7**. Unfortunately, isolation of these products proved impossible under the conditions explored. However, we discovered that the “open” macrocyclic structure **8a** can be obtained directly in quantitative yield upon reaction of ligand **7** with  $[\text{Rh}(\text{CO})_2(\text{Cl})]_2$ . The  $^31\text{P}\{^1\text{H}\}$  NMR spectrum of **8a** exhibits a single resonance at  $\delta$  21.6 (d,  $J_{\text{Rh-P}} = 124$  Hz), diagnostic of a highly symmetrical porphyrin complex with *trans*-phosphines and is consistent with the proposed structural formulation for **8a**.<sup>50</sup> ESIMS analysis of **8a** shows a parent ion at  $m/z$  1349.3, indicating the loss of each  $\text{Cl}^-$  ligand bound to the  $\text{Rh}^{\text{I}}$  centers.  $\text{Rh}^{\text{I}}$  complexes with analogous coordination environments often lose these ligands during ESIMS.<sup>50</sup>

Similar to the reactivity exhibited with  $\text{Rh}^{\text{I}}$ , ligand **7** forms the analogous “open” macrocyclic product **8b** when reacted with  $[\text{Cu}(\text{CH}_3\text{CN})_4]\text{PF}_6$ . The  $^31\text{P}\{^1\text{H}\}$  NMR spectrum of **8b** exhibits a singlet at  $\delta$  −11.5 consistent with a tetrahedral  $\text{Cu}^{\text{I}}$ –P environment.<sup>51</sup> The resonance is substantially shifted downfield from the one observed for free ligand at  $\delta$  −21.2. Characterization of this product via ESIMS leads to the observation of a parent ion at  $m/z$  1282.4 which corresponds to the loss of two  $\text{CH}_3\text{CN}$  molecules from each  $\text{Cu}^{\text{I}}$  metal center. This is consistent with the known lability of the bound  $\text{CH}_3\text{CN}$  molecules, which has been observed in analogous systems.<sup>51</sup>

**X-ray Structure Determination of  $8a\text{cDABCO}$  and  $8c\text{cDABCO}$ .** Initial attempts to grow crystals of **8a** in a variety of solvent mixtures proved unsuccessful, presumably due to the large free volume and flexibility of the cavity. We hypothesized that crystals of **8a** could be obtained in the presence of a ligand that can span and coordinate to the two Zn-centers within the macrocycle, filling the free volume and rigidifying the overall structure. Indeed, X-ray quality crystals were obtained from vapor diffusion of pentane/diethyl ether (1:1 v/v) into a THF/ $\text{CH}_2\text{Cl}_2$  (1:1 v/v) solution of **8a** containing DABCO which was layered with acetonitrile. The 56-membered macrocyclic product **8acDABCO** contains two porphyrin moieties aligned and locked into a cofacial arrangement in the solid-state by a DABCO ligand (Figure 1). The Zn–Zn distance of 7.09 Å is significantly longer than the P–Rh–P distance of 4.64 which



**Figure 1.** Graphical representations of the X-ray crystal structure of **8acDABCO** as viewed (A) from the side and (B) from the top containing a molecule of DABCO bridging both Zn atoms. Hydrogen atoms, disordered DABCO carbon atoms, and solvent molecules have been omitted for clarity. Pink = Rh, Red = O, Yellow = Cl, Green = P, Blue = N, Light Blue = Zn.

is made possible by the flexible  $(\text{CH}_2)_2\text{O}$  linkages between the porphyrin moieties and the  $\text{Rh}^{\text{I}}$  hinges. This large porphyrin–porphyrin distance allows for the mesityl arms of the porphyrin moieties to align in a near superimposable fashion.

Attempts to crystallize **8b** in a similar manner led to an unexpected reaction with the halogenated solvent. Because the solubility of **8b** in pure THF is poor, a mixture of solvents proved to be the most effective for crystallization. Analytically pure samples of **8b** were dissolved in a THF/ $\text{CH}_2\text{Cl}_2$  solution containing DABCO which was first layered with acetonitrile following by a slow second layer of a solution of pentane/diethyl ether (1:1 v/v). X-ray quality crystals were obtained from this mixture after 1 day and subjected to single-crystal diffraction analysis at the Advanced Photon Source (APS, Argonne National Laboratory). Although **8b** has been shown to contain four molecules of  $\text{CH}_3\text{CN}$  and two  $\text{PF}_6^-$  counterions according to elemental analysis, the diffraction experiment yielded a structure with DABCO bridging both Zn-atoms and two  $\text{Cl}^-$  anions bound to each tetrahedral  $\text{Cu}^{\text{I}}$  metal center, indicating an oxidation of the original  $\text{Cu}^{\text{I}}$  center to  $\text{Cu}^{\text{II}}$  forming **8ccDABCO** (see Supporting Information). Such a compound could be formed from the dehalogenation of alkyl halide solvents (i.e.,  $\text{CH}_2\text{Cl}_2$  and  $\text{CHCl}_3$ ) by the  $\text{Cu}^{\text{I}}$  center in **8b**, similar to that observed by Karlin et al. for  $(\text{TMPA})\text{Cu}^{\text{I}}$  ( $\text{TMPA} = (\text{tris}-(2\text{-pyridylmethyl})\text{amine})$ ) in  $\text{CH}_2\text{Cl}_2$ .<sup>52–54</sup> That **8ccDABCO** possesses the framework expected for **8b** supports our assignment of an open macrocyclic structure for the product isolated from the reaction of **7** and  $[\text{Cu}(\text{CH}_3\text{CN})_4]\text{PF}_6$ .


(50) Dixon, F. M.; Eisenberg, A. E.; Farrell, J. R.; Mirkin, C. A.; Liable-Sands, L. M.; Rheingold, A. L. *Inorg. Chem.* **2000**, *39*, 3432–3433.

(51) Masar III, M. S.; Mirkin, C. A.; Stern, C. L.; Zakharov, L. N.; Rheingold, A. L. *Inorg. Chem.* **2004**, *43*, 4693–4701.

(52) Lucchese, B.; Humphreys, K. J.; Lee, D. H.; Incarvito, C. D.; Sommer, R. D.; Rheingold, A. L.; Karlin, K. D. *Inorg. Chem.* **2004**, *43*, 5987–5998.

(53) Tyeklar, Z.; Richard, R. R.; Wei, N.; Murthy, N. N.; Zubieta, J.; Karlin, K. D. *J. Am. Chem. Soc.* **1994**, *116*, 5987–5988.

(54) Wei, N.; Murthy, N. N.; Tyeklar, Z.; Karlin, K. D. *Inorg. Chem.* **1994**, *33*, 1177–1183.

**Scheme 3.** Synthesis of Thioether-Based Ligand **13** and Macrocycles **14a–b**, **15a–b**<sup>a</sup>

<sup>a</sup> (i)  $\text{ClCH}_2\text{CH}_2\text{PPh}_2$ ,  $\text{Cs}_2\text{CO}_3$ ,  $\text{CH}_3\text{CN}$ , Reflux; (ii)  $\text{S}_8$ , THF; (iii)  $n\text{-BuLi}$ , DMF, THF,  $-78^\circ\text{C}$ ; (iv) 5-mesityldipyrromethane,  $\text{BF}_3\cdot\text{OEt}_2$ , DDQ,  $\text{NEt}_3$ ,  $\text{CHCl}_3$ , 4 Å Molecular Sieves; (v)  $\text{Zn}(\text{OAc})_2\cdot 2\text{H}_2\text{O}$ , 4:1  $\text{CHCl}_3/\text{MeOH}$ , Reflux; (vi)  $\text{Cp}_2\text{ZrHCl}$ , THF,  $60^\circ\text{C}$ ; (vii) for **14a**:  $[\text{Rh}(\text{NBD})\text{Cl}]_2$ ,  $\text{AgBF}_4$ ,  $\text{CH}_2\text{Cl}_2/\text{THF}$ ; (viii) for **14b**:  $[\text{Cu}(\text{CH}_3\text{CN})_4]\text{PF}_6$ ,  $\text{CH}_2\text{Cl}_2/\text{THF}$ ; (ix) for **15a**:  $\text{PPNCl}/\text{CO}$  (1 atm); (x) for **15b**:  $\text{C}_5\text{D}_5\text{N}$ .

### Synthesis of Thioether–Phosphine Hemilabile Ligand **13**.

Because type **I** structures are difficult to access from the ether–phosphine ligand **7**, we hypothesized that isostructural ligands containing thioether linkages would be more effective at forming condensed structures as has been observed previously with analogous smaller molecules.<sup>50</sup> To this end, we modified the synthesis of **7** to obtain the thioether-based porphyrin ligand **13** in six steps (Scheme 3). First, 4-bromothiophenol was stoichiometrically alkylated with 1-chloro-2-diphenylphosphinoethane to yield 2-(4-bromophenylsulfanyl)ethyl diphenyl phosphine. To protect the phosphine moiety from oxidation during the porphyrin synthesis, reaction with elemental sulfur yields phosphine sulfide **9**, which was then formylated with  $n\text{-BuLi}$  and DMF. The resulting aldehyde **10** was subsequently condensed with 5-mesityldipyrromethane in the presence of  $\text{BF}_3\cdot\text{OEt}_2$  to yield porphyrin **11** in 44% yield after a very simple chromatographic separation. Freebase porphyrin **11** was metallated with  $\text{Zn}(\text{OAc})_2\cdot 2\text{H}_2\text{O}$  to give **12** and followed by deprotection of the phosphine with Schwartz's reagent to yield the phosphine derivative **13** in 33% overall yield.

### Reactivity of Hemilabile Ligand **13** with $\text{Rh}^I$ and $\text{Cu}^I$ .


In contrast to the results obtained from ether ligand **7**, thioether ligand **13** reacts cleanly with “ $\text{Rh}(\text{NBD})\text{BF}_4$ ”<sup>55</sup> to form the condensed intermediate **14a**. The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of **14a** exhibits a doublet at  $\delta$  64.5 ( $J_{\text{Rh-P}} = 162$  Hz) which is highly diagnostic of its symmetrical structure and *cis*-Rh–P coordination centers.<sup>50</sup> Additionally, ESIMS analysis shows a peak corresponding to the  $\text{M}^{2+}$  ion at 1354.1  $m/z$ , indicating the formation of the desired condensed structure. Significantly, compound **14a** can be opened into macrocycle **15a** by introduction of benzyltriethyl ammonium chloride or by  $\text{PPNCl}$  (where  $\text{PPNCl}$  = bis(triphenylphosphoranylidene)ammonium chloride) and CO. This transformation can be followed by  $^{31}\text{P}\{^1\text{H}\}$  NMR through the appearance of a doublet at  $\delta$  25.1 (d,  $J_{\text{P-Rh}} = 123$  Hz), significantly upfield from the resonance for **14a**. This resonance is characteristic of a *trans*-phosphine environment about the  $\text{Rh}^I$  metal center and indicates a conversion from a condensed macrocyclic intermediate to the “open” macrocycle

(55) Formed by the reaction between  $[\text{Rh}(\text{NBD})\text{Cl}]_2$  and  $\text{AgBF}_4$  (NBD = norbornadiene).



**Table 1.** X-ray Crystallographic Data for **8a**⊂**DABCO** and **15a**⊂**DABCO**

	<b>8a</b> ⊂ <b>DABCO</b>	<b>15a</b> ⊂ <b>DABCO</b>
empirical formula	C <sub>181</sub> H <sub>166</sub> Cl <sub>4</sub> N <sub>12</sub> O <sub>8</sub> P <sub>4</sub> Rh <sub>2</sub> Zn <sub>2</sub>	C <sub>172</sub> H <sub>142</sub> Cl <sub>2</sub> N <sub>10</sub> O <sub>4</sub> P <sub>4</sub> Rh <sub>2</sub> S <sub>4</sub> Zn <sub>2</sub>
formula weight	3239.5	3072.54
temperature	153(2) K	100(2) K
wavelength	0.71073 Å	0.48595 Å
crystal system, space group	triclinic, <i>P</i> $\bar{1}$	monoclinic, <i>C</i> 2/ <i>c</i>
unit cell dimensions	<i>a</i> = 13.9395(13) Å $\alpha$ = 83.044(2)° <i>b</i> = 17.1623(17) Å $\beta$ = 73.368(2)° <i>c</i> = 20.894(2) Å $\gamma$ = 74.250(2)°	<i>a</i> = 17.0086(10) Å <i>b</i> = 44.661(3) Å $\beta$ = 104.252(2)° <i>c</i> = 23.0053(13) Å
volume	4604.5(8) Å <sup>3</sup>	16937.3(17) Å <sup>3</sup>
Z, calculated density	1, 1.168 Mg/m <sup>3</sup>	4, 1.205 Mg/m <sup>3</sup>
absorption coefficient	0.581 mm <sup>−1</sup>	0.342 mm <sup>−1</sup>
<i>F</i> (000)	1678	6336
crystal size	0.170 × 0.170 × 0.50 mm	0.150 × 0.025 × 0.010 mm
theta range for data collection	1.02 to 28.83°	0.90 to 15.70°
reflections collected/unique	42335/21291 [ <i>R</i> (int) = 0.0712]	123297/12229 [ <i>R</i> (int) = 0.0780]
absorption correction	integration	empirical
max. and min. transmission	0.9695 and 0.8265	1.000000 and 0.649500
refinement method	full-matrix least-squares on <i>F</i> <sup>2</sup>	full-matrix least-squares on <i>F</i> <sup>2</sup>
data/restraints/parameters	21291/0/915	12229/0/921
goodness-of-fit on <i>F</i> Δ <sup>2</sup>	0.912	1.063
final <i>R</i> indices [ <i>I</i> > 2σ( <i>I</i> )]	<i>R</i> 1 = 0.0910, <i>wR</i> 2 = 0.2301	<i>R</i> 1 = 0.0937, <i>wR</i> 2 = 0.2816
<i>R</i> indices (all data)	<i>R</i> 1 = 0.1924, <i>wR</i> 2 = 0.2591	<i>R</i> 1 = 0.1199, <i>wR</i> 2 = 0.3081
largest diff. peak and hole	1.378 and −0.747 e <sup>−</sup> /Å <sup>−3</sup>	2.236 and −0.800 e <sup>−</sup> /Å <sup>−3</sup>

**Figure 2.** Graphical representations of the X-ray crystal structure of **15a**⊂**DABCO** as viewed (A) from the side and (B) from the top containing a molecule of DABCO bridging both Zn atoms. Hydrogen atoms, disordered DABCO carbon atoms and solvent molecules have been omitted for clarity. Gray = Carbon, Pink = Rh, Red = O, Orange = S, Yellow = Cl, Green = P, Dark Blue = N, Light Blue = Zn.

**15a.**<sup>50</sup> The ESIMS spectrum of **15a** exhibits a peak at 1353.7 *m/z*, corresponding to the M<sup>2+</sup> ion without the Cl<sup>−</sup> and CO ligands and is consistent with ESIMS data for analogous structures.<sup>26</sup>

We observed condensed intermediate **14b** as the sole product from the reaction between thioether-based ligand **13** and [Cu(CH<sub>3</sub>CN)<sub>4</sub>][PF<sub>6</sub>]. The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of **14b** exhibits a singlet at δ 1.0 which is highly diagnostic of a tetrahedral environment at the Cu<sup>I</sup> metal center.<sup>51</sup> Additionally, ESIMS analysis yields a peak corresponding to the M<sup>2+</sup> ion at 1314.5


**Table 2.** Selected Distances for **8a**⊂**DABCO** and **15a**⊂**DABCO**

selected distance	<b>8a</b> ⊂ <b>DABCO</b> (Å)	<b>15a</b> ⊂ <b>DABCO</b> (Å)
Rh–Rh	24.8	22.5
Rh–P	2.32, 2.31	2.30
Zn–Zn	7.09	7.02


*m/z*, indicating formation of the desired condensed structure. Notably, compound **14b** can be converted into macrocycle **15b** upon the addition of stoichiometric quantities of pyridine. Once again, the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum allows us to monitor this transformation through the appearance of a singlet at δ −8.0, which has shifted upfield from the resonance observed for **14b**. The ESIMS analysis of **15b** yields a peak at 1314.9 *m/z* which corresponds to the pyridine-free molecular ion.

**X-ray Structure Determination of 15a**⊂**DABCO** and **15c**⊂**DABCO**. Similar to the problems encountered in crystallizing **8a**, our attempts to isolate X-ray quality single crystals of **14a** and **15a** proved unfruitful. Based upon the more promising strategy of adding DABCO to pure samples of **8a**, we hypothesized that crystals of **15a**⊂**DABCO** could be obtained in a similar fashion. X-ray quality crystals were obtained from vapor diffusion of pentane/diethyl ether (1:1 v/v) into a THF/CH<sub>2</sub>Cl<sub>2</sub> solution of **15a** containing DABCO, which was initially layered with acetonitrile. Similar to the analogous structure **8a**⊂**DABCO**, the 56-membered macrocyclic product **15a**⊂**DABCO** represents one of the largest macrocycles ever prepared via the Weak-Link Approach. In this structure, the Zn atoms are separated by a distance of distance of 7.02 Å and are bridged by a DABCO ligand (Figure 2). The structure of **15a**⊂**DABCO** is slightly puckered toward the Zn(porphyrin) centers, leading to a slightly staggered porphyrin–porphyrin geometry. As expected, the Rh–P and Zn–Zn distance in **8a**⊂**DABCO** are comparable to those found for **15a**⊂**DABCO** while the Rh–Rh distance differs significantly, presumably due to the presence of the four S atoms which have been incorporated in **15a**⊂**DABCO** (Table 2).

Contrary to **8b**, **15b** was quite soluble in pure THF; however, our efforts to crystallize **15b** from THF only yielded micro-



**Figure 3.** Acyl transfer reactions catalyzed by (left) a closed macrocycle vs (right) the corresponding open macrocycle. The open macrocycle can preorganize the substrates within the cavity, thereby increasing the rate of the reaction ( $k_2$ ) in comparison to that ( $k_1$ ) observed in the presence of the closed macrocycle.



**Figure 4.** Formation of the three *X*-(acetoxymethyl)pyridine ( $X = 2, 3$ , or  $4$ ) isomers by an acyl transfer reaction between 1-acetylimidazole and *X*-pyridylcarbinol, as catalyzed by Zn-porphyrin complexes **14a** and **15a**. Concentration vs time plots are shown for the formation of 4-(acetoxymethyl)pyridine (**A**, 4-AMP), 3-(acetoxymethyl)pyridine (**B**, 3-AMP), and 2-(acetoxymethyl)pyridine (**C**, 2-AMP). All data were corrected for background reactions (see Supporting Information). Conditions:  $\text{CH}_2\text{Cl}_2$ , rt, 9 mM *X*-pyridylcarbinol, 6 mM 1-acetylimidazole, 2.5 mM biphenyl (GC reference standard), and 0.3 mM supramolecular catalyst (**14a** and **15a**). CO (1 atm) and appropriate amounts of benzyltriethylammonium chloride when indicated.

crystalline powder. Attempts to crystallize **15b** using the same conditions as used for **8b** led to the formation of magenta single crystals of **15c**DABCO, which were confirmed by X-ray diffraction analysis at APS (see Supporting Information). Macrocycle **15c**DABCO encapsulates a bridging DABCO ligand between two Zn porphyrins, flanked by tetrahedral  $\text{CuCl}_2$  centers, again indicating the oxidation of  $\text{Cu}^{\text{I}}$  to  $\text{Cu}^{\text{II}}$  via solvent dehalogenation as has been observed previously. As for **8c**DABCO, isolation of **15c**DABCO strongly suggests an open macrocyclic structure for **15b** as shown in Scheme 3.

**Acyl Transfer Catalytic Experiments.** To demonstrate the ability of the  $\text{Zn}^{\text{II}}$ –porphyrin moieties in **14a** and **15a** to act cooperatively in an allosterically controlled fashion, we employed a catalytic acyl transfer reaction that has been shown by Sanders and co-workers to accelerate in the presence of trimetallic Lewis acidic porphyrin assemblies.<sup>56</sup> We hypothesized that the pyridylcarbinol and acetylimidazole substrates could be brought together within the cavity of **15a** by cofacial

$\text{Zn}^{\text{II}}$  metal centers and converted to the products in a catalytic fashion by virtue of their proximity (Figure 3). Furthermore, reactions involving 1-acetylimidazole and differentially substituted *X*-pyridylcarbinol (where  $X = 2, 3$ , or  $4$ ) can be used to evaluate the ability of **15a** to act allosterically: only the combination of substrates with the right distance can span the cavity and react at an accelerated rate.

The efficiency of both the “closed” (**14a**) and “open” (**15a**) supramolecular catalysts in the acyl transfer reaction were evaluated against a control reaction consisting of the monomer **Zn(TPP)** and analogous monomeric  $\text{Rh}^{\text{I}}$  complexes **16a** and **16b** (see Supporting Information). For 4-pyridylcarbinol (4-PC), both **14a** and **15a** significantly accelerate the reaction rate, with the open macrocycle **15a** being almost 14 times more active than the monomers and twice as fast as the closed macrocycle **14a** (Figure 4a). While **14a** is depicted as a rigid entity in Figure 3, its structure is probably dynamic when in solution and the observed catalytic activity may originate from the conforma-

tional flexibility around the S atoms and the low rotational barrier of the porphyrin ligand about the carbon-sulfur bond in the backbone of the ligand. For 3-pyridylcarbinol (3-PC), the background-corrected  $\text{rate}_{\text{macrocycle}}/\text{rate}_{\text{monomers}}$  ratio ( $\text{rate}_{\text{monomers}} = \text{rate of the reaction in the presence of the } [\text{Rh}^{\text{I}} \text{ monomer} + \text{Zn}(\text{TPP})] \text{ mixture}$ ) and the observed allosteric effect ( $\text{rate}_{15\text{a}}/\text{rate}_{14\text{a}}$ ) only drops slightly (Figure 4b), suggesting that the cavities of **14a** and **15a** are still flexible enough to accommodate the change in transition state distance for acyl transfer from acetylimidazole upon binding. For 2-pyridylcarbinol (2-PC), both the  $\text{rate}_{\text{macrocycle}}/\text{rate}_{\text{monomers}}$  ratio and the observed allosteric effect ( $\text{rate}_{15\text{a}}/\text{rate}_{14\text{a}}$ ) drop significantly with respect to 3- and 4-pyridylcarbinol and are similar to those observed for the monomers (Figure 4c). This may be explained by a simple geometry argument: if 2-pyridylcarbinol is bound to one of the Zn centers, the carbinol group will be pointed away from the imidazole *N*-acetyl group bound to the other side, resulting in an unfavorable transition state (in comparison to those for 4- and 3-pyridylcarbinol) for productive acyl transfer.

Our catalytic data offers strong evidence that  $\text{Cl}^-/\text{CO}$  can act as positive allosteric effectors in a  $\text{Rh}^{\text{I}}$ -based hemilabile supramolecular catalyst system, enhancing the efficiency of bimolecular acyl transfer reactions when the geometry is optimized. Significantly, these effectors operate cooperatively with each other and are incapable of effecting shape change if added independently to the  $\text{Rh}^{\text{I}}$  species. That 2-pyridylcarbinol can be selectively discriminated against 3- and 4-pyridylcarbinol provides an impetus for employing these supramolecular assemblies for chemical sensing and shape-selective recognition when coupled to catalytic processes that provide signal amplification.

## Conclusion

In conclusion, we have developed a coordination chemistry-based synthetic approach for the quantitative preparation of flexible cofacial porphyrin assemblies in which the porphyrins act as functional sites within an allosteric framework that is tunable via modulation of peripheral structure control domains.

(56) Mackay, L. G.; Wylie, R. S.; Sanders, J. K. M. *J. Am. Chem. Soc.* **1994**, *116*, 3141–3142.

Importantly, these architectures possess cavities whose sizes can be directly controlled *in situ* via the introduction of simple, cooperative ancillary ligands that bond to the structure control domains. This capability enables the cofacial porphyrin structures to act as allosteric catalysts capable of discriminating different substrate combinations and selectively transforming them into the desired products, two key steps in developing new biomimetic supramolecular systems. Most notably, our synthetic approach should allow for facile access to a range of systems with tunable cofacial porphyrin–porphyrin distances that are useful in studying distance-dependent electron-transfer phenomena, molecular switches, host–guest interactions, and catalysis. Efforts toward elaboration of these areas are currently underway.

**Acknowledgment.** C.A.M acknowledges the NSF and ARO for support of this research and is grateful for a NIH Director's Pioneer Award. This work was also supported, in part, by the NSF-NSEC program under NSF Award Number EEC-0118025. Portions of this work were performed at the DuPont-Northwestern-Dow Collaborative Access Team (DND-CAT) Synchrotron Research Center located at Sector 5 of the Advanced Photon Source. DND-CAT is supported by the E.I. DuPont de Nemours & Co., the Dow Chemical Company, the U.S. National Science Foundation through Grant DMR-9304725 and the State of Illinois through the Department of Commerce and the Board of Higher Education Grant IBHE HECA NWU 96. ChemMat-CARS Sector 15 is principally supported by the National Science Foundation/Department of Energy under grant number CHE0087817 and by the Illinois Board of Higher Education. The Advanced Photon Source is supported by the U.S. Department of Energy, Basic Energy Sciences, Office of Science, under Contract No. W-31-109-Eng-38.

**Supporting Information Available:** Crystallographic data, X-ray crystal structures of **8c**⊂**DABCO** and **15c**⊂**DABCO**, and uncorrected and background reactions for 2-PC, 3-PC, and 4-PC. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA0661010