The 'one-pot' syntheses of α, α' -diphosphino-substituted imines: a unique reaction of bulky bis(dialkylamino)chlorophosphines

Antoine Baceiredo,^{*a*} Guy Bertrand,^{*a*} Philip W. Dyer,^{*b} John Fawcett,^{*b*} Nina Griep-Raming,^{*b*} Olivier Guerret,^{*a*} Martin J. Hanton,^{*b*} David R. Russell^{*b*} and Anna-Maria Williamson^{*c*}

- ^a Université Paul Sabatier, Laboratoire d'Hétérochimie Fondamentale et Appliquée, ESA 5069–Bât. 2R1, 118 route de Narbonne, 31062 Toulouse cedex, France
- ^b Chemistry Department, University of Leicester, University Road, Leicester, UK LE1 7RH. E-mail: p.dyer@le.ac.uk
- ^c Chemistry Department, Imperial College of Science, Technology and Medicine, South Kensington, London, UK SW7 2AY

Received (in Cambridge, UK) 7th December 2000, Accepted 22nd January 2001 First published as an Advance Article on the web 14th March 2001

The reaction between various bis(dialkylamino)chlorophosphines and N-isopropyl-substituted lithium amides afforded bis(α, α' -phosphino)imines in a 'one-pot' procedure. These compounds react with sulfur to generate intramolecularly hydrogen-bonded ylide derivatives. The molecular structure of $(Pr_2^iN)_2PCH_2C(=NBu^i)CH_2P(NPr^i)_2$ has been determined by X-ray crystallography.

Introduction

Currently there is a considerable resurgence of interest in the use of heteroatom substituents at phosphorus to 'tune' both the steric requirements and basicity of phosphine-containing ligands. This is due, in part, to the ease and considerable scope associated with the synthesis of phosphorus-heteroatom bonds, exemplified by P–O and P–N bond formation.¹ It is therefore somewhat surprising that despite both the continued use of bidentate P,P'-ligands in many industrially important catalytic processes,² and the on-going development of so-called 'hemi-labile' phosphorus-containing systems (*e.g.* bidentate $P,O^{-3} P,N^{-4}$ and P,S-ligands⁵),⁶ there remain few examples of chelating ligands which possess a diaminophosphine component.^{1a,c}

As the application of phosphorus-based donors in catalysis continues apace, simple, high-yielding preparative routes to novel, easily tailored ligand systems are of growing commercial importance. During studies into the use of bis(dialkylamino)phosphino substituents, the simple 'one-pot' synthesis of bis(α,α' -phosphino)-substituted imines has been studied in detail, while the basis of a more general route to compounds containing the R₂PCH₂C=NR' motif has been established.

Results and discussion

Treating a solution of LDA (lithium diisopropylamide) with a solution of bis(dialkylamino)chlorophosphines 1 (Et₂O, 0 °C) in a 1 : 1 molar ratio cleanly affords bis(α, α' -phosphino)imines 2 and the corresponding secondary phosphines 3^{7} in equimolar amounts [³¹P NMR spectroscopy] (Scheme 1, Table 1).8 The addition of less than stoichiometric amounts of LDA leads to the formation of mixtures of 2, 3 and unchanged 1. Similarly, the reaction of the phosphenium salt [(Pr₂ⁱN)₂P][CF₃SO₃] 5 with LDA (Et₂O) also forms 2a in good yield (73%) with concomitant formation of secondary phosphine 3a and elimination of lithium triflate. It is noteworthy that although a number of phosphorus-containing products were generated from the reaction between 5 and $HNPr_2^i$ in the presence of NEt_3 or 1,4-diazabicyclo[2.2.2] octane (dabco), none was readily identifiable. This suggests that the formation of 2 does not necessitate the generation of an intermediate $[(Pr_2^iN)_2P]^+$ moiety, but favours a concerted mechanism as outlined below.

It seems most likely that the imine $Me_2C=NPr^i$ 4 is generated during the reaction between 1 and LDA via an Nanalogue of the Meerwein–Ponndorf–Verley reaction (Scheme 2), similar reactions having been reported previously for

 Table 1
 Selected ³¹P NMR and IR spectroscopic data

Product	R ¹	R ²	R ³	$\delta^{31} P ({}^{4} I_{} / H_{7})^{a}$	Vield ^b (%)	IR¢
Tioduct	ĸ	K	K	0 I (9 pp/112)	Tield (70)	ш
2a	NPr ⁱ ₂	NPr ⁱ ₂	$\mathbf{Pr^{i}}$	45.4, 52.9 (11.0)	80	1625
2b	NCy ₂	NCy ₂	Pr ⁱ	53.9, 59.3 (25.0)	75	1621
2c	NPr ⁱ ₂	NCy ₂	Pr ⁱ	49.6, 56.6 $(13.0)^d$	е	1625 ^f
	-			49.8, 56.1 $(20.0)^d$		
2d	NPr ⁱ ₂	NPr ⁱ ₂	Су	46.5, 53.5 (12.0)	70	1619
2e	$NPr_2^{\overline{i}}$	$NPr_2^{\overline{i}}$	Bu ^t	47.6, 55.8 (37.0)	83	1629

^{*a*} In CDCl₃, 300 MHz. ^{*b*} Isolated yield after work-up. ^{*c*} Solution cell, KBr windows, CH_2Cl_2 , cm⁻¹. ^{*d*} In Et₂O. ^{*e*} Product isolated as a mixture of two inseparable diastereomers, formed quantitatively according to ³¹P NMR spectroscopy. ^{*f*} In Et₂O, severely broadened.



Scheme 1 Reagents and conditions: i, 0 °C to r.t., Et₂O, 12 h; ii, $R^1 = R^2 = NPr_2^i$, 0 °C to r.t., Et₂O, 12 h; iii, $R^1 = R^2 = NPr_2^i$, BuⁿLi/tmeda, -78 °C, 40 min, hexane; iv, 1a -78 °C to r.t., hexane. OTf = O₃SCF₃ (triflate).

certain transformations involving this amide base.⁹ It would appear that imine **4** is then rapidly deprotonated by LDA to generate an intermediate aza-allyl anion, which subsequently undergoes substitution by chlorophosphines **1**, a process that is repeated to form **2**.

Low temperature ³¹P NMR spectroscopic studies (Et₂O) of the reaction between LDA and compound **1a** indicate that the secondary phosphine **3a** and phosphino-imine **2a** are formed simultaneously in a 1 : 1 molar ratio as the only phosphoruscontaining products, on the NMR timescale at 243 K (no reaction is observed below this temperature). There was no evidence to support the claim that $(Pr_2^iN)_3P 6$ is involved in the formation of **2a** from LDA and **1a**, despite a previous report to the contrary.^{8†}

Attempts to identify compound 4 directly during reactions of 1a and LDA by GC-MS were unsuccessful. However, its intermediacy is supported by the observation that 2a can also be synthesized (30% yield) by treating an authentic sample of 4 with two equivalents of BuⁿLi/tmeda (hexane, -78 °C)



[†] It is important to note the coincidence between the reported ³¹P NMR chemical shift for $(Pr_2^iN)_3P$ 6, δ +133.6 (solvent not specified),^{8,10} and that for its most convenient precursor, namely **1a** {³¹P NMR (Et₂O, 273 K): δ +133.6}, suggesting a previously incorrect assignment of the former.

followed by the addition of two equivalents of diaminochlorophosphine **1a** (hexane, 0°C) [Scheme 1]. An extension to this methodology can be used to prepare variously substituted α -phosphino-imines, *e.g.* 7, from R(Me)C=NR' and the appropriate chlorophosphine (Scheme 3).¹¹ A variation on this procedure has recently been utilised for the synthesis of unsymmetrical β -diimines.¹²

It has proved straightforward to introduce other alkyl substituents at the imine terminus of diphosphines 2 using the 'one-pot' methodology. For example, treatment of diaminochlorophosphine 1a with unsymmetrical lithium *N*alkyl-*N*-isopropyl amides [LiNPrⁱ(R³)] (Et₂O, 0 °C) cleanly affords the corresponding bis(phosphino)imines 2d,2e (Scheme 4, Table 1). In each case, the 'sacrificial' role of one *N*-isopropyl group is readily apparent, further supporting the intermediacy of an imine such as Me₂C=NR (*e.g.* R = Prⁱ, 4). These ideas are endorsed by the observation that the reaction of LiN(Prⁱ)(Me) with 1a affords H₂C=NPrⁱ (GC-MS, IR v(CN) 1649 cm⁻¹)¹³ and (Prⁱ₂N)₂PH (GC-MS/³¹P NMR)⁷ rather than {(Prⁱ₂N)₂PCH₂}₂C=NMe, as a result of hydride abstraction from the more hydridic NCH₃ group.

Overall, it would appear that the initial hydride transfer between compound 1a and LDA outlined in Scheme 2 occurs because simple substitution, which would lead to formation of $(Pr_2^iN)_3P 6$, is blocked due to unfavourable steric interactions between the bulky amide substituents. Despite an unverified report of the ³¹P NMR spectrum of 6 (*vide supra*) and two brief accounts of its generation, little convincing data have been reported which support the existence of what would be an extremely sterically encumbered tris(amino)phosphine.^{8,14}

Notably, phosphino-imine formation only occurs when bulky dialkylamino substituents (e.g. NPr₂ⁱ or NCy₂) are present on the chlorophosphines **1**. The less bulky derivatives, $(R'_2N)_2PCl$ (R' = Me or Et), react with LDA to afford intractable mixtures of phosphorus-containing products, which include trace amounts of the corresponding tris(amino)phosphines, $(R'_2N)_2PNPr_2^i$, according to ³¹P NMR spectroscopy, but show no evidence of imine formation.¹⁵ Similarly, the direct reaction between Ph₂PCl and LDA, irrespective of the nature of the solvent, affords a mixture of the product from reductive coupling, Ph₂PPPh₂ ($\delta - 14.2$),¹⁶ and



Scheme 3 Reagents and conditions: i, BuⁿLi/tmeda, -78 °C to r.t., hexane, 12 h (R = H, Me or Ph; R' = Ph, C₆H₃Pr₂ⁱ-2,6); ii, R''_PCl (R'' = NPr₂ⁱ or Ph), 0 °C, reflux, 3 h.



the aminophosphine $Ph_2P(NPr_2^i)$ (δ +37.1),¹⁷ resulting from substitution, in an approximately 1 : 2 ratio (Scheme 4).[‡]

It is noteworthy that when the reaction between compound 1a and LDA is undertaken in THF instead of Et₂O the major product becomes $(Pr_2^iN)_2P-P(NPr_2^i)_2$,¹⁸ with only trace amounts of 2 being detectable (<5% by ³¹P NMR spectroscopy). This solvent dependency is presumed to reflect small changes in the degree of chloride dissociation from 1a that occurs in each of the solvents.^{19,20,§} The formation of products resulting from reductive coupling is perhaps not surprising since LDA is known to act as a one-electron donor to species with favourable reduction potentials.²¹ However, as expected neither nucleophilic substitution nor imine formation was observed to occur when the corresponding $\sigma^4 \lambda^5$ thioxophosphine (Pr₂ⁱN)₂P(S)Cl was treated with LDA in Et₂O, presumably due to a combination of steric hindrance and the lack of a vacant orbital at phosphorus to accommodate initial hydride transfer (as required by the mechanism outlined in Scheme 2).

All the phosphino-imine derivatives 2 (with the exception of 2c) are readily isolated after appropriate work-up as colourless solids. Crystals of 2e suitable for an X-ray crystallographic study were grown from a concentrated hexane solution at -30 °C.¶ The structural parameters associated with the planar imine moiety (sum of angles about $C(2) = 360.0^{\circ}$) are typical for this type of functional group (Fig. 1).²² The phosphorus atom P(1) is orientated such that its lone pair is directed away from both the lone pair on the imine nitrogen N(1) and the second phosphorus P(2). This induces a staggered conformation along the PCCCP backbone such that P(2) and P(1) lie, respectively, 1.5447 Å above and -0.8824 Å below the plane defined by [N(1)-C(3)-C(2)-C(1)]. Changes in the extent of this 'twist' are believed to be the cause of the considerable variation in the ${}^{4}J_{PP}$ coupling constants observed (2a-2e, Table 1), although no significant difference in this parameter was observed for 2a at either reduced or elevated temperatures. The bond distances P(2)-C(3) and P(1)-C(1) are unremarkable and consistent with P-C single bonds.^{1f} As is observed for many aminophosphines, the four P-N bond distances are significantly shorter than that associated with a true P-N single bond, while the sums of angles about nitrogen atoms N(2) to N(5) [358.4, 359.9, 360.0, 360.0°, respectively] indicate significant sp² character.²³

Diphosphino-imine **2a** readily reacts with sulfur, initially forming the corresponding $bis(\sigma^4\lambda^5$ -thiophosphoryl)imine **8** [AB system: δ +70.9 and +72.6 (⁴J_{PP} 10.0 Hz)] (Scheme 5) which rearranges on standing (~12 hours, 25 °C) to afford the new hydrogen-bonded ylide tautomer **9**. This can be isolated after recrystallisation from dichloromethane (-30 °C) as an almost colourless solid in 61% yield. Its identity is clearly apparent from NMR spectroscopy with characteristic signals observed for the C=N, P=CH (confirmed from J-modulated



Fig. 1 Molecular structure of compound 2e. Displacement ellipsoids are drawn at the 30% probability level. Selected bond distances (Å) and angles (°): P(1)–N(2) 1.698(4), P(1)–N(3) 1.689(4), P(2)–N(4) 1.699(4), P(2)–N(5) 1.704(4), N(1)–C(2) 1.274(5), C(1)–P(1) 1.871(4), C(1)–C(2) 1.530(6), C(2)–C(3) 1.498(6), and C(3)–P(2) 1.861(5); C(2)–N(1)–C(4) 127.9(4), N(1)–C(2)–C(3) 116.5(4), N(1)–C(2)–C(1) 127.2(4), C(1)–P(1)–N(2) 100.03(19), C(1)–P(1)–N(3) 102.04(19), N(2)–P(1)–N(3) 109.1(2), C(3)–P(2)–N(4) 100.05(19), C(3)–P(2)–N(5) 102.4(2), N(4)–P(2)–N(5) 110.2(2).

¹³C NMR spectra) and N–H fragments. Unlike previous examples of this type of tautomerisation,²⁴ structure **9** is believed to be favoured over the C–H acid form as a result of the formation of an intramolecular S–H–N hydrogen bond, affording a six-membered ring system. The stability of this arrangement is demonstrated by the lack of reactivity of **9** towards strong bases {NaN(SiMe₃)₂ in C₆D₆ or LDA · tmeda in Et₂O} and to transition metals {Mo(CO)₄(pip)₂ (pip = piperidine) or RhCl(PPh₃)₃}.

Conclusion

It has been shown that sterically encumbered bis(dialkylamino)chlorophosphines exhibit unusual, yet understandable reactivity towards isopropyl-substituted lithium amides undergoing hydride transfer rather than nucleophilic substitution. This observation has led to the synthesis of a family of bulky phosphino-imines 2 in good yields. Compounds 2 make interesting bulky ligands for transition metals. They present both hard imino- and soft phosphino-donor sites making these species potentially 'hemi-labile'. Furthermore, the pres-



Scheme 5 Reagents and conditions: i, $S_8,\ CH_2Cl_2,\ r.t.,\ 1$ h; ii, $CH_2Cl_2,\ r.t.,\ 12$ h.

[‡] These two phosphines were isolated as their sulfides following treatment of the reaction mixture with an excess of sulfur and subsequent separation and purification by flash column chromatography on silica gel using toluene as eluent.

gel using toluene as eluent. § The ³¹P NMR chemical shift for chlorophosphine **1a** exhibits a significant solvent dependency (162.0 MHz, 300 K): δ +133.6 (Et₂O), +135.8 (THF) and +140.0 (CDCl₃).

[¶] Crystal data for compound **2e**: $C_{31}H_{69}N_5P_2$, M = 573.85, orthorhombic, space group *Pbca*, a = 17.676(6), b = 18.784(8), c = 22.185(11) Å, V = 7366(5) Å³, Z = 8, μ (Mo-K α) = 0.143 mm⁻¹, T = 200(2) K; clear colourless plate, Siemens P4 diffractometer, ω scans, 6459 measured, 6032 independent reflections ($R_{int} = 0.0304$). The structure was solved by direct methods and the non-hydrogen atoms were refined anisotropically using full matrix least squares based on F^2 to give R1 = 0.0776, wR2 = 0.2000 (all data) for 3052 independent observed reflections [$I > 2\sigma(I)$] and 343 parameters. CCDC reference number 156002. See http://www.rsc.org/suppdata/nj/b0/b0098671/ for crystal-lographic data in CIF or other electronic format.

ence of σ -electron withdrawing and π -electron donating substituents at phosphorus could have significant implications in catalysis.¹⁹ A study of the coordination chemistry of ligands **2** has been initiated and preliminary results will be reported elsewhere;²⁵ their coordination geometry should contrast with the many examples of related *ortho*-phenylene-bridged P–N ligands that have been reported.⁴

Experimental

All manipulations were performed under an atmosphere of dry argon or nitrogen. Solvents were distilled from the appropriate drying agent and stored under argon or nitrogen prior to use. ¹H, ¹³C and ³¹P NMR spectra were recorded on Bruker AC200, WM250, ARX 300 or JEOL 270 spectrometers at ambient probe temperatures. C₆D₆ and CDCl₃ were distilled from P_2O_5 and degassed prior to use. $^1\dot{H}$ and ^{13}C chemical shifts in ppm were referenced using the partially deuteriated solvent as internal reference. ³¹P downfield shifts are expressed with a positive sign, in ppm, relative to external 85% H₃PO₄; where appropriate a sealed capillary tube containing C₆D₆-PPh₃ or C₆D₅CD₃-PPh₃ was placed inside a normal 5 mm NMR tube to provide a deuterium lock and an internal reference. Coupling constants and linewidths are reported in Hertz. Infrared spectra were obtained in a solution cell with KBr windows on either a Perkin-Elmer FT-IR1725X or Mattson Research Series 1 spectrometer, mass spectra on a Ribermag R10 10E, Kratos Concept, or AutospecQ instrument. GC-MS were performed using a Perkin-Elmer Turbomass spectrometer/Perkin-Elmer Autosystem XL gas chromatograph. Satisfactory elemental analyses have been obtained for all isolated products. Melting points are uncorrected.

1a,²⁶ 1c,²⁷ 4,²⁸ and 5^{29} were prepared according to literature procedures, or minor modifications thereof. BuⁿLi (1.6 M, hexanes) was obtained from Aldrich. Amines (Aldrich and Lancaster) used in the preparation of the lithium amides were pre-dried over KOH pellets and subsequently stored under nitrogen over activated 4 Å molecular sieves.

Preparations

Lithium amides. These reagents were prepared by treating a solution of the appropriate amine in diethyl ether with a stoichiometric amount of BuⁿLi at -78 °C and used without further purification.

($Pr_2^iN)_2P(S)Cl.$ A Schlenk was charged with compound 1a (1.0 g, 3.7 mmol) and CH_2Cl_2 (10 cm³). An excess of sulfur (0.15 g, 4.6 mmol) was added slowly as a solid under a flow of nitrogen. The solution was stirred at r.t. for 48 h. Volatile components were removed under reduced pressure and the product extracted from any unchanged sulfur using pentane (3 × 10 cm³). Removal of pentane *in vacuo* afforded ($Pr_2^iN)_2P(S)Cl$ as a white solid which was sufficiently pure for further use (0.99 g, 98%). ³¹P NMR (109.4 MHz, CH_2Cl_2): δ + 68.7 (pent., ³J_{PH} 22.0 Hz).

 $(Pr_2^iN)_2PCH_2C(=NPr^i)CH_2P(NPr_2^i)_2$ 2a. A suspension of compound 1a (10 g, 37.0 mmol) in ether (20 cm³) was added to a cold (0 °C) solution of LDA (prepared as above). The mixture was allowed to warm to room temperature, where it became progressively pale yellow and the formation of a white precipitate occurred. Subsequently, the reaction was stirred for 12 h at room temperature. Removal of volatiles *in vacuo* afforded an orange solid. Extraction with pentane (3 × 20 cm³) afforded a clear pale yellow solution. Concentration and recrystallisation from the pentane solution at -30 °C afforded colourless crystals of 2a (5.52 g, 80%), mp 130–132 °C (Found: C, 62.1; H, 12.5; N, 12.1. C₃₀H₆₇N₅P₂ requires C, 64.3; H,

12.1; N, 12.5%); ¹H NMR (270.0 MHz; C₆D₆) δ 3.98 (1 H, sept., ³J_{HH} 6.0, CNCH(CH₃)₂), 3.35 (8 H, sept. d, ³J_{HH} 7.0, ³J_{PH} 12.0, PNCH), 3.10 (2 H, s, CH₂), 3.08 (2 H, s, CH₂), 1.37 (6 H, d, ³J_{HH} 6.0, C=NCH(CH₃)₂), 1.26 (24 H, dd, ³J_{HH} 7.0, ⁴J_{PH} 12.0, PNCH(CH₃)₂) and 1.19 (24 H, d, ³J_{HH} 7.0 Hz, PNCH(CH₃)₂); ¹³C NMR (62.9 MHz; CDCl₃) δ 166.0 (dd, ²J_{PC} 15.0 and 9.5, C=N), 50.2 (s, C=NCH), 46.6 (d, ²J_{PC} 11.0, PNCH), 46.5 (d, ²J_{PC} 11.0, PNCH), 44.4 (dd, ¹J_{PC} 12.0, ³J_{PC} 6.0, CH₂), 33.7 (dd, ¹J_{PC} 19.0, ³J_{PC} 8.0, CH₂), 24.3 (br d, ³J_{PC} 6.0, PNCH(CH₃)₂), 24.0 (br d, ³J_{PC} 6.0, PNCH(CH₃)₂), 23.7 (d, ³J_{PC} 7.0 Hz, PNCH(CH₃)₂) and 23.6 (s, C=NCHMe₂); *m*/z (EI) 245 (9.3%, [Prⁱ₂N]₂PCH₂).

From imine 4. BuⁿLi (23.0 cm³, 1.6 M, 36.6 mmol) was added dropwise to a cooled (-78 °C) solution of imine 4 (2.05 g 18.3 mmol) and tmeda (5.6 cm³, 37.0 mmol) in hexane (20 cm³) and subsequently stirred at this temperature for 40 min. A suspension of compound 1a (9.76 g, 36.6 mmol) in hexane (50 cm³) was then added dropwise at -78 °C over a period of 5 min. The mixture was stirred at room temperature overnight, the volatiles were removed under reduced pressure and the product was extracted with hexane (3 × 10 cm³). Prolonged cooling of a concentrated pentane solution (-30 °C) afforded 2a as a colourless solid (3.0 g, 30%).

From compound 5. An ether solution (10 cm^3) of compound 5 (0.25 g, 0.66 mmol) was treated with a solution of LDA (0.66 mmol) in ether (10 cm³) at 0°C, allowed to warm to room temperature and stirred overnight. Subsequent removal of solvent *in vacuo* and extraction of the product with hexane $(3 \times 2 \text{ cm}^3)$ followed by removal of solvent afforded **2a** as a white solid (0.09 g, 73% yield).

 $(Cy_2N)_2PCH_2C(=NPr^i)CH_2P(NCy_2)_2$ (2b). An analogous procedure to that used for the preparation of compound 2a was employed using 1b (0.5 g, 1.17 mmol) suspended in ether (30 cm³) and LDA (1.17 mmol) in ether (10 cm³). After extraction and removal of volatile components *in vacuo*, diphosphine-imine 2b was isolated as an off-white solid after washing with CH₃CN (3 × 10 cm³) (0.26 g, 75%), mp 185– 188 °C (Found: C, 73.1; H, 11.0; N, 7.9. C₅₄H₉₉N₅P₂ requires C, 74.0; H, 11.4; N, 8.0%); *m/z* (FAB+) 877 ([M – H]⁺).

Reaction of $(Pr_2^iN)(Cy_2N)PCl$ **1c with LDA.** A similar experimental procedure to that used for the preparation of compound **2a** was employed, using $(Pr_2^iN)(Cy_2N)PCl$ **1c** (5.91 g, 17.04 mmol) in ether (20 cm³) and LDA (17.04 mmol) in ether (50 cm³) at 0 °C. After stirring overnight at r.t. the solvent was removed *in vacuo* and the product extracted off the inorganic salts using pentane (3 × 20 cm³). On removal of the solvent under reduced pressure an intractable mixture of diastereomeric products (**2c**) was isolated as a waxy orange solid. Direct attempts at purification *via* column chromatography (silica) or thiolation with S₈ followed by chromatography (silica) led to decomposition, while recrystallisation from a variety of solvents was unsuccessful.

(Pr¹₂N)₂PCH₂C(=NCy)CH₂P(NPr¹)₂ 2d. An identical experimental procedure to that used for the preparation of compound 2a was employed. A suspension of 1a (2.0 g, 7.5 mmol) in ether (30 cm³) was added to a cold (0 °C) solution of LiN(Pr¹)(Cy) [prepared as above using BuⁿLi (1.6 M, 7.5 mmol, 4.7 cm³) and HN(Pr¹)(Cy) (7.5 mmol, 1.2 cm³)]. After stirring at r.t. for 12 h, removal of all volatile components under vacuum followed by extraction with hexane (20 cm³) and its subsequent removal *in vacuo* gave a viscous yellow oil. Washing the oil with CH₃CN (3 × 10 cm³) followed by drying *in vacuo* afforded 2d as a white solid (3.2 g, 70%), mp 114–118 °C (Found: C, 65.0; H, 12.1; N, 11.2. C₃₃H₇₁N₅P₂ requires C, 66.1; H, 11.9; N, 11.7%); ¹H NMR (250.1 MHz; CDCl₃) δ 3.41 (9 H, m, C=NCH and PNCH), 2.92 (2 H, s,

PCH₂), 1.74 (2 H, br s, PCH₂), 1.45–1.08 (58 H, m, CH₂ and CH₃); ¹³C NMR (62.9 MHz; CDCl₃) δ 166.4 (dd, ²J_{PC} 15.0 and 9.0, C=N), 59.8 (d, ⁴J_{PC} 4.0, C=NCH), 47.0 (d, ²J_{PC} 11.0, PNCH), 46. 8 (d, ²J_{PC} 11.0, PNCH), 45.0 (dd, ¹J_{PC} 12.0, ³J_{PC} 5.0, CH₂), 34.5 (s, CH₂), 34.3 (dd, ¹J_{PC} 20.0, ³J_{PC} 8.0, CH₂), 26.1 (s, CH₂), 25.5 (s, CH₂), 24.7 (pseudo t, ³J_{PC} 7.0, PNCH(CH₃)₂), 24.4 (d, ³J_{PC} 7.0, PNCH(CH₃)₂) and 24.1 (d, ³J_{PC} 7.0 Hz, PNCH(CH₃)₂).

(Pr₂ⁱN)₂PCH₂C(=NBu^t)CH₂P(NPrⁱ)₂ 2e. An identical experimental procedure to that used for the preparation of compound 2a was employed. A suspension of 1a (2 g, 7.5 mmol) in ether (10 cm³) was added to a cold (0 $^{\circ}$ C) solution of LiN(Prⁱ)(Bu^t) [prepared as above using BuⁿLi (1.6 M, 7.5 mmol, 4.7 cm³) and HN(Prⁱ)(Bu^t) (7.5 mmol, 1.2 cm³)]. Recrystallisation from pentane at -30 °C afforded a first batch of colourless crystals of 2e (1.0 g); subsequent concentration and cooling to -30 °C gave a further 0.2 g (total yield 1.2 g, 83%), mp 142–144 °C (Found: C, 64.8; H, 12.0; N, 12.1. $C_{31}H_{69}N_5P_2$ requires C, 65.0; H, 12.2; N, 12.2%); ¹H NMR (270.0 MHz; C₆D₆) δ 3.36 (8 H, m, PNCH), 3.14 (2 H, s, CH₂), 3.07 (2 H, s, CH₂), 1.54 (9 H, s, CNC(CH₃)₃), 1.27 (24 H, dd, ${}^{3}J_{\text{HH}}$ 7.0, ${}^{3}J_{\text{PH}}$ 17.0, PCH(CH₃)₂) and 1.21 (24 H, dd, ${}^{3}J_{\text{HH}}$ 7.0, ${}^{4}J_{\text{PH}}$ 3.2 Hz, PNCH(CH₃)₂); 13 C NMR (100.6 MHz; C₆D₆) δ 163.1 (dd, ²J_{PC} 17.0 and 10.0, C=N), 55.6 (s, C=NC), 47.1 (m, PNCH), 45.9 (dd, ${}^{1}J_{PC}$ 14.0, ${}^{3}J_{PC}$ 6.0, CH₂), 37.5 (dd, ${}^{1}J_{PC}$ 21.0, ${}^{3}J_{PC}$ 4.5, CH₂), 31.7 (s, C(CH₃)₃), 24.8 (d, ${}^{3}J_{PC}$ 8.0, PNCH(CH₃)₂), 24.8 (d, ${}^{3}J_{PC}$ 8.0, PNCH(CH₃)₂), 24.6 (d, ${}^{3}J_{PC}$ 6.0, PNCH($(CH_3)_2$) and 24.0 (d, ${}^{3}J_{PC}$ 6.0 Hz, PNCH($(CH_3)_2$); m/z (EI) 573 (M⁺).

Reaction of Ph₂PCl with LDA. To a cold (0 °C) ether solution (20 cm³) of LDA (5.6 mmol) was added a cold ether solution (10 cm³) of Ph₂PCl (1.0 cm³, 5.6 mmol). The mixture was allowed to warm to room temperature and stirred overnight. Volatiles were removed *in vacuo* and the mixture extracted with pentane (3 × 10 cm³) that was subsequently removed under reduced pressure. Ether (10 cm³) and an excess of sulfur (0.25 g) were added before stirring overnight. Volatiles were removed *in vacuo* and the mixture was separated by column chromatography on silica gel using toluene as eluent to afford Ph₂P(S)NPrⁱ₂ (R_f 0.5; 0.88 g, 52%) and Ph₂P(S)=P(S)Ph₂ (R_f 0.3; 0.61 g, 25%).

Reaction of (Pr_2^iN)_2PCl with LiN(Me)(Prⁱ). A suspension of compound 1a (2.0 g, 7.5 mmol) in ether (10 cm³) was added to an ether solution (10 cm³) of LiN(Me)(Prⁱ) (7.5 mmol) at 0 °C. After stirring at room temperature overnight a ³¹P NMR spectrum of the mixture showed a single resonance corresponding to $(Pr_2^iN)_2PH$ (³¹P: δ +42.1{¹J_{PH} 254.0 Hz}). All volatile components were isolated by vacuum distillation and subjected to analysis by GC-MS. *m*/z (EI) 71 (M⁺) and 132 (M⁺ - NPr_2).

 $(Pr_2^iN)_2PCH_2C(=NC_6H_3Pr_2^i-2,6)CH_3$ 7. A solution of BuⁿLi (2.4 mmol) was added dropwise from a syringe to a cold $(-78 \,^{\circ}\text{C})$ solution of Me₂C=N(C₆H₃Pr₂ⁱ-2,6) (2.4 mmol, 0.52 g) and tmeda (2.5 mmol, 0.4 cm³) in hexane (10 cm³) to afford a clear solution and a bright yellow precipitate which was stirred at r.t. for 16 h. A solution of compound 1a (2.4 mmol, 0.64 g) in hexane was then added slowly at 0 °C. On warming to r.t. the mixture was heated to reflux for 3 h. The resulting clear yellow solution was isolated by filtration. The white solid was washed with hexane $(3 \times 5 \text{ cm}^3)$ and the washings were combined. A dark yellow viscous oil was obtained on removing all volatiles in vacuo. Recrystallisation from the minimum of hot acetonitrile afforded colourless crystals of 7 (0.49 g); subsequent concentration and cooling to -30 °C afforded a further 0.4 g, a total yield of 0.89 g, 83% (Found: C, 72.0; H, 11.0; N, 9.8. C₂₇H₅₀N₃P requires C, 72.4; H, 11.3; N, 9.4%); v_{max} (KBr, CH₂Cl₂)/cm⁻¹ 1635 (C=N); ¹H NMR (301.5 MHz, CDCl₃) δ 7.00 (3H, m, C₆H₃), 3.46 (4 H, d. sept., ${}^{3}J_{HH}$ 7.0, ${}^{3}J_{PH}$ 11.0, PNCH), 2.99 (2 H, s, PCH₂), 2.79 (2 H, sept., ${}^{3}J_{HH}$ 7.0, CCHMe₂), 1.75 (3 H, s, CH₃), 1.14 (24 H, dd, ${}^{3}J_{PH}$ 7.0, ${}^{4}J_{PH}$ 17.0, PNCH(CH₃)₂) and 1.04 (12 H, d, ${}^{3}J_{HH}$ 7.0 Hz, CCHMe₂); 13 C NMR (75.8 MHz, CDCl₃) δ 171.0 (s, C=N), 146.9 (s, C_{ipso} of C₆H₃), 137.2 (s, o-C of C₆H₃), 123.3 (s, p-C of C₆H₃), 123.3 (s, m-C of C₆H₃), 47.6 (d, ${}^{2}J_{PC}$ 11.0, PNCH), 45.6 (d, ${}^{1}J_{PC}$ 15.5, CH₂), 28.0 (s, CCHMe₂), 24.5 (d, ${}^{3}J_{PC}$ 7.0, PNCMe₂), 24.2 (s, CCHMe₂), 23.8 (s, CCHMe₂) and 21.9 (d, ${}^{3}J_{PC}$ 10.0 Hz, CH₃); 31 P NMR (122.1 MHz): δ +47.5 (s).

1,3-Bis[**bis**(**diisopropylamino**)**thiophosphoryl**]-2-**isopropyliminopropane 8.** To a solution of compound 2a (0.10 g, 0.18 mmol) in CDCl₃ (0.5 cm³) was added an excess of sulfur (13.0 mg, 0.40 mmol). The solution was monitored by ³¹P NMR spectroscopy until complete conversion into 8 had occurred. v_{max} (KBr, CDCl₃)/cm⁻¹ 1620; ¹H NMR (301.5 MHz, CDCl₃) δ 3.90 (13H, m, PNCH and CNCH(CH₃)₂), 3.73 (2H, s, CH₂), 3.68 (2H, s, CH₂), 1.41 (12H, d, ³J_{HH} 7.0, PNCH(CH₃)₂), 1.40 (12H, d, ³J_{HH} 7.0, PNCH(CH₃)₂) and 1.14 (6H, d, ³J_{HH} 6.0 Hz, CNCH(CH₃)₂); ¹³C NMR (50.3 MHz; CDCl₃) δ 156.5 (pseudo triplet, ²J_{PC} 7.0), 51.2 (br t, ⁴J_{PC} 2.0, C=NCH), 46.8 (d, ²J_{PC} 5.0, PNCH), 46.2 (d, ²J_{PC} 5.0, PNCH), 45.2 (d, ¹J_{PC} 79.0, CH₂), 41.3 (d, ¹J_{PC} 79.0, CH₂), 24.0 (d, ³J_{PC} 2.5, PNCH(CH₃)₂); ³¹P NMR (81.0 MHz; CDCl₃) δ +70.9 (⁴J_{PP} 10.0, ³J_{PH} 16.0) and +72.6 (⁴J_{PP} 10.0, ³J_{PH} 16.0 Hz).

Hydrogen-bonded thio tautomer 9. An excess of elemental sulfur (0.03 g, 0.94 mmol) was added directly to a stirred dichloromethane solution (10 cm³) of compound 2a (0.20 g, 0.36 mmol) at room temperature. The solution became vellow almost immediately and gradually turned red upon stirring for approximately 12 h. After 24 h at room temperature the mixture was filtered to remove unchanged sulfur, concentrated and placed at -30 °C. Pale yellow cubic crystals of 9 (0.14 g, 61%) were isolated after prolonged cooling, mp 132-134 °C (Found: C, 56.9; H, 8.9; N, 11.0. C₃₀H₆₇N₅P₂S₂ requires C, 57.8; H, 10.9; N, 11.2%); v_{max} (KBr, CDCl₃ solution)/cm⁻¹ 3264mbr, 2475w, 1610wsh, 1584m and 1538m; ¹H NMR (270 MHz; C₆D₆) δ 8.21 (1 H, br t, ⁴J_{PH} 6.0, NH), 4.35 (2 H, br d, ²J_{PH} 17.0, PCH₂), 4.18 (1 H, dd, ²J_{PH} 7.0, ⁴J_{PH} 2.5, PCH), 3.93 (8 H, m, PNCH), 3.39 (1 H, sept., ³J_{HH} 6.0, CNCH(CH₃)₂), 1.45 (24 H, d, ${}^{3}J_{HH}$ 7.0, PNCH(CH₃)₂), 1.39 (24 H, d, ${}^{3}J_{HH}$ 7.0, PNCH(CH_3)₂) and 1.13 (6 H, d, ${}^{3}J_{HH}$ 6.0 Hz, CNCH(CH_3)₂); ¹³C NMR (67.9 MHz; C_6D_6) δ 152.2 (dd, ² J_{PC} 18.0, 7.0, CN), 83.1 (dd, ¹ J_{PC} 152.0, ³ J_{PC} 8.0, PCH), 47.9 (d, ² J_{PC} 5.0, P(S)NCH), 46.3 (d, ² J_{PC} 6.0, P(S)NCH), 43.8 (s, C=NCH), 37.2 F(S)(CH), 46.3 (d, J_{PC} 6.0, F(S)(CH), 45.8 (s, C-NCH), 57.2 (dd, $^{1}J_{PC}$ 85.0, $^{3}J_{PC}$ 4.0, PCH₂), 24.1 (d, $^{3}J_{PC}$ 5.0, PNCH(CH₃)₂), 24.0 (d, $^{3}J_{PC}$ 6.0, PNCH(CH₃)₂), 23.9 (d, $^{3}J_{PC}$ 2.0, PNCH(CH₃)₂), 23.5 (d, $^{3}J_{PC}$ 3.0 Hz, PNCH(CH₃)₂) and 22.0 (s, CNCH₃); ³¹P NMR (81.0 MHz; CDCl₃) δ + 66.7 (s) and +66.6 (s); m/z (EI) 623 (M⁺), 523 (M – N{Pr₂ⁱ}₂) and 262 $([{Pr_2^iN}_2P=S]^+).$

Acknowledgements

The Royal Society is thanked for a European Science Exchange Fellowship (PWD), The Nuffield Foundation for both an Undergraduate Summer Bursary (MJH) and an Award to Newly Appointed Science Lecturers (PWD). Financial and equipment support from the CNRS, the Departments of Chemistry at Imperial College and the University of Leicester, and The Royal Society are gratefully acknowledged.

References

 See for example: (a) S. C. van der Slot, P. C. J. Kamer, P. W. N. M van Leeuwen, J. Fraanje, K. Goubitz, M. Lutz and A. L. Spek, Organometallics, 2000, 19, 2504; (b) K. G. Moloy and J. L. Petersen, J. Am. Chem. Soc., 1995, 117, 7696; (c) A. Huang, J. E. Marcone, K. L. Mason, W. J. Marshall and K. G. Moloy, Organometallics, 1997, 16, 3377; (d) S. M. Aucott, A. M. Z. Slawin and J. D. Woollins, J. Chem. Soc., Dalton Trans., 2000, 2559; (e) H. Tye, D. Smyth, C. Eldred and M. Wills, Chem. Commun., 1997, 1053; (f) D. E. C. Corbridge, Phosphorus: An Outline of Its Chemistry, Biochemistry and Technology, Elsevier, Amsterdam, 5th edn., 1995, p. 51; (g) J. M. Brunel, A. Heumann and G. Buono, Angew. Chem., Int. Ed., 2000, 39, 1946; (h) M. L. Clarke, D. J. Cole-Hamilton, A. M. Z. Slawin and J. D. Woollins, Chem. Commun., 2000, 2065.

- 2 F. A. Cotton and B. Hang, Prog. Inorg. Chem., 1992, 40, 179.
- 3 A. M. Z. Slawin, M. B. Smith and J. D. Woollins, J. Chem. Soc., Dalton Trans., 1996, 1283; J. C. Jeffrey and T. B. Rauchfuss, Inorg. Chem., 1979, 18, 2658.
- 4 See for example: E. K. van den Beuken, W. J. J. Smeets, A. L. Spek and B. L. Feringa, *Chem. Commun.*, 1998, 223; G. P. C. M. Dekker, A. Buijs, C. J. Elsevier, K. Vrieze, P. W. N. M. van Leeuwen, W. J. J. Smeets, A. L. Spek, Y. F. Wang and C. H. Stam, *Organometallics*, 1992, 11, 1937; A. Pfaltz, *Acta Chem. Scand.*, 1996, 50, 189; K. R. Reddy, K. Surekha, G.-H. Lee, S.-M. Peng and S.-T. Liu, *Organometallics*, 2000, 19, 2637.
- 5 M. J. Baker, M. F. Giles, A. G. Orpen, M. J. Taylor and R. J. Watt, J. Chem. Soc., Chem. Commun., 1995, 197.
- 6 P. Molina, A. Arques, A. García and M. C. Ramírez de Arellano, Eur. J. Inorg. Chem., 1998, 1359; A. Bader and E. Lindner, Coord. Chem. Rev., 1991, 108, 27.
- 7 R. B. King, N. D. Sadanani and P. M. Sundaram, J. Chem. Soc., Chem. Commun., 1983, 477.
- 8 The formation of compound 2a has very briefly been reported, previously: H. Hund, R.-D. Hund, S. Has and G.-V. Röschenthaler, *Phosphorus, Sulfur Silicon Relat. Elem.*, 1996, 111, 193.
- 9 C. Kowalski, X. Creary, A. J. Rollin and M. C. Burke, J. Org. Chem., 1978, 43, 2601; N. De Kimpe, M. Palamareva and N. Schamp, J. Org. Chem., 1985, 50, 2993; E. C. Ashby, A. B. Goel and R. N. DePriest, Tetrahedron Lett., 1981, 22, 4355.
- 10 V. Mark, C. H. Dungan, M. V. Crutchfield and J. R. Vanwazer, Top. Phosphorus Chem., 1967, 5, 1.

- 11 P. W. Dyer, N. Griep-Raming and M. J. Hanton, unpublished results.
- 12 P. H. M. Budzelaar, A. B. van Oort and A. G. Orpen, *Eur. J. Inorg. Chem.*, 1998, 1485.
- 13 J.-C. Guillemin and J.-M. Denis, Tetrahedron, 1988, 44, 4431.
- 14 A. D. Sinitsa, L. I. Nesterova, D. M. Malenko and S. N. Luk'yanenko, Russ. J. Gen. Chem., 1997, 67, 62.
- 15 J.-P. Gouesnard, J. Dorie and G. J. Martin, Can. J. Chem., 1980, 58, 1295.
- 16 CRC Handbook of Phosphorus-31 Nuclear Magnetic Data, ed. J. C. Tebby, CRC Press, Boston, 1991.
- 17 H.-J. Cristau, A. Chêne and H. Christol, Synthesis, 1980, 551.
- 18 H. R. G. Bender, E. Niecke, M. Nieger and H. Westermann, Z. Anorg. Allg. Chem., 1994, 620, 1194; H. Westermann, M. Nieger and E. Niecke, Chem. Ber., 1991, 124, 13.
- 19 A. P. Abbott, A. V. Malkov, N. Zimmermann, J. B. Raynor, G. Ahmed, J. Steele and P. Kocovski, *Organometallics*, 1997, 16, 3690.
- 20 D. Gudat, A. Haghverdi, H. Hupfer and M. Nieger, Chem. Eur. J., 2000, 6, 3414.
- G. R. Newkome and D. C. Hager, J. Org. Chem., 1982, 47, 599; E. C. Ashby, A. B. Goel and R. N. DePriest, J. Org. Chem., 1981, 46, 2429.
- 22 C. Sandorfy, in *The Chemistry of the Carbon–Nitrogen Double* Bond, ed. S. Patai, Wiley, New York, 1970, p. 2.
- 23 J. C. Clardy, R. L. Kolpa and J. G. Verkade, *Phosphorus*, 1974, 4, 133; C. Rømming and J. Songstad, *Acta Chem. Scand. Ser. A*, 1978, 32, 689.
- 24 O. I. Kolodiazhnyi, Tetrahedron Lett., 1982, 23, 499.
- 25 F. Dahan, P. W. Dyer, M. J. Hanton, M. Jones, D. M. P. Mingos, A. J. P. White, D. J. Williams and A. M. Williamson, in the press.
- 26 R. B. King and N. D. Sadanani, Synth. React. Org. Met.-Org. Chem., 1985, 15, 149.
 27 I-L. Fauré R. Réau, M. W. Wong, R. Koch, C. Wentrup and G.
- 27 J.-L. Fauré, R. Réau, M. W. Wong, R. Koch, C. Wentrup and G. Bertrand, J. Am. Chem. Soc., 1997, 119, 2819.
- 28 D. G. Norton, V. E. Haury, F. C. Davis, L. J. Mitchell and S. A. Ballard, J. Org. Chem., 1954, 19, 1054.
- 29 M. R. Mazières, C. Roques, M. Sanchez, J. P. Majoral and R. Wolf, *Tetrahedron*, 1987, 43, 2109.