

Organic & Biomolecular Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: Y. Bilgi, M. Kus and L. Artok, *Org. Biomol. Chem.*, 2020, DOI: 10.1039/D0OB01226B.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Palladium-catalysed regio- and stereoselective arylative substitution of γ,δ -epoxy- α,β -unsaturated esters and amides by sodium tetraaryl borates†Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Yasemin Bilgi, Melih Kuş‡ and Levent Artok *

Palladium-catalysed reactions of γ,δ -epoxy- α,β -unsaturated esters and amides with NaBAR_4 reagents proceeded regio- and stereoselectively, producing allylic homoallyl alcohols with aryl-substituents in the allylic position for a wide range of substrates. AsPh_3 was found to be a competent ligand for the arylation reaction, whereas phosphine ligand/Lewis acidic organoboron combinations favoured the substitution reaction by oxygen nucleophiles (e.g. H_2O , ROH).


Introduction

Stereoselective ring opening reactions of epoxides constitute one of the most important methods in organic chemistry.¹ Specifically, epoxy-functionalized acrylic acid derivatives are versatile reagents in epoxide ring-opening substitution reactions with oxygen,² nitrogen,³ hydride,⁴ and carbon-centered^{5,6} nucleophiles, which mainly take place in the form of 1,2-addition ($\text{S}_{\text{N}}2$).

To date, arylative ring-opening reactions of γ,δ -epoxy- α,β -unsaturated carboxylic acid derivatives have been possible using methods involving π -electron-rich aromatics in the presence of a Lewis acid catalyst (Scheme 1a),^{5d–g} arylcoppers (Scheme 1b),^{5h,i} or Grignard reagents in the presence of an iron catalyst (Scheme 1c)^{5j}. The first two methods usually suffer from occasionally low regioselectivity or unsatisfactory yields. Furthermore, any method using hard organometal nucleophiles, such as organocopper or Grignard reagents, requires an absolutely moisture-free environment and has low functional group tolerance. Nevertheless, soft arylborons are generally preferable reagents for C–C bond formation because of their low toxicity, availability, high functional group tolerance, longer shelf lives, and moisture insensitivity.⁷

Interestingly, to the best of our knowledge, there is no reported method in the literature utilizing arylboron reagents in the C–C bond formation of epoxy acrylic acid derivatives, except for those methods involving simple vinyl epoxides⁸ and styrenyl epoxides⁹, wherein the reactions took place by 1,4-addition ($\text{S}_{\text{N}}2'$) and α -aryl substitution, respectively. Perhaps, the most important reason for this is that exposure of the allylpalladium, a putative intermediate in palladium-catalysed reactions of allylic reagents, to the Lewis acidic boron-promoted attack of oxygen nucleophiles, which may be

present in the form of boronic acid, water, or alcohol additives.² Nevertheless, herein,



Scheme 1 Previous studies on the arylation of γ,δ -epoxy- α,β -unsaturated carboxylic acid derivatives.

we demonstrate that stereoselective $\text{S}_{\text{N}}2$ -type arylation of γ,δ -epoxy- α,β -unsaturated carboxylic acid derivatives can be successfully performed by the use of NaBAR_4 as the non-acidic

* Department of Chemistry, Faculty of Science, Izmir Institute of Technology, Urla, Izmir 35430, Turkey. E-mail: leventartok@iyte.edu.tr

† Electronic Supplementary Information (ESI) available: General procedures, tables for optimisation studies and robustness screening data, spectroscopic data for new compounds. See DOI: 10.1039/x0xx00000x

‡ Present address: Integrated Research Centres, Izmir Institute of Technology, Urla, Izmir 35430, Turkey

ARTICLE


Journal Name

organoboron reagent and AsPh_3 ligand under palladium catalysis (Scheme 1d).

View Article Online
DOI: 10.1039/D0OB01226B

Results and discussion

As the first attempt, the reaction of ester-functionalized vinyl epoxide **1a** and PhBneop was carried out at room temperature (rt), in the presence of $(i\text{-Pr})_2\text{NH}$ base and the $\text{Pd}_2(\text{dba})_3\text{-CHCl}_3/\text{PPh}_3$ (P/Pd = 4.1:1) catalyst combination, in a THF/water (4:1) mixture, in a Schlenk flask, and under Ar gas. In line with the reactivity trend mentioned above, this reaction proceeded with complete conversion, producing a vicinal diol, structure **4a**, in exclusively the *syn*-configuration as the major product, albeit formed in low yield (Table 1, entry 1).¹⁰ No detectable amount

Table 1. Reaction optimization with vinyl epoxide **1a**.^a

entry	organoboron	Pd/ligand	solvent (mL)	T (°C)	time	yield% ^b
1	PhBneop	Pd ₂ (dba) ₃ -CHCl ₃ /PPh ₃	THF/H ₂ O (4:1)	rt	45 min	34 4a
2	PhBneop	Pd ₂ (dba) ₃ -CHCl ₃ /dppe	THF/H ₂ O (4:1)	rt	O.N.	77 (69) ^c 4a
3	NaBPh ₄	Pd ₂ (dba) ₃ -CHCl ₃ /dppe	THF/H ₂ O (4:1)	rt	O.N.	N.D.
4	PhBneop	Pd ₂ (dba) ₃ -CHCl ₃ /AsPh ₃	THF/H ₂ O (4:0.5)	rt	48 h	55 3aa
5	PhBneop	Pd ₂ (dba) ₃ -CHCl ₃	THF/H ₂ O (4:0.5)	rt	48 h	N.C.
6	PhBneop	Pd ₂ (dba) ₃ -CHCl ₃ /AsPh ₃	THF/H ₂ O (4:0.5)	50	O.N.	55 3aa
7	PhBneop	Pd ₂ (dba) ₃ -CHCl ₃ /AsPh ₃	THF/MeOH (4:0.5)	50	3 h	55 3aa
8	NaBPh ₄	Pd ₂ (dba) ₃ -CHCl ₃ /AsPh ₃	THF/MeOH (4:0.5)	50	3.5 h	76 3aa
9 ^d	NaBPh ₄	Pd ₂ (dba) ₃ -CHCl ₃ /AsPh ₃	THF/MeOH (4:0.5)	50	5 h	48 3aa
10	NaBPh ₄	Pd(OAc) ₂ /AsPh ₃	THF/MeOH (4:0.5)	50	4 h	78 3aa
11	NaBPh ₄	Pd(OAc) ₂ /AsPh ₃	THF/MeOH (4:0.5)	70	1 h	82 3aa
12	NaBPh ₄	Pd(OAc) ₂ /AsPh ₃	dioxane/MeOH (4:0.5)	70	40 min	85 3aa
13	NaBPh ₄	Pd(OAc) ₂ /AsPh ₃	dioxane/MeOH (4:0.5)	110	2 min	90 (87) 3aa
14	NaBPh ₄ ^e	Pd(OAc) ₂ /AsPh ₃	dioxane/MeOH (4:0.5)	110	2.5 min	90 3aa
15	NaBPh ₄ ^e	Pd(OAc) ₂ /AsPh ₃ ^f	dioxane/MeOH (4:0.5)	110	4 min	50 3aa

^aEpoxide **1a**: 0.2 mmol; Phneop: phenylboronic acid neopentylglycol ester; N.D.: not determined; N.C.: no conversion; dppe: 1,2-bis(diphenylphosphino)ethane. ^bDetermined by ¹H-NMR using benzaldehyde as the internal standard. ^cIsolated yield. ^dWithout added base. ^e2.0 equiv. ^f1.0% Pd.

of any arylated product could be detected to form by GC/MS and ¹H-NMR techniques.

Numerous electron-rich and poor monodentate, as well as bidentate ligands were screened in the reaction (see the supporting file). Generally, phosphine ligands favoured the formation of **4a**, among which dppe provided the highest yield of **4a** (entry 2). In the presence of dppe, a non-acidic borate reagent, NaBPh₄, was also tried, but a complex mixture formed as a result of the reaction (entry 3).

Nevertheless, it was intriguing to find that the use of a more labile ligand, such as AsPh₃, for the reaction of **1a** and PhBneop rendered the arylative ring-opening reaction from the allylic position, thereby leading to γ -aryl-substituted homoallyl alcohols **3aa** in a moderate yield with almost clean stereoselectivity,¹¹ and no trace of **4a** was detected (entry 4). It is apparent that the presence of a ligand is essential for the reaction because no conversion occurred under a ligand-free conditions (entry 5).


Substitution of water with MeOH co-solvent resulted in considerable acceleration of the reaction at 50 °C (entries 6 and 7). Contrary to the result obtained under the conditions of entry 3, a good yield of **3aa** was obtained by using NaBPh₄ (**2a**) instead of PhBneop under the conditions of entry 8;¹² however, the presence of a base is required for a reasonable yield (entry 9). Gradual increments in the yield were

observed using 1,4-dioxane, Pd(OAc)₂, and elevated reaction temperatures (entries 10-12).

To maintain volatile MeOH and (*i*-Pr)₂NH in solution with high proportion, all reactions at ≥ 70 °C were carried out in a sealed Schlenk tube. The reaction was incredibly fast at 110 °C, completed within just 2 minutes, as judged by the appearance of palladium black (entry 13). The amount of **2a** could be reduced to 2 equivalents without sacrificing the yield (entry 14).¹³ On the other hand, reducing the amount of palladium catalyst resulted in a lower yield of the product (entry 15).















Having established effective conditions, an array of vinyl epoxide substrates was then subjected to the reactions with a number of NaBAR₄ reagents and the related results are listed in Table 2. When reacted with **2a**, high yields of homoallyl alcohols were able to be achieved with epoxy alkenyl ester substrates having a *primary* alkyl-substituted and *trans*-configured epoxy moiety, regardless of the structure of the ester group (entries 1–3).

The relative stereoisomeric configuration of homoallyl alcohol **3da** was decidedly verified by NOE interaction studies and the ¹H-NMR coupling constant (10 Hz) of the vicinal *tertiary* hydrogen signals of its δ -lactone derivative (eq 1); these data clearly indicated that **3da** possessed 4,5-*anti*-configuration.^{5,c,f,j,14}




This protocol was also be successfully implemented on NaBAR₄ reagents having electron-rich or moderately electron-poor phenyl groups (entries 4–6). The desired arylated product **3de**, was also be obtained in a good yield (72%) from the reaction of **1d** and the highly electron deficient NaB(*p*-CF₃C₆H₄)₄ (**2e**) reagent. However, its formation was accompanied by an amount of methoxylated side product **6d** (entry 7).¹⁵

Table 2. Allylic arylation of γ,δ -epoxy- α,β -unsaturated esters with NaBAR₄ reagents.^a


Entry	Epoxide	Product	Yield%
1			89
2			82
			
3		Ar: Ph 3da	90
4		<i>p</i> -Me-C ₆ H ₄ 3db	79
5		<i>m</i> -MeOC ₆ H ₄ 3dc	81
6		<i>m</i> -FC ₆ H ₄ 3dd	87
7		<i>p</i> -CF ₃ C ₆ H ₄ 3de	72 ^b
8			67
9			56 ^c
10			N.D.
11			50 ^d

^aThe reaction conditions of entry 14 in Table 1 were applied in the reactions. The reactions were complete within 2–5 minutes. ^bThe reaction also gave rise to an allylic methoxylated product (**6d**, 12%). ^cdr = 89:11. ^dThe reaction also gave rise to a phenyl-substituted unsaturated lactone (**7ha**, 32%).




The reaction of the substrate **1e**, containing a cyclohexyl group on the epoxy terminus, and **2a** led to a relatively lower yield (**3ea**). This result may imply that the method is sensitive to sterical effects at the epoxy site for ester compounds (entry 8). Moreover, when the epoxy group is tri-substituted, the reaction proceeded with a lower level of stereoselectivity (dr = 89:11), and furnished the homoallyl product **3fa** in a moderate yield (entry 9). Aryl-substituted substrates do not appear to be suitable for the method; the reaction of the epoxy ester **1g** led to an intricate mixture (entry 10).

It was surprising that the reaction of the substrate **1h**, having a *cis*-configured epoxy moiety, gave rise to an unsaturated lactone product, **7ha**, with *trans*-configuration in a reasonable yield, along with the expected homoallyl product **3ha**, which is the diastereomer of **3ba**. The formation of **7ha** requires a homoallyl structure, **3ha'**, with *Z*- and *anti*-4,5-configurations (eq 2), because the inherent structures of all the *E*-configured **3** structures do not allow the hydroxyl and ester groups to provide sufficient distance to undergo cyclative condensation.



The reaction mechanism should be similar to its iron-catalysed counterpart using Grignard reagents as the nucleophilic source (Scheme 1c).^{5j} The reaction cycle should begin with ring-opening by the attack of a palladium complex to **1** in *anti*-mode, leading to the formation of π -allylpalladium complex **A** (Scheme 2).¹⁶ Then, in the case where the ligand is AsPh₃, sequential transmetalation (**B**) and reductive elimination steps should furnish the arylated homoallyl alcohol structure **3**. The formation of **6d** from the



Scheme 2 General reaction mechanisms.

reaction of **1d** and **2e**, even in the presence of AsPh₃ (Table 2, entry 7), could be ascribed to the lower transmetalation activity of highly electron deficient **2e** with **A**. At this stage,


Journal Name

probably with a promotive effect of the by-product BPh_3 after reaching a critical concentration, **A** was exposed, to some extent, to nucleophilic attack by MeOH .^{2,3}

It has been previously shown that AsPh_3 is an activating palladium ligand that is superior to the most phosphines in accelerating the rate determining transmetalation step in the Stille coupling reaction.¹⁷ Consistent with this finding, it seems that the transmetalation step was the preferred pathway in the presence of the softer ligand, AsPh_3 , whereas the presence of the more strongly coordinating dppe ligand probably led to a deceleration of the transmetalation step. In turn, **A** was more exposed to nucleophilic oxygen attack in the presence of a Lewis acidic boron compound.

The formation of **3ha'**, which is presumed to be an intermediate structure for the formation of cyclic unsaturated lactone **7ha**, should necessitate π -allylpalladium intermediates **E** or **F**, in which planar allyl ligands bear ester groups oriented *anti* with respect to the middle allylic C-H bond. Further, since the reaction of **1h**, under the standard conditions, primarily yields π -allylpalladium intermediates **C** and **D**, with ester groups oriented *syn*, the formation of **C** or **D** should be followed by a π - σ - π interconversion. For the underlying reasons, it is difficult for us to explain this configurational transition at this time.

The versatility of this protocol has been further demonstrated by the reaction of an optically active ester reagent and a scale-up experiment. The asymmetric epoxidation of the corresponding diene ester by Shi's method and subsequent reaction of the produced epoxide (**2R,3R**)-**1d** and **2a** under standard conditions afforded (**4R,5R**)-**3da** with a high level of ee%, which proves that, combined with the Shi's procedure,¹⁸ the present method would be a convenient strategy for the synthesis of optically active aryl-functionalized homoallyl compounds (Scheme 3a).¹⁹ The method was also successfully applied on a gram scale (Scheme 3b); the reaction of 1.00 g of **1b** with 2 equivalents of **2a** was complete within 9 minutes, providing the desired product, **3da**, in a yield comparable to the counterpart mg scale reaction.



Scheme 3 Synthesis of an optically active product and scale-up experiment.

Scheme 4 Allylic arylation of γ,δ -epoxy- α,β -unsaturated amides (**8**) with NaBAr_4 (**2**) reagents.^a

Entry	Epoxide	Product	Yield%
1		Ar: Ph 9aa	90
2		<i>p</i> -Me-C ₆ H ₄ 9ab	79
3		<i>m</i> -FC ₆ H ₄ 9ad	87
4			92
5			73
6			88
7			86
8			91

^aThe reaction conditions of entry 14 in Table 1 were applied, but with 1.5 equivalent of **2**.

Finally, a robustness screen, which was formulated by Glorius and co-workers,²⁰ was performed with 9 different functional group additives (1 equivalent), including aldehyde, terminal alkyne, terminal alkene, primary alcohol, primary alkyl chloride, phenyl chloride, nitrile, and aniline (see Supporting Information), to the optimized conditions though it should be qualified that these results only precisely diagnose intermolecular competitions. Subsequently, we followed the yield of **3da** and recovery of the additive. While the protocol generally well-tolerated these additives in terms of **3da** formation, more or less reduction in additive recovery was observed.

After revealing that γ,δ -epoxy- α,β -unsaturated esters are perfectly amenable reagents toward allylic arylation, we then extended our studies to their amide derivatives (**8**). Gladly, the use of a relatively smaller amount of **2** (1.5 equiv) was sufficient for effective conversion of **8**, generally yielding the desired products in good to high yields (Scheme 4). The method operates well with both electron-rich and

ARTICLE

poor reagents **2** (entries 1–3). Although the structure of the amide group does not seem too critical for the process (entries 4–6), **9ca** could be isolated at a somewhat lower corresponding yield from the reaction of **8c**, containing relatively bulkier diisopropyl amide group (entry 5). Having larger groups on the oxirane terminus, such as propyl (**8e**) or cyclohexyl (**8f**), had no influence on the behaviour of the amide substrate (entries 7 and 8). It should be noted that an inferior result was obtained with its ester counterpart **1f** (Table 2, entry 8).

Conclusions

In conclusion, the palladium-catalysed reaction of γ,δ -epoxy- α,β -unsaturated esters or amides with NaBAR_4 reagents proceeded regio- and stereoselectively, specifically in *anti*-mode to produce allylic aryl-substituted homoallyl alcohols. While phosphine ligands showed very poor performance in terms of producing the desired product, AsPh_3 was found to be more suitable for the method, most likely because it increased the activity of the palladium at transmetalation step.

Experimental

General procedure for catalytic reactions.

The reactions at $<70^\circ\text{C}$ were carried out in a two-necked Schlenk attached to a condenser and inert gas-line, while those at $\geq 70^\circ\text{C}$ were performed in a sealed Schlenk tube under an inert gas.

The Pd-catalyst, ligand, and dry solvent (half of the volume required for reaction) were added, successively, into the Schlenk (dried in an oven and cooled under Ar gas) connected to an inert gas-line. The mixture was stirred for 5 min at rt, and then, organoboron, epoxide compound in a dry solvent (another quarter volume required for the reaction), additive, and base (in remaining last quarter volume of the dry solvent) were added successively. The mixture was stirred magnetically in a preheated oil bath. The reaction period was monitored by TLC when carried out in a two-necked Schlenk, while completion of the reaction was judged by the formation of palladium black when a sealed Schlenk tube was used. After completion of the reaction, the mixture was filtered through a short column of silica gel (height: 10 cm and width: 2 cm), washed with Et_2O , and concentrated under reduced pressure. The crude product was analysed by ^1H NMR using benzaldehyde as the internal standard. The residue was purified using silica gel column chromatography to afford the target product, the homoallylic alcohol **3** or **9** usually as a colourless oil.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Dedicated in memory of Prof. D. Nüket Öcal-Sunguroğlu. The financial support of the Scientific and Technological Research Council of Turkey (216Z094) and IZTECH (2017IYTE85) is gratefully acknowledged. We thank Mr. S. Sakrak of Redoks Analytical Laboratory Equipments Ltd. Co., Dr. O. N. Aslan of Atatürk University, and Ms. T. Tunceli of Trakya University for HRMS analyses.

Notes and references

- (a) A. Gansäuer, In *Aziridines and Epoxides in Organic Synthesis*; A. Yudin, Ed.; Wiley-VCH: Weinheim, 2006; p. 492; for reviews on the metal-catalysed reactions of epoxides, see: (b) J. Muzart, *Eur. J. Org. Chem.*, 2011, 4717; (c) C. Wang, L. Luo, H. Yamamoto, *Acc. Chem. Res.* 2016, **49**, 193; for reviews on the reactions of vinyl epoxides, see: (d) A. Kar, N. P. Argade, *Synthesis*, 2005, 2995; (e) M. Pineschi, F. Bertolini, V. D. Bussolo, P. Crotti, *Curr. Org. Chem.*, 2009, **13**, 290; (f) J. He, J. Ling, P. Chiu, *Chem. Rev.*, 2014, **114**, 8037.
- (a) A. Hirai, X.-Q. Yu, T. Tonooka, M. Miyashita, *Chem. Commun.*, 2003, 2482; (b) X.-Q. Yu, A. Hirai, M. Miyashita, *Chem. Lett.*, 2004, **33**, 764; (c) X.-Q. Yu, Y. Fumihiko, K. Tanino, M. Miyashita, *Tetrahedron Lett.*, 2008, **49**, 7442; (d) X.-Q. Yu, F. Yoshimura, F. Ito, M. Sasaki, A. Hirai, K. Tanino, M. Miyashita, *Angew. Chem., Int. Ed.*, 2008, **47**, 750.
- (a) T. Tsuda, Y. Horii, Y. Nakagawa, T. Ishida, T. Saegusa, *J. Org. Chem.*, 1989, **54**, 977; (b) M. Miyashita, T. Mizutani, G. Tadano, Y. Iwata, M. Miyazawa, K. Tanino, *Angew. Chem. Int. Ed.*, 2005, **44**, 5094.
- M. Oshima, H. Yamazaki, I. Shimizu, M. Nisar, J. Tsuji, *J. Am. Chem. Soc.*, 1989, **111**, 6280.
- (a) N. Ishibashi, M. Miyazawa, M. Miyashita, *Tetrahedron Lett.* 1998, **39**, 3775; (b) M. Miyashita, M. Hoshino, A. Yoshikoshi, *J. Org. Chem.*, 1991, **56**, 6483; (c) T. Ibuka, M. Tanaka, H. Nemoto, Y. Yamamoto, *Tetrahedron*, 1989, **45**, 435; (d) H. Akita, I. Umezawa, M. Takano, C. Ohyama, H. Matsukura, T. Oishi, *Chem. Pharm. Bull.*, 1993, **41**, 55; (e) M. Ono, R. Todoriki, Y. Yamamoto, H. Akita, *Chem. Pharm. Bull.*, 1994, **42**, 1590; (f) M. Ono, T. Ehara, H. Yokomoya, N. Ohtani, Y. Hoshino, H. Akita, *Chem. Pharm. Bull.*, 2005, **53**, 1259; (g) T. Ehara, S. Takinawa, M. Ono, H. Akita, *Chem. Pharm. Bull.*, 2007, **55**, 1361; (h) S. Nagumo, S. Irie, H. Akita, *J. Chem. Soc., Chem. Commun.*, 1995, 2001; (i) S. Nagumo, S. Irie, H. Akita, *Chem. Pharm. Bull.* 1996, **44**, 675; (j) T. Hata, R. Bannai, M. Otsuki, H. Urabe, *Org. Lett.*, 2010, **12**, 1012.
- For examples of C-C coupling reactions proceeded with 1,4-addition selectivity to form allyl alcohols: (a) J. Tsuji, H. Kataoka, Y. Kobayashi, *Tetrahedron Lett.*, 1981, **22**, 2575; (b) H. Nemoto, T. Ibaragi, M. Bando, M. Kido, M. Shibuya, *Tetrahedron Lett.*, 1999, **40**, 1319; (c) A. Hirai, A. Matsui, K. Komatsu, K. Tanino, M. Miyashita, *Chem. Commun.*, 2002, 1970.
- Hall, D. *Boronic Acids: Preparation and Applications in Organic Synthesis and Medicine*; Wiley-VCH: Weinheim, Germany, 2005.
- (a) N. Miyaoura, Y. Tanabe, H. Sugimoto, *J. Organomet. Chem.*, 1982, **233**, C13; (b) J. Kjellgren, J. Aydin, O. A.

Journal Name


- Wallner, I. V. Saltanova, K. J. Szabó, *Chem. Eur. J.*, 2005, **11**, 5260.
- 9 D. K. Nielsen, A. G. Doyle, *Angew. Chem. Int. Ed.*, 2011, **50**, 6056.
- 10 Diastereomeric form of diol compound **4a** was determined by NOE measurement of its acetonide derivative as described elsewhere (see the reference 2a,b).
- 11 NMR analyses of crude and isolated products in CDCl₃ or C₆D₆ revealed the presence of only single diastereomer of **3**, unless otherwise stated.
- 12 For selected examples of metal-mediated reactions using NaBAR₄ reagents as the transmetalation partner, see: (a) P. G. Ciattini, E. Morera, G. Ortá, *Tetrahedron Lett.*, 1992, **33**, 4815; (b) K. Ueura, S. Miyamura, T. Satoh, M. Miura, *J. Organomet. Chem.*, 2006, **691**, 2821; (c) H. Zeng, R. Hua, *J. Org. Chem.*, 2008, **73**, 558; (d) R. Shintani, Y. Tsutsumi, M. Nagaosa, T. Nishimura, T. Hayashi, *J. Am. Chem. Soc.*, 2009, **131**, 13588; (e) R. Shintani, S. Isobe, M. Takeda, T. Hayashi, *Angew. Chem. Int. Ed.* 2010, **49**, 3795.
- 13 No trans-esterification or amide by-products were found which may occur by the reaction of **1** or **3** with MeOH and (*i*-Pr)₂NH additives, respectively.
- 14 S. Nagumo, M. Ono, Y.-i. Kakimoto, T. Furukawa, T. Hisano, M. Mizukami, N. Kawahara, H. Akita, *J. Org. Chem.*, 2002, **67**, 6618.
- 15 The diastereomeric form of **6d** was tentatively estimated, assuming that its occurrence was similar to that of formation of **4a**.
- 16 (a) Hayashi, T., Hagihara, T., Konishi, M., Kumada, M., *J. Am. Chem. Soc.*, 1983, **105**, 7767; (b) M. Oshima, H. Yamazaki, I. Shimizu, M. Nisar, J. Tsuji, *J. Am. Chem. Soc.*, 1989, **111**, 6280.
- 17 V. Farina, B. Krishnan, *J. Am. Chem. Soc.*, 1991, **113**, 9585.
- 18 (a) Z.-X. Wang, Y. Tu, M. Frohn, J.-R. Zhang, Y. Shi, *J. Am. Chem. Soc.*, 1997, **119**, 11224; (b) K. Uchida, K. Ishigami, H. Watanabe, T. Kitahara, *Tetrahedron*, 2007, **63**, 1281.
- 19 Although we do not have a precise description for the slight erosion at the ee level, it can be caused by a minor equilibrium pathway that cleaves the non-allylic C-O bond.
- 20 K. D. Collins, F. Glorius, *Nat. Chem.*, 2013, **5**, 597.

Organic & Biomolecular Chemistry

View Article Online
DOI: 10.1039/D0OB01226B

ARTICLE

Organic & Biomolecular Chemistry Accepted Manuscript



View Article Online
DOI: 10.1039/D0OB01226B

The palladium-catalysed reaction of γ,δ -epoxy- α,β -unsaturated esters or amides with NaBAr_4 reagents proceeded regio- and stereo-selectively to produce aryl-substituted homoallyl alcohols