

Tetrahedron Letters 40 (1999) 6451-6454

TETRAHEDRON LETTERS

Homogeneous Diels-Alder catalysis by organotungsten Lewis acid containing *tris*(2-pyridyl)phosphine ligand

Cheng-Yi Kuo, Yaw-Shyan Fuh, Ming-Cheng Chaen and Shuchun Joyce Yu * Department of Chemistry, National Chung Cheng University, Ming Hsiung, Chia Yi, Taiwan 621, People's Republic of China

Received 27 May 1999; revised 22 June 1999; accepted 25 June 1999

Abstract

The complex $[P(2-py)_3W(CO)(NO)_2](BF_4)_2$ (1) is an excellent catalyst precursor for Diels-Alder reactions of 1,3-diene and α , β -unsaturated ketones and/or aldehydes with a catalyst loading as low as 0.3 mol%. © 1999 Elsevier Science Ltd. All rights reserved.

Keywords: Diels-Alder reaction; homogeneous catalysis; organotungsten Lewis acid; tris(2-pyridyl)phosphine complex.

The Diels-Alder (D-A) reaction is the most commonly used method for the synthesis of six-membered rings.¹ Lewis acid catalyzed D-A reactions generally provide better regio- and stereoselectivity than their corresponding thermal reactions.² However, most conventional Lewis acids are generally employed in large catalyst loading due to their extreme water-sensitivity and their potential in product binding. These Lewis acids are sometimes too active to prevent undesired side reactions. Thus, the study of low-valent transition metal complexes as Lewis acid catalysts for D-A reactions has become a subject of intense current interest.³⁻⁶

Recent reports⁷ from this laboratory have demonstrated that the complex [P(2py)₃W(CO)(NO)₂](BF₄)₂ (1) possesses strong Lewis acidity upon loss of the CO ligand. The relative Lewis acid strength⁸ of 1 is found to be comparable to that of BF₃.⁷ With a view to studying the catalytic aspect of 1, we report here the effective reactivity of 1 in catalyzing the D-A reactions of 1,3-dienes with α , β -unsaturated ketones and aldehydes (Scheme 1).

The catalyst precursor $[P(2-py)_3W(CO)(NO)_2](BF_4)_2$ (1) can be easily synthesized in two steps in high yield from the commercially available $W(CO)_6$. In addition, the complex 1 can be stored as a crystalline solid in air for months without significant decomposition.

The isolated yields of D-A adducts from 1-catalyzed D-A reactions were summarized in Table 1. The catalytic reactions⁹ were usually conducted in CH_3NO_2 using a catalyst loading as low as 0.3 mol% (catalyst-to-dienophile mole ratio) and an initial diene-to-dienophile ratio of 1.2:1.0.

^{*} Corresponding author. Fax: (886) 5-2721040; e-mail: chejyy@ccunix.ccu.edu.tw

^{0040-4039/99/\$ -} see front matter © 1999 Elsevier Science Ltd. All rights reserved. *P11:* S0040-4039(99)01284-8



 Table 1

 Diels-Alder reactions catalyzed by complex 1 at various temperatures

entry ^a	diene	dienophile R'	cat. loading (mol %) ^b	solvent	temp. (°C)	time ^c	endo : exo ^d	product n	yield (%)
1	3	CH ₃	0.3	CH ₂ Cl ₂	0	10 hr	93:7	1	91
2	3	CH ₃	0.3	CH ₃ NO ₂	0	1 hr	94 : 6	1	92
3	3	CH ₃	0.3	CH ₃ NO ₂	RT	30 min	94 : 6	1	87
4	3	CH ₃	0.3	CD ₃ NO ₂	70	< 3 min	93:7	1	93 ⁸
5	3	н	0.3	CH ₂ Cl ₂	0	9 hr	93 : 7	1	90
6	4	CH ₃	0.3	CH ₂ Cl ₂	0	30 hr	99:1	2	85 ^k
7	4	CH ₃	0.3	CH ₃ NO ₂	0	15 hr	99 :1	2	81 ^{<i>h</i>}
8	4	CH ₃	0.3	CH ₃ NO ₂	RT	12 hr	97:3	2	83
9	4	CH ₃	0.3	CH ₃ NO ₂	70	55 min	96:4	2	88
10	4	CH ₃	2	CH ₃ NO ₂	RT	2 hr	98:2	2	82
11	2	CH ₃	0.3	CH ₂ Cl ₂	0	35 hr	99 :1	0	79
12	2	CH ₃	0.3	CH ₃ NO ₂	0	72 hr		Polymer	91

^aInitial diene-to-dienophile ratio of 1.2 : 1.0 for each entry. ^bCatalyst-to-dienophile mole ratio. ^cTime were for greater than 90% completion. ^dDetermined by ¹H NMR. ^ePolymerization products were analyzed by ¹H and ¹³C NMR in C₆D₆. ^fIsolated yield. ^gAnalytical yield. ^hwork from reference 4.

The 1-catalyzed reaction of cyclopentadiene (3) and methyl vinyl ketone (MVK) at ambient temperature was complete in 30 minutes, however the corresponding thermal reaction was 83% complete in 20 hours (entries 3, Tables 1 and 2). Under the same conditions, the 1-catalyzed reaction of cyclohexadiene (4) and MVK was complete in 12 hours (entry 8, Table 1). In contrast, the thermal reaction was only 5% complete in 24 hours (entry 7, Table 2). Higher reaction rates can be obtained by increasing the catalyst loading. The reaction of 4 and MVK was complete in 2 hours using a catalyst loading of 2 mol% (entry 10, Table 1). The rates of both 1-catalyzed and uncatalyzed systems can also be enhanced by increasing reaction temperature. At 70°C and 0.3 mol% 1, the reactions of 3 with MVK and 4 with MVK were complete in <3 and 55 minutes, respectively (entries 4 and 9, Table 1). The corresponding uncatalyzed reactions were complete in 90 minutes and 7% complete in 6 hours, respectively (entries 4 and 8, Table 2). The observed rates of 3 and MVK are higher than those of 4 and MVK in both catalyzed and uncatalyzed systems. From ab initio calculations,¹⁰ the estimated intrinsic barrier for the gas phase uncatalyzed D–A reaction of 3 and MVK is 5 kcal/mol lower than that of uncatalyzed 4 and MVK system (this translates to a ratio of approximate 4000 in rate constants at room temperature). This result is, at least qualitatively, consistent with the experimental rate differences between 3/MVK and 4/MVK systems.

In order to minimize the thermal interference, the catalytic reactions were also carried out at milder

entry [#]	diene	dienophile R'	solvent	temp. (°C)	time (hours)	endo : exo ^b	product n	conversion ^c (%)
1	3	CH ₃	CH ₂ Cl ₂	0	24	85 : 15	1	86
2	3	CH3	CH ₃ NO ₂	0	22	86 : 14	1	82
3	3	CH3	CH ₃ NO ₂	RT	20	88 : 12	1	83
4	3	CH3	CH ₃ NO ₂	70	1.5	81 : 19	1	91
5	3	н	CH ₂ Cl ₂	0	24	81 : 19	1	83
6	4	CH ₃	CH ₂ Cl ₂	0	24			< 5
7	4	CH ₃	CH ₂ Cl ₂	RT	24			5
8	4	CH3	CH ₃ NO ₂	70	6			7
9	2	CH3	CH ₂ Cl ₂	0	24			< 5

 Table 2

 Uncatalyzed Diels-Alder reactions at various temperatures

^aInitial diene-to-dienophile ratio of 1.2 : 1.0 for each entry. ^bDetermined by ¹H NMR and GC. ^cThe percentage conversion is determined by ¹H NMR.

conditions. At 0°C, the catalytic reaction of 3 and MVK was complete in an hour, while the thermal reactions were 82% complete in 22 hours (entries 2, Tables 1 and 2). Under the same conditions, the catalytic reaction of 4 and MVK was complete in 15 hours, however, the thermal reactions were <5% complete in 24 hours in CH₂Cl₂ (entry 7, Table 1; entry 6, Table 2). Furthermore, the 1-catalyzed reaction of 3 and acrolein in CH₂Cl₂ was complete in 9 hours, while the uncatalyzed reaction took 24 hours to complete (entry 5, Table 1).

It is noteworthy that complex 1 provides not only efficiencies but also excellent *endo*-selectivity even at very mild conditions. In addition, 1 can be washed away from the reaction mixture¹¹ after the reaction to allow an easier process of product isolation.

The rates of most D–A catalyses are generally unaffected by solvents.¹² However, we have noticed in our study that the rates are, to a certain degree, influenced by the media.¹³ The 1-catalyzed reaction of 3 and MVK was complete in 10 hours in CH₂Cl₂, while the same reaction was completed within 1 hour in CH₃NO₂ (entries 1 and 2, Table 1). Also, the reactions of 4 and MVK were complete in 30 hours in CH₂Cl₂, and 15 hours in CH₃NO₂ (entries 6 and 7, Table 1). These results might imply that either a certain degree of change in charge separation of substrates takes place ongoing from the initial state to the transition structure or the solvent effect induces the change of Lewis acidity of the catalyst.

The diene polymerization reaction is often observed as a side reaction during the course of Lewis acid catalyzed D-A reaction. We have demonstrated in our system¹⁴ that the degree of polymerization can be excluded by reducing the solvent polarity. When the catalytic reaction of isoprene (2) and MVK was carried out in CH₂Cl₂ (μ =1.60), the product was dominated by the D-A adduct (entry 11, Table 1). In contrast, polyisoprene became the only detectable product when the catalysis was conducted in CH₃NO₂ (μ =3.73) (entry 12, Table 1).

In conclusion, the stable complex $[P(2-py)_3W(CO)(NO)_2](BF_4)_2$ (1) is an effective Lewis acid catalyst precursor for the D-A reactions of 1,3-diene and α , β -unsaturated ketones and aldehydes. Both the rates and the branching ratio of diene polymerization versus D-A reaction are found to be strongly solvent dependent. However, the actual mechanisms based on reactivity and selectivity are still speculative. The study of reaction mechanisms and reaction kinetics as well as a theoretical modeling of the catalytic system is currently in progress.

Acknowledgements

Financial supports from the National Science Council (NSC88-2113-M-194-018) and National Chung Cheng University are gratefully acknowledged.

References

- 1. Carruthers, W. Cycloaddition Reactions in Organic Synthesis; Pergamon Press: New York, 1990.
- For reviews see: (a) Kagan, H. B.; Riant, O. Chem. Rev. 1992, 92, 1007. (b) Pindur, U.; Lutz, G.; Otto, C. Chem. Rev. 1993, 932, 741.
- (a) Honeychuck, R. V.; Bonnesen, P. V.; Farahi, J.; Hersh, W. H. J. Org. Chem. 1987, 52, 5293. (b) Bonnesen, P. V.; Puckett, C. L.; Honeychuck, R. V.; Hersh, W. H. J. Am. Chem. Soc. 1989, 111, 6070. (c) Faller, J. W.; Smart, C. J. Tetrahedron Lett. 1989, 30, 1189. (d) Faller, J. W.; Ma, Y. J. Am. Chem. Soc. 1991, 113, 1579. (e) Olson, A. S.; Seitz, W. J.; Hossain, M. M. Tetrahedron Lett. 1991, 32, 5299.
- 4. (a) Odenkirk, W.; Rheingold, A. L.; Bosnich, B. J. Am. Chem. Soc. 1992, 114, 6392. (b) Hollis, T. K.; Robinson, N. P.; Bosnich, B. J. Am. Chem. Soc. 1992, 114, 5464. (c) Odenkirk, W.; Bosnich, B. J. Chem. Soc., Chem. Commun. 1995, 1181. (d) Hollis, T. K.; Odenkirk, W.; Robinson, N. P.; Whelan, J.; Bosnich, B. Tetrahedron 1993, 49, 5415.
- (a) Künding, E. P.; Bourdin, B.; Bernardinelli, G. Angew. Chem., Int. Ed. Engl. 1994, 33, 1856. (b) Bruin, M. E.; Künding, E. P. Chem. Commun. 1998, 2635. (c) Künding, E. P.; Saudan, C. M.; Bernardinelli, G. Angew. Chem., Int. Ed. Engl. 1999, 38, 1220.
- (a) Evans, D. A.; Lectka, T.; Miller, S. J. Tetrahedron Lett. 1993, 34, 7027. (b) Evans, D. A.; Lectka, T.; Miller, S. J. J. Am. Chem. Soc. 1993, 115, 6460. (c) Evans, D. A.; Barnes, D. M. Tetrahedron Lett. 1997, 38, 57. (d) Nie, J.; Kobayashi, H.; Sonoda, T. Catal. Today 1997, 36, 81. (e) Kanemasa, S.; Oderaotoshi, Y.; Sakaguchi, S.; Yamamoto, H.; Tanaka, J.; Wada, E.; Curran, D. P. J. Am. Chem. Soc. 1998, 120, 3074.
- 7. Kuo, C.-Y.; Fuh, Y.-S.; Shie, J.-Y.; Lee, G.-H.; Peng, S.-M.; Yu, S. J. J. Organomet. Chem. in press.
- 8. The complexation of crotonaldehyde to a Lewis acid provides a downfield shift of the H atom at C-3 position in the ¹H NMR spectrum and this correlation has been used to compare the relative acid strength of various Lewis acids. For details, see: Childs, R. F.; Mulholland, D. L.; Nixon, A. Can. J. Chem. 1982, 60, 801 and Ref. 3b.
- 9. A representative procedure follows: 10 mg (14.0 μmol) of 1 in CH₂Cl₂ (10 mL) was placed in a 25 mL flask. Methyl vinyl ketone (0.4 mL, 4.8 mmol) and cyclohexadiene (0.53 mL, 5.6 mmol) were then added to the stirred reaction mixture. When TLC showed the reaction to be complete (defined as >90% of MVK conversion), the mixture was dried under vacuum. The crude was then purified by flash chromatography on SiO₂ using hexane:EA (20:1) as eluent. Yield=0.60 g (83%).
- 10. Fuh, Y.-S.; Chaen, M.-C.; Tsai, L.-C.; Hu, W.-P.; Yu, S. J. in preparation.
- 11. A possible product [P(2-py)₃W(H₂O)(NO)₂](BF₄)₂ might be formed from the reaction of 1 and water. The resulting metal species was slightly soluble in water. For examples of documented transition metal aqua complexes as efficient D-A catalysts, see Refs. 4a-c and 6e.
- (a) Sauer, J.; Sustmann, R. Angew. Chem., Int. Ed. Engl. 1980, 19, 779. (b) Huisgen, R. Pure Appl. Chem. 1980, 52, 2283.
 (c) Reichardt, C. Solvents and Solvent Effects in Organic Chemistry; VCH: Cambridge, 1990.
- (a) Breslow, R. Acc. Chem. Res. 1991, 24, 150, and references cited therein. (b) Desimoni, G.; Faita, G.; Pasini, D.; Righetti, P. P. Tetrahedron 1992, 48, 1667, and references cited therein. (c) Otto, S.; Bertoncin, F.; Engberts, J. B. F. N. J. Am. Chem. Soc. 1996, 118, 7702, and references cited therein. (f) Otto, S.; Engberts, J. B. F. N.; Kwak, J. C. T. J. Am. Chem. Soc. 1998, 120, 9517.
- 14. There was no polymeric materials observed by the NMR nor detectable precipitates formed during the course of 1catalyzed D-A reactions of cyclopentadiene and cyclohexadiene systems.