Accepted Manuscript

In vitro and in silico insights into tyrosinase inhibitors with (E)benzylidene-1-indanone derivatives



Hee Jin Jung, Sang Gyun Noh, Yujin Park, Dongwan Kang, Pusoon Chun, Hae Young Chung, Hyung Ryong Moon

PII:	S2001-0370(19)30175-8
DOI:	https://doi.org/10.1016/j.csbj.2019.07.017
Reference:	CSBJ 375
To appear in:	Computational and Structural Biotechnology Journal
Received date:	12 April 2019
Revised date:	26 July 2019
Accepted date:	27 July 2019

Please cite this article as: H.J. Jung, S.G. Noh, Y. Park, et al., In vitro and in silico insights into tyrosinase inhibitors with (E)-benzylidene-1-indanone derivatives, Computational and Structural Biotechnology Journal, https://doi.org/10.1016/j.csbj.2019.07.017

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

In vitro and in silico insights into tyrosinase inhibitors with (E)-benzylidene-1-indanone derivatives

Hee Jin Jung^{1,2,3}, Sang Gyun Noh^{1,2,3}, Yujin Park¹, Dongwan Kang¹, Pusoon Chun⁴, Hae Young Chung^{1,2,3,*} hyjung@pusan.ac.kr, Hyung Ryong Moon^{1,*} mhr108@pusan.ac.kr ¹College of Pharmacy, Pusan National University, Busan 46241, Republic of Korea ²Longevity life Science and Technology Institutes, Pusan National University, Busan 46241, Republic of Korea

³Aging Tissue Bank, College of Pharmacy, Pusan National University, Busan 46241, Republic of Korea

⁴College of Pharmacy, Inje University, Gimhae, 47392, Republic of Korea *Corresponding authors.

Abstract

Tyrosinase is a key enzyme responsible for melanin biosynthesis and is effective in protecting skin damage caused by ultraviolet radiation. As part of ongoing efforts to discover potent tyrosinase inhibitors, we systematically designed and synthesized thirteen (*E*)-benzylidene-1-indanone derivatives (**BID1** – **13**) and determined their inhibitory activities against tyrosinase. Among the compounds evaluated, **BID3** was the most potent inhibitor of mushroom tyrosinase (IC₅₀ = 0.034 μ M, monophenolase activity; IC₅₀ = 1.39 μ M, diphenolase activity). Kinetic studies revealed that **BID3** demonstrated a mixed type of tyrosinase inhibition with *K*_i value of 2.4 μ M using L-DOPA as a substrate. *In silico* molecular docking simulations demonstrated that **BID3** can bind to the catalytic and allosteric sites of tyrosinase to inhibit enzyme activity which confirmed *in vitro* experimental studies between **BID3** and tyrosinase. Furthermore, melanin contents decreased and cellular tyrosinase activity was inhibited after **BID3** treatment. These observations revealed that **BID3** is a potent tyrosinase inhibitor and potentially could be used as a whitening agent for the treatment of pigmentation-related disorders.

Keywords: 1-Indanone; 2,4-Dihydroxy benzylidene; Melanin; Tyrosinase inhibitor

1. Introduction

Melanin is synthesized by melanocytes in the vassal layer of the epidermis generally refers to the family of pigments commonly known for protecting mammalian skin from damage caused by harmful UV radiation by scavenging free radicals or dispersing incoming UV light [1]. However, abundant generation of melanin can cause visible hyperpigmentation of the epidermis, which may be obvious as melasma, freckles, age spots, or senile lentigines [2].

Tyrosinase (EC 1.14.18.1), a copper-containing metalloenzyme, is a key enzyme involved in melanogenic processes. Tyrosinase catalyzes the two rate-limiting steps in melanogenesis; hydroxylation of tyrosine (cresolase or monophenolase activity) to produce 3,4dihydroxyphenylalanine (L-DOPA) and the subsequent oxidation of L-DOPA (catecholase or diphenolase activity) to the corresponding DOPA-quinone. When L-tyrosine is the substrate, the product of tyrosinase catalyzed reaction is DOPA-quinone, which is then converted to melanin [3]. These steps are crucial for the protection of skin against UV damage. Mushroom *Agaricus bisporus* is used to its commercially available and high homology with mammalian tyrosinase enzyme that renders it well suited as a model for studies on melanogenesis [4]. In humans, melanin helps defend skin form the damage caused by UV [5]. However, the excess level of melanin can cause various dermatological disorders including hyperpigmentations, melisma, freckles, and age spots [6]. Therefore, new tyrosinase inhibitors exhibiting drug-like and skin-whitening properties are required to inhibit excessive skin pigmentation.

1-Indanone, with a benzocyclopentanone skeleton, is considered as the rigid cousins of chalcones, incorporating the α , β -unsaturated ketone system of chalcones forming a cyclic 5 membered ring, which is a naturally occurring component present in various edible plant sources [7]. Several studies have shown that compounds possessing a 1-indanone moiety have pharmacological importance, as they exhibit various beneficial biological activities,

including anti-inflammatory [8-10], anticancer [11, 12], antioxidant [13], anti-Parkinson's disease [14], anti-Alzheimer disease [15-17], antimicrobial [18], anti-immune suppressive [19], and anti-tyrosinase [20] properties.

Chalcones (1,3-diaryl-2-propen-1-ones) are aromatic ketones consisting of two aromatic rings linked to a three carbon α,β -unsaturated carbonyl system as a central core [21, 22]. These are naturally occurring component found in various edible natural plants and considered as precursor compounds for synthetic route of flavonoids and isoflavonoids such as pyrazolines, pyrimidine, flavanol, flavones, flavanones, isollavone, aurones, antocianidin, dihydroflavanol and dihydrochalcone [23-25]. The α,β -unsaturated carbonyl system is paramount to the type of biological activity, since these system are distribute in many natural products like chalcones as well as indanone, where the molecular are relatively more planar, and the transmission of the electronic effects of the aryl substituents are directed to the carbonyl group [7]. Previous many researcher reported that chalcones were highlighted as a new class of tyrosinase inhibitor in several publication [26-29]. Recently, Kim and co-worker [30, 31] designed and synthesized a series of chalcone derivatives and evaluated them for their *in vitro* anti-tyrosinse and anti-melanogenic activities against a murine melanocytes.

According to our previously reported data, derivatives with an (E)- β -phenyl- α , β -unsaturated carbonyl scaffold showed potent tyrosinase inhibition [32-34]. (*E*)-Benzylidene-1-indanones can be an attractive anti-melanogenic agent, since the compounds have the (E)- β -phenyl- α , β -unsaturated carbonyl scaffold in their chemical structure. Notably, Radhakrishnan et al. (2015)[20] designed and synthesized a series of benzylidene-1-indanone derivatives bearing a (*Z*)-configuration and evaluated them for anti-tyrosinase activity. Although these (*Z*)-benzylidene-1-indanone derivatives have been investigated anti-tyrosinase and anti-melanogenic activities of (*E*)-benzylidene-1-indanone derivatives have yet to be scrutinized.

Therefore, we designed and synthesized thirteen compounds (BID1 - 13) of the 1-indanone

skeleton bearing a (*E*)-form benzylidene. Among these synthetic compounds, with 2,4dihydroxy group on the B ring of 1-indanone showed the greatest tyrosinase inhibition (approximately 400-fold more effective than kojic acid, positive control). Moreover, we further evaluate **BID3** tyrosinase inhibitory activity as well as characterize its regulatory role in melanin synthesis using B16F10 melanoma cells. In extended experiments, we evaluated anti-melanogenic potential of **BID3**, and sought to identify the enzyme kinetics and molecular docking studies. In successive experiments **BID3** cellular tyrosinase potential and melanin production in α -MSH and IBMX-induced B16F10 melanoma cells was also explored.

2. Materials and Methods

2.1. Reagents

Mushroom tyrosinase (EC 1.14.18.1), alpha-melanocyte stimulating hormone (α -MSH), 3isobutyl-1-methylxanthine (IBMX), L-tyrosine, L-3,4-dihydroxyphenylalanine (L-DOPA), dimethyl sulfoxide (DMSO), kojic acid, phthalic acid, and *trans*-cinnamic acid were purchased from Sigma-Aldrich (St. Louis, MO, USA). Dulbecco's modified Eagle's medium (DMEM), fetal bovine serum (FBS), streptomycin, and amphotericin were purchased from WELGENE Inc. (Gyeongsan-si, South Korea). All other reagents were purchased from Sigma-Aldrich.

2.2. Chemistry

2.2.1. General methods

All reagents were obtained commercially and used without further purification. Thin layer chromatography (TLC) and column chromatography were conducted on Merck precoated 60F245 plates and MP Silica 40-63, 60 Å, respectively. High resolution (HR) mass spectroscopy data were obtained on an Agilent Accurate Mass quadruple-time of flight (Q-TOF) liquid chromatography (LC) mass spectrometer (Agilent, Santa Clara, CA, USA) in ESI positive mode. Nuclear magnetic resonance (NMR) spectra were recorded on a Varian

Unity INOVA 400 spectrometer or a Varian Unity AS500 spectrometer (Agilent Technologies, Santa Clara, CA, USA) for ¹H NMR (400 and 500 MHz) and for ¹³C NMR (100 MHz), respectively. DMSO- d_6 , CD₃OD, and CDCl₃ were used as NMR solvents for NMR samples. Coupling constant (*J*) and chemical shift values were measured in hertz (Hz) and parts per million (ppm), respectively. The abbreviations used in the analysis of ¹H NMR data are as follows: s (singlet), brs (broad singlet), d (doublet), dd (doublet of doublets), t (triplet), td (triplet of doublets), q (quartet), and m (multiplet).

2.2.2. Procedure for the synthesis of BID1 - 13

A solution of benzaldehyde (1.1 - 1.2 equiv.) and 1-indanone (100 mg, 0.76 mmol) in 1M in HCl acetic acid (0.4 mL) was stirred at room temperature for 4 – 48 h. After the addition of water, the precipitates were filtered and washed with water and hexane:ethyl acetate (1:1), hexane:dichloromethane (4:1 – 1:1), or a mixture of dichloromethane and methanol, depending on the properties of the remaining benzaldehydes to give (*E*)-benzylidene-1-indanones (**BID1–13**) in yields of 32.8–99.1%. Structural characterization (¹H and ¹³C NMR and mass data of all BIDs) of synthesized compounds is provided in Supplementary information.

(*E*)-2-(4-Hydroxybenzylidene)-2,3-dihydro-1*H*-inden-1-one (**BID1**)

Yellow solid; yield, 93.4%; reaction time, 6 h; molecular formula, $C_{16}H_{12}O_2$; mp, 224.7-225.9 °C; ¹H NMR (400 MHz, DMSO- d_6) δ 10.10 (s, 1H, OH), 7.72 (d, 1H, J = 7.6 Hz, 7-H), 7.64–7.58 (m, 4H, 4-H, 5-H, 2'-H, 6'-H), 7.43 (s, 1H, vinylic H), 7.39 (t, 1H, J = 7.2 Hz, 6-H), 6.87 (d, 2H, J = 8.0 Hz, 3'-H, 5'-H), 3.99 (s, 2H, CH₂); ¹³C NMR (100 MHz, DMSO- d_6) δ 193.9, 160.1, 150.4, 138.3, 135.1, 134.0, 133.6, 132.2, 128.2, 127.2, 126.7, 124.1, 116.7, 32.6; HRMS (ESI+) $m/z C_{16}H_{13}O_2$ (M+H)⁺ calcd 237.0910, obsd 237.0911.

(*E*)-2-(3,4-Dihydroxybenzylidene)-2,3-dihydro-1*H*-inden-1-one (**BID2**)

Yellow solid; yield, 86.6%; reaction time, 5 h; molecular formula, C₁₆H₁₂O₃; mp, 251.2-

252.4 °C; ¹H NMR (400 MHz, DMSO- d_6) δ 9.65 (s, 1H, OH), 9.24 (s, 1H, OH), 7.72 (d, 1H, J = 7.6 Hz, 7-H), 7.66–7.61 (m, 2H, 4-H, 5-H), 7.42 (t, 1H, J = 7.2 Hz, 6-H), 7.35 (s, 1H, vinylic H), 7.19 (s, 1H, 2'-H), 7.08 (d, 1H, J = 8.0 Hz, 6'-H), 6.83 (d, 1H, J = 8.0 Hz, 5'-H), 3.98 (s, 2H, CH₂); ¹³C NMR (100 MHz, DMSO- d_6) δ 193.8, 150.4, 148.7, 146.3, 138.3, 135.1, 134.5, 132.0, 128.2, 127.3, 127.1, 125.0, 124.0, 118.2, 116.7, 32.7; HRMS (ESI+) m/z C₁₆H₁₃O₃ (M+H)⁺ calcd 253.0859, obsd 253.0857.

(E)-2-(2,4-Dihydroxybenzylidene)-2,3-dihydro-1H-inden-1-one (BID3)

Yellow solid; yield, 50.9%; reaction time, 10 h; molecular formula, $C_{16}H_{12}O_3$; mp, 198.5-199.9 °C (dec.); ¹H NMR (500 MHz, CD₃OD) δ 8.17 (s, 1H, vinylic H), 7.82 (d, 1H, J = 8.0 Hz, 6'-H), 7.67–7.63 (m, 3H, 4-H, 5-H, 7-H), 7.45 (t, 1H, J = 7.0 Hz, 6-H), 6.45 (d, 1H, J = 8.5 Hz, 5'-H), 6.39 (s, 1H, 3'-H), 4.01 (s, 2H, CH₂); ¹³C NMR (100 MHz, DMSO- d_6) δ 193.9, 161.6, 160.4, 150.3, 138.6, 134.8, 131.7, 130.3, 128.7, 128.1, 127.2, 123.9, 114.3, 108.6, 103.1, 32.7; HRMS (ESI+) $m/z C_{16}H_{13}O_3(M+H)^+$ calcd 253.0859, obsd 253.0858.

(*E*)-2-(4-Hydroxy-3-methoxybenzylidene)-2,3-dihydro-1*H*-inden-1-one (**BID4**)

Yellow solid; yield, 52.0%; reaction time, 4 h; molecular formula, $C_{17}H_{14}O_3$; mp, 186.9-187.4 °C;¹H NMR (400 MHz, DMSO- d_6) δ 9.71 (s, 1H, OH), 7.73 (d, 1H, J = 7.6 Hz, 7-H), 7.67–7.62 (m, 2H, 4-H, 5-H), 7.45 (d, 1H, J = 2.0 Hz, 2'-H), 7.43 (t, 1H, J = 7.2 Hz, 6-H), 7.31 (s, 1H, vinylic H), 7.24 (dd, 1H, J = 8.0, 2.0 Hz, 6'-H), 6.87 (d, 1H, J = 8.0 Hz, 5'-H), 4.05 (s, 2H, CH₂), 3.84 (s, 3H, OCH₃); ¹³C NMR (100 MHz, DMSO- d_6) δ 193.9, 150.5, 149.7, 148.5, 138.3, 135.2, 134.4, 132.3, 128.2, 127.3, 127.1, 125.7, 124.1, 116.6, 115.4, 56.3, 32.5; HRMS (ESI+) m/z C₁₇H₁₅O₃ (M+H)⁺ calcd 267.1016, obsd 267.1016.

(*E*)-2-(3-Ethoxy-4-hydroxybenzylidene)-2,3-dihydro-1*H*-inden-1-one (**BID5**)

Yellow solid; yield, 60.1%; reaction time, 2 d; molecular formula, $C_{18}H_{16}O_3$; mp, 135.8-136.4 °C;1H NMR (500 MHz, CDC13) δ 7.90 (d, 1H, J = 7.5 Hz, 7-H), 7.61 (s, 1H, vinylic H), 7.60 (td, 1H, J = 7.5, 1.0 Hz, 5-H), 7.55 (d, 1H, J = 7.5 Hz, 4-H), 7.42 (td, 1H, J = 7.5, 1.0

Hz, 6-H), 7.28 (dd, 1H, J = 8.0, 2.0 Hz, 6'-H), 7.14 (d, 1H, J = 2.0 Hz, 2'-H), 7.01 (d, 1H, J = 8.0 Hz, 5'-H), 6.05 (s, 1H, OH), 4.20 (q, 2H, J = 7.0 Hz, CH₂CH₃), 4.00 (s, 2H, CH₂), 1.50 (t, 3H, J = 7.0 Hz, CH₂CH₃); ¹³C NMR (100 MHz, CDCl₃) δ 194.2, 149.6, 147.8, 146.2, 138.5, 134.6, 132.5, 128.2, 127.8, 126.3, 125.0, 124.5, 115.2, 114.4, 64.9, 32.6, 15.0; HRMS (ESI+) m/z C₁₈H₁₇O₃ (M+H)⁺ calcd 281.1172, obsd 281.1172.

(*E*)-2-(3-Hydroxy-4-methoxybenzylidene)-2,3-dihydro-1*H*-inden-1-one (**BID6**)

Yellow solid; yield, 85.9%; reaction time, 4 h; molecular formula, $C_{17}H_{14}O_3$; mp, 186.1-186.9 °C; ¹H NMR (400 MHz, DMSO- d_6) δ 9.27 (s, 1H, OH), 7.72 (d, 1H, J = 7.2 Hz, 7-H), 7.65–7.59 (m, 2H, 4-H, 5-H), 7.41 (t, 1H, J = 7.2 Hz, 6-H), 7.37 (s, 1H, vinylic H), 7.21 (s, 1H, 2'-H), 7.18 (d, 1H, J = 8.4 Hz, 6'-H), 6.98 (d, 1H, J = 8.4 Hz, 5'-H), 3.98 (s, 2H, CH₂), 3.79 (s, 3H, CH₃); ¹³C NMR (100 MHz, DMSO- d_6) δ 193.9, 150.4, 150.3, 147.4, 138.2, 135.2, 134.0, 133.0, 128.4, 128.3, 127.3, 124.6, 124.1, 117.6, 112.8, 56.3, 32.6; HRMS (ESI+) $m/z C_{17}H_{15}O_3 (M+H)^+$ calcd 267.1016, obsd 267.1009.

(*E*)-2-(4-Methoxybenzylidene)-2,3-dihydro-1*H*-inden-1-one (**BID7**)

White solid; yield, 32.8%; reaction time, 6 h; molecular formula, $C_{17}H_{14}O_2$; mp, 138.9-139.5 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.88 (d, 1H, J = 7.6 Hz, 7-H), 7.63 (s, 1H, vinylic H), 7.63–7.56 (m, 3H, 5-H, 2'-H, 6'-H), 7.53 (d, 1H, J = 8.0 Hz, 4-H), 7.40 (t, 1H, J = 7.2 Hz, 6-H), 6.96 (d, 2H, J = 8.4 Hz, 3'-H, 5'-H), 3.98 (s, 2H, CH₂), 3.84 (s, 3H, OCH₃); ¹³C NMR (100 MHz, CDCl₃) δ 194.6, 161.1, 149.7, 138.5, 134.5, 134.0, 132.8, 132.6, 128.4, 127.8, 126.3, 124.5, 114.7, 55.6, 32.7; HRMS (ESI+) m/z C₁₇H₁₅O₂ (M+H)⁺ calcd 251.1067, obsd 251.1064.

(*E*)-2-(3,4-Dimethoxybenzylidene)-2,3-dihydro-1*H*-inden-1-one (**BID8**)

Yellowish solid; yield, 80.5%; reaction time, 2 d; molecular formula, $C_{18}H_{16}O_3$; mp, 176.5-177.2 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.90 (d, 1H, J = 7.5 Hz, 7-H), 7.62 (s, 1H, vinylic H), 7.60 (td, 1H, J = 7.0, 1.0 Hz, 5-H), 7.55 (d, 1H, J = 7.5 Hz, 4-H), 7.42 (t, 1H, J = 7.5 Hz,

6-H), 7.30 (dd, 1H, J = 8.0, 2.0 Hz, 6'-H), 7.18 (d, 1H, J = 2.0 Hz, 2'-H), 6.95 (d, 1H, J = 8.0 Hz, 5'-H), 4.01 (s, 2H, CH₂), 3.96 (s, 3H, OCH₃), 3.94 (s, 3H, OCH₃); ¹³C NMR (100 MHz, CDCl₃) δ 194.5, 150.8, 149.6, 149.3, 138.4, 134.6, 134.3, 132.8, 128.6, 127.8, 126.3, 124.8, 124.5, 113.7, 111.5, 56.2, 56.2, 32.6.

(*E*)-2-(2,4-Dimethoxybenzylidene)-2,3-dihydro-1*H*-inden-1-one (**BID9**)

Yellow solid; yield, 99.1%; reaction time, 24 h; molecular formula, $C_{18}H_{16}O_3$; mp, 74.0.-74.6 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.12 (s, 1H, vinylic H), 7.91 (d, 1H, *J* = 7.5 Hz, 7-H), 7.66 (d, 1H, *J* = 8.5 Hz, 6'-H), 7.58 (t, 1H, *J* = 7.5 Hz, 5-H), 7.53 (d, 1H, *J* = 7.5 Hz, 4-H), 7.41 (t, 1H, J = 7.5 Hz, 6-H), 6.58 (dd, 1H, *J* = 8.5, 2.5 Hz, 5'-H), 6.49 (d, 1H, *J* = 2.5 Hz, 3'-H), 3.97 (s, 2H, CH₂), 3.89 (s, 3H, OCH₃), 3.87 (s, 3H, OCH₃); ¹³C NMR (100 MHz, CDCl₃) δ 194.6, 162.7, 161.0, 149.8, 138.8, 134.3, 132.4, 131.0, 128.6, 127.7, 126.2, 124.5, 117.9, 105.4, 98.5, 55.8, 55.7, 32.8; HRMS (ESI+) *m*/*z* C₁₈H₁₇O₃ (M+H)⁺ calcd 281.1172, obsd 281.1180.

(*E*)-2-(3,4,5-Trimethoxybenzylidene)-2,3-dihydro-1*H*-inden-1-one (**BID10**)

Beige solid; yield, 63.2%; reaction time, 6 h; molecular formula, $C_{19}H_{18}O_4$; mp, 164.4-165.2 ^oC; ¹H NMR (400 MHz, CDCl₃) δ 7.87 (d, 1H, J = 7.2 Hz, 7-H), 7.59–7.52 (m, 3H, 4-H, 5-H, vinylic H), 7.39 (t, 1H, J = 7.2 Hz, 6-H), 6.86 (s, 2H, 2'-H, 6'-H), 3.99 (s, 2H, CH₂), 3.91 (s, 6H, 2 x OCH₃), 3.89 (s, 3H, OCH₃); ¹³C NMR (100 MHz, CDCl₃) δ 194.4, 153.6, 149.5, 140.0, 138.3, 134.8, 134.3, 134.0, 131.1, 127.9, 126.3, 124.6, 108.3, 61.2, 56.5, 32.4; HRMS (ESI+) $m/z C_{19}H_{18}O_4$ (M+H)⁺ calcd 311.1278, obsd 311.1282.

(E)-2-(4-Hydroxy-3,5-dimethoxybenzylidene)-2,3-dihydro-1H-inden-1-one (BID11)

Yellow solid; yield, 84.0%; reaction time, 5 h; molecular formula, $C_{18}H_{16}O_4$; mp, 190.2-191.1 °C;¹H NMR (500 MHz, DMSO- d_6) δ 9.12 (s, 1H, OH), 7.76 (d, 1H, J = 7.5 Hz, 7-H), 7.69–7.66 (m, 2H, 4-H, 5-H), 7.49 (s, 1H, vinylic H), 7.47–7.44 (m, 1H, 6-H), 7.07 (s, 2H, 2'-H, 6'-H), 4.12 (s, 2H, CH₂), 3.86 (s, 6H, 2 x OCH₃); ¹³C NMR (100 MHz, DMSO- d_6) δ

193.9, 150.5, 148.7, 138.9, 138.2, 135.2, 134.8, 132.6, 128.2, 127.3, 125.9, 124.1, 109.5, 56.8, 32.4; HRMS (ESI+) *m*/*z* C₁₈H₁₇O₄ (M+H)⁺ calcd 297.1121, obsd 297.1128.

(*E*)-2-(3-Bromo-4-hydroxybenzylidene)-2,3-dihydro-1*H*-inden-1-one (**BID12**)

Yellowish beige solid; yield, 54.4%; reaction time, 10 h; molecular formula, $C_{16}H_{11}BrO_{2}$; mp, 225.0-226.2 °C; ¹H NMR (400 MHz, DMSO- d_{6}) δ 10.92 (s, 1H, OH), 7.89 (s, 1H, 2'-H), 7.72 (d, 1H, J = 7.6 Hz, 7-H), 7.67–7.64 (m, 2H, 4-H, 5-H), 7.61 (d, 1H, J = 8.4 Hz, 6'-H), 7.42 (t, 1H, J = 7.2 Hz, 6-H), 7.40 (s, 1H, vinylic H), 7.03 (d, 1H, J = 8.4 Hz, 5'-H), 4.02 (s, 2H, CH₂); ¹³C NMR (100 MHz, DMSO- d_{6}) δ 193.8, 156.4, 150.5, 138.0, 136.0, 135.4, 133.6, 132.4, 132.4, 128.4, 128.3, 127.3, 124.1, 117.3, 110.7, 32.4; HRMS (ESI+) m/z C₁₆H₁₂BrO₂ (M+H)⁺ calcd 315.0015, obsd 315.0008, m/z C₁₆H₁₂BrO₂ (M+2+H)⁺ calcd 316.9996, obsd 316.9996.

(E)-2-(3,5-Dibromo-4-hydroxybenzylidene)-2,3-dihydro-1H-inden-1-one (BID13)

White solid; yield, 63.7%; reaction time, 24 h; molecular formula, $C_{16}H_{10}Br_2O_2$; mp, 267.4-268.1 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.55 (brs, 1H, OH), 7.92 (s, 2H, 2'-H, 6'-H), 7.73 (d, 1H, J = 7.2 Hz, 7-H), 7.67–7.63 (m, 2H, 4-H, 5-H), 7.43 (t, 1H, J = 7.2 Hz, 6-H), 7.38 (s, 1H, vinylic H), 4.05 (s, 2H, CH₂); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 193.8, 152.7, 150.7, 137.8, 135.6, 135.4, 135.1, 130.9, 130.2, 128.4, 127.4, 124.2, 112.8, 32.2; HRMS (ESI+) m/z C₁₆H₁₁Br₂O₂ (M+H)⁺ calcd 392.9120, obsd 392.9124, C₁₆H₁₁Br₂O₂ (M+2+H)⁺ calcd 394.9101, obsd 394.9101, C₁₆H₁₁Br₂O₂ (M+H)⁺ calcd 396.9082, obsd 396.9087.

2.3. Biological evaluation

2.3.1. Mushroom tyrosinase inhibitory assay

Tyrosinase inhibitory activity assay was performed using mushroom tyrosinase as described previously, with minor modification [35, 36]. Briefly, 10 μ L of a specified concentrations of each compound (0.0005 – 50 μ M) and 20 μ L of mushroom tyrosinase (1000 units/mL) in a 50 mM phosphate buffer (pH 6.5) were added to 170 μ L reaction mixture in a 96-well

microplate (Corning, USA). The ratio of 1mM L-tyrosine or L-DOPA solution, 50 mM potassium phosphate buffer (pH 6.5), and distilled water was 10:10:9. The reaction mixtures were incubated at 37 °C for 30 min. After that, the amount of dopachrome produced in each well was measured by spectrophotometric analysis at 492 nm (OD₄₉₂) using a microplate reader. The tyrosinase inhibition rate (%) was calculated as $(1 - Abs_{sample} / Abs_{control}) \times 100\%$. The degree of sample inhibition was expressed as the concentration required for 50% inhibition (IC₅₀).

2.3.2. Enzyme kinetic analysis with tyrosinase

2.3.3. Tyrosinase molecular docking simulations

To determine the structure of the enzyme-inhibitor complex and to ensure accuracy, repeatability, and reliability of the docking results, we employed AutoDock4.2 software. For dicking studies, the crystal structures of mushroom tyrosinase protein target was obtained from the protein sequence alignment (Protein Data Bank (PDB ID: 2Y9X) (http://www.rcsb.org/adb) [40]. Automated docking simulations were performed between tyrosinase and kojic acid, phthalic acid, cinnamic acid, or **BID3**. For the docking procedure: conversed 2D into 3D stuctures, calculated charges, and added hydrogen atoms using the

ChemOffice program (http://www.cambridgesoft.com) [41]. LigandScout 4.1.5 was used for the prediction of possible interactions between ligands and tyrosinase and the identification of pharmacophores.

2.3.4. Cell culture and cell viability assay

Murine melanoma B16F10 cells were purchased from the Korean Cell Line Bank (KCLB, Seoul, Korea) and cultured in DMEM supplemented with 10% FBS and 1% streptomycin, and then incubated at 37 °C, humidified with 5% atmospheric CO₂. Cell viability analyses were performed as previously described [42]. Briefly, cells seeded at a density of 1×10^4 cells/well in a 96-well plate for 24 h. On the following day, the cells were exposed to different concentrations of **BID3** and incubated for 24 or 48 h, respectively. Then, 10 µL EZ-Cytox solution was added to each well and the cells were incubated for 2 - 4 h. Absorbance was determined using ELISA at a wavelength of 450 nm. Each assay was performed in triplicate.

2.3.5. Melanin contents assay

Melanin content was determined as described previously [43]. B16F10 melanoma cells (2 × 10^5 cells/well) were seeded in 6-well culture plates. To determine the inhibitory effect of **BID3** on melanogenesis, fresh medium was replaced with medium containing **BID3** (1, 5 and 10 µM) or kojic acid (10 µM) as a positive control for 1 hr, and then stimulated with α-MSH (5 µM) and IBMX (200 µM) for 48 h. After washed twice with PBS, the cells were detached by incubation in Trypsin/EDTA, and the pellets were dissolved in 100 µL of 1N NaOH, and then incubated at 60 °C for 1 h and mixed to solubilize the melanin. The melanin contents were determined by measuring absorbance at 405 nm by using the ELISA reader (TECAN, Sunrise, Austria). The melanin content was calculated using the following equation: ($\Delta_{sample}/\Delta_{control}$) × 100%. All determinations were performed in triplicate.

2.3.6. Cellular tyrosinase activity

Cellular tyrosinase activity was assessed as described previously, with slight modifications [44, 45]. Briefly, 2×10^5 /cells were plated in 6-well dishes and incubated overnight. The cells were exposed to various concentrations of **BID3** (1, 5, and 10 µM) or kojic acid (10 µM) for 1 h, and subsequently stimulated with α-MSH (5 µM) and IBMX (200 µM) for 48 h. After being rinsed twice with PBS, the cells were lysed with 100 µL lysis solution containing 50 mM phosphate buffer (pH 6.5), 0.1 mM phenylmethylsulfonylfluoride (PMSF), and 1% Triton X-100 and frozen at -80 °C for 30 min. Lysates were thawed and centrifuged at 12,000 rpm for 30 min at 4 °C. Then, supernatant (80 µL) were combined with and 2 mg/mL L-DOPA (20 µL) in a 96-well plate. After incubation at 37 °C for 30 min, the optical density at 492 nm was measured using an ELISA reader (TECAN, Salzburg, Austria). The protein concentration was determined using BCA protein assay reagent using BCA as a standard (Thermo Scientific, Rockford, IL, USA).

2.3.7. Statistical analysis

All data are presented as the mean \pm standard error of the mean (S.E.M). The data were analyzed by one-way analysis of variance (ANOVA) to determined differences between treatments, followed by the Bonferroni post-hoc test. A value of p < 0.05 was considered statistically significant.

3. Results and discussion

3.1. Chemistry

The (*E*)-benzylidene-1-indanones derivatives (**BID1** – **13**) were synthesized according to the general scheme 1. In this present study, thirteen (*E*)-2-benzylidene-1-indanone derivatives were synthesized and their tyrosinase inhibitory properties were examined. Target (*E*)-2benzylidene-1-indanone derivatives were synthesized employing acidic (1M HC1 in acetic acid) conditions. These reactions generated the target (*E*)-2-benzylidene-1-indanone derivatives in yields of 32.8 - 99.1%. It appears that only the *E*-isomers of 2-benzylidene-1-

indanones were generated due to the A-strain known as the 1,3-allylic strain. The A-strain (steric hindrance) between the phenyl ring and the carbonyl group in the Z-isomer is apparently larger than that between the phenyl ring and the methylene group in the *E*-isomer. The structures of synthetic compounds were verified by ¹H and ¹³C NMR and mass spectrometry as described in the experimental section (Fig. S1-S38). In particular, the chemical shift of olefinic proton is deshielded in the *E*-isomer of 2-benzylidene due to the anisotropic effect of the carbonyl group and appears downfield (above 7 ppm) compared to the Z isomer (less than 7 ppm) [46]. The chemical shifts of olefinic protons in the benzylidene-1-indanones appeared in the range of 7.31 – 8.17 ppm. Therefore, the benzylidene-1-indanone derivatives was determined to be (*E*)-stereoisomers. We found that the presence of hydroxyl or methoxyl group at the 2-position of the phenyl ring moves the chemical shift of the olefinic proton to 0.5 - 0.8 ppm downfield.



Scheme 1. Synthesis of (E)-2-benzylidene-1-indanones.

3.2. Tyrosinase inhibitory properties of compounds

The tyrosinase inhibitory potential of the (*E*)-benzylidene-1-indanones derivatives were examined using mushroom tyrosinase. The monophenolase (L-tyrosine) and diphenolase (L-DOPA) were used as substrates for these experiments [35]. Enzyme reactions were performed with tyrosinase, substrate, and test inhibitors. All tested compounds demonstrated a concentration-dependent inhibition. The IC₅₀ values of the (*E*)-benzylidene-1-indanones derivatives are shown in Table 1.

Table 1. Mushroom tyrosinase inhibitory potential of the substituted 2,3-dihydro-1*H*-inden-1one (1-indanone) chalcone-like derivatives (**BID1** – **13**)



Compounds	R ₁	R ₂	R ₃	R ₄	$IC_{50} (\mu M)^{a}$	IC ₅₀ (μM) ^D
BID1	Н	Н	ОН	Н	39.74 ± 3.71	130.0 ± 5.39
BID2	Н	OH	ОН	Н	41.27 ± 3.03	52.93 ± 2.62
BID3	ОН	Н	ОН	Н	0.034 ± 0.00224	1.39 ± 0.00004
BID4	Н	OMe	ОН	Н	73.51 ± 5.78	> 200
BID5	Н	OEt	OH	Н	>200	> 200
BID6	Н	OH	OMe	Н	5.00 ± 0.91	50.78 ± 2.16
BID7	Н	Н	OMe	Н	> 200	> 200
BID8	Н	OMe	OMe	Н	> 200	> 200
BID9	OMe	Н	OMe	Н	> 200	96.85 ± 9.78
BID10	Н	OMe	OMe	OMe	> 200	> 200
BID11	Н	OMe	OH	OMe	75.15 ± 2.24	> 200
BID12	Н	Br	OH	Н	>200	> 200
BID13	Н	Br	OH	Br	> 200	75.31 ± 7.94
Kojic acid ^c					13.77 ± 0.20	33.14 ± 0.93
Phthalic acid ^a					> 200	> 200
Cinnamic acid ^a					> 200	> 200

The IC₅₀ values (μ M) were calculated from a log dose inhibition curve using L-tyrosine ^a and L-DOPA ^b as a substrate, respectively and are as means ± SEM of triplicate experiments. ^c Reported competitive type inhibitor. ^d Reported mixed type inhibitor.

In both L-tyrosine and L-DOPA substrates, **BID3** exhibited strong inhibitory activitt with IC_{50} values of 0.034 \pm 0.00224 μ M and 1.39 \pm 0.00004 μ M, respectively, compared with the positive control kojic acid, which had IC_{50} values of 13.77 \pm 0.20 μ M and 33.14 \pm 0.93 μ M, respectively (Fig. 1A). Furthermore, **BID6** showed potent activity toward L-tyrosine and L-DOPA with IC_{50} values of 5.00 \pm 0.91 μ M and 53.11 \pm 2.79 μ M, respectively. **BID1** showed

significant inhibition against tyrosinase via L-tyrosine and L-DOPA pathways, with IC₅₀ values of $39.74 \pm 3.71 \ \mu\text{M}$ and $130.0 \pm 5.39 \ \mu\text{M}$, respectively. **BID4** and **BID11** displayed moderate tyrosinase inhibitory activity. Other derivatives were inactive. Likewise, **BID2** showed significant activity toward L-tyrosine and L-DOPA, with IC₅₀ values of $41.27 \pm 3.03 \ \mu\text{M}$ and $52.93 \pm 2.62 \ \mu\text{M}$. In addition, reported mixed type inhibitors, phthalic acid and cinnamic acid showed no inhibitory effect at the concentration of 200 μ M [47].

Consistent with the findings of structure-activity relationship (SAR) studies, derivatives containing (E)-benzylidene-1-indanone substituents on the phenyl ring noticeably influenced the potential tyrosinase inhibition. **BID3**, which inhibited tyrosinase with the highest potency, bear a 2-hydroxy group in the presence of a 4-hydroxy group and greatly inhibited tyrosinase activity (BID1 vs BID3). These results suggested that the 2,4-dihydroxy group (resorcinol moiety) in the phenyl ring structure may be responsible for the enhanced inhibition of tyrosinase. Interestingly, in a previous study on potential synthetic tyrosinase inhibitors, we showed that the 2,4-dihydroxy substitution potently inhibited tyrosinase activity [48, 49]. In addition, our SAR data also indicated that the introduction of a 3-hydroxy group in the presence of a 4-hydroxy group affects tyrosinase inhibition. These phenomena were demonstrated by the finding that a 4-methoxy-3-hydroxy phenyl ring increased inhibitory activity against tyrosinase, whereas the introduction of a 3,4-dihydroxy group (catechol moiety) decreased tyrosinase inhibitory activity (BID2 vs BID6) [35]. There results suggest that the 3-hydroxy-4-methoxy group is also responsible for tyrosinase inhibitory activity. Nevertheless, in our present study, BID9 and BID11, which contain 2,4-dimethoxyphenyl or 3,5-dimethoxyphenyl groups, showed moderate tyrosinase inhibitory activity. Based on IC_{50} values via the L-tyrosine and L-DOPA pathway, BID3 demonstrated the most potent tyrosinase inhibitory activity therefore, we further studied the mechanism underlying this inhibitory effect. Radhakrishnan et al. (2015) [20] reported that hydroxyl substituted (Z)-

benzylidene-1-indanone compounds potent tyrosinase inhibitors. Although previous studies have shown that hydroxyl-substituted (Z)-benzylidene-1-indanone derivatives possess anti-tyrosinase activity, to the best of our knowledge, there have been no reports of tyrosinase inhibition by (E)-benzylidene-1-indanone derivatives using cell-based experiments.



Fig. 1. Concentration-dependent inhibition of mushroom tyrosinase activity by **BID3** and kojic acid (positive control) (n = 3).

3.3. Enzyme kinetic analysis of tyrosinase inhibition

Since **BID3** was the most potent inhibitor, we further studied the mechanism underlying its inhibitory effect. In an attempt to explain the mode of enzyme inhibition, kinetic analyses were performed at various concentrations of L-DOPA and various **BID3** concentrations. Dixon and Lineweaver-Burk plots were drawn using the data obtained from the kinetic studies in order to confirm the inhibition pattern and the inhibition constants (K_i) were determined by interpretation of Dixon plots (Fig. 2A and 2B). The concentration of L-DOPA is denoted as follows: 1, 2, 4 mM. The intersection lies on the left side, indicating mixed-type inhibition against tyrosinase with a K_i value 2.4 μ M (Fig. 2B). The K_i value represent the concentrations required to form an enzyme inhibitor complex, so inhibitor with lower K_i value indicate greater tyrosinase inhibition activity.



Fig. 2. Lineweaver-Burk (A) and Dixon plots (B) for the inhibition of mushroom tyrosinase by **BID3** using L-DOPA as the substrate. The effects are shown in the presence of different concentrations of **BID3** [0 μ M (*closed circle*), 1 μ M (*open circle*), 2 μ M (*closed triangle*), and 4 μ M (*open triangle*)] and in the presence of different concentrations of substrate [0.125 mM (*closed circle*), 0.25 mM (*open circle*), 0.5 mM (*closed triangle*), and 1 mM (*open triangle*)] of L-DOPA. Error bars indicate the standard error of the mean (SEM).

Molecular docking has contributed important insights into drug discovery over many years. However, docking procedures aim to identify the correct positions of ligands in the binding pocket of a protein and to predict the affinity between ligand and protein. In other words, docking describes a process through which two molecules fit together in a three-dimensional space. To determine whether **BID3** binds to the active site of tyrosinase, molecular docking analysis was performed to identify the possible binding positions of **BID3** in the crystal structure of mushroom tyrosinase (PDB ID: 2Y9X) (Fig. 3A). These results were calculated using the AutoDock4.2. program. The most stable binding position was that with the lowest score [50]. Recently, Hassani et al. [47] reported that cinnamic acid and phthalic acid were mixed-mode tyrosinase inhibitor. Thus, kojic aid, cinnamic acid and phthalic acid were used to validate docking results [51].

The molecular docking models of **BID3** and kojic acid (well-known competitive type inhibitor) in the catalytic site of tyrosinase are illustrated in Fig. 3B [52]. The tyrosinase-**BID3** inhibitor complex presented –6.28 kcal/mol binding energy including two hydrogen

bonds with the ASN260 and MET280 residue of tyrosinase, and hydrophobic interactions were observed between **BID3** and tyrosinase residues VAL248, PHE264, and VAL283, which further stabilized the interaction in the catalytic site for mushroom tyrosinase (Table 2, and Fig. 3E and H). Moreover, molecular docking models of BID3, and of phthalic acid (allosteric inhibitor 1) and cinnamic acid (allosteric inhibitor 2) in the two allosteric sites are illustrated in Fig. 3C and 3D. Phthalic acid and cinnamic acid were taken as reference ligands at allosteric sites 1 and 2, respectively [47, 51]. As shown in Fig. 3F and I, BID3 and phthalic acid exhibited one hydrogen bond with the TRY140 residue of tyrosinase, and three hydrophobic interactions residues with the LEU24, PHE147, and ILE148 of tyrosinase at allosteric site 1. The corresponding ligand interactions of BID3 and cinnamic acid in the allosteric site 2 of tyrosinase included hydrogen bonds between ASP312 and LYS376 of tyrosinase, whereas THR308 interacted hydrophobically at allosteric site 2 (Fig. 3G and 3J). Furthermore, **BID3**-tyrosinase binding was found to exert binding energy in both allosteric sites of tyrosinase (-4.38 and -5.68 kcal/mol, respectively), indicating high binding affinity with both allosteric sites. Based on enzyme kinetic studies, **BID3** exerted a mixed-type inhibition, binding ability of BID3 with both the catalytic site and two allosteric sites confirmed its mixed-type inhibition of tyrosinase.



Fig. 3. Molecular docking models of tyrosinase inhibition by kojic acid, phthalic acid, cinnamic acid, and **BID3** (blue color) (A). Inhibition mode of **BID3** at the tyrosinase catalytic site with the catalytic inhibitor, kojic acid (pink color) (B) and at allosteric sites 1 and 2 with the allosteric inhibitor, phthalic acid (violet color) (C) and cinnamic acid (orange color) (D), respectively. 2D ligand interaction diagrams of the catalytic (E) and allosteric (F and G) of tyrosinase by **BID3** and kojic acid (H), phthalic acid (I), and cinnamic acid (J) as catalytic and allosteric inhibitors, respectively. Green and red arrow: hydrogen-bonding, yellow: hydrophobic interaction.

K	Compounds	Binding energy (kcal/mol) ^a	Binding residues ^b
Catalytic inhibitor	BID3	-6.28	VAL248, ASN260, PHE264, MET280, VAL283
	Kojic acid [°]	-5.23	HIS61, HIS85, HIS263, MET280
Allosteric inhibitor1	BID3	-4.38	LEU24, TRY140, PHE147, ILE148
	Phthalic acid ^d	-3.19	TRP136, GLY149, ILE217, ALA220, ALA221, PHE224, LEU265
Allosteric inhibitor2	BID3	-5.68	THR308, ASP312, LYS376

Table 2. Binding sites and docking scores of **BID3** in mushroom tyrosinase (PDB ID: 2Y9X) as determined using AutoDock4.2 program

|--|

^a Binding energy indicate binding affinity and capacity for the active site of tyrosinase enzyme. ^b All amino acid residues from the enzyme–inhibitor complex were determined using AutoDock4.2 program^c Reported competitive type inhibitor. ^d Reported mixed type inhibitors. **3.2. Biological activity**

To exploit the anti-tyrosinase effects of **BID3** in the cell culture model, we examined its cytotoxic effects on B16F10 cells. Cells were treated with different concentration of **BID3** (1 – 20 μ M) for 48 h and assessed using the EZ-Cytox assay. The results indicated that **BID3** had no significant cytotoxic effect in B16F10 melanoma cells up to 20 μ M (Fig. 4A and B). Therefore, subsequent experiments were performed using **BID3** up to 20 μ M.

Melanogenesis is regulated by a tyrosinase enzymatic cascade. For this reason, several skin-whitening compounds have been developed to decrease tyrosinase activity. Kojic acid, a tyrosinase inhibitor, was used as positive control [53]. Thus, to investigate the effects of **BID3** on cAMP elevation induced hyperpigmentation, B16F10 cells were treated with α -MSH and IBMX in the presence of **BID3** (1, 5, and 20 μ M) for 48h, and melanin contents, and cellular tyrosinase activities were determined (Fig. 4C and D). Treatment with α -MSH and IBMX induced a significant increase ($P < 0.001^{\#}$) in melanin contents (Fig. 4C) and, cellular tyrosinase activity (Fig. 4D) in B16F10 cells compared with the untreated control. However, **BID3** treatment significantly ($P < 0.01^{**}$; $P < 0.001^{***}$) and concentration dependently decrease melanin contents and cellular tyrosinase activity of B16F10 cells compared with α -MSH or IBMX treated cells, indicating that the decrease in cellular melanin might be due to the inhibition of tyrosinase activity. Thus, these findings clearly shown that **BID3** exerted anti-melanogenic effects inhibiting tyrosinase activity and melanin synthesis in B16F10 cells without inducing cytotoxicity.

Indanone is one of the privileged structures of medicinal chemistry and is generally

associated with variety of pharmacologically active compounds [54]. The indanone moiety is present in variety of physiologically active natural products. In view of this, Okpekon et al., [55] reported a novel 1-indanone compound, afzeliindanone, is isolated from *Uvaria afzelii* roots. This compound is the first 1-indanone derivative isolated from plants. Nagle et al., [56] reported a new methylated indane-aldehyde was isolated from marine cyanobacterium as tumor angiogenesis regulator. Likewise, since the biological importance of indanone core, past several years researchers have been focused on developing of pharmacologically active indanone analogs as therapeutic agents. The structural diversity of 1-indanones implies variety of biological responses and these compounds may be applied in agriculture and medicine. Various indanone based compounds have been developed as anti-Alzheimer's disease [16, 57-59], anticancer [60-62], antimicrobial [63, 64], and antiviral agents [65, 66]. Due to the wide application potential, 1-indanones were interesting objects for further research and it is desirable to design new classes for their synthesis.



Fig. 4. Effects of BID3 on melanogenesis in B16F10 melanoma cells. Effect of BID3 on

B16F10 cell viability. Viability of cells treated with **BID3** (1 – 20 μ M) for 24 h (A) and 48 h (B). Cell viability was determined using EZ-Cytox solution. The melanin contents (C) and intracellular tyrosinase activity (D) of B16F10 cells were determined after incubation with **BID3** or kojic acid (10 μ M) for 48 h. Each experiment was conducted in triplicate, and the data represent the as mean ± SEM. [#]*P* < 0.05 compared with the control; ^{*}*p* < 0.05, ^{**}*p* < 0.01 and ^{***}*p* < 0.001, compared with the α -MSH and IBMX-treated.

4. Conclusion

To identify new potent tyrosinase inhibitors, we have designed and synthesized thirteen (E)benzylidene-1-indanone derivatives. compounds, Among these (E)-2-(2,4dihydroxybenzylidene)-2,3-dihydro-1H-inden-1-one (BID3) potently inhibited tyrosinase activity (IC₅₀ = 0.034 and 1.39 μ M, for L-tyrosine and L-DOPA, respectively), which was better than reference compound, kojic acid (IC₅₀ = 13.77 μ M and 33.14 μ M, respectively). Structure activity relationship revealed that presence of the dihydroxyl group at the 2 and 4position of the (E)-benzylidene-1-indanone skeleton improved the activity against tyrosinase. The mode of enzymatic inhibition of the enzyme, determined by Lineweaver-Burk and Dixon plots, indicated that **BID3** is a mixed-type inhibitor. Molecular docking studies demonstrates that binding at the active site and at allosteric sites are an important mechanisms towards tyrosinase inhibitory activity. Based on these results, **BID3** seemed to be a potent tyrosinase inhibitor for the treatment of skin disorders such as hyperpigmentation.

Conflicts of interest

The authors declare no competing interest.

Acknowledgements

This work was supported by the National Research Foundation of Korea (NRF) funded by the Korean government (Grant Nos. 2018R1A6A3A11047399 and 2018R1A2A3075425). We thank Aging Tissue Bank for providing us with research materials for the study.

Author contributions

HJJ wrote the manuscript and performed the experiments. HRM and HYC designed and interpreted data. HRM, DK, and YJ synthesized samples. SGN performed the *in silico* docking simulation.

References

[1] Boissy RE. Melanosome transfer to and translocation in the keratinocyte, Exp Dermatol,12 Suppl 2, 2003: 5–12.

[2] Gilchrest BA, Park HY, Eller MS, Yaar M. Mechanisms of ultraviolet light-induced pigmentation, Photochem Photobiol 1996;63:1–10.

[3] Lai X, Wichers HJ, Soler-Lopez M, Dijkstra BW. Structure and Function of Human Tyrosinase and Tyrosinase-Related Proteins. Chem Eur J 2018;24:47–55.

[4] Song KK, Chen QX, Wang Q, Qiu L, Huang H. Inhibitory effects of 4-vinylbenzaldehyde and 4-vinylbenzoic acid on the activity of mushroom tyrosinase. J Enzyme Inhib Med Chem 2005;20:239–43.

[5] Abu Ubeid A, Zhao L, Wang Y, Hantash BM. Short-sequence oligopeptides with inhibitory activity against mushroom and human tyrosinase. J Invest Dermatol 2009;129:2242–9.

[6] Solimine J, Garo E, Wedler J, Rusanov K, Fertig O. Hamburger M, et al. Tyrosinase inhibitory constituents from a polyphenol enriched fraction of rose oil distillation wastewater. Fitoterapia 2016;108:13–9.

[7] Menezes JCJMDS. Arylidene indanone scaffold: medicinal chemistry and structureactivity relationship view. RSC Advances 2017;7:9357–72.

[8] Li CS, Black WC, Chan CC, Ford-Hutchinson AW, Gauthier JY, Gordon R, et al. Cyclooxygenase-2 Inhibitors. Synthesis and Pharmacological Activities of 5-Methanesulfonamido-1-indanone Derivatives. J Med Chem 1995;38:4897–905.

[9] Shrestha A, Jin OH, Kim MJ, Pun NT, Magar TBT, Bist G, et al. Design, synthesis, and

structure-activity relationship study of halogen containing 2-benzylidene-1-indanone derivatives for inhibition of LPS-stimulated ROS production in RAW 264.7 macrophages. Eur J Med Chem 2017;133:121–38.

[10] Xiao S, Zhang W, Chen H, Fang B, Qiu Y, Chen X, et al. Design, synthesis, and structure-activity relationships of 2-benzylidene-1-indanone derivatives as anti-inflammatory agents for treatment of acute lung injury. Drug Des Devel Ther 2018;12:887–99.

[11] Chanda D, Bhushan S, Guru SK, Shanker K, Wani ZA, Rah BA, et al. Anticancer activity, toxicity and pharmacokinetic profile of an indanone derivative. Eur J Pharm Sci 2012;47:988–95.

[12] Singh A, Fatima K, Singh A, Behl A, Mintoo MJ, Hasanain M, et al. Anticancer activity and toxicity profiles of 2-benzylidene indanone lead molecule. Eur J Pharm Sci 2015;76:57–67.

[13] Tang ML, Zhong C, Liu ZY, Peng P, Liu XH, Sun X. Discovery of novel sesquistilbene indanone analogues as potent anti-inflammatory agents. Eur J Med Chem 2016;113:63–74.

[14] Nel MS, Petzer A, Petzer JP, Legoabe LJ. 2-Benzylidene-1-indanone derivatives as inhibitors of monoamine oxidase. Bioorg Med Chem Lett 2016;26:4599–605.

[15] Huang L, Miao H, Sun Y, Meng F, Li X. Discovery of indanone derivatives as multitarget-directed ligands against Alzheimer's disease. Eur J Med Chem 2014;87:429–39.

[16] Rampa A, Mancini F, De Simone A, Falchi F, Belluti F, Di Martino RM, et al. From AChE to BACE1 inhibitors: The role of the amine on the indanone scaffold. Bioorg Med Chem Lett 2015;25:2804–8.

[17] Piplani P, Jain A, Devi D, Anjali, Sharma A, Silakari P. Design, synthesis and pharmacological evaluation of some novel indanone derivatives as acetylcholinesterase inhibitors for the management of cognitive dysfunction. Bioorg Med Chem 2018;26:215–24.
[18] Butkevich AN, Ranieri B, Meerpoel L, Stansfield I, Angibaud P, Corbu A, et al.

Synthesis of substituted indenones and indanones by a Suzuki-Miyaura coupling/acidpromoted cyclisation sequence. Org Biomol Chem 2014;12:728–31.

[19] Gao D, Li Y. Identification and preliminary structure-activity relationships of 1-Indanone derivatives as novel indoleamine-2,3-dioxygenase 1 (IDO1) inhibitors. Bioorg Med Chem 2017;25:3780–91.

[20] Radhakrishnan S, Shimmon R, Conn C, Baker A. Inhibitory kinetics of novel 2,3dihydro-1*H*-inden-1-one chalcone-like derivatives on mushroom tyrosinase. Bioorg Med Chem Lett 2015;25:5495–9.

[21] Bukhari SN, Jasamai M, Jantan I. Synthesis and biological evaluation of chalcone derivatives (mini review). Med Chem 2012;12:1394–1403.

[22] Matos MJ, Vazquez-Rodriguez S, Uriarte E, Santana L. Potential pharmacological uses of chalcones: a patent review (from June 2011 - 2014). Expert Opin Ther Pat 2015;25:351–66.
[23] Wong E. The role of chalcones and flavanones in flavonoid biosynthesis. Phytochemistry 1968;7:1751–8.

[24] Zhang Y, Shi S, Zhao M, Jiang Y, Tu P. A novel chalcone from *Coreopsis tinctoria* Nutt.Biochem Syst Ecol 2006;34:766–9.

[25] Singh P, Anand A, Kumar V. Recent developments in biological activities of chalcones:A mini review. Eur J Med Chem 2014;85:758–77.

[26] Nerya O, Vaya J, Musa R, Izrael S, Ben-Arie R, Tamir S. Glabrene and Isoliquiritigenin as Tyrosinase Inhibitors from Licorice Roots. J Agric Food Chem 2003;51:1201–7.

[27] Nerya O, Musa R, Khatib S, Tamir S, Vaya J. Chalcones as potent tyrosinase inhibitors: the effect of hydroxyl positions and numbers. Phytochemistry 2004;65:1389–95.

[28] Seo WD, Ryu YB, Curtis-Long MJ, Lee CW, Ryu HW, Jang KC, et al. Evaluation of anti-pigmentary effect of synthetic sulfonylamino chalcone. Eur J Med Chem 2010;45:2010–

7.

[29] Liu J, Chen C, Wu F, Zhao L. Microwave-Assisted Synthesis and Tyrosinase Inhibitory Activity of Chalcone Derivatives. Chem Biol Drug Des 2013;82:39–47.

[30] Kim BH, Park KC, Park JH, Lee CG, Ye SK, Park JY. Inhibition of tyrosinase activity and melanin production by the chalcone derivative 1-(2-cyclohexylmethoxy-6-hydroxyphenyl)-3-(4-hydroxymethyl-phenyl)-propenone. Biochem Biophys Res Commun 2016;480:648–54.

[31] Kim BH, Hong SN, Ye SK, Park JY. Evaluation and Optimization of the Anti-Melanogenic Activity of 1-(2-Cyclohexylmethoxy-6-hydroxy-phenyl)-3-(4-hydroxymethylphenyl)-propenone Derivatives. Molecules 2019;24:1372.

[32] Kim HR, Lee HJ, Choi YJ, Park YJ, Woo Y, Kim SJ. Benzylidene-linked thiohydantoin derivatives as inhibitors of tyrosinase and melanogenesis: importance of the β -phenyl- α , β -unsaturated carbonyl functionality. Med Chem Comm 2014:5:1410–7.

[33] Ullah S, Park Y, Park C, Lee S, Kang D, Yang J, et al. Antioxidant, anti-tyrosinase and anti-melanogenic effects of (*E*)-2,3-diphenylacrylic acid derivatives. Bioorg Med Chem 2019;27:2192–200.

[34] Ullah S, Kang D, Lee S, Ikram M, Park C, Park Y, et al. Synthesis of cinnamic amide derivatives and their anti-melanogenic effect in alpha-MSH-stimulated B16F10 melanoma cells. Eur J Med Chem 2019;161:78–92.

[35] Jung HJ, Lee MJ, Park YJ, Noh SG, Lee AK, Moon KM, et al. A novel synthetic compound, (*Z*)-5-(3-hydroxy-4-methoxybenzylidene)-2-iminothiazolidin-4-one (MHY773) inhibits mushroom tyrosinase. Biosci Biotechnol Biochem 2018;82:759–67.

[36] Jung HJ, Lee AK, Park YJ, Lee S, Kang D, Jung YS, et al. (2*E*,5*E*)-2,5-Bis(3-hydroxy-4methoxybenzylidene) cyclopentanone Exerts Anti-Melanogenesis and Anti-Wrinkle Activities in B16F10 Melanoma and Hs27 Fibroblast Cells. Molecules 2018;23:1415.

[37] Lineweaver H, Burk D. The Determination of Enzyme Dissociation Constants. J Am

Chem Soc 1934;56:658–66.

[38] Dixon M. The determination of enzyme inhibitor constants. Biochem J 1953;55:170-1.

[39] Cornish-Bowden A. A simple graphical method for determining the inhibition constants of mixed, uncompetitive and non-competitive inhibitors. Biochem J 1974;137:143–4.

[40] Ismaya WT, Rozeboom HJ, Weijn A, Mes JJ, Fusetti F, Wichers HJ, et al. Crystal structure of *Agaricus bisporus* mushroom tyrosinase: identity of the tetramer subunits and interaction with tropolone. Biochemistry 2011;50:5477–86.

[41] Morris GM, Huey R, Lindstrom W, Sanner MF, Belew RK, Goodsell DS, et al. AutoDock4 and AutoDockTools4: Automated docking with selective receptor flexibility. J Comput Chem 2009;30:2785–91.

[42] Kim H, Roh HS, Kim JE, Park SD, Park WH, Moon JY. Compound K attenuates stromal cell-derived growth factor 1 (SDF-1)-induced migration of C6 glioma cells. Nutr Res Pract 2016;10:259–64.

[43] Bilodeau ML, Greulich JD, Hullinger RL, Bertolotto C, Ballotti R, Andrisani OM. BMP-2 stimulates tyrosinase gene expression and melanogenesis in differentiated melanocytes.
Pigmet Cell Res 2001;14:328–36.

[44] Kim MJ, Kim DS, Yoon HS, Lee WJ, Lee NH, Hyun CG. Melanogenesis inhibitory activity of Korean Undaria pinnatifida in mouse B16 melanoma cells. Interdiscip Toxicol 2014;7:89–92.

[45] Seo GY, Ha Y, Park AH, Kwon OW, Kim YJ. *Leathesia difformis* Extract Inhibits alpha-MSH-Induced Melanogenesis in B16F10 Cells via Down-Regulation of CREB Signaling Pathway. Int J Mol Sci 2019; 20:536.

[46] Kadayat TM, Banskota S, Gurung P, Bist G, Thapa Magar TB, Shrestha A, et al. Discovery and structure-activity relationship studies of 2-benzylidene-2,3-dihydro-1*H*-inden-1-one and benzofuran-3(2H)-one derivatives as a novel class of potential therapeutics for

inflammatory bowel disease. Eur J Med Chem 2017;137:575-97.

[47] Hassani S, Haghbeen K. Fazli M. Non-specific binding sites help to explain mixed inhibition in mushroom tyrosinase activities. Eur J Med Chem 2016;122:138–48.

[48] Kim CS, Noh SG, Park Y, Kang D, Chun P, Chung HY, et al. A Potent Tyrosinase Inhibitor, (*E*)-3-(2,4-Dihydroxyphenyl)-1-(thiophen-2-yl)prop-2-en-1-one, with Anti-Melanogenesis Properties in alpha-MSH and IBMX-Induced B16F10 Melanoma Cells. Molecules 2018;23:2715.

[49] Kim SJ, Yang J, Lee S, Park C, Kang D, Akter J, et al. The tyrosinase inhibitory effects of isoxazolone derivatives with a (*Z*)-beta-phenyl-alpha, beta-unsaturated carbonyl scaffold. Bioorg Med Chem 2018;26:3882–9.

[50] Lee GY, Kim JH, Choi SK, Kim YH. Constituents of the seeds of *Cassia tora* with inhibitory activity on soluble expoxide hydrolease. Bioorg Med Chem Lett 2015;25:5097–101.

[51] Yin SJ, Si YX, Qian GY. Inhibitory effect of phthalic Acid on tyrosinase: the mixed-type inhibition and docking simulations. Enzyme Res, 2011;2011:294724–30.

[52] Sohretoglu D, Sari S, Barut B, Ozel A. Tyrosinase inhibition by some flavonoids: Inhibitory activity, mechanism by *in vitro* and *in silico* studies. Bioorg Chem 2018;81:168–74.
[53] Cabanes J, Chazarra S, Garcia-Carmona F. Kojic acid, a cosmetic skin whitening agent, is a slow-binding inhibitor of catecholase activity of tyrosinase. J Pharm Pharmacol 1994;46:982–5.

[54] Patil SA, Patil R, Patil SA. Recent developments in biological activities of indanones.Eur J Med Chem 2017;138:182–98.

[55] Okpekon T, Millot M, Champy P, Gleye C, Yolou S, Bories C, et al. A novel 1-indanone isolated from *Uvaria afzelii* roots. Nat Prod Res 2009;23:909–15.

[56] Nagle DG, Zhou YD, Park PU, Paul VJ, Rajbhandari I, Duncan CJG, et al. A New

Indanone from the Marine Cyanobacterium *Lyngbya majuscula* That Inhibits Hypoxia-Induced Activation of the VEGF Promoter in Hep3B Cells. Nat Prod Res 2000;63:1431–3.

[57] Qiao JP, Gan CS, Wang CW, Ge JF, Nan DD, Pan J, et al. Novel Indanone Derivatives as Potential Imaging Probes for β -Amyloid Plaques in the Brain. Chem Bio Chem 2012;13:1652–62.

[58] Sağlık BN, Ilgın S, Özkay Y. Synthesis of new donepezil analogues and investigation of their effects on cholinesterase enzymes. Eur J Med Chem 2016;124:1026–40.

[59] van Greunen DG, Cordier W, Nell M, van der Westhuyzen C, Steenkamp V, Panayides JL, et al. Targeting Alzheimer's disease by investigating previously unexplored chemical space surrounding the cholinesterase inhibitor donepezil. Eur J Med Chem 2017;127:671–90.
[60] Gomez N, Santos D, Vazquez R, Suescun L, Mombru A, Vermeulen M, et al. Synthesis, structural characterization, and pro-apoptotic activity of 1-indanone thiosemicarbazone platinum(II) and palladium(II) complexes: potential as antileukemic agents. Chem Med Chem 2011;6:1485–94.

[61] Prakasham AP, Saxena AK, Luqman S, Chanda D, Kaur T, Gupta A, et al. Synthesis and anticancer activity of 2-benzylidene indanones through inhibiting tubulin polymerization. Bioorg Med Chem 2012;20:3049–57.

[62] Singh A, Fatima K, Srivastava A, Khwaja S, Priya D, Singh A, et al. Anticancer activity of gallic acid template-based benzylidene indanone derivative as microtubule destabilizer. Chem Biol Drug Des 2016;88:625–34.

[63] Finkielsztein LM, Castro EF, Fabian LE, Moltrasio GY, Campos RH, Cavallaro LV, et al. New 1-indanone thiosemicarbazone derivatives active against BVDV. Eur J Med Chem 2008;43:1767–73.

[64] Patel VM, Bhatt ND, Bhatt PV, Joshi HD. Novel derivatives of 5,6-dimethoxy-1indanone coupled with substituted pyridine as potential antimicrobial agents. Arab J Chem

2018;11:137-42.

[65] Patil V, Barragan E, Patil SA, Patil SA, Bugarin A. Direct Synthesis and Antimicrobial Evaluation of Structurally Complex Chalcones. ChemistrySelect 206;1:3647–50.
[66] Patil SA, Patil V, Patil R, Beaman K, Patil SA. Identification of Novel 5,6-Dimethoxyindan-1-one Derivatives as Antiviral Agents. Med Chem 2017;13:787–95.

