

Beneficial Participation of the Polymer : Improvement in Polymer-Supported Oligosaccharide Synthesis

Seema Mehta and Dennis Whitfield*

National Research Council of Canada, Institute for Biological Sciences, Ottawa, Ontario, Canada K1A 0R6.


Received 16 February 1998; revised 27 March 1998; accepted 9 June 1998

Abstract: The reagent combination scandium(III) trifluoromethanesulfonate and acetic anhydride efficiently cleaves oligosaccharides bound to the polyethylene glycol (*PEG*) polymer-support via the dioxyxylene (*DOX*) linker. Unexpectedly, the site of cleavage is between the *DOX* linker and the *PEG* polymer. This method of cleavage allows for the use of *DOX* as a linker for synthesis on polystyrene-polyethylene glycol, *PS-PEG* beads. Crown copyright © 1998 Published by Elsevier Science Ltd. All rights reserved.

Polymer-supported synthesis is an emerging tool for the production of oligosaccharides.^{1,2,3} The effective use of the polymer-support polyethylene glycol (*PEG*) for such syntheses is well demonstrated.^{1,3} The dioxyxylene (*DOX*) linker, introduced by Krepinsky *et al.*, has been efficiently employed for the synthesis of up to a pentasaccharide.⁴ This linker enables the use of base labile acyl groups as cleavable protecting groups. In spite of the considerable efforts to improve *PEG*-based oligosaccharide synthesis, a number of challenges remain. An increase in the repertoire of orthogonal protecting groups, as well as an increase in the efficiency of the glycosylation reactions are required.^{1,3}

The propensity of *PEG* derivatives to complex metal cations has been demonstrated.⁵ It is known that a number of conventional metal based reagents render *PEG* into an unfilterable slurry. This makes the usual purification procedure of precipitation and reprecipitation impossible.⁶ Oligosaccharides linked via the benzyl-like *DOX* linker, to *PEG* can be released by Pd, Pt or Ni catalyzed hydrogenation.⁴ However, hydrogenation is known to be inhibited by *PEG* derivatives and is substrate specific.⁷ This paper describes the novel use of Sc(OTf)₃ for the cleavage of the *DOX* linker from the *PEG* polymer-support. This reaction likely proceeds via the coordination of the Sc³⁺ ions to the *PEG*, and represents a new type of reaction for polymer-supported chemistry, that is based on reagent-polymer interactions.

As a part of the process towards developing vaccines against Group B *Streptococcus* bacteria, we are involved in a program to synthesize oligosaccharide fragments of the serotype specific capsular polysaccharides.⁸ In order to take advantage of the ease of purification of polymer bound products, we have employed polymer-supported methods based on the soluble polymer monomethyl polyethylene glycol (MeO-(CH₂CH₂O)_nOH MW 5,000), *MPEG*.¹ Our initial test substrate was the β1,3-disaccharide bound to the *MPEG* polymer-support via the *DOX* linker **1** (Scheme 1). In this example the *DOX* linker has been modified by mono-α-methyl substitution to give *MDOX*. Stimulated by the report that FeCl₃ promoted benzyl ether cleavage,⁹ we tested the cleavage of **1** with FeCl₃ and Ac₂O.¹⁰ Although these conditions did lead to the cleavage of the disaccharide from the polymer, they resulted in decomposition. The use of the hard Lewis acid Sc(OTf)₃ was next examined. The combination Sc(OTf)₃/Ac₂O did promote the cleavage of the disaccharide from the polymer-support. To our surprise, the site of bond cleavage was between the *MPEG* terminal oxygen and the benzylic carbon of the *DOX* linker, and not the expected site between the oligosaccharide and the *DOX* linker (Scheme 1). Thus, the free β1,3-disaccharide **2** was obtained. The cleavage was complete as only traces of the sugar bound to *MPEG* could be detected by ¹H NMR. Hydrogenation of the diastereomers **2** afforded the reducing disaccharide **3**,¹¹ a derivative that can be




Scheme 1.

easily converted into a disaccharide donor.

Since this reagent combination has been reported to be an effective acylating system, the free hydroxyl group at the 4 position was acetylated under these reaction conditions.¹²


This unexpected position of cleavage led us to experiment with these cleavage conditions on the polymer-supported disaccharide 4. The reaction of 4 with $\text{Sc}(\text{OTf})_3/\text{Ac}_2\text{O}$ effected cleavage at the *MPEG-DOX*, C-O bond, and not at the anomeric position, to afford the free disaccharide 5.¹³



Scheme 2.

These results led us to examine oligosaccharide synthesis using polystyrene-polyethylene glycol (*PS-PEG*) beads. Thus, α,α -dichloroxylylene 6, was attached to *PS-PEG-OH* with NaH in tetrahydrofuran, followed by hydrolysis to give 7. The alcohol 7 was glycosylated with 5 equivalents of the glycosyl donor 8¹⁴ under TESOTf promotion, in dichloromethane to afford the resin bound monosaccharide 9. Upon cleavage with $\text{Sc}(\text{OTf})_3/\text{Ac}_2\text{O}$, the expected free monosaccharide 10 and the diacetate of *DOX* 11, were obtained (about 3:1). The formation of the diacetate 11 suggested that no side reactions had occurred but that some sites on the beads had remained unreacted. Repetitive treatments or even more vigorous glycosylation conditions did not improve the yield, suggesting that some sites on the beads are not accessible to the glycosyl donor. Deacetylation of 13 in the presence of 1,8-diazabicyclo[5.4.0] undec-7-ene afforded the *PS-PEG* bound triol 12. The glycosylation of the triol 12 with glycosyl donor 13, under promotion by *N*-iodosuccinimide / triflic acid, afforded the resin


bound disaccharide 14.¹⁵ Upon cleavage from the resin with $\text{Sc}(\text{OTf})_3/\text{Ac}_2\text{O}$, the free disaccharide 15 was obtained in 10% yield for 6 steps, based on 100% derivatization of the beads.¹⁶ Traces of monosaccharide 10 and the diacetate 11 were obtained as well as traces of other regio- and stereoisomeric disaccharides. Thus, this new method of cleavage allows each step of the solid-phase synthesis to be easily monitored. It also allows the combination of the stable DOX linker, and base labile acyl protecting groups to be used with PS-PEG beads.



Scheme 3.

Typically 0.5 equivalents of $\text{Sc}(\text{OTf})_3$ were used. Some cleavage could be detected even after 1 min, however 2-4 hours were necessary to obtain complete cleavage.¹⁷ In the case of soluble MPEGDOX bound substrates, MPEGOAc was the major by-product.

The dominant reaction at the MPEGDOX ether linkage is highly suggestive of a complexation mechanism,¹⁸ since other benzylic ethers, such as the electronically activated MDOX (Scheme 1) were stable. The observation of the cleavage of the DOX linker from sites that are inaccessible to the glycosyl donor suggest that steric effects are also not important. Furthermore, two control experiments support the premise of a complexation driven mechanism. Firstly, $\text{Sc}(\text{OTf})_3$ is insoluble in dichloromethane but is freely soluble in the presence of MPEG. Secondly, the subjection of 1,6-di-*O*-acetyl-2,3,4-tri-*O*-benzyl-D-glucopyranoside to $\text{Sc}(\text{OTf})_3/\text{Ac}_2\text{O}$ conditions for 90 hours, in the absence of MPEG, led to its complete decomposition. However, in an analogous reaction conducted in the presence of 1 equivalent of MPEG, the sugar was recovered intact. The benzylic protecting groups were not cleaved under these reaction conditions. These observations suggest that the Sc(III) ion forms a complex with the PEG in such a way that its reactivity is tuned to the cleavage of the DOX-PEG, C-O linkage only (Scheme 4).



Scheme 4.

Thus, we have not only discovered a useful reagent for polymer-supported oligosaccharide synthesis, but we have discovered a new type of reaction that utilizes the complexation properties of the polymer-support to promote a useful reaction. It can be envisaged that the deliberate design of polymeric-supports that complex reagents is now feasible. Furthermore, this reaction opens up the use of PEGDOX combinations for so called "traceless" linker synthetic applications to compounds other than oligosaccharides.¹⁹

References

- Douglas, S. P.; Whitfield, D. M.; Krepinsky, J. J. *J. Am. Chem. Soc.* **1991**, *113*, 5095-5097.
- (a) Liang, R.; Yan, L.; Loebach, J.; Ge, M.; Uozumi, Y.; Sekanina, K.; Horan, N.; Gildersleeve, J.; Thompson, C.; Smith, A.; Biswas, K.; Still, W. C.; Kahne, D. *Science* **1996**, *274*, 1520-1522. (b) Rademann J.; Schmidt, R. R. *Tetrahedron Lett.* **1996**, *37*, 3989-3990. (c) Halcomb, R. L.; Huang, H.; Wong, C. H. *J. Am. Chem. Soc.* **1994**, *116*, 11315-11322. (d) Danishefsky, S. J.; McClure, K. F.; Randolph, J. T.; Ruggeri, R. B. *Science* **1993**, *260*, 1307-1309. (e) Nicolau, K. C.; Winssinger N.; Pastor, J.; DeRoose F. *J. Am. Chem. Soc.* **1997**, *119*, 449-450. (f) Rodebaugh, R.; Joshi, S.; Fraser-Reid, B.; Geysen, H. M. *J. Org. Chem.* **1997**, *62*, 5660-5661.
- (a) Ito, Y.; Ogawa, T. *J. Am. Chem. Soc.* **1997**, *119*, 5562-5566. (b) Dreef-Tromp, C. M.; Willems, H. A. M.; Westerduin, P.; van Veelen, P.; van Boeckel, C. A. A. *Bioorganic Med.Chem. Lett.* **1997**, *7*, 1175-1180. (c) Wang, Z. G.; Douglas, S. P.; Krepinsky, J. J. *Tetrahedron Lett.* **1996**, *39*, 6985-6988.
- Douglas, S. P.; Whitfield, D. M.; Krepinsky, J. J. *J. Am. Chem. Soc.* **1995**, *117*, 2116-2117.
- (a) Whitfield, D. M.; Douglas, S. P.; Krepinsky, J. J. *Tetrahedron Lett.* **1992**, *33*, 6795-6798. (b) Bernson, A.; Lindgren, J.; Huang, W.; Frech, R. *Polymer*, **1995**, *36*, 4471-4478. (c) Rogers, R. D.; Zhang, J.; Bauer, C. B. *J. Alloys Compounds* **1997**, *249*, 41-48, and references therein.
- Harris, J. M. *Rev. Macromol. Chem. Phys.* **1985**, *C25*, 325-373.
- Bergbreiter, D. E.; Kimmel, T.; Caraway, J. W. *Tetrahedron Lett.* **1995**, *36*, 4757-4760.
- Jennings H. J. *Curr. Top. Microbiol. Immunol.* **1990**, *150*, 97-127.
- (a) Ganem, B.; Small Jr., V. R. *J. Org. Chem.* **1974**, *39*, 3728-3730. (b) Bhar, S.; Ranu, B. S. *J. Org. Chem.* **1995**, *60*, 745-754. (c) Rodebaugh, R.; Debenham, J. S.; Fraser-Reid, B. *Tetrahedron Lett.* **1996**, *37*, 5477-5478.
- Ranu, B. C.; Bhar, S. *Org. Prep. Proc. Int.* **1996**, *28*, 371-409.
- 3**: [α]²⁰_D: 19.2° (c 0.13, CHCl₃); ¹H NMR (CDCl₃): δ 5.67 (1H, brd, *J* = 2.7 Hz, H-4), 5.53 (1H, d, *J* = 8.2 Hz, H-1'), 5.45 (1H, d, *J* = 3.8 Hz, H-1), 4.15 (1H, m, H-3); HRMS (FAB) calcd for C₄₂H₄₀NO₁₇: 830.2294, found *m/z* 830.2309 (M+ - OH).
- Ishihara, K.; Kubota, M.; Kurihara, H.; Yamamoto, H. *J. Org. Chem.* **1996**, *61*, 4560-4567.
- 5**: [α]²⁰_D: -25° (c 0.1, CHCl₃). ¹H NMR (CDCl₃): δ 4.98 (2H, s, CH₂ of DOX), 4.89 (1H, d, *J* = 7.9 Hz, H-1'), 4.41-4.47 (4H, m, H-1), 4.07- 4.12 (2H, m, H-4, 6'), 3.78 (3H, s, OCH₃), 3.55-3.58 (2H, m, H-5, 3'), ¹³C NMR (CDCl₃): 100.7 (C-1'), 99.1 (C-1); MS(MALDI- TOF) calcd for C₅₂H₅₆O₂₀Na: 1024, found *m/z*: 1024.6 (M + Na+).
- Lemieux, R. U.; Takeda, T.; Chung, B. Y. *Synthetic Methods for Carbohydrates, ACS Symp. Series* **1976**, *39*, 90-100.
- Veeneman, G. H.; van Leeuwen, S. H.; van Boom, J. H. *Tetrahedron Lett.* **1990**, *31*, 1331-1334.
- 15**: [α]²⁰_D: 58.7° (c 0.53, CHCl₃); ¹H NMR (CDCl₃): δ 5.70 (1H, dd, *J* = 8.9, 10.4 Hz, H-3), 5.21 (1H, d, *J* = 8.5 Hz, H-1), 4.94 (2H, m, H-1', 4), 4.40 (1H, d, *J* = 12.5 Hz, CHH of DOX), 4.12 (1H, d, *J* = 12.5 Hz, CHH of DOX); ¹³C NMR (CDCl₃): δ 101.5 (C-1'), 96.8 (C-1); HRMS (FAB+) calcd for C₆₂H₅₅NO₁₉Na: 1156.3215, found *m/z* 1156.3181 (M + Na+).
- On an analytically scale to 15-30 mg of resin is added sequentially CH₂Cl₂ (0.5 mL), Ac₂O (0.5 mL) and Sc(OTf)₃ (2-3 mg). This mixture is stirred for 3-4 hours, cooled, quenched with TBME, and filtered. The filtrate is evaporated and analysed by TLC, NMR, MS etc. On a larger scale silica gel chromatography can be used to purify the products.
- (a) Castellani, C. B.; Carugo, O.; Giusti, M.; Leopizzi, C.; Perotti, A.; Invernizzi, A.G.; Vidari, G. *Tetrahedron* **1996**, *52*, 11045-11052. (b) Kobayashi, S.; Moriwaki, M.; Hachiya, I. *Bull. Chem. Soc. Jpn.* **1997**, *70*, 267-273. (c) Tsuchimoto, T.; Tobita, K.; Hiyama, T.; Fukuzawa, S. J. *J. Org. Chem.* **1997**, *62*, 6997-7005.
- (a) Plunkett, M. J.; Ellman, J. A. *J. Org. Chem.* **1997**, *62*, 2885-2893. (b) Deegan, T. L.; Gooding, O. W.; Baudart, S.; Porco Jr., J. A. *Tetrahedron Lett.* **1997**, *38*, 4973-4976. (c) Atrash, B.; Bradley, M. *Chem. Commun.* **1997**, 1397-1398.