

A convenient route to vinylsiloxane-tertiary phosphine-nickel(0) complexes; the molecular structure of $[(\text{Ni}\{\text{P}(\text{C}_6\text{H}_4\text{Me-4})_3\})_2\{\mu\text{-(L''L'')}\}_2]\{\text{(L''L'')}_2 = [\text{CH}_2=\text{CH}(\text{Me})\text{Si}(\mu\text{-O})]_4\}$

Peter B. Hitchcock, Michael F. Lappert *, Hieronim Maciejewski ¹

The Chemical Laboratory, University of Sussex, Brighton BN1 9QJ, UK

Received 18 April 2000; accepted 18 May 2000

Abstract

A simple one pot synthesis in Et₂O at ambient temperature, from the readily available starting materials [Ni(cod)₂], PR₃ and LL or (L''L'')₂, provides an essentially quantitative route to the known vinylsiloxanenickel(0) complexes $[\text{Ni}(\overline{\text{LL}})\text{PR}_3]$ and the new binuclear analogues $[(\text{Ni}(\text{PR}_3))_2\{\mu\text{-(L''L'')}\}_2]$ (**5**) $\{\text{LL} = [\text{CH}_2=\text{CH}(\text{Me})_2\text{Si}]_2\text{O}$, (L''L'')₂ = $[\text{CH}_2=\text{CH}(\text{Me})\text{Si}(\mu\text{-O})]_4$ and R = Ph, C₆H₄Me-4 or C₆H₁₁-c}. The X-ray crystal structure of **5b** (R = C₆H₄Me-4) shows it to be a centrosymmetric binuclear complex containing (L''L'')₂ as a chair-shaped bridging ligand. It is bound to each Ni(PR₃) moiety by a pair of μ²-vinyl groups (having M and M' as centroids) and each NiMSiOSi'M' metallacycle is also of chair conformation. © 2000 Elsevier Science S.A. All rights reserved.

Keywords: Nickel; Vinylsiloxane complexes

1. Introduction

Hydrosilylation is widely used in the silicone industry for the synthesis of monomers containing Si–C bonds, the cross-linking of polymers and as a source of various silane coupling reagents, while in organic chemistry it has a role for the synthesis of alcohols by reduction of carbonyl compounds [1]. Such processes generally require the use of a highly active platinum-containing catalyst. The silicone-soluble Karstedt catalyst is commercially important for hydrosilylation. It is obtained, in presence of the silane, by reaction of chloroplatinic acid H₂[PtCl₆]_xH₂O (**1**) (Speier's catalyst) with a vinyl-silicon-containing compound, such as sym-divinyltetra(methyl)disiloxane $[\text{CH}_2=\text{CH}(\text{Me})_2\text{Si}]_2\text{O}$ (≡LL) [2,3].

Our earlier studies had established that **1** and LL in the absence of a silane yielded a solution **A** [4], from which the crystalline platinum(0) complex $[(\text{Pt}(\text{LL}))_2(\mu\text{-(L''L'')})_2]$ (**2**) was isolated [5]. Complex **2** was shown to be a Karstedt catalyst; with styrene as substrate, transient 14-electron intermediates $[\text{Pt}(\text{LL})(\text{LL})]$ and $[\text{Pt}(\text{LL})(\eta^2\text{-CH}_2=\text{CHPh})]$ were identified by NMR spectroscopy [6]. Complex **2** was also shown to be a convenient source of further well characterised siloxaneplatinum(0) complexes, such as $[\text{Pt}(\text{LL})(\text{PR}_3)]$ [4,7], $[(\text{Pt}(\text{LL}))_2(\mu\text{-dppe})]$ [8] and $[(\text{Pt}(\text{LL}))_3(\mu_3\text{-triphos})]$ [8] [dppe = (Ph₂PCH₂)₂, triphos = (Ph₂PCH₂)₃CMe or (Ph₂PCH₂CH₂)₂PPh]. A series of palladium(0) complexes $[(\text{Pd}(\text{LL}))_2(\mu\text{-LL})]$ (from $[\text{PdCl}_2(\text{cod})_2]$, Li₂cot and LL) and $[\text{Pd}(\text{LL})\text{D}]$ (D = C₂H₄ or PR₃; R = Me, ^tPr, ^tBu, Ph or C₆H₄Me-2) has also been reported [9].

The high cost of platinum has led to much research on various nickel complexes as potential hydrosilylation catalysts [1]. Also relevant to the present study is the fact that tertiary phosphine ligands have a pervasive place in many industrially important noble metal-catalysed organic reactions, including hydrosilylation; complexes of Rh(I), Pd(0 or II) and Pt(0) are particularly prominent [10]. The other pertinent ligand component for this report relates to a vinylsilane. Previous

The high cost of platinum has led to much research on various nickel complexes as potential hydrosilylation catalysts [1]. Also relevant to the present study is the fact that tertiary phosphine ligands have a pervasive place in many industrially important noble metal-catalysed organic reactions, including hydrosilylation; complexes of Rh(I), Pd(0 or II) and Pt(0) are particularly prominent [10]. The other pertinent ligand component for this report relates to a vinylsilane. Previous

* Corresponding author. Fax: +44-1273-677196.

E-mail address: m.f.lappert@sussex.ac.uk (M.F. Lappert).

¹ Present address. Adam Mickiewicz University, 60–780 Poznan, Poland.

work dealing with these parameters concerned the formation of (a) $[\text{Ni}(\text{LL})(\text{PR}_3)]$ ($\text{R} = \text{Ph}$, $\text{C}_6\text{H}_4\text{Me-4}$ or $\text{C}_6\text{H}_{11}\text{-c}$) (**3**) from *trans,trans,trans*-cyclododecatrienickel(0) $\{\equiv[\text{Ni}(\text{CDDT})]\}$ and LL [7,11], and (b) $[\text{Ni}(\text{LL})(\text{PPh}_3)]$ or $[\text{Ni}(\text{LPhL}^{\text{Ph}})(\text{PPh}_3)]$ from $[\text{NiCl}_2(\text{PPh}_3)_2]$, zinc dust and LL or $\text{L}^{\text{Ph}}\text{L}^{\text{Ph}}$ $\{\equiv[\text{CH}_2=\text{CH}(\text{Ph})_2\text{Si}]_2\text{O}\}$ [7]. The compound $[\text{Ni}\{\text{CH}_2=\text{CH}(\text{Me})_2\text{SiOSi}(\text{Me})_2\text{CH}=\text{CHSi}(\text{Me})_2\text{OSi}(\text{Me})_2\text{CH}=\text{CH}_2\}]$ was obtained from nickel atoms and LL under metal vapour synthesis conditions [12]. Other vinylsilanes which have been employed as ligands in Group 10 metal chemistry include $(\text{CH}_2=\text{CH})_2\text{SiMe}_2$ ($\equiv\text{L}'\text{L}'$) and $[\text{CH}_2=\text{CH}(\text{Me})\text{Si}(\mu\text{-O})]_4$ [$\equiv(\text{L}'\text{L}')_2$] (**4**). Thus, $[\text{NiCl}_2(\text{PPh}_3)_2]$, zinc dust and $\text{L}'\text{L}'$ afforded $[\{\text{Ni}(\mu\text{-L}'\text{L}')(\text{PPh}_3)_2\}_2]$ [8]; and **4** has been used in platinum chemistry [7,13] to generate hydrosilylation catalysts related to **2** [13].



2. Results and discussion

2.1. Synthetic studies

Our objective was to develop convenient syntheses for nickel(0) complexes having η^2 -vinylsilicon and ter-

tiary phosphine ligands, study their structures and explore their potential as hydrosilylation catalysts; catalytic aspects will be considered in a later paper.

An improved synthesis (cf. [7] and [11]) of the complexes **3** was accomplished, using the readily available $[\text{Ni}(\text{cod})_2]$ as precursor (Eq. 1). The displaced volatile cyclooctadiene was easily removed in vacuo, providing the essentially pure residue of the appropriate complex **3** in quantitative yield. Each of the crystalline complexes **3a**, **3b** and **3c** was authenticated by showing it to have identical NMR spectra to those of the previously characterised specimens [7].



In a similar fashion, but using cyclotetrakis[vinyl(methyl)siloxane] (**4**) [$\equiv(\text{L}'\text{L}')_2$] in place of LL, there were obtained the binuclear nickel(0) complexes **5**, Eq. (2). Each of the crystalline, yellow complexes **5a**, **5b** and **5c** gave satisfactory analyses and appropriate NMR spectra. The various $^1\text{H}\text{-}^1\text{H}$, and $^1\text{H}\text{-}^{31}\text{P}$ and $^{31}\text{P}\text{-}^{29}\text{Si}$ coupling constants, Fig. 2, were similar for each of **5a–5c**, and also to those found for **3a–3c** [7]. Like their analogues **3** [7], it is interesting that **5c** is the rac-diastereoisomer in the solid state.








Fig. 1. The molecular structure of $[(\text{Ni}\{\text{P}(\text{C}_6\text{H}_4\text{Me-4})_3\}_2[\text{CH}_2=\text{CH}(\text{Me})\text{Si}(\mu\text{-O})]_4$ (**5b**) with atom labelling.




Fig. 2. Schematic representation of $^1\text{H}\text{-}^{31}\text{P}$ and $^{31}\text{P}\text{-}^{29}\text{Si}$ coupling constants (Hz), from ^1H - and ^{29}Si -NMR spectra, for **5a**, **5b**, **5c**, respectively in C_6D_6 at 298 K.

Table 1
Selected bond lengths (Å) and angles (°) for **5b**^a

<i>Bond lengths</i>			
Ni–P	2.180(1)	Ni–C(1)	2.008(3)
Ni–C(2)	2.005(3)	Ni–C(3)	2.023(3)
Ni–C(4)	2.000(3)	Ni–M(1)	1.882(3)
Ni–M(2)	1.887(3)	C(1)–C(2)	1.393(4)
C(3)–C(4)	1.392(4)		
<i>Bond angles</i>			
M(1)–Ni–M(2)	131.85(11)	M(1)–Ni–P	111.18(9)
M(2)–Ni–P	116.00(9)	Si(1)–O(1)–Si(2)	129.08(11)
Si(1)–O(2)–Si(2) ^c	148.67(12)	C(2)–C(1)–Si(1)	122.4(2)
C(2)–C(1)–Ni	69.6(2)	Si(1)–C(1)–Ni	113.05(13)
C(1)–C(2)–Ni	69.8(2)	C(4)–C(3)–Si(2)	123.4(2)
C(4)–C(3)–Ni	68.9(2)	Si(2)–C(3)–Ni	113.03(13)
C(3)–C(4)–Ni	70.7(2)	C(12)–C(7)–C(8)	117.5(2)

^a M(1) and M(2) are the centres of the C(1)–C(2) and C(3)–C(4) bonds, respectively. Symmetry transformations used to generate equivalent atoms: $-x, -y, -z$.

A platinum analogue of **5c**, prepared from $[\text{Pt}(\text{LL})\{\text{P}(\text{C}_6\text{H}_{11}\text{-c})_3\}]$ and **4**, had been reported [4]; its structure was conjectured to be similar to that now established for **5b** (see Section 2.2).

2.2. The X-ray molecular structure of $[\{\text{Ni}\{\text{P}(\text{C}_6\text{H}_4\text{Me-4})_3\}_2\{\mu\text{-(L''L'')}_2\}]$ (**5b**)

Crystalline **5b**, obtained from benzene, is a centrosymmetric binuclear nickel(0) complex, the inversion centre being the mid-point of the chair-shaped $(\text{SiO})_4$ ring, Fig. 1. Selected bond lengths and angles are listed in Table 1. There is a molecule of benzene solvate in a general position.

The local geometry at each three-coordinate nickel atom is planar, taking its vertices to be the phosphorus atom and the centroids M of adjacent η^2 -vinyl groups. The M(1)–Ni–M(2) angle is wider [131.8(1)°] than M(1)–Ni–P [111.2(1)°] or M(2)–Ni–P [116.0(1)°]. This situation is similar to that in $[\text{Ni}(\text{LL})\{\text{P}(\text{C}_6\text{H}_{11}\text{-c})_3\}]$, the corresponding angles being 130.1(1), 113.9(1) and 115.1(1)° [7]. Likewise, the Ni–P and average Ni–C_α and Ni–C_β bond lengths in that compound of 2.206(1), 2.000(3) and 2.027(3) Å are close to the 2.179(1), 2.003(3) and 2.015(3) Å in **5b**. Each six-membered metallacyclic ring $[\text{NiM}(1)\text{Si}(1)\text{OSi}(2)\text{M}(2)]$ (**5b**) adopts a chair conformation, as had previously been observed for (Met)M(1)Si(1)OSi(2)M(2) rings in $\text{Ni}(\text{LL})$ [7], $\text{Pt}(\text{LL})$ [4,5,7,8], $\text{Rh}(\text{LL})$ [14] and $\text{Rh}(\text{LPhL}^{\text{Ph}})$ [14] complexes (Met = Ni, Pt or Rh).

3. Experimental

3.1. General procedures and starting silicon reagents

All manipulations and instruments were as described earlier [7,8,14]. The silicon reagents used in Sections 3.2, 3.3, 3.4 and 3.5 were kindly provided by Dow Corning, Ltd.

3.2. Synthesis of

$[\{\text{Ni}(\text{PPh}_3)_2\{\mu\text{-(}\eta\text{-CH}_2\text{=CH(Me)Si}(\mu\text{-O))}_4\}]$ (**5a**)

Cyclotetakis[vinyl(methyl)siloxane] (1.0 ml) was added slowly to a stirred red suspension of $[\text{Ni}(\text{cod})_2]$ (0.50 g, 1.8 mmol) and PPh_3 (0.47 g, 1.8 mmol) in diethyl ether (10 ml) at ambient temperature, yielding a yellow solution. The reaction mixture was allowed to stir overnight. The volatiles were removed in vacuo to give a yellow oil, which was extracted into pentane (2 × 2 ml). The extract was filtered through Celite. The filtrate was concentrated, then set aside at -30°C to yield the yellow solid compound **5a** (0.77 g, 0.8 mmol, 86%). Yellow crystals of **5a**, m.p. 212–214°C (dec) were obtained by recrystallisation from benzene, at ambient temperature. Anal. Found: C, 58.3; H, 5.49. $\text{C}_{48}\text{H}_{54}\text{Ni}_2\text{O}_4\text{P}_2\text{Si}_4$ Calc.: C, 58.4; H, 5.52%. $^1\text{H-NMR}$ (C_6D_6 , 298 K, 500 MHz): δ 0.39 (s, 12H), 2.56 (td, 4H), 3.07 (dd, 8H), 7.03–7.64 (m, 30H), $^3J(^1\text{H}_1\text{-}^1\text{H}_2) = 12.6$ Hz, $^3J(^1\text{H}_1\text{-}^1\text{H}_3) = 16.5$ Hz, $^3J(^1\text{H}_1\text{-}^{31}\text{P}) = 4.8$ Hz, $^3J(^1\text{H}_2\text{-}^{31}\text{P}) = 7.9$ Hz, $^3J(^1\text{H}_3\text{-}^{31}\text{P}) = 6.4$ Hz. $^{13}\text{C}\{^1\text{H}\}\text{-NMR}$ (C_6D_6 , 298 K, 125.8 MHz): δ 0.61 (s, Me), 58.17 (s, =CH–), 63.15 (s, =CH₂), 127.81–136.60 (m, Ph). $^{29}\text{Si}\{^1\text{H}\}\text{-NMR}$ (C_6D_6 , 298 K, 99.4 MHz): δ –24.0 (d), $^3J(^{29}\text{Si}\text{-}^{31}\text{P}) = 4.0$ Hz. $^{31}\text{P}\{^1\text{H}\}\text{-NMR}$ (C_6D_6 , 298 K, 101.2 MHz): δ 41.2 (s).

3.3. Synthesis of $[\{\text{NiP}(\text{C}_6\text{H}_4\text{Me-4})_3\}_2\{\mu\text{-(}\eta\text{-CH}_2\text{=CH(Me)Si}(\mu\text{-O))}_4\}]$ (**5b**)

Compound **5b** (0.83 g, 0.8 mmol, 85%) was prepared in a similar manner as **5a**, except that $\text{P}(\text{C}_6\text{H}_4\text{Me-4})_3$ was used in place of PPh_3 . Yellow crystals of **5b**, m.p. 224–226°C (dec) suitable for X-ray diffraction, were grown from benzene at ambient temperature over a period of 24 h. Anal. Found: C, 60.7; H, 6.54. $\text{C}_{54}\text{H}_{66}\text{Ni}_2\text{O}_4\text{P}_2\text{Si}_4$ Calc.: C, 60.6; H, 6.21%. $^1\text{H-NMR}$ (C_6D_6 , 298 K, 500 MHz): δ 0.48 (s, 12H), 2.03 (s, 9H), 2.61 (td, 4H), 3.16 (dd, 8H), 6.93–7.65 (m, 24H), $^3J(^1\text{H}_1\text{-}^1\text{H}_2) = 12.1$ Hz, $^3J(^1\text{H}_1\text{-}^1\text{H}_3) = 16.4$ Hz, $^3J(^1\text{H}_1\text{-}^{31}\text{P}) = 4.6$ Hz, $^3J(^1\text{H}_2\text{-}^{31}\text{P}) = 6.0$ Hz, $^3J(^1\text{H}_3\text{-}^{31}\text{P}) = 10.5$ Hz. $^{13}\text{C}\{^1\text{H}\}\text{-NMR}$ (C_6D_6 , 298 K, 125.8 MHz): δ 0.69 (s, Me), 21.01 (s, Me), 57.68 (s, =CH–), 63.00 (s, =CH₂), 127.80–139.22 (m, tol). $^{29}\text{Si}\{^1\text{H}\}\text{-NMR}$ (C_6D_6 , 298 K, 99.4 MHz): δ –23.9 (d), $^3J(^{29}\text{Si}\text{-}^{31}\text{P}) = 3.8$ Hz. $^{31}\text{P}\{^1\text{H}\}\text{-NMR}$ (C_6D_6 , 298 K, 101.2 MHz): δ 39.1 (s)

3.4. Synthesis of $[\{Ni(P(C_6H_{11}-c)_3)_2-\mu-(\eta-CH_2=\overline{CH(Me)Si}(\mu-O))_4\}]$ (**5c**)

Compound **5c** (0.70 g, 0.7 mmol, 76%) was prepared in similar manner as **5a**, except that $P(C_6H_{11}-c)_3$ was used in place of PPh_3 . Yellow crystals of **5c**, m.p. 235–237°C (dec) were obtained by recrystallisation from benzene, at ambient temperature. Anal. Found: C, 57.3; H, 9.01. $C_{48}H_{90}Ni_2O_4P_2Si_4$ Calc.: C, 56.4; H, 8.87%. 1H -NMR (C_6D_6 , 298 K, 500 MHz): δ 0.51 (s, 12H), 1.1–1.8 (m, 33H), 2.53 (td, 4H), 3.06 (dd, 8H), $^3J(^1H_1-^1H_2)$ = 12.6 Hz, $^3J(^1H_1-^1H_3)$ = 16.3 Hz, $^3J(^1H_1-^{31}P)$ = 3.4 Hz, $^3J(^1H_2-^{31}P)$ = 5.6 Hz, $^3J(^1H_3-^{31}P)$ = 8.2 Hz. $^{13}C\{^1H\}$ -NMR (C_6D_6 , 298 K, 125.8 MHz): δ 0.65 (s, Me), 26.99–36.54 (m, $C_6H_{11}-c$), 55.60 (s, $=CH-$), 58.95 (s, $=CH_2$). $^{29}Si\{^1H\}$ -NMR (C_6D_6 , 298 K, 99.4 MHz): δ –23.6 (d), $^3J(^{29}Si-^{31}P)$ = 3.2 Hz. $^{31}P\{^1H\}$ -NMR (C_6D_6 , 298 K, 101.2 MHz): δ 39.4 (s).

3.5. Typical synthesis of

$[Ni\{\mu-CH_2=\overline{CH(Me)_2Si}\}_2O(PR_3)]$ (**3**)

Divinyltetramethyldisiloxane (1.0 ml) was added slowly to a stirred red suspension of $[Ni(cod)_2]$ (0.56 g, 2.5 mmol) and PR_3 (2.5 mmol; R = Ph, C_6H_4Me-4 or $C_6H_{11}-c$) in diethyl ether (10 ml) at ambient temperature, yielding a yellow solution. The reaction mixture was allowed to stir overnight and the volatiles were removed under reduced pressure to yield a yellow oil. This was taken up into pentane (2×2 ml) and filtered through Celite. Concentration of the filtrate and cooling to $-30^\circ C$ yielded yellow crystals of **3**. Each of the crystalline complexes $[Ni\{\mu-CH_2=\overline{CH(Me)_2Si}\}_2O(PPh_3)]$, (**3b**), $[Ni\{\mu-CH_2=\overline{CH(Me)_2Si}\}_2O\{P(C_6H_4Me-4)_3\}]$ **3a** and $[Ni\{\mu-CH_2=\overline{CH(Me)_2Si}\}_2-$

$O\{P(C_6H_{11}-c)_3\}]$ (**3c**) was authenticated by showing it to have identical NMR spectra to those of the previously characterised specimens [7].

3.6. X-ray structure determination of **5b**

Intensities were measured on a Enraf–Nonius CAD 4 diffractometer with monochromated Mo– K_α radiation ($\lambda = 0.71073$ Å). Structure solution was by SHELXS-86 [15]. Refinement was by full-matrix least-squares on F^2 using all reflections and SHELXL-93 [16]. Non-hydrogen atoms were refined anisotropically. Hydrogen atoms were included in riding mode with $U_{iso}(H)$ equal to $1.2U_{eq}(C)$ or $1.5U_{eq}(C)$ for methyl groups. The 4-tolyl methyl H atoms were included as disordered equally over two sets of positions related by a 60° rotation. Further details are in Table 2.

4. Supplementary material

Crystallographic data for the structural analysis have been deposited at the Cambridge Crystallographic Data Centre: CCDC no. 142037. Copies of this information may be obtained free of charge from The Director, CCDC, Cambridge CB2 1EZ, UK (fax: +44-1223-336-033; e-mail: deposit@ccdc.cam.ac.uk or www: http://www.ccdc.cam.ac.uk).

Acknowledgements

We thank Dow Corning Ltd. (Barry) for gifts of chemicals, the Royal Society for the award of a R.S.-NATO postdoctoral fellowship to H.M. and EPSRC for other support

Table 2
Crystal data and structural refinement parameters for **5b**

Formula	$C_{54}H_{66}Ni_2O_4P_2Si_4 \cdot 2(C_6H_6)$
Formula weight	1227.0
Crystal system	Triclinic
<i>a</i> (Å)	10.045(5)
<i>b</i> (Å)	11.529(3)
<i>c</i> (Å)	14.179(9)
α (°)	80.65(4)
β (°)	76.86(4)
γ (°)	85.47(3)
<i>U</i> (Å ³)	1576.3(13)
<i>Z</i>	1
<i>D</i> _{calc} (g cm ^{−3})	1.29
Space group	$P\bar{1}$ (No.2)
θ_{max} for data collection (°)	25
Unique reflections	5527
Reflections with $[I > 2\sigma(I)]$	4543
<i>R</i> ₁ [for $I > 2\sigma(I)$]	0.035
<i>wR</i> ₂ (for all data)	0.083

References

- [1] B. Marciniec (Ed.), Comprehensive Handbook on Hydrosilylation, Pergamon, Oxford, 1992.
- [2] D.N. Willing, US Patent 34 19 593 (1968).
- [3] B.D. Karstedt, US Patent 37 75 452 (1973).
- [4] G. Chandra, P.Y. Lo, P.B. Hitchcock, M.F. Lappert, Organometallics 6 (1987) 191.
- [5] P.B. Hitchcock, M.F. Lappert, N.J.W. Warhurst, Angew Chem. Int. Ed. Engl. 30 (1991) 438.
- [6] M.F. Lappert, F.P.E. Scott, J. Organomet. Chem. 492 (1995) C11.
- [7] P.B. Hitchcock, M.F. Lappert, C. MacBeath, F.P.E. Scott, N.J.W. Warhurst, J. Organomet. Chem. 528 (1997) 185.
- [8] P.B. Hitchcock, M.F. Lappert, C. MacBeath, F.P.E. Scott, N.J.W. Warhurst, J. Organomet. Chem. 534 (1997) 139.
- [9] J. Krause, G. Cestarić, K.-J. Haack, K. Seevogel, W. Storm, K.R. Pörschke, J. Am. Chem. Soc. 121 (1999) 9807.
- [10] B. Cornils, W.A. Herrmann, Applied Homogeneous Catalysis with Organometallic Compounds, VCH, Weinheim, 1996.
- [11] B. Proft, PhD. Dissertation, University of Düsseldorf, 1993.

- [12] F.G.N. Cloke, P.B. Hitchcock, M.F. Lappert, C. MacBeath, G.O. Mepsted, *J. Chem. Soc. Chem. Commun.* (1995) 87.
- [13] J. Stein, L.N. Lewis, Y. Gao, R.A. Scott, *J. Am. Chem. Soc.* 121 (1999) 3693.
- [14] C.J. Cardin, P.B. Hitchcock, M.F. Lappert, C. MacBeath, N.J.W. Warhurst, *J. Organomet. Chem.* 584 (1999) 366.
- [15] G.M. Sheldrick, in: G.M. Sheldrick, C. Krüger, R. Goddard (Eds.), *Crystallographic Computing*, vol. 3, Oxford University, Oxford, 1985, p. 175.
- [16] G.M. Sheldrick, *SHELXL-93*, A Program for Crystal Structure Refinement, University of Göttingen, 1993.