Internally Phosphine-Stabilized Zirconocene Cations Employing Substituted ((Diarylphosphino)methyl)cyclopentadienyl Ligand Systems

Boris E. Bosch, Gerhard Erker,* Roland Fröhlich, and Oliver Meyer

Organisch-Chemisches Institut der Universität Münster, Corrensstrasse 40, D-48149 Münster, Germany

Received July 8, 1997[®]

The dimethylzirconocene complex $[(Cp-CMe_2-PAr_2)_2ZrMe_2]$, **2a** (Ar = *p*-tolyl), was treated with 1 molar equiv of $B(C_6F_5)_3$ to yield the salt $[(Cp-CMe_2-PAr_2)_2ZrMe^+MeB(C_6F_5)_3^-]$, **3a**. The X-ray crystal structure analysis of **3a** shows that both $-PAr_2$ units are intramolecularly coordinated to zirconium in a close to C_2 -symmetric arrangement with the $[Zr]-CH_3$ group placed in the central position in the bent metallocene σ -ligand plane. Treatment of **2a** with 2 equiv of $B(C_6F_5)_3$ generates the highly reactive dication system $[(C_P-CMe_2-PAr_2)_2Zr^{2+}]$ (**4** with two MeB(C_6F_5)₃⁻ anions). The highly electrophilic cation **4** abstracts chloride from, e.g., dichloromethane solvent to yield $[(Cp-CMe_2-PAr_2)_2Zr-Cl^+]$ (5, with MeB(C₆F₅)₃⁻ anion). The same cation (5', with $ClB(C_6F_5)_3^-$ anion) was obtained from the reaction of [(Cp-CMe₂- $PAr_2_2ZrCl_2$ (1a) with B(C₆F₅)₃. The C₂-symmetric, internally –PAr₂-stabilized dication adds acetonitrile or 2,6-dimethylphenyl isocyanide to give the respective C_2 -symmetric donorligand adducts in which the $-PAr_2$ coordination to zirconium is retained. The complexes **3a** ($\Delta G^{\ddagger}_{enant}$ (300 K) = 14.0 ± 0.5 kcal/mol) and **5** ($\Delta G^{\ddagger}_{enant}$ (360 K) = 17.5 ± 0.5 kcal/mol) show dynamic NMR spectra due to an intramolecular enantiomerization process proceeding with a rate-determining cleavage of the Zr-P linkages. Complex 5 was also characterized by an X-ray crystal structure analysis.

Introduction

The chemistry of zirconocene cation systems has become of great interest recently. The donor-ligand-free Cp_2ZrR^+ species and especially the olefin complexes of the related modified *ansa*-metallocene cations probably serve as essential intermediates in homogeneous metallocene Ziegler catalysis.¹ Also, the donor-ligandstabilized systems $Cp_2ZrR(L)^+$ or $Cp_2ZrR(L)_2^+$ are of importance.² Early examples have paved the way toward an understanding of group 4 metallocene cation chemistry related to homogeneous Ziegler catalysis,³ and postulated single steps of catalytic cycles could be studied in detail using such stabilized metallocene cations as suitable model systems.⁴ Also, the metallocene cation serves as an activating organometallic Lewis acid in these adducts that may lead to novel stoichiometric reactions.⁵

Donor ligands used to stabilize the Cp₂ZrR⁺ moiety in these adduct systems are often nitriles or cyclic ethers.^{2,3} Phosphorus-containing systems are quite rare in this metallocene cation chemistry.⁶ After we had gained some experience with internal amine stabilization of alkylzirconocene cation-type systems, making use of substituted ((dimethylamino)methyl)cyclopentadienyl

[®] Abstract published in Advance ACS Abstracts, November 1, 1997. (1) Reviews: Sinn, H.; Kaminsky, W. Adv. Organomet. Chem. **1980**, 18, 99. Aulbach, M.; Küber, F. Chem. Unserer Zeit **1994**, 28, 197. Brintzinger, H.-H.; Fischer, D.; Mülhaupt, R.; Rieger, B.; Waymouth, R. Angew. Chem. **1995**, 107, 1255; Angew. Chem., Int. Ed. Engl. **1995**, 34, 1143. Bochmann, M. J. Chem. Soc., Dalton Trans. **1996**, 255 and references cited in these articles.

⁽²⁾ Jordan, R. F. Adv. Organomet. Chem. **1991**, 32, 325. Jordan, R. F.; Guram, A. S. Comprehensive Organometallic Chemistry; Abel, E. W., Stone, F. G. A., Wilkinson G., Eds.; Pergamon: New York, 1995; Vol. 4, pp 589–625.

⁽³⁾ Jordan, R. F.; Bajgur, C. S.; Willett, R.; Scott, B. J. Am. Chem. Soc. 1986, 108, 7410. Jordan, R. F.; LaPointe, R. E.; Bajgur, C. S.; Echols, S. F.; Willett, R. J. Am. Chem. Soc. 1987, 109, 4111. Jordan, R. F.; LaPointe, R. E.; Baenziger, N.; Hinch, G. D. Organometallics 1990, 9, 1539. Eshuis, J. J. W.; Tan, Y. Y.; Renkema, J.; Teuben, J. H. J. Mol. Catal. 1990, 62, 277. Borkowsky, S. L.; Jordan, R. F.; Hinch, G. D. Organometallics 1991, 10, 1268. Alelyunas, Y. W.; Jordan, R. F.; Echols, S. F.; Borkowsky, S. L.; Bradley, P. K. Organometallics 1991, 10, 2191. Eshuis, J. J. W.; Tan, Y. Y.; Meetsma, A.; Teuben, J. H.; Renkema, J.; Evens, G. G. Organometallics 1992, 11, 362.

^{(4) (}a) Temme, B.; Karl, J.; Erker, G. Chem. Eur. J. 1996, 2, 919.
Karl, J.; Erker, G. J. Mol. Catal., in press. Karl, J.; Erker, G. Chem. Ber., in press. (b) Pellechia, C.; Immirzi, A.; Grassi, A.; Zambelli, A. Organometallics 1993, 12, 4473. Pellechia, C.; Grassi, A.; Immirzi, A. J. Am. Chem. Soc. 1993, 115, 1160. Pellechia, C.; Immirzi, A.; Pappalardo, D.; Peluso, A. Organometallics 1994, 13, 3773. Pellechia, C.; Grassi, A.; Zambelli, A. Organometallics 1994, 13, 3773. Pellechia, C.; Grassi, A.; Zambelli, A. Organometallics 1994, 13, 3773. Pellechia, C.; Grassi, A.; Zambelli, A. Organometallics 1994, 13, 3773. Pellechia, C.; Grassi, A.; Zambelli, A. Organometallics 1994, 13, 388. Immirzi, A.; Zambelli, A. J. Organomet. Chem. 1977, 178, 1077. Akita, M.; Yasuda, H.; Nakamura, A. Bull. Chem. Soc. Jpn. 1984, 57, 480. Eisch, J. J.; Piotrowski, A. M.; Brownstein, S. K.; Gabe, E. J.; Florence, L. L. J. Am. Chem. Soc. 1985, 107, 7219. Thompson, M. E.; Baxter, S. M.; Bulls, A. R.; Burger, B. J.; Nolan, M. C.; Santarsiero, B. D.; Schaefer, W. P.; Bercaw, J. E. J. Am. Chem. Soc. 1987, 109, 203. den Haan, K. H.; Wielstra, Y.; Teuben, J. H. Organometallics 1987, 6, 2053. Ishikawa, M.; Ohskita, J.; Ito, Y.; Minato, A. J. Organometallics 1991, 10, 1980. Guram, A. S.; Jordan, R. F.; Taylor, D. F. J. Am. Chem. Soc. 1991, 113, 1833. Horton, A. D.; Orpen, A. G. Angew. Chem. 1992, 104, 912; Angew. Chem., Int. Ed. Engl. 1992, 31, 876; Organometallics 1992, 11, 8. Horton, A. D. J. Chem. Soc., Chem. Commun. 1992, 185. Röttger, D.; Erker, G.; Fröhlich, R.; Kotila, S. Chem. Ber. 1996, 129, 5. (5) Jordan, R. F.; La Pointe, R. E.; Bradley, P. K.; Baenziger, N. Organometallics 1998, 2080. Baye, E. L. Petersen, L. J. Organometallics 1998, 7199.

⁽⁵⁾ Jordan, R. F.; La Pointe, R. E.; Bradley, P. K.; Baenziger, N. Organometallics **1989**, *8*, 2892. Berg, F. J.; Petersen, J. L. Organometallics **1989**, *8*, 2461. Jordan, R. F.; Guram, A. S. Organometallics **1990**, *9*, 2116. Guram, A. S.; Jordan, R. F. Organometallics **1991**, *10*, 3470. Guram, A. S.; Guo, Z.; Jordan, R. F. J. Am. Chem. Soc. **1993**, *115*, 4902. Guo, Z.; Swenson, D. C.; Guram, A. S.; Jordan, R. F. Organometallics **1994**, *13*, 766.



Figure 1. View of the molecular geometry of **2b**. Selected bond lengths (Å) and angles (deg): Zr-C8 2.277(3), $Zr-C_{Cp}$ 2.544(3), C70-C1 1.524(4), C1-C2 1.542(4), C1-C3 1.521(4), C1-P4 1.904(3), P4-C50 1.840(3), P4-C60 1.844(3); C8-Zr-C8* 92.2(2), C2-C1-C3 109.9(3), C2-C1-P4 103.2(2), C2-C1-C70 111.2(2), C3-C1-P4 114.6(2), C3-C1-C70 111.5(3), P4-C1-C70 106.0(2), C1-P4-C50 107.85(14), C1-P4-C60 102.38(13), C50-P4-C60 100.84-(13).

ligand systems,⁷ we developed synthetic entries into related ((diarylphosphino)methyl)cyclopentadienyl-derived zirconocenes suited for conversion into the respective internally phosphine-stabilized zirconocene cation systems.⁸ This has led to the development of straightforward synthetic routes to novel mono- and dicationic phosphine-stabilized zirconocene systems, some of which constitute very interesting new organometallic Lewis acid systems.⁹ This synthetic development along with a chemical and structural characterization of a variety of key examples is described in this article.

Results and Discussion

The synthesis of the [1-methyl-1-((diarylphosphino)ethyl)]cyclopentadienyl ligand systems used in this study was carried out by means of the fulvene route.¹⁰ 6,6-Dimethylfulvene was treated with lithiodiarylphosphide (Ar = *p*-tolyl or phenyl), Scheme 1. The resulting ligand system was then treated with 0.5 molar equiv of ZrCl₄ to yield the metallocene complexes **1a** (Ar = *p*-tolyl) and **1b** (Ar = phenyl), as was previously described by us.⁸ The complexes **1a,b** were each subsequently treated with 2 molar equiv of methyllithium to furnish the dimethylzirconocene complexes **2a** (Ar = *p*-tolyl) and **2b** (Ar = phenyl), respectively. Out of each



series of these starting materials, one representative example was crystallized and characterized by an X-ray crystal structure analysis. The molecular structures of the metallocene dichloride (1b) and dimethyl (2b) complexes are very similar. Therefore, the structural data of **1b** are provided in the Supporting Information. A view of the molecular geometry of **2b** is given in Figure 1. Figure 1 shows that the bonding parameters around the pseudotetrahedrally coordinated zirconium atom in **2b** are unexceptionalⁱ¹ (Zr–C8 = 2.277(3) Å, $C8-Zr-C8^* = 92.2(2)^\circ$, $D1-Zr-D1^* = 132.0^\circ$, D1 and D2 denote the centroids of the two Cp ring systems). The conformational properties of the metallocene system **2b** are noteworthy though. In the crystal, **2b** is C_2 symmetric. The system shows a rotameric structure where both -CMe₂-PPh₂ substituents at the Cp rings are oriented in opposite lateral sectors at the bent metallocene wedge. The (Cp-CMe₂PPh₂)₂ZrMe₂ complex 2b, thus, exhibits a bis(lateral):anti-metallocene

⁽⁶⁾ Jordan, R. F.; Bajgur, C. S.; Dasher, W. E.; Rheingold, A. L. *Organometallics* **1987**, *6*, 1041. Jordan, R. F.; Bradley, P. K.; Baenziger, N. C.; LaPointe, R. E. J. Am. Chem. Soc. **1990**, *112*, 1289. Razavi, A.; Thewalt, U. J. Organomet. Chem. **1993**, *445*, 111. Alelyunas, Y. W.; Guo, Z.; LaPointe, R. E.; Jordan, R. F. Organometallics **1993**, *12*, 544. Guo, Z.; Swenson, D. C.; Jordan, R. F. Organometallics **1994**, *13*, 1424.

⁽⁷⁾ Bertuleit, A.; Fritze, C.; Erker, G.; Fröhlich, R. Organometallics 1997, 16, 2891. and references cited therein. See also (reviews): Jutzi, P.; Dahlhaus, J. Phosphorus, Sulfur Silicon Relat. Elem. 1994, 87, 73. Jutzi, P.; Dahlhaus, J. Coord. Chem. Rev. 1994, 137, 179. Okuda, J. Comm. Inorg. Chem. 1994, 16, 185. Jutzi, P.; Siemeling, U. J. Organomet. Chem. 1995, 500, 175.

⁽⁸⁾ Bosch, B.; Erker, G.; Fröhlich, R. *Inorg. Chim. Acta*, in press. (9) For the use of specific organometallic Lewis acids in organic catalysis see e.g.: Erker, G.; Schamberger, J.; van der Zeijden, A. A. H.; Dehnicke, S.; Krüger, C.; Goddard, R.; Nolte, M. *J. Organomet. Chem.* **1993**, *459*, 107 and references cited therein.

⁽¹⁰⁾ Ziegler, K.; Schäfer, W. Justus Liebigs Ann. Chem. 1934, 511,
(10) Ziegler, K.; Gellert, H.-G.; Martin, H.; Nagel, K.; Schneider, J. Justus Liebigs Ann. Chem. 1954, 589, 91. Sullivan, M. F.; Little, W. F. J. Organomet. Chem. 1967, 8, 277. Renaut, P.; Tainturier, G.; Gautheron, B. J. Organomet. Chem. 1978, 148, 43. Brickhouse, M. D.; Squires, R. R. J. Am. Chem. Soc. 1988, 110, 2706. Okuda, J. Chem. Ber. 1989, 122, 1075. Squires, R. R. Acc. Chem. Res. 1992, 25, 461. Clark, T. J.; Killian, C. M.; Luthra, S.; Nile, T. A. J. Organomet. Chem. 1993, 462, 247.

⁽¹¹⁾ Bochmann, M. Comprehensive Organometallic Chemistry, Abel, E. W., Stone, F. G. A., Wilkinson, G., Eds.; Pergamon: New York, 1995; Vol. 4, p 273. Ryan, E. J. *Ibid.* p 483. Hey-Hawkins, E. *Ibid.* p 501. Gambarotta, S.; Jubb, J.; Song, J.; Richeson, D. *Ibid.*, p. 543. Cardin, D. J.; Lappert, M. F.; Raston, C. L. Chemistry of Organo-Zirconium and -Hafnium Compounds; Wiley: New York, 1986.

Phosphine-Stabilized Zirconocene Cations

conformation.¹² One of the methyl groups at C1, namely C³H₃, is oriented close to the Cp-ligand plane (dihedral angle θ C3-C1-C70-C71 = -3.9(4)°), whereas the phosphorus atom P4 and the remaining C²H₃ group are located out of this plane (θ P4-C1-C70-C71 121.5(3)°, C2-C1-C70-C71 = -127.0(3)°).

The [Cp-CMe₂-P(p-tolyl)₂]₂Zr(CH₃)₂ complex **2a** was used as a starting material for generating the metallocene cation complexes. Complex 2a was therefore treated with 1 molar equiv of tris(pentafluorophenyl)borane in toluene.¹³ The $[(Cp-CMe_2-P(p-tolyl)_2)_2Zr (CH_3)^+$ $CH_3B(C_6F_5)_3^-$] salt **3a** was formed instantaneously and isolated in >90% yield. In dichloromethane d_2 , it exhibits a C_2 -symmetric structure in solution, as apparent from the NMR spectra at low temperature (1H/ 13 C NMR (600/150 MHz at 253 K) δ 6.44, 6.40, 6.26, 5.94/111.9, 106.92, 106.79, 97.8 (Cp-); δ 1.82, 1.28/30.3, 25.5 (C(CH_3)₂); δ 2.37 (12H)/21.38, 21.15 (p-tolyl- CH_3), $\delta - 0.39/17.3$ (Zr-CH₃)). The ¹H NMR signal of the Zr- CH_3 group is split as a binominal triplet due to coupling with the two symmetry-equivalent phosphorus nuclei $({}^{3}J_{PH} = 15 \text{ Hz})$, as is the corresponding ${}^{13}C$ NMR resonance (${}^{2}J_{PC} = 10.5$ Hz). The C(CH_{3})₂ methyl-group ¹H NMR resonances also show coupling to two ³¹P nuclei (as part of an $A_3A_3'XX'$ -type system¹⁴). The cation complex 3a exhibits a single ³¹P NMR resonance as expected, but with a very unusual chemical shift of δ -22.0 (for comparison, the corresponding ³¹P NMR resonance of the starting material **2a** occurs at δ +31.2). This enormous shifting of the ³¹P NMR resonance of the system upon cation formation and establishing the phosphine-to-metal coordination was not expected per se. It may be explained by either a very pronounced $\Delta_{\rm R}$ -effect¹⁵ or by assuming that the phosphorus of the -P(aryl)₂ moiety has entered the magnetically deshielding anisotropy cone of the adjacent Cp-ring system in the unusual overall coordination geometry that has resulted from the formation of the internally phosphinecoordinated cationic metallocene system (further structural evidence on this point is given below).

Complex **3a** exhibits dynamic NMR spectra. Raising the monitoring temperature of the ¹H NMR experiment (at 200 MHz) above 280 K results in a broadening of the C_5H_4 resonances and the C(CH₃)₂ signals. Eventually, the δ 6.44/5.94 (273 K) pair of resonances reaches coalescence ($T_c = 300$ K), as does the 6.39/6.24 pair ($T_c = 295$ K) and the δ 1.82/1.31 pair of methyl signals ($T_c = 300$ K). The dynamic process that leads to a symmetry exchange of these pairs of signals must be described as an intramolecular enantiomerization process of **3a**. Its Gibbs activation energy was estimated



at ΔG^{\ddagger} (300 K) = 14.0 ± 0.5 kcal/mol from the temperature-dependent dynamic ¹H NMR spectra described above.¹⁶ The dynamic symmetrization process requires a "left/right-exchange" of the phosphorus ligands bonded to zirconium in the σ -ligand plane of **3a**. Due to the specific stereoelectronic features of the bent metallocene unit,¹⁷ this requires (reversible) rupture of *both* zirconium-phosphorus bonds (see Scheme 2). The magnitude of the activation barrier is, thus, determined by a process leading to the unsaturated intermediate 8 that is devoid of any stabilization by coordination of electrondonating –PAr₂ ligands. Qualitatively, the activation barrier of this dynamic automerization process of **3a**, thus, provides a measure of the stabilization energy that a very electrophilic "naked" [Zr]-CH₃ cation gains by (intramolecularly) coordinating two R-PAr₂ donor ligands to it. The diphenylphosphino-substituted derivative 3b shows an analogous dynamic behavior. From the temperature-dependent ¹H NMR spectra, an almost identical $\Delta G^{\dagger}_{\text{enant}} = 14.1 \pm 0.5$ kcal/mol was obtained.

Complex 3a was characterized by an X-ray crystal structure analysis. In the crystal, the $CH_3B(C_6F_5)_3^{-1}$ anion is independent of the **3a** cation. The structural parameters of the anion are unexceptional.¹³ The cation shows a pseudopenta-coordinated zirconium atom with two substituted Cp ligands, a methyl group, and both $-P(p-tolyl)_2$ moieties bonded to the metal center. This leads to a (noncrystallographic) very close to C2-symmetric molecular geometry of the 3a cation. The D1-Zr-D2 angle is 137.2° (D1 and D2 denote the centroids of the two Cp ligands). The Zr-C10 bond length is 2.350(4) Å.¹⁸ The C¹⁰H₃ group is bonded to zirconium in the front of the bent metallocene wedge in the central position of the σ -ligand plane. The phosphorus atoms P1 and P11 are bonded to Zr laterally adjacent to the $Zr-CH_3$ vector (Zr-P1 = 2.8370(14) Å, Zr-P11 =2.8447(14) Å, $P1-Zr-C10 = 77.55(10)^\circ$, P11-Zr-C10 $= 77.31(10)^{\circ}$, P1-Zr-P11 = 154.78(4)°). Both P atoms

⁽¹²⁾ For leading references on the conformational features of group 4 bent metallocene complex chemistry, see, e.g.: Krüger, C.; Nolte, M.; Erker, G.; Thiele, S. Z. Naturforsch. **1992**, 47b, 995. Erker, G.; Temme, B. J. Am. Chem. Soc. **1992**, 114, 4004. Krüger, C.; Lutz, F.; Nolte, M.; Erker, G.; Aulbach, M. J. Organomet. Chem. **1993**, 452, 79. Erker, G.; Aulbach, M.; Knickmeier, M.; Wingbermühle, D.; Krüger, C.; Nolte, M.; Werner, S. J. Am. Chem. Soc. **1993**, 115, 4590. Fritze, C.; Knickmeier, M.; Erker, G.; Zaegel, F.; Gautheron, B.; Meunier, P.; Paquette, L. A. Organometallics **1995**, 14, 5446. Knickmeier, M.; Erker, G.; Fox, T. J. Am. Chem. Soc. **1996**, 118, 9623 and references cited therein.

⁽¹³⁾ For leading references on this method of generating group 4 bent metallocene cation complexes, see: Yang, X.; Stern, C. L.; Marks, T. J. *J. Am. Chem. Soc.* **1994**, *116*, 10015 and references cited therein.

⁽¹⁴⁾ Leoni, P.; Marchetti, F.; Paoletti, M. Organometallics **1997**, *16*, 2146.

⁽¹⁵⁾ Garrou, P. E. *Inorg. Chem.* **1975**, *14*, 1435. Garrou, P. E. *Chem. Rev.* **1981**, *81*, 229.

⁽¹⁶⁾ Gutowsky, H. S.; Holm, C. H. *J. Chem. Phys.* **1956**, *25*, 1228. Green, M. L. H.; Wong, L.-L.; Seela, A. *Organometallics* **1992**, *11*, 2660 and references cited therein.

⁽¹⁷⁾ Brintzinger, H. H.; Bartell, L. S. J. Am. Chem. Soc. **1970**, *92*, 1105. Lauher, J. W.; Hoffmann, R. J. Am. Chem. Soc. **1976**, *98*, 1729.



Figure 2. Two projections of the molecular structure of **3a** (cation only) in the crystal. Selected bond lengths (Å) and angles (deg): $Zr-C10 \ 2.350(4)$, $Zr-C_{Cp} \ 2.512(5)$, $Zr-P1 \ 2.8370(14)$, $Zr-P11 \ 2.8447(14)$, $P1-C2 \ 1.894(5)$, $P11-C12 \ 1.892(5)$, $P1-C21 \ 1.809(5)$, $P11-C51 \ 1.808(5)$, $P1-C31 \ 1.850(5)$, $P1-C41 \ 1.835(5)$, $C2-C3 \ 1.540(7)$, $C12-C13 \ 1.543(7)$, $C2-C4 \ 1.538(7)$, $C12-C14 \ 1.529(7)$, $C2-C5 \ 1.518(7)$, $C12-C15 \ 1.516(7)$; C10-Zr-P177.55(10), $C10-Zr-P11 \ 77.31(10)$, $P1-Zr-P11 \ 154.78(4)$, $Zr-P1-C2 \ 89.0-(2)$, $Zr-P11-C12 \ 89.2(2)$, $Zr-P1-C21 \ 125.7(2)$, $Zr-P11-C51 \ 123.8(2)$, $Zr-P1-C31 \ 120.0(2)$, $Zr-P11-C41 \ 120.7(2)$, $P1-C2-C3 \ 114.9(4)$, $P11-C12-C13 \ 113.7(3)$, $P1-C2-C4 \ 111.8(4)$, $P11-C12-C14 \ 113.5(4)$, $P1-C2-C5 \ 99.5(3)$, $P11-C12-C15 \ 98.9(3)$.

are located markedly outside the bent metallocene σ -ligand plane (i.e., C_2 -symmetrically above and below, see Figure 2). The angle between the D1-Zr-D2 and P1-Zr-P11 planes is 80.6°. This deviation from an electronically favored arrangement of the P1.Zr.P11 unit in the Cp-Zr-Cp bisecting major plane of the bent metallocene unit is enforced geometrically by the attachment of the -PAr₂ groups at the Cp ligands by the CMe_2 linkages (C2-P1 = 1.894(5) Å, C12-P11 = 1.892-(5) Å, $C5-C2-P1 = 99.5(3)^\circ$, $C2-P1-Zr = 89.0(2)^\circ$, $C15-C12-P11 = 98.9(3)^{\circ}, C12-P11-Zr = 89.2(2)^{\circ}.$ This geometric arrangement brings each phosphorus atom rather close to one Cp edge ($C5 \cdots P1 = 2.615$ Å, $C15 \cdots P11 = 2.600$ Å). The methyl groups at C2 (and C12) are in a markedly different environment. The C⁴H₃ group points toward the front side of the bent metallocene wedge, whereas the $C^{3}H_{3}$ substituent is oriented toward a hind lateral sector. The relative orientation of the aryl groups at the phosphorus atoms



Figure 3. View of the molecular geometry of **5** $(CH_3B(C_6F_5)_3^-$ anion not depicted). Selected bond lengths (Å) and angles (deg): Zr-Cl 2.453(2), Zr-C_{Cp} 2.499(7), Zr-P1 2.846(2), Zr-P11 2.846(2), P1-C2 1.885(7), P11-C12 1.891(7), P1-C21 1.798(7), P11-C51 1.814(7), P1-C31 1.833(7), P11-C41 1.827(7), C2-C3 1.534(10), C12-C13 1.536(10), C2-C4 1.549(10), C12-C14 1.529(9), C2-C5 1.516(10), C12-C15 1.532(10); Cl-Zr-P177.83(6), Cl-Zr-P11 78.03(6), P1-Zr-P11 155.81(6), Zr-P1-C2 88.3(2), Zr-P11-C12 88.8(2), Zr-P1-C21 124.6(2), Zr-P11-C51 122.6(3), Zr-P1-C31 120.8(2), Zr-P11-C41 121.9(2), P1-C2-C3 115.3(6), P11-C12-C13 113.9(5), P1-C2-C4 111.7-(5), P11-C12-C14 112.5(5), P1-C2-C5 99.5(4), P11-C12-C15 98.3(4).

(P1, P11) is similar; this leads to some lateral shielding of the central $Zr-CH_3$ group by the bulky substituents at phosphorus (see Figure 2).

Next, we tried to generate the internally phosphanestabilized dication 4 (Ar = *p*-tolyl, see Scheme 1).¹⁹ In the first series of experiments, we treated the dimethylzirconocene complex 2a (Ar = *p*-tolyl) with 2 molar equiv of $B(C_6F_5)_3$ in dichloromethane. The reaction rapidly goes to completion when carried out at room temperature. A single reaction product was obtained almost quantitatively. Recrystallization from dichloromethane/pentane gave single crystals suited for the X-ray crystal structure analysis. This revealed that instead of the expected dication complex [(Cp-CMe₂- $PAr_{2}_{2}Zr^{2+} \cdot 2$ MeB(C₆F₅)₃⁻] (4, see below) the [(Cp-CMe₃PAr₂)₂Zr-Cl⁺MeB(C₆F₅)₃⁻] salt 5 had been formed and isolated in excellent yield. The structure of the chloro-containing monocation 5 (see Figure 3) is very similar to the methyl-containing cation **3a** (see above). The overall molecular arrangement is again close to C_2 symmetric with two $-P(p-tolyl)_2$ ligands internally coordinating to zirconium as close to the σ -ligand plane of the bent metallocene wedge as the rigid system allows. The Zr-Cl vector is located in the central position in the σ -ligand plane (Zr-Cl = 2.453(2) Å, Zr-P1 = 2.846(2) Å, Zr-P11 = 2.846(2) Å, D1-Zr-D2 =134.9°, P1-Zr-Cl = 77.83(6)°, P11-Zr-Cl = 78.03(6)°, P1-Zr-P11 = 155.81(6)°, C2-P1-Zr = 88.3(2)°, C12- $P11-Zr = 88.8(2)^{\circ}$).

Complex **5** exhibits the very characteristic NMR spectra of the chiral C_2 -symmetric cation at ambient

⁽¹⁸⁾ Atwood, J. L.; Barker, G. K.; Holton, J.; Hunter, W. E.; Lappert, M. F.; Pearce, R. J. Am. Chem. Soc. **1977**, 99, 6645. Jeffrey, J.; Lappert, M. F.; Luong-Thi, N. T.; Webb, M.; Atwood, J. L.; Hunter, W. E. J. Chem. Soc., Dalton Trans. **1981**, 1593. Atwood, J. L.; Hunter, W. E.; Hrncir, D. C.; Bynum, V.; Penttila, R. A. Organometallics **1983**, 2, 750. Krüger, C.; Müller, G.; Erker, G.; Dorf, U.; Engel, K. Organometallics **1985**, 4, 215. See also: Orpen, G. A.; Brammer, L.; Allen, F. H.; Kennard, O.; Watson, D. G.; Taylor, R. J. Chem. Soc., Dalton Trans. **1989**, S1.

^{(19) (}a) Zirconium: Jordan, R. F.; Echols, S. F. Inorg. Chem. **1987**, 26, 383. Billinger, P. N.; Claire, P. P. K.; Collins, H.; Willey, G. R. Inorg. Chim. Acta **1988**, 149, 63. Thewalt, U.; Lasser, W. J. Organomet. Chem. **1984**, 276, 341. Lasser, W.; Thewalt, U. J. Organomet. Chem. **1986**, 311, 69. (b) Titanium: Thewalt, U.; Klein, H.-P. J. Organomet. Chem. **1980**, 194, 297. Thewalt, U.; Berhalter, K. J. Organomet. Chem. **1986**, 316, 291. Berhalter, K.; Thewalt, U. J. Organomet. Chem. **1987**, 332, 123.

Phosphine-Stabilized Zirconocene Cations

temperature (¹³C NMR (CD₂Cl₂, 150 MHz) δ 116.3, 112.8, 108.4, 100.9 (Cp), 29.8, 26.4 (C(CH₃)₂), 21.7, 21.4 (tolyl-CH₃); ¹H NMR (C₆D₅Br, 200 MHz, 303 K) δ 6.23, 6.22, 6.01, 5.86 (Cp), 1.49, 1.32 (C(CH₃)₂). The ³¹P NMR resonance of **5** is at δ –33.0 ppm (in CD₂Cl₂).

Complex 5 shows a dynamic NMR behavior at higher temperatures similar to that previously observed for 3a. The ¹H NMR spectrum of **5** in bromobenzene- d_5 shows a broadening of the lines at temperatures above ca. 340 K (200 MHz) with subsequent coalescence of the former δ 6.23/6.01 (T_c = 363 K) and δ 6.22/5.86 (T_c = 370 K) pairs of Cp resonances as well as the former δ 1.49/1.32 ¹H NMR C(CH₃)₂ resonance pair ($T_c = 358$ K). The enantiomerization process of 5 is analogous to that observed and described for 3 (see above and Scheme 2) and is also likely to proceed through a nondonor-ligandstabilized intermediate (8), however, in the case of the [Zr]-Cl⁺ complex this should be even further destabilized due to the presence of the electron-withdrawing σ -chloride ligand at zirconium instead of $-CH_3$. Thus, the $\mathbf{5} \rightarrow \mathbf{8}$ transition should have a higher activation barrier compared to the corresponding $3 \rightarrow 8$ reaction, which it has. The Gibbs activation energy of the thermally induced intramolecular $5 \Rightarrow \text{ent-}5$ automerization process was derived at $\Delta G^{\ddagger}_{ent}$ (360 K) = 17.5 \pm 0.5 kcal/mol from the dynamic ¹H NMR spectra.¹⁶

The cation 5 was prepared independently. We have treated the substituted zirconocene dichloride 1a (Ar = p-tolyl) directly with 1 molar equiv of the Lewis acid $B(C_6F_5)_3$ in dichloromethane. At ambient temperature one chloride is rapidly removed from zirconium with formation of the [(Cp-CMe₂-P(p-tolyl)₂)₂Zr-Cl⁺ClB- $(C_6F_5)_3^{-1}$ salt 5'. The ClB $(C_6F_5)_3^{-1}$ anion was identified by LDI-TOF-MS (m/z = 547) in the negative ion mode and by an X-ray crystal structure analysis (B-Cl2 =1.907(8) Å; for details of this structure, see the Supporting Information). The NMR spectra of the independently prepared compound 5' are very similar to those of 5 (for details, see the Experimental Section).

It is very likely that the chloride-containing complex [(Cp-CMe₂-PAr₂)₂Zr-Cl⁺CH₃B(C₆F₅)₃⁻] **5** was formed in the reaction between 2a and 2 equiv of $B(C_6F_5)_3$ via the reactive $[(Cp-CMe_2-PAr_2)_2Zr^{2+}\cdot 2CH_3B(C_6F_5)_3^-]$ dication intermediate 4, which is not stable under the applied reaction conditions. It must be assumed that the ultimately obtained reaction products originate from 4, which is converted to 5 by a subsequent chloride abstraction reaction from the halocarbon solvent.

This was shown to be true, although the organic stoichiometric reaction product could not be identified. The reaction of **2a** with 2 equiv of $B(C_6F_5)_3$ in CD_2Cl_2 was carried out at variable temperatures under direct ¹H and ³¹P NMR control. The reaction was started at 213 K, a temperature where the formation of the monocation 3a is very rapid. After 30 min at 213 K, a mixture containing 3a (65%), 5 (9%), and the new complex 4 (i.e., the donor-ligand-stabilized dication, 26%) was obtained. Slowly raising the temperature resulted in a predominant conversion of 3a to 4 (by means of intermolecular CH₃ abstraction by the B(C₆F₅)₃ reagent) without very much subsequent chloride abstraction. At 228 K after a total reaction time of 60 min, a 3a:5:4 ratio of 27:11:62 was obtained, and at 253 K after a total reaction time of 125 min, the primary product 3a was no longer observed, the 5:4 ratio was

now 20:80. Raising the temperature further then led to an increased consumption of the dication 4 with formation of 5. The characteristic NMR data of 4 were obtained from a ca. 1:1 mixture of 4 and 5 at 275 K in dichloromethane- d_2 (δ 2.39, 2.29 (*p*-tolyl CH₃); δ 1.93, 1.59 (C(CH₃)₂)). The ³¹P NMR resonance of **4** (CD₂Cl₂) is at δ –48.5 ppm. Complex **4** was also obtained as an insoluble oil by treatment of **2a** with ca. 2 molar equiv of $B(C_6F_5)_3$ in toluene (see Experimental Section). Since **4** is insoluble in the aromatic hydrocarbon solvent, the product was spectroscopically characterized using CD₂- Cl_2 or C_6D_5Br solutions (see above).

We have characterized the dication 4 by its reaction with external donor ligands. The resulting [(Cp- $CMe_2 - PAr_2 Zr - L^{2+} Z CH_3 B(C_6F_5)_3^{-1}$ complexes are more stable than the very electrophilic dication 4 and, thus, are easier to handle and isolate.²⁰

The reaction between **3a**, $B(C_6F_5)_3$ (2 molar equiv), and acetonitrile was carried out in toluene. The reaction rapidly turns heterogeneous, and the reaction product (6, see Scheme 1) separates as an oil. The isolated (89% yield) complex [(Cp-CMe₂-PAr₂)₂Zr- $(N \equiv C - CH_3)^{2+2} CH_3B(C_6F_5)_3^{-1}]$ (6) is stable in dichloromethane. It shows the typical NMR spectra of a C_2 symmetric cationic species in solution ($^{1}H/^{13}C$ NMR δ 6.73, 6.46, 6.38, 6.20/117.1, 108.5, 107.9, 101.2 (Cp); δ 2.43, 2.41/21.53, 21.42 (CH₃ of p-tolyl), 1.78, 1.44/29.4, 26.3 (C(CH_3)₂)). The ¹H NMR resonance of the coordinated acetonitrile ligand appears at δ 1.32 as a triplet due to coupling with the two adjacent symmetryequivalent ³¹P NMR nuclei (${}^{5}J_{PH} = 3.3$ Hz). The coordinated $-C \equiv N^{13}C$ NMR resonance is at δ 141.5 ppm, and the corresponding acetonitrile methyl ¹³C NMR signal is at δ 2.92. Complex **6** exhibits a ³¹P NMR resonance at δ –44.7 ppm. The typical IR bands of the coordinated acetonitrile ligand are found at 2307 (combination) and 2273 cm⁻¹ (C=N) (free acetonitrile shows the corresponding IR bands at $\tilde{\nu} \approx 2290$ and 2255 cm^{-1}).^{20,21}

We have also trapped the in-situ-generated metallocene dication 4 by addition of 2,6-dimethylphenyl isocyanide (reaction carried out in toluene). The [(Cp- $CMe_2 - PAr_2)_2Zr(C \equiv N-Ar')^{2+} \cdot 2CH_3B(C_6F_5)_3^{-}]$ salt 7 (see Scheme 1) was isolated in ca. 80% yield. It again exhibits the typical NMR spectra of a C₂-symmetric adduct (for details see the Experimental Section). The $\tilde{\nu}$ (C=N-R) IR band of 7 was monitored at $\tilde{\nu} = 2170$ cm^{-1} (free C=N-C₆H₃Me₂ 2122 cm⁻¹).

Conclusions

This study shows that phosphine stabilization of the very electrophilic zirconium center in the mono- and dicationic bent metallocene complexes can become very pronounced and dominant if the P-donor ligands become attached to the Cp rings by suitable linkages and, thus, can coordinate intramolecularly. In this manner, even the very electrophilic 16-electron dication 4 has attained a moderate stability, although it apparently is so electrophilic that it rapidly starts to abstract chloride

⁽²⁰⁾ See, for a comparison: Brackemeyer, T.; Erker, G.; Fröhlich, R. Organometallics 1997, 16, 531. Brackemeyer, T.; Erker, G.; Fröhlich, R.; Prigge, J.; Peuchert, U. Chem. Ber. 1997, 130, 899.
(21) Bruce, M. R. M.; Tyler, D. R. Organometallics 1985, 4, 528. Jordan, R. F.; Dasher, W. E.; Echols, S. F. J. Am. Chem. Soc. 1986, 108, 1718.

^{108, 1718.}

from the solvent at or above ca. -20 °C to cleanly form the C_2 -symmetric chloride containing 18-electron monocation 5. Similarly, there is an apparently large thermodynamic driving force of 4 to add neutral donor ligands (nitrile, isonitrile) to give the respective very stable 18-electron complexes (6,7) which have the third donor ligand bonded at zirconium in the central position in the σ -ligand plane. This study shows that the selected synthetic route $(1 \rightarrow 2 \rightarrow 3 \rightarrow 4$, see Scheme 1) is suited and easy to use for generating the reactive chiral internally donor-ligand-stabilized metallocene dication system 4.²² Whether this rather electrophilic dication system can be used as a catalyst or a stoichiometric reagent in organic transformations and whether its inherent chirality can be utilized in such reactions is currently being studied in our laboratory.

Experimental Section

All compounds described are air and moisture sensitive. They were prepared and handled in an argon atmosphere using Schlenk-type glassware or in a glovebox. Solvents (including deuterated solvents) were dried and distilled under argon prior to use. NMR spectra were recorded on a Bruker AC 200 P (¹H, 200 MHz; ³¹P, 81 MHz, H₃PO₄ external standard; ¹³C, 50.3 MHz) or a Varian Unity plus (¹H, 600 MHz; ¹³C, 150 MHz) NMR spectrometer. IR spectroscopy: Nicolet 5 DXC FT-IR spectrometer (KBr). Elemental analyses: Foss Heraeus CHNO-Rapid. DSC: Dupont 2910 DSC (STA instruments). X-ray crystal structure analyses: Enraf-Nonius MACH 3 diffractometer (programs used, SHELX 86, SHELX 93, DIAMOND, and SCHAKAL). The complexes **1a** and **1b** were prepared as recently described by us.⁸ B(C₆F₅)₃ was prepared according to a literature procedure.²³

X-ray Crystal Structure Analysis of 1b. Formula $C_{40}H_{40}$ -Cl₂P₂Zr; M = 744.78; $0.20 \times 0.20 \times 0.10$ mm; a = 31.058(6) Å, b = 6.537(1) Å, c = 18.167(4) Å, $\beta = 108.57(2)^{\circ}$, V = 3496.2-(12) Å³, $\rho_{calc} = 1.415$ g cm⁻³, $\mu = 5.87$ cm⁻¹, empirical absorption correction via φ scan data ($0.925 \leq C \leq 0.998$), Z = 4, monoclinic, space group C2/c (No. 15), $\lambda = 0.710$ 73 Å, T =223 K, ω scans, 3660 reflections collected ($\pm h$, +k, -h), [(sin $\theta)/\lambda$] = 0.62 Å⁻¹, 3528 independent and 2112 observed reflections [$I \geq 2 \sigma(I)$], 207 refined parameters, R1 = 0.048, wR2 = 0.106, max (min) residual electron density 0.76 (-1.15) e Å⁻³, hydrogens calculated and refined as riding atoms.

Bis[[1-methyl-1-(di-p-tolylphosphino)ethyl]cyclopentadienyl]dimethylzirconium (2a). 1a (16.0 g, 20 mmol) was suspended in 150 mL of diethyl ether, and 24 mL (40 mmol) of methyllithium (1.68 M in Et₂O) was added at -78 °C. The suspension was allowed to warm to room temperature overnight. Celite filtration and washings of the filter pad with 100 mL of diethyl ether produced a red solution. The solvent was evaporated, the residue taken up in 60 mL of pentane, and the product collected on a Schlenk frit, washed with two 20 mL portions of pentane, and dried in vacuo. Yield: 8.61 g, 57%. Mp 112 °C. Anal. Calcd for C46H54P2Zr (760.1): C, 72.69; H, 7.16. Found: C, 70.25; H, 7.21. IR (KBr): v 3080, 3069, 3022, 3013, 2971, 2960, 2911, 2864, 1598, 1558, 1498 (vs), 1474, 1446, 1409, 1397, 1377, 1358, 1307, 1261, 1187, 1120 (vs), 1089, 1038, 1020, 932, 874, 862, 853, 810 (br, vs), 716, 710, 699, 662, 649, 623, 611 cm⁻¹. ¹H NMR (benzene- d_6 , 200.1 MHz): 8 7.47 (m, 8H, o-tol-H), 6.99 (m, 8H, m-tol-H), 5.73 (s, 8H, C_5H_4), 2.08 (s, 12H, *p*-tol-CH₃), 1.42 (d, ${}^3J_{PH} = 13.6$ Hz, 12H, CMe₂), -0.03 (s, 6H, Zr-CH₃). ¹³C NMR (benzene*d*₆, 50.3 MHz): δ 138.8 (*p*-tol), 135.7 (²*J*_{PC} = 21 Hz, *o*-tol), 135.0 $(^{2}J_{PC} = 7 \text{ Hz}, C1 \text{ of } Cp), 133.4 (^{1}J_{PC} = 21 \text{ Hz}, ipso$ C of tol), 128.9 (${}^{3}J_{PC} = 7$ Hz, *m*-tol), 110.3 (${}^{3}J_{PC} = 5$ Hz, C2 of Cp), 109.6 (Cp), 37.9 (${}^{1}J_{PC} = 21$ Hz, *C*Me₂), 32.3 (Zr–*CH*₃), 26.4 (${}^{2}J_{PC} = 18$ Hz, C(*C*H₃)₂), 21.2 (tol-CH₃). 31 P NMR (benzene-*d*₆, 81.0 MHz): δ 31.2.

Bis[[1-methyl-1-(diphenylphosphino)ethyl]cyclopentadienyl]dimethylzirconium (2b). To a suspension of 3.11 g (4.17 mmol) of bis[[1-methyl-1-(diphenylphosphino)ethyl]cyclopentadienyl]dichlorozirconium (1b) in 80 mL of pentane was added 5.0 mL of methyllithium (1.68 M in diethyl ether, 8.4 mmol) at -78 °C. A yellow suspension was obtained after stirring overnight. Subsequently, the solvent was evaporated into a cold trap, and the residue was taken up in 150 mL of toluene and filtered through a 2 cm bed of Celite. The toluene was evaporated, and the crude product 2b was washed with two 20 mL portions of pentane, leaving 2.35 g (80%) of 2b as a white solid. A sample of 2b was dissolved in a 3:1 mixture of toluene and pentane. Cooling this mixture to -20°C vielded white crystals suitable for the X-ray diffraction analysis. Mp 178 °C. Anal. Calcd for C42H46P2Zr (704.0): C, 71.66; H, 6.59. Found: C, 70.17; H, 6.47. IR (KBr): $\tilde{\nu} = 3075$, 3058, 2971, 2964, 2932, 2915, 2868, 1583 (s), 1481, 1461, 1435 (s), 1429 (s), 1412, 1377, 1358, 1223, 1091, 1068, 1051, 1039, 1025, 1000, 931, 875, 865, 854, 810 (vs), 747, 739 (s), 698 (s) cm⁻¹. ¹H NMR (benzene-*d*₆, 200.1 MHz): δ 7.53–7.44 (m, 8H, o-Ph), 7.19-7.14 (m, 12H, p,m-Ph), 5.73 (s, 8H, C₅H₄), 1.39 (d, ${}^{3}J_{PH} = 13.5$ Hz, 12H, CMe₂), -0.07 (s, 6H, Zr-CH₃). ${}^{13}C$ NMR (APT, benzene- d_6 , 50.3 MHz): δ 136.7 ($^1J_{PC}$ = 23 Hz, ipso C of Ph), 135.5 (²J_{PC} = 20 Hz, o-Ph), 129.0 (p-Ph), 128.2 $({}^{3}J_{PC} = 7 \text{ Hz}, \text{ m-Ph}), 110.2 ({}^{3}J_{PC} = 6 \text{ Hz}, C2 \text{ of Cp}), 109.6 (Cp),$ 37.9 (${}^{1}J_{PC} = 21$ Hz, *C*Me₃), 32.3 (Zr-C*H*₃), 26.3 (${}^{2}J_{PC} = 18$ Hz, $C(CH_3)_2$; C1 of Cp not observed. ³¹P NMR (benzene- d_6 , 81.0 MHz): δ 32.5.

X-ray Crystal Structure Analysis of 2b. Formula $C_{42}H_{46}P_2Zr; M = 703.95; 0.50 \times 0.30 \times 0.20 \text{ mm}; a = 31.554-(3) Å; b = 7.028(1) Å; c = 16.735(2) Å, \beta = 105.72(1)^\circ, V = 3572.4(7) Å^3, \rho_{calc} = 1.309 \text{ g cm}^{-3}, \mu = 4.26 \text{ cm}^{-1}, \text{ empirical absorption correction via } \varphi \text{ scan data } (0.962 \le C \le 0.999), Z = 4, \text{ monoclinic, space group } C2/c \text{ (No. 15), } \lambda = 0.710 \text{ 73 Å}, T = 223 \text{ K}, \omega \text{ scans, } 3771 \text{ reflections collected } (\pm h, -k, -l), [(sin <math>\theta)/\lambda] = 0.62 \text{ Å}^{-1}$, 3636 independent and 2622 observed reflections [$I \ge 2 \sigma(I)$], 207 refined parameters, R1 = 0.039, wR2 = 0.083, max (min) residual electron density 0.41 (-0.32) e Å^{-3}, hydrogens calculated and refined as riding atoms.

Methylbis{[[1-methyl-1-(di-*p*-tolylphosphino)ethyl]-η⁵cyclopentadienyl]- κ^2 -P,P'}zirconium(IV) Methyltris(pentafluorophenyl)borate (3a). 2a (760 mg, 1.0 mmol) and 512 mg (1.0 mmol) of $B(C_6F_5)_3$ were mixed as solids and suspended in 25 mL of toluene. The orange suspension was stirred for 1 h, and the mixture was concentrated in vacuo to 2 mL. Pentane (20 mL) was added, and the mixture was stirred for 1 h. The pale yellow product was isolated on a Schlenk frit, washed with two 10 mL portions of pentane, and dried in vacuo to yield 1.15 g (91%) of the pale yellow monocation complex 3a. It was possible to grow large crystals of 3a by layering a dichloromethane solution with an equal amount of pentane at -20 °C. Single crystals suitable for the X-ray diffraction analysis were obtained by slow diffusion of pentane into a dichloromethane solution of 3a at 4 °C. Mp 160 °C (dec). Anal. Calcd for C₆₄H₅₄BF₁₅P₂Zr (1272.08): C, 60.43; H, 4.28. Found: C, 59.26; H, 4.09. IR (KBr): $\tilde{\nu} = 3028$, 2994, 2974, 2956, 2923, 2869, 1641 (B(C₆F₅)₃), 1600, 1510, 1498, 1458 (br, vs), 1265 (B(C₆F₅)₃), 1190, 1117, 1088, 1081, 994, 975, 965, 947, 934, 838, 822, 800, 658 cm⁻¹. ¹H NMR (dichloromethane-d₂, 200.1 MHz, 253 K): & 7.27 (m, 16H, p-tol-H), 6.43 (m, 4H, C₅H₄), 6.27, 5.94 (each m, each 2H, C₅H₄), 2.36 (s, 12H, tol-CH₃), 1.81 (m, 6H, CMe₂), 1.28 (m, 6H, CMe₂), 0.43 (br s, 3H, [B]-CH₃), -0.42 (t, ${}^{3}J_{PH} = 15.1$ Hz, 3H, Zr-CH₃). ${}^{13}C$ NMR (dichloromethane- d_2 , 150 MHz, 253 K): δ 148.1 ($^1J_{\rm FC} = 228$ Hz, o-C₆F₅), 142.05, 141.92 (each *p*-tol), 137.4 (${}^{1}J_{FC} = 235$ Hz, p-C₆F₅), 137.1 (ipso-C of Cp), 136.3 (C₆F₅), 135.38 (o-tol), 134.43 (o-tol), 130.1 (m-tol), 129.9 (m-tol), 125.8 (ipso-C of tol), 121.7 (ipso-C of tol), 111.9, 106.92, 106.79, 97.8 (Cp), 34.9 (CMe2),

⁽²²⁾ For a structurally related neutral system, see: Christoffers, J.; Bergman, R. G. Angew. Chem. **1995**, 107, 2423; Angew. Chem., Int. Ed. Engl. **1995**, 34, 2266.

⁽²³⁾ Massey, A. G.; Park, A. J. J. Organomet. Chem. 1964, 2, 245.

Phosphine-Stabilized Zirconocene Cations

30.3, 25.5 (C(*C*H₃)₂), 21.38, 21.15 (tol-CH₃) 17.3 (t, ${}^{2}J_{PC} = 10.5$ Hz, Zr–C*H*₃), 9.8 (br, H₃*C*[B]). 31 P NMR (dichloromethane- d_{2} , 81.0 MHz): δ –22.0. 11 B NMR (dichloromethane- d_{2} , 64.2 MHz): δ –15.

X-ray Crystal Structure Analysis of 3a. Formula $C_{64}H_{54}$ -BF₁₅P₂Zr; M = 1272.04; $0.40 \times 0.25 \times 0.10$ mm; a = 14.139(2) Å, b = 14.976(2) Å, c = 16.145(2) Å, $\alpha = 113.62(1)^{\circ}$, $\beta = 109.24$ - $(1)^{\circ}$, $\gamma = 95.84(1)^{\circ}$, V = 2847.2(7) Å³, $\rho_{calc} = 1.484$ g cm⁻³, $\mu = 3.40$ cm⁻¹, empirical absorption correction via φ scan data (0.918 $\leq C \leq 0.999$), Z = 2, triclinic, space group $P\overline{1}$ (No. 2), $\lambda = 0.710$ 73 Å, T = 223 K, $\omega/2\theta$ scans, 10 412 reflections collected ($\pm h, \pm k, -h$), [(sin $\theta)/\lambda$] = 0.59 Å⁻¹, 10 024 independent and 6956 observed reflections [$I \geq 2 \sigma(h)$], 758 refined parameters, R1 = 0.063, wR2 = 0.182, max (min) residual electron density 0.99 (-1.25) e Å⁻³, hydrogens calculated and refined as riding atoms.

Methylbis{[[1-methyl-1-(diphenylphosphino)ethyl]- η^{5} cyclopentadienyl]- κ^2 -P,P'}zirconium(IV) Methyltris(pentafluorophenyl)borate (3b). In a similar procedure, the analogous cationic system with Ar = Ph was obtained, starting from 211 mg (0.3 mmol) of 2b and 154 mg (0.3 mmol) of B(C₆F₅)₃. Yield of **3b**: 350 mg (95%). Large colorless crystals were grown by diffusion of pentane into a dichloromethane solution of 3a at -20 °C. Mp 201 °C (dec). Anal. Calcd for C₆₀H₄₆BF₁₅P₂Zr (1215.97): C, 59.27; H, 3.81. Found: C, 59.30; H, 3.81. IR (KBr): v 3121, 3064, 2996, 2964, 2941, 2929, 2903, 2844, 1641, 1511, 1458 (br, vs), 1436, 1265, 1096 (br, vs), 995, 978, 965, 949, 935, 828, 802, 743, 707, 696, 659, 640 cm⁻¹. ¹H NMR (dichloromethane-d₂, 200.1 MHz, 243 K): δ 7.7-7.2 (m, 20H, Ph), 6.47 (m, 4H, C₅H₄), 6.33, 5.97 (each m, each 2H, C₅H₄), 1.86 (m, 6H, CMe₂), 1.29 (m, 6H, CMe₂), 0.45 (br s, 3H, [B]-CH₃), -0.39 (t, ${}^{3}J_{PH} = 15.1$ Hz, 3H, Zr-CH₃). ${}^{13}C$ NMR (dichloromethane- d_2 , 50.3 MHz, 238 K): δ 147.8 (${}^1J_{\rm FC} = 229$ Hz, o-C₆F₅), 137.3 (${}^{1}J_{FC} = 230$ Hz, p-C₆F₅), 136.9 (ipso-C of Cp), 136.0 (${}^{1}J_{FC} = 250 \text{ Hz}, m \cdot C_{6}F_{5}$), 135.3 (o-Ph), 134.5 (o-Ph), 131.13, 131.05 (p-Ph), 129.3 (m-Ph), 129.0 (m-Ph), 124.9 (ipso-C of Ph), 111.5, 106.9 (double intensity), 97.7 (Cp), 34.8 (CMe₂), 30.3, 25.2 (C(CH₃)₂), 17.3 (t, ²J_{PC} = 10 Hz, Zr-CH₃), 9.7 (br, [B]-CH₃); one ipso-C of Ph not observed. ³¹P NMR (dichloromethane- d_2 , 81.0 MHz): δ -22.3.

Chlorobis{[[1-methyl-1-(di-*p*-tolylphosphino)ethyl]-η⁵cyclopentadienyl]-k²-P,P }zirconium(IV) Methyltris(pentafluorophenyl)borate (5). A solution of 408 mg (0.80 mmol) of $B(C_6F_5)_3$ in 20 mL of dichloromethane was added to a solution of 304 mg (0.43 mmol) of 2a in 20 mL of dichloromethane at 0 °C. A yellow solution was formed immediately. Removal of the solvent and washing of the product with pentane and drying in vacuo yielded 670 mg of crude 5. Mp. 222 °C (dec). By layering a dichloromethane solution of 5 with pentane at -20 °C, big crystals covered with some oily product could be isolated. Repetition of the crystallization process and washing of the so-formed crystals with cold dichloromethane yielded crystals suitable for an X-ray-diffraction analysis. IR (KBr): $\tilde{\nu}$ 3029, 2993, 2973, 2957, 2924, 2864, 1641, 1598, 1510, 1498, 1459 (br, vs), 1411, 1398, 1384, 1370, 1311, 1295, 1266, 1191, 1088, 1081 993, 976, 965, 947, 935, 830, 802, 766, 757, 736, 707, 659, 631, 607 cm⁻¹. ¹H NMR (dichloromethane-d₂, 200.1 MHz): δ 7.5–7.2 (m, 16H, tol-H), 6.5–6.4 (m, 8H, C₅H₄), 6.22 (m, 4H, C₅H₄), 6.12 (m, 4H, C₅H₄), 2.41 (s, 12H, tol-CH₃), 1.75 (m, 6H, CMe₂), 1.55 (m, 6H, CMe₂), 0.50 (br s, 6H, [B]–CH₃). ¹³C NMR (dichloromethane- d_2 , 150 MHz): δ 148.7 $({}^{1}J_{\text{FC}} = 221 \text{ Hz}, o \cdot C_{6}F_{5}), 142.88, 142.33 \text{ (each ipso-C of p-tol)},$ 139.5 (br s, C1 of Cp), 137.9 (p-C₆F₅), 136.8 (${}^{1}J_{FC} = 249$ Hz, m-C₆F₅), 136.89 (o-tol), 134.2 (o-tol), 130.7 (m-tol), 129.9 (mtol), 129.2 (br, ipso-C of H₃CB(C₆F₅)₃⁻), 126.3 (ipso-C of tol), 122.1 (ipso-C of tol), 116.3, 112.7, 108.3, 100.8 (Cp), 35.73 (CMe₂), 29.75, 26.38 (C(CH₃)₂), 21.70, 21.35 (CH₃) 10.6 ([B]-CH₃). ³¹P NMR (dichloromethane- d_2 , 81.0 MHz): δ -33.0. ¹¹B NMR: δ –15. LDI-TOF MS (negative): m/z = 527Calcd for C₆₃H₅₁BClF₁₅P₂Zr $(H_3CB(C_6F_5)_3^-)$. Anal. (1292.50): C, 58.54; H, 3.98. Found: C, 57.41; H, 3.97.

X-ray Crystal Structure Analysis of 5. Formula $C_{63}H_{51}$ -BClF₁₅P₂Zr; M = 1292.46; $0.40 \times 0.30 \times 0.30$ mm; a = 14.165-(3) Å, b = 14.950(2) Å, c = 16.165(3) Å, $\alpha = 113.94(2)^{\circ}$, $\beta = 109.62(2)^{\circ}$, $\gamma = 95.48(2)^{\circ}$, V = 2838.3(9) Å³, $\rho_{calc} = 1.512$ g cm⁻³, $\mu = 3.88$ cm⁻¹, empirical absorption correction via φ scan data ($0.907 \le C \le 0.999$), Z = 2, triclinic, space group $P\overline{1}$ (No. 2), $\lambda = 0.710$ 73 Å, T = 223 K, $\omega/2\varphi$ scans, 10 438 reflections collected ($-h, \pm k, \pm l$), [(sin $\theta)/\lambda$] = 0.59 Å⁻¹, 9994 independent and 6378 observed reflections [$I \ge 2 \sigma(l)$], 753 refined parameters, R1 = 0.065, wR2 = 0.168, max (min) residual electron density 1.13 (-0.93) e Å⁻³, hydrogens calculated and refined as riding atoms.

Chlorobis{[[1-methyl-1-(di-*p*-tolylphosphino)ethyl]-η⁵cyclopentadienyl]-k²-P,P'}zirconium(IV) Chlorotris(pentafluorophenyl)borate (5'). 1a (200 mg, 0.25 mmol) and 1 equiv of $B(C_6F_5)_3$ (128 mg, 0.25 mmol) were mixed and then dissolved in 20 mL of dichloromethane at room temperature. The yellow solution was stirred for 15 min. The solvent was then removed in vacuo and the residue was washed with two 5 mL portions of pentane and dried in vacuo to yield 290 mg (88%) of 5' as an off-white solid. Crystalline material was obtained by layering a dichloromethane solution of 5' with an equal amount of pentane at -20 °C. Mp 99 °C (dec). IR (KBr): $\tilde{\nu}$ 3032, 2966, 2925, 2871, 1644, 1599, 1515, 1464 (br, vs), 1399, 1381, 1374, 1280, 1263, 1194, 1095 (br, s), 979 (br, s), 825, 806 (s), 772, 765, 668 cm⁻¹. ¹H NMR (dichloromethaned₂, 200.1 MHz): δ 7.50, 7.33, 7.14 (each m, 16H, tol-H), 6.42 (2 m), 6.26, 6.21 (each m, each 2H, C₅H₄), 2.41 (s, 12H, tol-CH₃), 1.75 (m, 6H, CMe₂), 1.51 (m, 6H, CMe₂). ¹³C NMR (dichloromethane- d_2 , 150 MHz): δ 148.5 ($^1J_{FC}$ = 232 Hz, o-C₆F₅), 142.77, 142.27 (each ipso-C of p-tol), (m,p-C₆F₅ could not unambigously be assigned), 139.3 (br s, C1 of Cp), 136.9 (o-tol), 134.2 (o-tol), 130.6 (m-tol), 129.8 (m-tol), 126.2 (ipso-C of tol), 122.2 (ipso-C of tol), 116.2, 112.6, 108.7, 101.1 (Cp), 35.7 (CMe₃), 29.7, 26.4 (C(CH₃)₂), 21.7, 21.3 (each tol-CH₃). ³¹P NMR (dichloromethane- d_2 , 81.0 MHz): δ –33.0. ¹¹B NMR: δ -3. LDI-TOF (negative): m/z = 547 (ClB(C₆F₅)₃⁻). Anal. Calcd for C₆₂H₄₈BCl₂F₁₅P₂Zr (1312.92): C, 56.72; H, 3.68. Found: C, 54.29; H, 3.44.

X-ray Crystal Structure Analysis of 5'. Formula $C_{62}H_{48}$ -BCl₂F₁₅P₂Zr; M= 1312.87; 0.15 × 0.10 × 0.05 mm; a = 14.157-(4) Å, b = 14.964(5) Å, c = 16.122(4) Å, α = 114.37(2)°, β = 109.69(3)°, γ = 95.51(2)°, V = 2817.0(14) Å³, ρ_{calc} = 1.548 g cm⁻³, μ = 4.38 cm⁻¹, empirical absorption correction via φ scan data (0.924 $\leq C \leq$ 0.999), Z = 2, triclinic, space group PI (No. 2), λ = 0.710 73 Å, T = 223 K, ω scans, 9230 reflections collected ($\pm h$, +k, $\pm J$), [(sin θ)/ λ] = 0.57 Å⁻¹, 8846 independent and 4148 observed reflections [$I \geq 2 \sigma(I)$], 756 refined parameters, R1 = 0.057, wR2 = 0.109, max (min) residual electron density 1.05 (-0.48) e Å⁻³, hydrogens calculated and refined as riding atoms.

Low-Temperature Study on the Demethylation– **Chlorination Process in Dichloromethane.** 2a (15.2 mg) and B(C₆F₅)₃ (20.4 mg) were dissolved separately in dichloromethane- d_2 , and the B(C₆F₅)₃ solution was cannula-transferred to the **2a** solution at -78 °C. The first NMR spectrum was obtained at a temperature of -50 °C after keeping the solution for 30 min at -78 °C. ³¹P NMR (dichloromethane- d_2 , 81.0 MHz): δ -22.0 (**3a**), -33 (**5**), -48 (**4**). The following **3a:5:4** ratios were observed by ³¹P NMR spectroscopy subsequently after keeping the same sample for the time indicated at the temperatures noted: *T* = 213 K (30 min), 65:9:26; 223 K (45 min), 51:9:40; 228 K (60 min), 27:11:62; 228 K (90 min), 11:15:74; 238 K (105 min), 9:18:73; 253 K (125 min), 0:20:80; 275 K (170 min) and 295 K (30 min) 0:100:0.

Formation of the Dication 4. Toluene (20 mL) was vacuum-condensed from potassium/sodium alloy onto a mixture of 400 mg (0.57 mmol) of **2a** and 512 mg (1.00 mmol) of $B(C_6F_5)_3$ in a Schlenk tube. A red oily product formed as the reaction mixture was warmed to room temperature. After 15 min at this temperature, the toluene was decanted inside the

glovebox and the remaining red oil was washed twice with 10 mL portions of pentane and then dried in vacuo yielding 850 mg (93%) of **4** as an orange solid. The product is insoluble in toluene and slightly soluble in bromobenzene and was identified by low-temperature NMR in dichloromethane- d_2 and its dichloromethane-stable acetonitrile adduct (**6**), which could also be synthesized on a preparative scale. ¹H NMR of **4** (275 K, dichloromethane- d_2 (spectra were recorded using a 1:1 mixture of **4** and **5**)): 7.6–7.0 (m, 16H, tol-H), 6.85 (m, 4H, C₅H₄), 6.49 (m, 4H, C₅H₄), 2.39 (s, 6H, tol-CH₃), 2.29 (s, 6H, tol-CH₃), 1.93 (m, 6H, CMe₂), 1.59 (m, CMe₂), 0.45 ([B]-CH₃). ³¹P NMR (dichloromethane- d_2 , 81.0 MHz; δ –48.3. Anal. Calcd for C₈₂H₅₄B₂F₃₀P₂Zr (1784.06); C, 55.21; H, 3.05. Found: C, 55.18; H, 3.44.

Preparation of Acetonitrilobis{[[1-methyl-1-(di-ptolylphosphino)ethyl]- η^5 -cyclopentadienyl]- κ^2 -**P**,**P'**}zirconium(IV) Bis(methyltris(pentafluorophenyl)borate) 6. (a) Homogeneously in CH_2Cl_2 : To a solution of 152 mg (0.2 mmol) of 2a in 5 mL of dichloromethane was added a solution of 204 mg of B(C₆F₅)₃ (0.4 mmol) in 20 mL of dichloromethane at -78 °C. The solution turned bright yellow within minutes. The temperature was raised to -50 °C within 1 h. The reaction mixture was stirred for additional 90 min at temperatures around -40 °C. At this temperature, 10.4 μ L (0.2 mmol) of acetonitrile was added and the mixture kept overnight at -20 °C. The solvent was removed in vacuo, yielding 350 mg of a 70:30 mixture of the acetonitrile adduct 6 and the chlorocation 5. (b) Heterogeneously from toluene: A Schlenk flask was charged with 234 mg (0.3 mmol) of 2a and 308 mg (0.6 mmol) of $B(C_6F_5)_3$, and then 10 mL of toluene was added at room temperature in the glovebox. The mixture immediately turned orange with the dication **4** precipitating as a red oil. After 30 min, 16 μ L (0.3 mmol) of acetonitrile was added and the reaction mixture stirred for additional 15 min. The toluene solvent was decanted from the oily product. Washing with 5 mL of toluene and then 5 mL of pentane and drying in vacuo for 24 h led to the formation of a yellow solid product. Yield of 6: 490 mg (89%). Mp 192 °C (dec). IR (KBr): *ṽ* 3122, 2962, 2927, 2867, 2307 (*v*(C≡N), comb.), 2273 $(\nu(C=N))$, 1641, 1599, 1511 (vs), 1458 (br, vs), 1400, 1382, 1267, 1194, 1087 (br, vs), 995, 979, 966, 952, 934, 835, 805, 660 cm⁻¹. ¹H NMR (dichloromethane-d₂, 600 MHz): 7.38 (m, 12H, o, mtol-H), 7.18 (m, 4H, o-tol-H), 6.73, 6.46, 6.38, 6.20 (each m, each 2H, C₅H₄), 2.43, 2.41 (each s, each 6H, tol-CH₃), 1.78 (m, 6H, CMe₂), 1.44 (m, 6H, CMe₂), 1.32 (t, ${}^{5}J_{PH} = 3.3$ Hz, 3H, Zr-NC-CH₃), 0.50 (br, [B]-CH₃). ¹³C NMR (dichloromethane d_2 , 150 MHz): δ 148.7 (${}^1J_{\rm FC}$ = 240 Hz, o-C₆F₅), 145.1, 144.6 (each s, ipso-C of *p*-tol), 141.5 (s, C=N), 137.8 (${}^{1}J_{FC} = 242$ Hz, $p-C_6F_5$), 137.8 (C1 of Cp), 136.7 (${}^1J_{FC} = 245$ Hz, $m-C_6F_5$), 135.5 (o-tol), 134.4 (o-tol), 131.5 (2m-tol), 122.5 (ipso-C of tol), 119.6 (ipso-C of tol), 117.1, 108.5, 107.9, 101.2 (Cp), 36.0 (CMe₂), 29.4, 26.3 (C(CH₃)₂), 21.53, 21.42 (tol-CH₃), 10.5 ([B]-CH₃), 2.92 (s, CH₃CN). GHSQC-NMR²⁴ (dichloromethane- d_2): δ 135.5/7.38, 134.4/7.18, (each o-tol); 131.5/7.38 (m-tol); 117.1/6.73, 108.5/ 6.46, 107.9/6.20, 101.2/6.38, (each C_5H_4); 29.4/1.44, 26.33/1.78(CMe₂); 21.53, 21.42/2.43, 2.41 (each tol-CH₃); 10.5/0.50 ([B]-CH₃); 1.32/2.92 (CH₃CN). GHMBC-NMR²⁴ (dichloromethane- d_2): δ 145.1 (*p*-tol)/7.38 (*o*,*m*-tol-H), 2.43, 2.41 (tol-CH₃); 144.6 (p-tol)/7.18 (o-tol-H), 2.43, 2.41 (tol-CH₃); 141.5 (C=N)/ 1.32 (CH₃CN); 137.8 (C1 of Cp)/6.44, 6.38 (C₅H₄), 1.78, 1.44 (CMe2); 135.5 (o-tol)/7.38 (o,m-tol-H); 134.4 (o-tol)/7.18 (o-tol-H); 131.5 (2-m-tol)/7.38 (o,m-tol-H), 2.43, 2.41 (tol-CH₃); 122.5 (ipso-C of tol)/7.38 (o,m-tol-H); 119.6 (ipso-C of tol)/7.38 (o,mtol-H); 117.1/6.38, 108.5/6.20, 107.9/6.46, 6.38, 101.2/6.20 (each C_5H_4/C_5H_4 ; 36.0 (CMe₂)/1.78, 1.44 (C(CH₃)₂); 29.4 (C(CH₃)₂)/ 1.78 $(C(CH_3)_2)$; 26.3 $(C(CH_3)_2)/1.44$ $(C(CH_3)_2)$; 21.5, 21.4 (tol-CH₃)/7.38 (o,m-tol-H), 2.43, 2.41 (tol-CH₃). ³¹P NMR (dichloromethane- d_2 , 81.0 MHz): δ -44.7. Anal. Calcd for

(24) Braun, S.; Kalinowski, H.; Berger, S. In *100 and More Basic NMR Experiments*; VCH: Weinheim, 1996; and references cited therein.

 $C_{84}H_{57}NB_2F_{30}P_2Zr \ (1825.11); \ C, \ 55.28; \ H, \ 3.15; \ N, \ 0.77. Found: \ C, \ 55.40; \ H, \ 3.53; \ N, \ 0.70.$

((2,6-Dimethylphenyl)isonitrilo)bis{[[1-methyl-1-(di-ptolylphosphino)ethyl]- η^5 -cyclopentadienyl]- κ^2 -P,P'}zirconium(IV) Bis(methyltris(pentafluorophenyl)borate) (7). Formation of the dication 4 in toluene from 234 mg (0.3 mmol) of 2a and 308 mg (0.6 mmol) of B(C₆F₅)₃ followed by addition of 39 mg (0.30 mmol) of (2,6-dimethylphenyl)isonitrile, dissolved in 3 mL of toluene, resulted in the formation of adduct 7 as a red oil. The solvent was decanted, and the product was worked-up as described above. Yield: 470 mg (0.24 mmol, 81%). Mp 168 °C (dec). IR (KBr): v 3123, 2964, 2927, 2871, 2170 (C=N-R), 1641, 1599, 1511 (vs), 1458 (br, vs), 1400, 1382, 1267, 1087 (br, vs), 995, 979, 966, 952, 934, 833, 805, 660 cm⁻¹. ¹H NMR (dichloromethane- d_2 , 600 MHz): 7.37, 7.32, 7.07-7.03 (each m, 19H, tol-H, isonitrile-H), 6.54, 6.46, 6.44, 6.38 (m, each 2H, each C₅H₄), 2.42 (s, 6H, tol-CH₃), 2.06 (s, 6H, CH₃ of isonitrile), 2.02 (s, 6H, tol-CH₃), 1.97 (m, 6H, CMe₂), 1.42 (m, 6H, CMe₂), 0.51 (br, 6H, [B]–CH₃). ¹³C NMR (dichloromethane- d_2 , 150 MHz): δ 148.6 $({}^{1}J_{FC} = 235 \text{ Hz}, o-C_{6}F_{5}), 144.71, 144.57 (p-tol), 137.9 ({}^{1}J_{FC} =$ 242 Hz, p-C₆F₅), 136.7 (${}^{1}J_{FC} = 238$ Hz, m-C₆F₅), 136.4 (C1 of Cp), 135.5 (C2/C6 of isonitrile), 134.45 (o-tol), 134.19 (o-tol), 133.1 (CN-Ph tentatively), 131.4 (m-tol), 130.9 (m-tol), 129.3 (m-CH of isonitrile), 123.3 (ipso-C of tol), 122.8 (C1 of isonitrile), 120.7 (ipso-C of tol), 117.7, 107.3, 101.6, 100.3 (Cp), 36.0 (CMe₂), 28.9, 27.1 (C(CH₃)₂), 21.4, 21.2 (tol-CH₃), 19.2 (CH₃ of isonitrile), 10.5 (br, [B]–CH₃); p-CH of isonitrile not observed, probably hidden by o-tol-CH at 134.5 ppm. GHSQC-NMR²⁴ (dichloromethane- d_2): δ 134.5/7.32 (o-tol), 134.2/7.07 (o-tol), 131.4/7.37 (m-tol), 130.9/7.03 (m-tol), 129.3/7.07 (m-CH of isonitrile) 117.6/6.46, 107.3/6.54, 101.6/6.38, 100.3/6.44 (each C₅H₄), 28.9/1.42, 27.1/1.97 (C(CH₃)₂), 21.4/2.42, 21.2/2.02 (tol-CH₃), 19.2/2.06 (CH₃ of isonitrile), 10.5/0.51 ([B]-CH₃). GH-MBC-NMR²⁴ (dichloromethane- d_2): δ 144.71, 144.57 (p-tol)/ 7.32, 7.07-7.03 (o,m-tol-H), 2.42, 2.02 (tol-CH₃), 136.4 (C1 of Cp)/6.54, 6.46, 6.44, 6.38 (C₅H₄), 1.97, 1.42 (C(CH₃)₂), 135.5 (C2/C6 of isonitrile)/7.32 (CH of isonitrile), 2.06 (CH₃ of isonitrile), 134.5, 134.2 (o-tol)/7.37, 7.32, 7.07-7.03 (o,m-tol-H), 131.4, 130.9 (m-tol)/7.37, 7.03 (o,m-tol-H), 2.42, 2.02 (tol-CH₃), 129.3 (CH of isonitrile)/7.07 (CH of isonitrile), 2.06 (CH₃) of isonitrile), 123.3 (ipso-C of tol)/7.37 (m-tol-H), 1.97, 1.42 (C(CH₃)₂), 122.8 (C1 of isonitrile)/7.07 (m-CH of isonitrile), 120.7 (ipso-C of tol)/7.03 (m-tol-H), 1.42 (C(CH₃)₂), (117.7/6.44, 6.38), (107.3/6.44, 6.38), (101.6/6.54, 6.44), (100.3/6.38) (each C₅H₄/C₅H₄), 36.0 (CMe₂)/1.97, 1.42 (C(CH₃)₂), 28.9 (C(CH₃)₂)/ 1.97 (C(CH₃)₂), 27.08 (C(CH₃)₂)/1.42 (C(CH₃)₂), 21.4, 21.2 (tol-CH₃)/7.37, 7.07-7.03 (o,m-tol-H), 2.42, 2.02 (tol-CH₃), 19.2 (CH₃ of isonitrile)/2.06 (CH₃ of isonitrile). ³¹P NMR (dichloromethane- d_2 , 81.0 MHz): δ -51.7. Anal. Calcd for C₉₁H₆₃-NB₂F₃₀P₂Zr (1915.24): C, 57.07; H, 3.32; N, 0.73. Found: C, 56.70; H, 3.59; N, 0.80.

Acknowledgment. Financial support from the Fonds der Chemischen Industrie, the Ministerium für Wissenschaft und Forschung des Landes Nordrhein-Westfalen, and the Hoechst AG is gratefully acknowledged.

Supporting Information Available: Details about the X-ray crystal structures; including ORTEP diagrams and tables of crystal data and structure refinement, atomic coordinates, bond lengths and angles, anisotropic displacement parameters, and hydrogen coordinates for **1b**, **2b**, **3a**, **5**, and **5**' (52 pages). Ordering information is given on any current masthead page.

OM970573T