

Published online: 12 Aug 2010.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

BIS(DIAZASILAPHOSPHETIDINES) AND THEIR METAL COMPLEXES

Graham R. Lief, Ingo Schranz, and Lothar Stahl

University of North Dakota, Grand Forks, North Dakota, USA

(Received September 4, 2003; accepted October 3, 2003)

The syntheses and solid-state structures of bis(diazasilaphosphetidines) of the type $[\text{Me}_2\text{Si}(\mu\text{-N}^t\text{Bu})_2\text{P}]_2\text{NR}$, $\text{R} = \text{Ph}$, ^iPr , ^tBu , of the P-chiral analogue $[\text{Me}_2\text{Si}(\mu\text{-N}^t\text{Bu})(\mu\text{-N-2,6-}^i\text{PrPh})\text{P}]_2(\text{C}_2\text{H}_4)$, and of some of their metal complexes are reported. The highly electron-rich, modular bis(phosphines) are easy to synthesize and may have applications in catalysis.

Keywords: Bis(diazasilaphosphetidines); bis(phosphines); diazadiphosphetidines; P,N-heterocycles; Si,N,P-heterocycles

Diazadiphosphetidines, ambidentate phosphorus-nitrogen molecules, have been known for some time, but the comparatively low nucleophilicity of their phosphorus atoms has made them poor phosphine ligands, thereby limiting their role to building blocks for polymers¹ and scaffolds for chelating amido ligands (Chart 1, **1** and **2**).^{2,3} Recent studies on tethered bis(diazasilaphosphetidines) (**3**), however, have shown that the silicon-phosphorus analogues are much better phosphorus donors than diazadiphosphetidines.⁴ This electron-richness makes these bis(diazasilaphosphetidines) good candidates for reactions requiring electron-donating phosphines, e.g., homogenous hydrogenations and polyolefin catalysis by late-transition metal complexes.⁵ Thus, for example, $[(\text{Me}_2\text{Si}(\mu\text{-N}^t\text{Bu})_2\text{P})_2\text{C}_2\text{H}_4\text{Rh}]\text{BF}_4$ reduces alkenes to alkanes under mild conditions.

Enantioselective, homogeneous hydrogenations—key steps in the synthesis of fine chemicals—are typically catalyzed by transition-metal complexes bearing C_2 -symmetric ancillary ligands. For this reason we developed a synthetic route for a C_2 -symmetric bis(diazasilaphosphetidine) (Scheme 1) which yields a diphosphine with two

Address correspondence to Lothar Stahl, University of North Dakota, Department of Chemistry, Box 9024, Grand Forks, ND 58202. E-mail: lstahl@chem.und.edu




Downloaded by ["Queen's University Libraries, Kingston"] at 03:29 08 October 2014

Downloaded by ["Queen's University Libraries, Kingston"] at 03:29 08 October 2014



Downloaded by ["Queen's University Libraries, Kingston"] at 03:29 08 October 2014


Downloaded by ["Queen's University Libraries, Kingston"] at 03:29 08 October 2014



SCHEME 2

asymmetrical bis(phosphines) can be synthesized with relative ease. Even amine-linkers of appreciable length, such as 1,6-diaminohexane, may be used to synthesize tetrakis(diazasilaphosphetidino)phosphines for the construction of one-dimensional coordination polymers or for the synthesis of complexes with two remote catalytic sites.

To derivatize these organoaminobis(diazasilaphosphetidines) we have, thus far, synthesized mainly nickel dibromide and molybdenumtetracarbonyl derivatives, Scheme 3, but these ligands are equally well suited for the coordination of zero-valent nickel and other late transition metals.



SCHEME 3

REFERENCES

- [1] T. G. Hill, R. C. Haltiwanger, T. R. Prout, and A. D. Norman, *Inorg. Chem.*, **28**, 3461 (1989).
- [2] G. G. Briand, T. Chivers, and M. Krahn, *Coord. Chem. Rev.*, **233–234**, 237 (2002).
- [3] L. Stahl, *Coord. Chem. Rev.*, **210**, 203 (2000).
- [4] I. Schranz, G. R. Lief, S. J. Midstokke, and L. Stahl, *Inorg. Chem.*, **41**, 6919 (2002).
- [5] N. A. Cooley, S. M. Green, D. F. Wass, K. Heslop, A. G. Orpen, and P. G. Pringle, *Organometallics*, **20**, 4769 (2001).
- [6] M. L. Clark, D. J. Cole-Hamilton, A. M. Z. Slawin, and J. D. Woollins, *Chem. Commun.*, 2065 (2000).