

Published on Web 10/09/2007

Bisphosphine-Catalyzed Mixed Double-Michael Reactions: Asymmetric Synthesis of Oxazolidines, Thiazolidines, and Pyrrolidines

Vardhineedi Sriramurthy, Gregg A. Barcan, and Ohyun Kwon*

Department of Chemistry and Biochemistry, University of California, Los Angeles, California 90095-1569

Received May 24, 2007; E-mail: ohyun@chem.ucla.edu

Five-membered nitrogen-atom-containing heterocycles are structural components of many natural products and pharmaceuticals;¹ in addition, many of them-for example, enantiopure azolidine derivatives-have been employed as synthetic intermediates, auxiliaries, ligands, and catalysts for asymmetric synthesis.² Consequently, there is a high demand for new methods for the efficient construction of optically active azolidine derivatives.³ As part of a program aimed at developing phosphine-mediated annulation reactions,⁴ we sought a novel route toward highly substituted and functionalized five-membered-ring nitrogen-atom-containing heterocycles. In light of recent reports on the phosphine-catalyzed conjugate additions of electron-deficient olefins and acetylenes with alcohols,⁵ herein we report a bisphosphine-catalyzed mixed double-Michael process⁶ that generates azolidines (2; eq 1). Use of aminoacid-derived pronucleophiles (1) as Michael donors and electrondeficient acetylenes as Michael acceptors provides efficient access to azolidines containing both diversity and asymmetry.

$$R \xrightarrow{XH} + \equiv E \xrightarrow{DPPP (10 \text{ mol}\%)} R \xrightarrow{X} E (1)$$

$$1 \text{ XH = OH, SH}_{CH(CO_0Me)_0} 2$$

Our initial evaluation of the proposed double-Michael addition began with the reaction between amino alcohol 1a and methyl propiolate (Table 1). Employing PPh₃ as the catalyst gave the desired double-Michael adduct 2a in 35% yield in addition to a 40% yield of the mono-Michael adduct 3a (entry 1).7 Use of Ph₂-PEt led to a moderate improvement in the yield of the oxazolidine product 2a (entry 2), but none was formed from the reaction catalyzed by Me₃P (entry 3).⁸ In contrast, diphenylphosphinopropane (DPPP) catalysis increased the yield of the desired double-Michael adduct 2a to 71% (entry 4).9 Further increases in the yield and reaction rate were achieved when performing the reaction in a more polar solvent, CH₃CN (entry 5). On the basis of the encouraging results we obtained with DPPP as the catalyst, we also tested the applicability of the homologous bisphosphines diphenylphosphinomethane (DPPM), diphenylphosphinoethane (DPPE), diphenylphosphinobutane (DPPB), and diphenylphosphinopentane (DPPPent). The appreciably poorer yield (37%) of the DPPMmediated reaction, relative to those of the other bisphosphines (entries 6-9), provides a critical clue regarding the reaction mechanism (vide infra).

Scheme 1 presents a plausible mechanism for the dependence of the reaction on both the bidentate nature of the phosphine catalyst and the tether length between the two phosphine moieties. The reaction is triggered by the conjugate addition of the phosphine to the electron-deficient acetylene. The resulting vinyl anion 4 deprotonates the pronucleophile 1, which facilitates the first conjugate addition to form intermediate $6.^{5b,10}$ Upon β -elimination of the phosphine, the mono-Michael product 3 is formed. The presence of an additional phosphine moiety at the optimal distance,

Table 1. Evaluation of Catalysts for Double-Michael Additions ^a						
	+	ent 9 h 2a	CO ₂ Me	NH CO ₂ Me		
			isolated y	vields (%)		
entry	catalyst	solvent	2a	3a		
1^b	Ph ₃ P	toluene	35	40		
2^b	Ph ₂ PEt	toluene	42	30		
3^b	Me ₃ P	toluene	0	22		
4^b	DPPP	toluene	71	12		
5	$DPPP^{c}$	CH ₃ CN	92	0		
6	DPPM ^c	CH ₃ CN	37	42		
7	$DPPE^{c}$	CH ₃ CN	84	0		
8	$DPPB^{c}$	CH ₃ CN	82	0		
9	DPPPent ^c	CH ₃ CN	79	6		

^{*a*} All reactions were performed using 1 mmol of **1a**, 1.1 mmol of methylpropiolate, and 10 mol % of the catalyst. ^{*b*} These reactions were run for 48 h. ^{*c*} DPPM, DPPE, DPPP, DPPB, and DPPPent are acronyms for diphenylphosphinomethane, -ethane, -propane, -butane, and -pentane, respectively.

as in DPPP, provides additional stabilization to the intermediate phosphonium ions **6** and **7**. The latter undergoes S_N2 displacement to produce the cyclized product **2**.¹¹ In the absence of anchimeric assistance, as in the case of the monodentate phosphines, the decreased stability of the phosphonium ion led to an unfavorable equilibrium for the formation of **6** from **3**.¹² The relatively short tether of DPPM prohibits the orbital overlap required for anchimeric assistance because of geometrical constraints. The other phosphines for which intramolecular stabilization was possible, namely, DPPE, DPPB, and DPPPent, gave results similar to those obtained using DPPP. Note that intramolecular stabilization of phosphonium ions by nitrogen atoms has precedent in the literature.¹³

With the optimal reaction conditions in hand, that is, DPPP as catalyst and CH₃CN as solvent, we next explored the scope of the double-Michael reaction using a variety of amino-acid-derived pronucleophiles and electron-deficient acetylenes (Table 2). The formation of oxazolidines from β -amino alcohols and methyl propiolate proceeded smoothly, with high yields and diastereose-lectivities (entries 1 and 4). The Michael acceptors acetylacetylene and tosylacetylene also gave good results (entries 2 and 3). This methodology works well for the syntheses of thiazolidines from β -amino thiols (entries 5–7).¹⁴ All of the substrates provided similarly high yields and diastereoselectivities for the formation of thiazolidines.¹⁵

We further tested the generality of our reaction by using carbonucleophiles (entries 8–10) for the preparation of pyrrolidine derivatives, which are ubiquitous in natural products of pharma-cological interest.¹⁶ Under the optimized conditions, we generated the pyrrolidines **2h** and **2i** from the valine-derived γ -amino malonate **1h** (entries 8 and 9, respectively).¹⁷ Employing the cyclic γ -amino diester **1j** furnished the octahydroindole derivative **2j** as a single diastereoisomer in good yield (entry 10). Octahydroindoles, which

Scheme 1. Proposed Mechanism for the Formation of 2



Table 2. Syntheses of Various Azolidines^a

entry	substrate	product	yield $\%^{b}$
		0	(cis.traits)
1	^{i-Pr} OH NHTs 1a	i-Pr	92 (96:4)
2	1a		92 (94:6)
3	1a	iPr N Ts 2c	87 (97:3)
4	Bn OH NHTs 1d	$Bn \xrightarrow{N}_{Ts} 2d CO_2 Me$	91 (95:5)
5	^{i-Pr} SH NHTs 1e	i-Pr N S O Ts 2e	93 (95:5)
6	1e	HPr N S 2f	89 (96:4)
7	Ph SH NHTs 1g	Ph N $2g$ S 0 Ts $2g$	88 (96:4)
8	i-Pr TsHN CO ₂ Me 1h	MeO ₂ C i-Pr N Ts 2h	82 (94:6)
9	1h	MeO ₂ C CO ₂ Me	91 (95:5)
10	CO ₂ Me CO ₂ Me NHTs 1j	$\underbrace{MeO_{2}C}_{N} \underbrace{CO_{2}M}_{O} \underbrace{O}_{O} \underbrace{O} \underbrace{O}$	80 (100:0)

^{*a*} All reactions were performed using 1 mmol of the substrate, 1.1 equiv of the corresponding acetylene, and 10 mol % of DPPP in CH₃CN at 80 °C for 9 h. ^{*b*} Isolated yields after chromatographic purification. ^{*c*} Determined through ¹H NMR spectroscopic analysis.

are present in a large number of natural products, are often challenging synthetic targets.¹⁸

In summary, we have developed a remarkably simple protocol for the synthesis of oxazolidines, thiozolidines, pyrrolidines, and octahydroindoles. This mixed double-Michael process operates best under bisphosphine catalysis to provide β -amino carbonyl derivatives of azolidines¹⁹ in excellent yields and with high diastereoselectivities. Presumably, the use of bis(diphenylphosphine) derivatives allows intramolecular stabilization of the phosphonium ion intermediates. We are currently exploring the development of an enantioselective version of this ring-forming process from achiral starting materials and its application to the synthesis of selected drug candidates.

Acknowledgment. This study was funded by the NIH (R01GM071779). We thank Dr. Saeed Khan for performing the

crystallographic analyses. O.K. thanks Dr. Patrick J. Walsh, Richard P. Hsung, and Chulbom Lee for helpful discussions.

Supporting Information Available: Representative experimental procedures and spectral data for all new compounds (PDF). Crystallographic data for **2a** and **3a** (CIF). This material is available free of charge via the Internet at http://pubs.acs.org.

References

- (1) Butler, M. S. J. Nat. Prod. 2004, 67, 2141.
- (2) (a) Comprehensive Asymmetric Catalysis I-III; Jacobsen, E. N., Pfaltz, A., Yamamoto, H., Eds.; Springer: Berlin, 1999. (b) Asymmetric Synthesis, 2nd ed.; Ojima, I., Ed.; VCH: New York, 2000. (c) Fache, F.; Schultz, E.; Tommasino, M. L.; Lemaire, M. Chem. Rev. 2000, 100, 2159. (d) Chiral Reagents for Asymmetric Synthesis; Paquette, L. A., Ed.; Wiley: Chichester, 2003. (e) Seayad, J.; List, B. Org. Biomol. Chem. 2005, 3, 719.
- (3) Katritzky, A. R. *Chem. Rev.* 2004, 104, 2125–2812 (special issue on heterocycles).
- (4) (a) Zhu, X.-F.; Henry, C. E.; Kwon, O. J. Am. Chem. Soc. 2007, 129, 6722. (b) Castellano, S.; Fiji, H. D. G.; Kinderman, S. S.; Watanabe, M.; de Leon, P.; Tamanoi, F.; Kwon, O. J. Am. Chem. Soc. 2007, 129, 5843. (c) Mercier, E.; Fonovic, B.; Henry, C. E.; Kwon, O.; Dudding, T. Tetrahedron Lett. 2007, 48, 3617. (d) Dudding, T.; Kwon, O.; Mercier, E. Org. Lett. 2006, 8, 3643. (e) Tran, Y. S.; Kwon, O. Org. Lett. 2005, 7, 4289. (f) Zhu, X.-F.; Henry, C. E.; Wang, J.; Dudding, T.; Kwon, O. Org. Lett. 2005, 7, 1387. (g) Zhu, X.-F.; Schaffner, A.-P.; Li, R. C.; Kwon, O. Org. Lett. 2005, 61, 6276. (i) Zhu, X.-F.; Henry, C. E.; Kwon, O. Tetrahedron 2005, 61, 6276. (i) Zhu, X.-F.; Lan, J.; Kwon, O. J. Am. Chem. Soc. 2003, 125, 4716.
- (5) (a) Stewart, I. C.; Bergman, R. G.; Toste, F. D. J. Am. Chem. Soc. 2003, 125, 8696. (b) Inanaga, J.; Baba, Y.; Hanamoto, T. Chem. Lett. 1993, 2, 241.
- (6) Intramolecular double-Michael reactions of acetylenes. Double-carbo-Michael: (a) Grossman, R. B. Synlett 2001, 13. Double-thia-Michael: (b) Kuroda, H.; Tomita, I.; Endo, T. Synth. Commun 1996, 26, 1539. Double-oxa-Michael: (c) Ariza, X.; Costa, A. M.; Faja, M.; Pineda, O.; Vilarrasa, J. Org. Lett. 2000, 2, 2809. (d) Forsyth, C. J.; Hao, J.; Aiguade, J. Angew. Chem., Int. Ed. 2001, 40, 3663. Double-aza-Michael: (e) Diez-Barra, E.; Guerra, J.; Hornillos, V.; Merino, S.; Tejeda, J. Tetrahedron Lett. 2004, 45, 6937.
- (7) The structures of 2a and 3a were confirmed through X-ray crystallographic analysis. See the Supporting Information for details.
- (8) Other bases known to facilitate Michael addition of alcohols to alkynes, such as K₂CO₃ and morpholine, did not provide any cyclized product. (a) Eaton, P. E.; Stubbs, C. E. J. Am. Chem. Soc. **1967**, 89, 5722. (b) Lee, E.; Tae, J. S.; Lee, C.; Park, C. M. Tetrahedron Lett. **1993**, *34*, 4831.
- (9) For examples of the use of bisphosphines in organocatalysis, see: (a) Trost, B. M.; Li, C.-J. J. Am. Chem. Soc. 1994, 116, 10819. (b) Trost, B. M.; Dake, G. R. J. Am. Chem. Soc. 1997, 119, 7595. (c) Kamijo, S.; Kanazawa, C.; Yamamoto, Y. Tetrahedron Lett. 2005, 46, 2563. (d) Silva, F.; Sawicki, M.; Gouverneur, V. Org. Lett. 2006, 8, 5417.
 (10) Alternatively, the deprotonated nucleophile can add to the acetylene to
- (10) Alternatively, the deprotonated nucleophile can add to the acetylene to form a vinyl anion, which will deprotonate the pronucleophile and form 3. In this scenario, the phosphine is an initiator for the actual catalytic cycle. See: Grossman, R. B.; Comesse, S.; Rasne, R. M.; Hattori, K.; Delong, M. N. J. Org. Chem. 2003, 68, 871 and references therein.
- (11) Alternatively, one can invoke a mechanism where intermediate 6 functions as a base to deprotonate 3, which subsequently undergoes intramolecular Michael addition. The resulting enolate deprotonates another molecule of 3 to reiterate the productive cycle. Ion paring between the protonated 6 and deprotonated 3 (and the following intermediates) has been proposed previously; see: (a) ref 5a. (b) White, D. A.; Baizer, M. M. *Tetrahedron Lett.* **1973**, *14*, 3579.
- (12) The β -alkoxy acrylate intermediate **3a** smoothly converted to the cyclized product **2a** in 94% yield when treated with DPPP.
- (13) Although hexamethylphosphorous triamide (HMPT) did not facilitate any Michael reaction, Verkade's proazaphosphatrane (10 mol %) provided the desired cyclized product in 42% yield, indicating that other donators of electron lone pairs can provide similar anchimeric assistance. (a) Kisanga, P. B.; Ilankumaran, P.; Fetterly, B. M.; Verkade, J. G. J. Org. Chem. 2002, 67, 3555. (b) Verkade, J. G.; Kisanga, P. B. Aldrichimica Acta 2004, 37, 3.
- (14) Anderson, J. C.; Cubbon, R.; Harding, M.; James, D. S. Tetrahedron: Asymmetry 1998, 9, 3461.
- (15) 1,2-Diamine pronucleophiles are known to provide piperazines under phosphine catalysis; see: Lu, C.; Lu, X. Org. Lett. 2002, 4, 4677.
 (16) (a) O'Hagan, D. Nat. Prod. Rep. 2000, 17, 435. (b) Daly, J. W.; Spande,
- (16) (a) O Hagan, D. Nat. Prod. Rep. 2000, 17, 453. (b) Daly, J. W.; Spande, T. F.; Garraffo, H. M. J. Nat. Prod. 2005, 68, 1556.
 (17) W. H. H. G. J. Daly, J. W. H. F. G. L. Charles, M. F. F.
- (17) Mao, H.; Joly, G. J.; Peeters, K.; Hoornaert, G. J.; Compernolle, F. *Tetrahedron* 2001, 57, 6995.
 (18) (a) Kobayashi, L: Morita, H. In *The Alkalaids*: Cordell, G. A. Ed.;
- (18) (a) Kobayashi, J.; Morita, H. In *The Alkaloids*; Cordell, G. A., Ed.; Academic: New York, 2003; Vol. 60, p 165. (b) Hanessian, S.; Tremblay, M.; Petersen, J. F. W. *J. Am. Chem. Soc.* 2004, *126*, 6064. (c) Aron, Z. D.; Overman, L. E. Org. Lett, 2005, 7, 913.
- (19) Fustero, S.; Jimenez, D.; Sanchez-Rosello, M.; del Pozo, C. J. Am. Chem. Soc. 2007, 129, 6700.

JA073754N