

Published on Web 03/08/2006

## A Very Simple, Highly Stereoselective and Modular Synthesis of Ferrocene-Based P-Chiral Phosphine Ligands

Weiping Chen,\*,† William Mbafor, Stanley M. Roberts, and John Whittall

StylaCats Ltd., Chemistry Department, University of Liverpool, Liverpool L69 7ZD, United Kingdom

Received December 16, 2005; E-mail: weiping.chen@solvias.com

5

6

7

8

Ph

Ph

Cy

Ph

2-Np

Ph

Cy

2-Biph

Ferrocene-based ligands provide an important subset of chiral phosphines,<sup>1</sup> and several families with subtle structural variations, such as Josiphos,<sup>2</sup> Taniaphos,<sup>3</sup> Mandyphos,<sup>4</sup> Walphos,<sup>5</sup> Trap,<sup>6</sup> and BoPhoz,<sup>7</sup> have been developed in the past few years. Most of these ligands incorporate both carbon-centered chirality and planar chirality, and they have been proven to be highly effective in numerous asymmetric reactions. In stark contrast, much less attention has been paid to P-chiral phosphines bearing ferrocenyl groups, although, in theory, they are highly likely to provide a superior class of ligands for asymmetric catalysis by virtue of bringing the chiral environment into the closest possible proximity to the catalytic center. The paucity of examples undoubtedly reflects the difficulties in synthesis of P-chiral phosphines.<sup>8</sup>

Almost all of the documented ferrocene-based planar chiral phosphine ligands are synthesized by chiral group-directed diastereoselective ortho-lithiation9 or enantioselective lithiation10 followed by reaction with a monochlorophosphine. We envisioned that ferrocene-based P-chiral phosphines could be synthesized by reaction of a *dichlorophosphine* with a chiral lithiated ferrocene, followed by a second organometallic reagent. Herein we report a first example of highly stereoselective and modular synthesis of ferrocene-based P-chiral phosphine ligands using this simple and straightforward strategy.

Enantiopure *N*,*N*-dimethyl-1-ferrocenylethylamine (Ugi's amine) is readily available<sup>11</sup> and provided a suitable starting material for our investigations. Thus, (R)-Ugi's amine 1 was lithiated with t-BuLi, followed by reaction with PhPCl<sub>2</sub> and then reacted with o-anisyllithium unexpectedly (and most pleasingly!) to afford a single diastereomer ( $R_C$ , $S_{Fc}$ , $S_P$ )-2a (as judged by <sup>1</sup>H NMR and <sup>31</sup>P NMR spectroscopy) in 91% yield (Table 1).12 The absolute configuration of the product  $(R_C, S_{EC}, S_P)$ -2a was determined by single-crystal X-ray diffraction analysis. Similarly, the naphthalene derivative  $(R_{C}, S_{Fc}, S_{P})$ -2b was prepared as a single diastereomer in 89% yield. Recrystallization of  $(R_{\rm C}, S_{\rm Fc}, S_{\rm P})$ -2b proved to be difficult as it is very soluble in every solvent tested. By way of contrast, its epimer,  $(R_{\rm C}, S_{\rm Fc}, R_{\rm P})$ -2c, is almost insoluble in hot hexane and is easily prepared from  $(R_{\rm C}, S_{\rm Fc}, S_{\rm P})$ -2b by crystallization induced asymmetric transformation. The absolute configuration of  $(R_{\rm C},S_{\rm C})$  $_{\rm Fc}$ ,  $R_{\rm P}$ )-2c was determined by single-crystal X-ray diffraction analysis

This new synthetic methodology is highly modular. The scope of the above reaction was extended by using different dichlorophosphines R<sup>1</sup>PCl<sub>2</sub> and organometallic compounds R<sup>2</sup>M (Table 1) to provide a variety of ferrocene-based P-chiral phosphines each as a single diastereomer in high yield. Replacement of the lithium reagent with the corresponding Grignard reagent gave similar results (entry 2 vs entry 3). More importantly, the two diastereomers with the opposite configuration at the phosphorus atom can be prepared simply by exchanging  $R^1$  and  $R^2$  in  $R^1PCl_2$  and  $R^2M$  (entry 7 vs

	Fe (R)	Me NMe <sub>2</sub>	1) t-BuLi, -7 2) R1PCl2, - 3) R <sup>2</sup> M, -78	78 °C~rt -78 °C~rt ℃~rt ℃~rt		NMe <sub>2</sub> 1 2
entry	R1	R <sup>2</sup>	М	yield (%)	product	configuration
1	Ph	o-An	Li	91	2a	$(R_{\rm C}, S_{\rm Fc}, S_{\rm P})$
2	Ph	1-Np	Li	89	2b	$(R_{\rm C}, S_{\rm Fc}, S_{\rm P})$
3	Ph	1-Np	MgBr	83	2b	$(R_{\rm C}, S_{\rm Fc}, S_{\rm P})$
4	1-Np	Ph	•	78	$2c^a$	$(R_{\rm C}, S_{\rm Fc}, R_{\rm P})$

Table 1. Synthesis of Ferrocene-Based P-Chiral Phosphines

9	Ph	Me	MgB	r 89	2h	$(R_{\rm C}, S_{\rm Fc}, R_{\rm P})$ $(R_{\rm C}, S_{\rm Fc}, R_{\rm P})$
10	<i>t</i> -Bu	Me	Li	86	2i	
a (R	$C_{\rm C}, S_{\rm Fc}, R_{\rm P})$ -	2c was	prepared t	from $(R_{\rm C}, S_{\rm F})$	-c,S <sub>P</sub> )- <b>2b</b>	by crystallization

87

94

95

92

2d

2e

2f

2g

 $(R_{\rm C}, S_{\rm Fc}, S_{\rm P})$ 

 $(R_{\rm C}, S_{\rm Fc}, S_{\rm P})$ 

 $(R_{\rm C}, S_{\rm Fc}, S_{\rm P})$ 

 $(R_{\rm C}, S_{\rm Fc}, R_{\rm P})$ 

MgBr

MgCl

Li MgBr

induced asymmetric transformation (CIAT).

entry 8). The absolute configuration of each compound in the series 2d-2i was inferred from the established stereochemistry of 2a and 2c. We have also extended the scope of this reaction by using other ferrocenyllithium species generated using, initially, the classical chiral group directed ortho-lithiation, and the results will be reported elsewhere.

The stereochemistry and high diastereoselectivity of this reaction can be explained by the following plausible reaction mechanism (Scheme 1). For the substrates bearing groups which are strong Lewis bases (such as the dimethylamino unit in Ugi's amine), a five-membered ring quaternary ammonium salt intermediate, such as 3, is formed and then the organometallic compound RM attacks from the front (as portrayed) to give the products.





Compound 2 is useful as a starting point for expanding the range of novel compounds of potential utility. Thus several new families of ferrocene-based phosphine ligands, combining phosphorus chirality and other chiralities, such as P-chiral phosphine-aminophosphine ligands 4a-e, P-chiral phosphine-phosphoramidite ligands 5a,b,d, and P-chiral 1,3-bisphosphine ligands 6a-e (Scheme 2), can be prepared easily by similar methods to those used for the preparation of Josiphos<sup>4</sup> and BoPhoz.<sup>9</sup> It should be noted that epimerization at the phosphorus stereocenter occurred when the transformations were performed at elevated temperature. Fortu-

<sup>&</sup>lt;sup>†</sup> Present address: Solvias AG, WRO-1055.6.68B, Mattenstrasse 22, Postfach, CH-4002 Basel, Switzerland.

Scheme 2



Table 2.	Asymmetric Hydrogenation of $\alpha$ -Dehydroamino A	cid
Derivative	es l	

	NHAc	H <sub>2</sub>	, L*-Rh(CO	NHAc				
R	CO <sub>2</sub> Me		solvent, rt				CO <sub>2</sub> Me	
entry	L*	R	solvent	H <sub>2</sub> (psi)	time (h)	conv. (%)	ee (%)	
1	BoPhoz	Н	THF	50	2.5	99	94.5	
2	4b	Н	THF	50	2.5	100	98.3	
3	4c	Н	THF	50	2.5	100	92.3	
5	BoPhoz	Н	MeOH	50	2.5	91.4	81.1	
6	4b	Н	MeOH	50	2.5	100	>99	
7	4c	Н	MeOH	50	2.5	84.5	90.3	
8	BoPhoz	Ph	THF	100	2	100	95.7	
10	4b	Ph	THF	100	2	100	97.0	
11	<b>4</b> c	Ph	THF	100	2	100	92.4	

nately, most of transformations can be carried out at room temperature, with no evidence of any epimerization.

The preliminary results of asymmetric hydrogenation of two  $\alpha$ -dehydroamino acid derivatives catalyzed by Rh complexes prepared from the new ligands show that the introduction of P-chirality in ferrocene-based phosphine ligands enhances the enantioselective discrimination produced by the catalyst when a cooperative effect operates among the planar chirality, carbon chirality, and the chirality at phosphorus (Table 2). The activities and enantioselectivities of the matched ligand  $(R_C, S_{Fc}, S_P)$ -4b are higher than that of BoPhoz (entry 1 vs entry 2 and entry 8 vs entry 10), while the mismatched ligand  $(R_{\rm C}, S_{\rm Fc}, R_{\rm P})$ -4c is less active and enantioselective than BoPhoz (entry 1 vs entry 3 and entry 8 vs entry 11). Interestingly, the matched ligand  $(R_C, S_{Fc}, S_P)$ -4b seems to be particularly suitable for hydrogenation conducted in methanol (entry 5 vs entry 6).

In summary, a very simple, highly stereoselective synthesis of ferrocene-based P-chiral phosphine ligands has been developed. This new methodology offers modular construction of ligands and hence generates many possibilities for modifying the structure of these ferrocene-based phosphines. Indeed, on the basis of this new method, several new families of ferrocene-based phosphine ligands

combining phosphorus chirality and other chiralities have been prepared. The preliminary results from some asymmetric hydrogenation reactions show that the introduction of P-chirality into well-known ferrocene-based phosphine ligands enhances the enantioselective discrimination produced by the corresponding Rh catalyst when a matching among the planar chirality, carbon chirality, and the chirality at phosphorus exists.

Acknowledgment. We thank Dr. Jamie Bickley of Liverpool University for the crystal structure analysis, and Dr. David Justice for the synthesis of dehydrophenylalanine.

Supporting Information Available: Experimental procedures and confirmation of the absolute configuration of  $(R_C, S_{Fc}, S_P)$ -2a,  $(R_C, S_{Fc}, R_P)$ -**2c**,  $(R_C, S_{Fc}, S_P)$ -**2a**, and  $(R_C, S_{Fc}, S_P)$ -**6b**. This material is available free of charge via the Internet at http://pubs.acs.org.

## References

- For reviews, see: (a) Richards, C. J.; Locke, A. J. *Tetrahedron: Asymmetry* **1998**, *9*, 2377. (b) Colacot, T. J. *Chem. Rev.* **2003**, *103*, 3101. (c) Dai, L. X.; Tu, T.; You, S. L.; Deng, W. P.; Hou, X. L. Acc. Chem. Res. **2003**, *36*, 659. (d) Barbaro, P.; Bianchini, C.; Giambastiani, G.; Parisel, S. L. Coord. Chem. Rev. 2004, 248, 2131.
- (2) Togni, A.; Breutel, C.; Schnyder, A.; Spindler, F.; Landert, H.; Tijani, A. J. Am. Chem. Soc. 1994, 116, 4062.
- Lotz, M.; Polborn, K.; Knochel, P. Angew. Chem., Int. Ed. 2002, 41, 4708.
  Perea, A. J. J.; Borner, A.; Knochel, P. Tetrahedron Lett. 1998, 39, 8073.
- (5) Weissensteiner, W.; Sturm, T.; Spindler, F. Adv. Synth. Catal. 2003, 345, 160.
- (6) Sawamura, M.; Hamashima, H.; Sugawara, M.; Kuwano, N.; Ito, Y. Organometallics 1995, 14, 4549.
- Boaz, N. W.; Debenham, S. D.; Mackenzie, E. B.; Large, S. E. Org. Lett. (7)2002. 4. 2421
- (8) Some phosphines incorporating phosphorus chirality and other chiralities are known. For phosphines incorporating both planar and phosphorus chirality, see: (a) Nettekoven, U.; Widhalm, M.; Kamer, P. C. J.; van Leeuwen, P. W. N. M.; Mereiter, K.; Lutz, M.; Spek, A. L. Organome*tallics* **2000**, *19*, 2299. (b) Nettekoven, U.; Kamer, P. C. J.; Widhalm, M.; van Leeuwen, P. W. N. M. *Organometallics* **2000**, *19*, 4596. (c) Nettekoven, U.; Widhalm, M.; Kalchhauser, H.; Kamer, P. C. J.; van Leeuwen, P. W. N. M.; Lutz, M.; Spek, A. L. J. Org. Chem. 2001, 66, 759. For phosphines combining carbon, planar, and phosphorus chirality, see: (d) Barbaro, P.; Bianchini, C.; Giambastiani, G.; Togni, A. *Chem.* Commun. 2002, 2672
- (a) Hayashi, T.; Mise, T.; Fukushima, M.; Kagotani, M.; Nagashima, N.; Hamada, Y.; Matsumoto, A.; Kawakami, S.; Konishi, M. M.; Yamamoto, K.; Kumada, M. Bull. Chem. Soc. Jpn. 1980, 53, 1138. (b) Riant, O.; Argouarch, G.; Guillaneux, D.; Samuel, O.; Kagan, H. B. J. Org. Chem. 1998, 63, 3511. (c) Riant, O.; Samuel, O.; Flessner, T.; Taudien, S.; Kagan, H. B. J. Org. Chem. **1997**, 62, 6733. (d) Richards, J.; Damalidis, T.; Hibbs, D. E.; Hursthouse, M. B. Synlett **1995**, 74. (e) Sammakai, T.; Latham, H. D. E., Huisthouse, M. B. Synten 1995, 14: (c) Sammakai, T., Lamani, H.
   A.; Schaad, D. R. J. Org. Chem. 1995, 60, 10. (f) Nishibayashi, Y.;
   Uemura, S. Synlett 1995, 79. (g) Sammakai, T.; Latham, H. A. J. Org.
   Chem. 1995, 60, 6002. (h) Ganter, C.; Wagner, T. Chem. Ber. 1995, 128, 1157. (i) Enders, D.; Peters, R.; Lochtman, R.; Runsink, J. Eur. J. Org.
   Chem. 2000, 2839. (j) Lotz, M.; Ireland, T.; Tappe, K.; Knochel, P. Chirality 2000, 12, 389. (k) Kitzler, R.; Xiao, L.; Weissensteiner, W. Tetrahedron: Asymmetry 2000, 11, 3459. (1) Widhalm, M.; Mereiter, K.;
- Bourghida, M. *Tetrahedron: Asymmetry* 1998, 9, 2983.
  (10) (a) Nishibayashi, Y.; Arikawa, Y.; Ohe, K.; Uemura, S. J. Org. Chem. 1996, 61, 1172. (b) Tsukazaki, M.; Tinkl, M.; Roglans, A.; Chapell, B. J.; Taylor, N. J.; Snieckus, V. J. Am. Chem. Soc. 1996, 118, 685. (c) Jendralla, H.; Paulus, E. Synlett 1997, 471. (d) Price, D.; Simpkins, N. S. Tetrahedron Lett. 1995, 36, 6135
- (11) (a) Marquarding, D.; Klusacek, H.; Gokel, G.; Hoffmann, P.; Ugi, I. J. Am. Chem. Soc. **1970**, 92, 5389. (b) Boaz, N. W. Tetrahedron Lett. **1989**, 30, 2061.
- (12) Bulter, I. R.; Cullen, W. R.; Rettig, S. Organometallics 1986, 5, 1320. JA058531G