


This article was downloaded by: [The University of Manchester Library]

On: 01 December 2014, At: 23:33

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Nucleosides and Nucleotides

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn19>

### Synthesis of New Acyclic Nucleoside Vinyl Phosphonates

Annamaria Esposito<sup>a</sup> & Maurizio Taddei<sup>a</sup>

<sup>a</sup> Dipartimento di Chimica, Università di Sassari, Via Vienna 2, 07100, Sassari, Italy

Published online: 04 Oct 2006.

To cite this article: Annamaria Esposito & Maurizio Taddei (1999) Synthesis of New Acyclic Nucleoside Vinyl Phosphonates, *Nucleosides and Nucleotides*, 18:4-5, 985-986, DOI: [10.1080/15257779908041622](https://doi.org/10.1080/15257779908041622)

To link to this article: <http://dx.doi.org/10.1080/15257779908041622>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

## SYNTHESIS OF NEW ACYCLIC NUCLEOSIDE VINYL PHOSPHONATES


Annamaria Esposito and Maurizio Taddei\*

Dipartimento di Chimica, Università di Sassari, Via Vienna 2, 07100 Sassari Italy


**ABSTRACT:** Chiral vinyl phosphonates, homomorphous with natural nucleotides, have been prepared starting from (*S*)-malic acid as potential monomers for the synthesis of acyclic polynucleotides.

Nucleoside phosphonates have been considered for isosteric alternatives of nucleoside phosphates as antiviral agents.<sup>1</sup> Enantiomerically pure (*S*)-phosphonomethoxypropyl derivatives of purines exhibited a remarkable activity against several DNA and RNA viruses.<sup>2</sup> As a part of our interest in the field of DNA analogues,<sup>3</sup> we decided to prepare simple nucleoside phosphonates homomorphous with natural nucleotides. These new compounds would be acyclic and would contain an element of rigidity as a *trans* double bond. The synthesis started from (*S*)-malic acid **1** that was transformed into the corresponding diol **2** following standard conditions.<sup>4</sup> Product **2** was further protected as a cyclic acetal using benzaldehyde dimethyl acetal or *p*-methoxybenzaldehyde dimethyl acetal in the presence of canphorsulfonic acid (CSA). Dioxolanes **3** or **4** were reduced to the corresponding aldehydes with DIBAL-H in toluene at -78°C and the aldehydes **5** and **6** underwent Horner-Emmons type reaction with the anion of tetraisopropyl methylene-diphosphonate to give products **7** and **8**.

Scheme 1



We first tried to selectively deprotect the dioxolane to produce a diol having the secondary OH protected. Unfortunately any attempt to perform this reaction (using DIBAL-H at room temperature or NaCNBH<sub>3</sub>/ TMSCl)<sup>5</sup> did not give any result. Thus we decided to completely deprotect compound **7** to give diol **9** that was first protected at the primary OH as TBDMS ether (TBDMSCl, DMF, imidazole) then protected at the secondary OH as MOM ether (MOMCl, DIPEA, DMAP) and finally deprotected at the primary OH (TBAF, THF).



Compound **9** was transformed into the corresponding mesylate (MsCl, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>) and the reaction with adenine was carried out in DMF at 80°C in the presence of Cs<sub>2</sub>CO<sub>3</sub>. Product **10** was isolated by crystallisation and resulted a single isomer at <sup>1</sup>H and <sup>13</sup>C NMR analysis. Product **10** could be further completely deprotected at the hydroxyl and phosphonate functions to give a nucleoside phosphonate or could be selectively deprotected at one of these two functions for the preparation of modified oligonucleotides.

## REFERENCES

1. De Clercq E. *J. Med. Chem.* **1995**, *38*, 2491.
2. Holy, A.; Dvorakova, H.; Jindrich, J.; Masojidkova, M.; Budesinsky, M.; Balzarini, J.; Andrei, G.; De Clercq, E. *J. Med. Chem.* **1996**, *39*, 4073.
3. Ciapetti, P.; Taddei, M. *Tetrahedron* **1998**, *54*, 11305
4. Wunsch, B.; Diekmann, H., Höfner, G. *Liebigs Ann. Chem.* **1993**, 1273.
5. Johansson, R., Samuelsson, B. *J. Chem. Soc., Perkin Trans. I* **1984**, 2371.