

Pseudotetrahedral Manganese Complexes Supported by the Anionic Tris(phosphino)borate Ligand [PhBP<sup>Pr</sup><sub>3</sub>]

Connie C. Lu and Jonas C. Peters\*

*Contribution from the Division of Chemistry and Chemical Engineering, Arnold and Mabel Beckman Laboratories of Chemical Synthesis, California Institute of Technology, Pasadena, California 91125*

Received April 29, 2006

This paper presents aspects of the coordination chemistry of mono- and divalent manganese complexes supported by the anionic tris(phosphino)borate ligand, [PhBP<sup>Pr</sup><sub>3</sub>] (where [PhBP<sup>Pr</sup><sub>3</sub>] = [PhB(CH<sub>2</sub>P<sup>Pr</sup><sub>2</sub>)<sub>3</sub>]<sup>−</sup>). The Mn(II) halide complexes, [PhBP<sup>Pr</sup><sub>3</sub>]MnCl (**1**) and [PhBP<sup>Pr</sup><sub>3</sub>]MnI (**2**), have been characterized by X-ray diffraction, SQUID magnetometry, and EPR spectroscopy. Compound **2** serves as a precursor to a series of Mn azide, alkyl, and amide species: [PhBP<sup>Pr</sup><sub>3</sub>]Mn(N<sub>3</sub>) (**3**), [PhBP<sup>Pr</sup><sub>3</sub>]Mn(CH<sub>2</sub>Ph) (**4**), [PhBP<sup>Pr</sup><sub>3</sub>]Mn(Me) (**5**), [PhBP<sup>Pr</sup><sub>3</sub>]Mn(NH(2,6-*i*-Pr<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>)) (**6**), [PhBP<sup>Pr</sup><sub>3</sub>]Mn(dbabh) (**7**), and [PhBP<sup>Pr</sup><sub>3</sub>]Mn(1-Ph(isoindolate)) (**8**). The complexes **2–8** feature a divalent-metal center and are pseudotetrahedral. They collectively represent an uncommon structural motif for low-coordinate, polyphosphine-supported Mn complexes. Two Mn(I) species have also been prepared. These include the Ti–Mn adduct [PhBP<sup>Pr</sup><sub>3</sub>]Ti–MnBr(CO)<sub>4</sub> (**9**) and the octahedral complex [PhBP<sup>Pr</sup><sub>3</sub>]Mn(CN<sup>t</sup>Bu)<sub>3</sub> (**10**). Some of our initial synthetic efforts to generate [PhBP<sup>Pr</sup><sub>3</sub>]Mn≡N<sub>x</sub> species are briefly described, as are DFT studies that probe the electronic viability of these types of multiply bonded target structures.

## I. Introduction

Transition metal complexes containing metal–ligand multiple bonds figure prominently in coordination chemistry, especially in atom- and group-transfer processes.<sup>1,2</sup> Metal species of the types L<sub>n</sub>M=E and L<sub>n</sub>M≡E, where E = O, NR, CR<sub>2</sub>, can act as intermediates in numerous catalytic reactions including epoxidation,<sup>3–6</sup> aziridination,<sup>7–10</sup> cyclopropanation,<sup>11–13</sup> olefin metathesis,<sup>14–17</sup> and C–H bond inser-

tion.<sup>10,18</sup> Besides having broad synthetic utility, metal-to-ligand multiply bonded species also serve as biomimetic models of active sites in metalloenzymes.<sup>19–22</sup> Until a few years ago, examples of stable complexes featuring metal-to-ligand multiple bonds were rare for the later metals,<sup>23–25</sup> especially for those in the first-row.<sup>26</sup> While the paucity of these compounds was typically attributed to an inherent

\* To whom correspondence should be addressed. E-Mail: jpeters@caltech.edu.

- (1) Nugent, W. A.; Mayer, J. M. *Metal–Ligand Multiple Bonds*; Wiley: New York, 1988.
- (2) Holm, R. H. *Chem. Rev.* **1987**, *87*, 1401.
- (3) Jorgensen, K. A. *Chem. Rev.* **1989**, *89*, 431.
- (4) Gao, Y.; Hanson, R. M.; Klunder, J. M.; Ko, S. Y.; Masamune, H.; Sharpless, K. B. *J. Am. Chem. Soc.* **1987**, *109*, 5765.
- (5) Collman, J. P.; Zhang, X. M.; Lee, V. J.; Uffelman, E. S.; Brauman, J. I. *Science* **1993**, *261*, 1404.
- (6) Collman, J. P.; Brauman, J. I.; Meunier, B.; Raybuck, S. A.; Kodadek, T. *Proc. Natl. Acad. Sci. U.S.A.* **1984**, *81*, 3245.
- (7) Li, Z.; Quan, R. W.; Jacobsen, E. N. *J. Am. Chem. Soc.* **1995**, *117*, 5889.
- (8) Evans, D. A.; Woerpel, K. A.; Hinman, M. M.; Faul, M. M. *J. Am. Chem. Soc.* **1991**, *113*, 726.
- (9) Jacobsen, E. N. *Comprehensive Asymmetric Catalysis*; Springer: Hamburg, 1999.
- (10) Müller, P.; Fruit, C. *Chem. Rev.* **2003**, *103*, 2905.
- (11) Brookhart, M.; Studabaker, W. B. *Chem. Rev.* **1987**, *87*, 411.
- (12) Doyle, M. P. *Chem. Rev.* **1986**, *86*, 919.

- (13) Doyle, M. P.; McKervey, M. A.; Ye, T. *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds*; Wiley: New York, 1997.
- (14) Nguyen, S. T.; Grubbs, R. H.; Ziller, J. W. *J. Am. Chem. Soc.* **1993**, *115*, 9858.
- (15) Grubbs, R. H.; Chang, S. *Tetrahedron* **1998**, *54*, 4413.
- (16) Hoveyda, A. H.; Schrock, R. R. *Chem. Eur. J.* **2001**, *7*, 945.
- (17) Feldman, J.; Schrock, R. R. *Prog. Inorg. Chem.* **1991**, *39*, 1.
- (18) Davies, H. M. L.; Hansen, T.; Churchill, M. R. *J. Am. Chem. Soc.* **2000**, *122*, 3063.
- (19) Groves, J. T.; Han, Y.-Z. *Cytochrome P450: Structure, Mechanism, and Biochemistry*; Plenum Press: New York, 1995; p 3.
- (20) Abu-Omar, M. M.; Loaiza, A.; Hontzeas, N. *Chem. Rev.* **2005**, *105*, 2227.
- (21) Nakamoto, K. *Coord. Chem. Rev.* **2002**, *226*, 153.
- (22) Costas, M.; Mehn, M. P.; Jensen, M. P.; Que, L. *Chem. Rev.* **2004**, *104*, 939.
- (23) Glueck, D. S.; Wu, J. X.; Hollander, F. J.; Bergman, R. G. *J. Am. Chem. Soc.* **1991**, *113*, 2041.
- (24) Glueck, D. S.; Hollander, F. J.; Bergman, R. G. *J. Am. Chem. Soc.* **1989**, *111*, 2719.
- (25) Hay-Motherwell, R. S.; Wilkinson, G.; Hussain-Bates, B.; Hursthouse, M. B. *Polyhedron* **1993**, *12*, 2009.



incompatibility of high d-electron counts with multiply bonded  $\pi$ -donor substituents,<sup>1</sup> low-coordinate geometries have provided one strategic path to circumvent this limitation. The recent rise in literature reports of well-defined  $L_nM=E$  and  $L_nM\equiv E$  systems for the metals  $M = Fe$ ,<sup>27–33</sup>  $Co$ ,<sup>34–38</sup>  $Ni$ ,<sup>39–43</sup> and  $Cu$ ,<sup>44,45</sup> strongly attests to the viability of these types of species for the first-row, mid-to-late metals.

Manganese compounds featuring multiply bonded terminal functionalities such as nitrides and imides are far more abundant than their later first-row counterparts.<sup>46</sup> For example, Mn(V) nitrides are well known to be stabilized by various ligand auxiliaries, including porphyrins,<sup>47–51</sup> macrocyclic amines,<sup>52,53</sup> cyanides,<sup>54</sup> and Schiff bases.<sup>55–58</sup> In contrast, Mn(VI) and Mn(VII) nitrides are rare. Wieghardt and co-workers generated two examples of Mn(VI) nitrides in situ and assigned them from spectral data.<sup>52,54</sup> Hursthouse and co-workers isolated the sole example of a Mn(VII)

nitride,  $[Mn(N)(N'Bu)_3]^{2-}$ , which is supported by imido ligands.<sup>59</sup> Other relevant manganese imides from the Hursthouse group include species of the type  $Mn^{VII}(N'Bu)_3X$ , as well as the homoleptic compound  $[Mn^{VI}(N'Bu)_4]^{2-}$ .<sup>60–62</sup> Terminal Mn(V) imides are also stabilized by corroles, as recently reported by Abu-Omar and co-workers.<sup>63–65</sup> Finally, elegant work from Groves and Carreira strongly implicates Mn(V) acylimides (generated in situ from the nitrides) as effective group-transfer catalysts to olefin and enol ether substrates.<sup>47,55,66</sup>

Our group has been investigating the tris(phosphino)borate (generally denoted as  $[BP_3]$ ) and hybrid bis(phosphino)-pyrazolylborate ( $[BP_2(pz)]$ ) ligands as scaffolds for  $M\equiv E$  linkages. Our endeavors have led to the characterization of stable Co(III)<sup>36</sup> and Fe(II/III) imides,<sup>27,29</sup> as well as distinctive examples of Fe(IV) imides and nitrides.<sup>28,67</sup> Motivated by these results, we wish to extend this chemistry to other mid-to-late first-row metals such as Ni<sup>68</sup> and Mn. In particular, we are interested in whether  $[BP_3]Mn\equiv N_x$  species ( $d^3$  and  $d^4$ ) are electronically accessible and in defining synthetic methods for their generation. In pursuit of this task, we began to study aspects of the fundamental coordination chemistry of previously unexplored  $[BP_3]Mn$  systems. In this first study, we report on the synthesis and characterization of a family of monovalent- and divalent-manganese compounds supported by the  $[PhBP^{iPr}_3]$  ligand (where  $[PhBP^{iPr}_3] = [PhB(CH_2P^{iPr}_2)_3]^-$ ), some of which might ultimately serve as precursors for the preparation of  $[BP_3]Mn\equiv N_x$  species. Complementary theoretical studies are described for hypothetical imide  $[PhBP^{iPr}_3]Mn(N'Bu)$  and nitride  $[PhBP^{iPr}_3]Mn(N)$  systems. These species appear to be plausible target structures on the basis of electronic considerations. While studies toward well-defined  $[BP_3]Mn\equiv N_x$  systems are ongoing, the four-coordinate  $[PhBP^{iPr}_3]Mn(II)$  species that are described here represent structurally unusual examples of low-coordinate manganese complexes supported by polyphosphine ligands.


## II. Results and Discussion

**Synthesis and Characterization of  $[PhBP^{iPr}_3]Mn(II)$  Halides.** The Mn(II) halides  $[PhBP^{iPr}_3]MnCl$  (**1**) and  $[PhBP^{iPr}_3]MnI$  (**2**) were readily prepared by mixing the thallium reagent

- (26) Verma, A. K.; Nazif, T. N.; Achim, C.; Lee, S. C. *J. Am. Chem. Soc.* **2000**, *122*, 11013.
- (27) Brown, S. D.; Peters, J. C. *J. Am. Chem. Soc.* **2005**, *127*, 1913.
- (28) Betley, T. A.; Peters, J. C. *J. Am. Chem. Soc.* **2004**, *126*, 6252.
- (29) Brown, S. D.; Betley, T. A.; Peters, J. C. *J. Am. Chem. Soc.* **2003**, *125*, 322.
- (30) Klinker, E. J.; Kaizer, J.; Brennessel, W. W.; Woodrum, N. L.; Cramer, C. J.; Que, L. *Angew. Chem., Int. Ed.* **2005**, *44*, 3690.
- (31) Rohde, J. U.; In, J. H.; Lim, M. H.; Brennessel, W. W.; Bukowski, M. R.; Stubna, A.; Münck, E.; Nam, W.; Que, L. *Science* **2003**, *299*, 1037.
- (32) Kaizer, J.; Klinker, E. J.; Oh, N. Y.; Rohde, J. U.; Song, W. J.; Stubna, A.; Kim, J.; Münck, E.; Nam, W.; Que, L. *J. Am. Chem. Soc.* **2004**, *126*, 472.
- (33) Bart, S. C.; Lobkovsky, E.; Bill, E.; Chirik, P. J. *J. Am. Chem. Soc.* **2006**, *128*, 5302.
- (34) Shay, D. T.; Yap, G. P. A.; Zakharov, L. N.; Rheingold, A. L.; Theopold, K. H. *Angew. Chem., Int. Ed.* **2005**, *44*, 1508.
- (35) Dai, X. L.; Kapoor, P.; Warren, T. H. *J. Am. Chem. Soc.* **2004**, *126*, 4798.
- (36) Jenkins, D. M.; Betley, T. A.; Peters, J. C. *J. Am. Chem. Soc.* **2002**, *124*, 11238.
- (37) Termaten, A. T.; Aktas, H.; Schakel, M.; Ehlers, A. W.; Lutz, M.; Spek, A. L.; Lammertsma, K. *Organometallics* **2003**, *22*, 1827.
- (38) Sanchez-Nieves, J.; Sterenberg, B. T.; Udachin, K. A.; Carty, A. J. *J. Am. Chem. Soc.* **2003**, *125*, 2404.
- (39) Mindiola, D. J.; Hillhouse, G. L. *Chem. Commun.* **2002**, 1840.
- (40) Mindiola, D. J.; Hillhouse, G. L. *J. Am. Chem. Soc.* **2002**, *124*, 9976.
- (41) Melenkivitz, R.; Mindiola, D. J.; Hillhouse, G. L. *J. Am. Chem. Soc.* **2002**, *124*, 3846.
- (42) Mindiola, D. J.; Hillhouse, G. L. *J. Am. Chem. Soc.* **2001**, *123*, 4623.
- (43) Kogut, E.; Wiencko, H. L.; Zhang, L. B.; Cordeau, D. E.; Warren, T. H. *J. Am. Chem. Soc.* **2005**, *127*, 11248.
- (44) Dai, X. L.; Warren, T. H. *J. Am. Chem. Soc.* **2004**, *126*, 10085.
- (45) Straub, B. F.; Hofmann, P. *Angew. Chem., Int. Ed.* **2001**, *40*, 1288.
- (46) Eikey, R. A.; Abu-Omar, M. M. *Coord. Chem. Rev.* **2003**, *243*, 83.
- (47) Groves, J. T.; Takahashi, T. *J. Am. Chem. Soc.* **1983**, *105*, 2073.
- (48) Buchler, J. W.; Dreher, C.; Lay, K. L.; Lee, Y. J. A.; Scheidt, W. R. *Inorg. Chem.* **1983**, *22*, 888.
- (49) Hill, C. L.; Hollander, F. J. *J. Am. Chem. Soc.* **1982**, *104*, 7318.
- (50) Buchler, J. W.; Dreher, C.; Lay, K. L. *Z. Naturforsch., B: Chem. Sci.* **1982**, *37*, 1155.
- (51) Dehnicke, K.; Strahle, J. *Angew. Chem., Int. Ed. Engl.* **1981**, *20*, 413.
- (52) Meyer, K.; Bendix, J.; Metzler-Nolte, N.; Weyhermüller, T.; Wieghardt, K. *J. Am. Chem. Soc.* **1998**, *120*, 7260.
- (53) Niemann, A.; Bossek, U.; Haselhorst, G.; Wieghardt, K.; Nuber, B. *Inorg. Chem.* **1996**, *35*, 906.
- (54) Bendix, J.; Meyer, K.; Weyhermüller, T.; Bill, E.; Metzler-Nolte, N.; Wieghardt, K. *Inorg. Chem.* **1998**, *37*, 1767.
- (55) Du Bois, J.; Tomooka, C. S.; Hong, J.; Carreira, E. M. *Acc. Chem. Res.* **1997**, *30*, 364.
- (56) Du Bois, J.; Tomooka, C. S.; Hong, J.; Carreira, E. M.; Day, M. W. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 1645.
- (57) Chang, C. J.; Connick, W. B.; Low, D. W.; Day, M. W.; Gray, H. B. *Inorg. Chem.* **1998**, *37*, 3107.
- (58) Chang, C. J.; Low, D. W.; Gray, H. B. *Inorg. Chem.* **1997**, *36*, 270.


- (59) Danopoulos, A. A.; Wilkinson, G.; Sweet, T. K. N.; Hursthouse, M. B. *J. Chem. Soc., Dalton Trans.* **1995**, 205.
- (60) Danopoulos, A. A.; Green, J. C.; Hursthouse, M. B. *J. Organomet. Chem.* **1999**, *591*, 36.
- (61) Danopoulos, A. A.; Wilkinson, G.; Sweet, T. K. N.; Hursthouse, M. B. *J. Chem. Soc., Dalton Trans.* **1995**, 937.
- (62) Danopoulos, A. A.; Wilkinson, G.; Sweet, T. K. N.; Hursthouse, M. B. *J. Chem. Soc., Dalton Trans.* **1994**, 1037.
- (63) Abu-Omar, M. M.; Shields, C. E.; Edwards, N. Y.; Eikey, R. A. *Angew. Chem., Int. Ed.* **2005**, *44*, 6203.
- (64) Edwards, N. Y.; Eikey, R. A.; Loring, M. I.; Abu-Omar, M. M. *Inorg. Chem.* **2005**, *44*, 3700.
- (65) Eikey, R. A.; Khan, S. I.; Abu-Omar, M. M. *Angew. Chem., Int. Ed.* **2002**, *41*, 3592.
- (66) Du Bois, J.; Hong, J.; Carreira, E. M.; Day, M. W. *J. Am. Chem. Soc.* **1996**, *118*, 915.
- (67) Thomas, C. M.; Mankad, N. P.; Peters, J. C. *J. Am. Chem. Soc.* **2006**, *128*, 4956.
- (68) MacBeth, C. E.; Thomas, J. C.; Betley, T. A.; Peters, J. C. *Inorg. Chem.* **2004**, *43*, 4645.






**Figure 1.** Displacement ellipsoid representations (50%) of (A) [PhBP<sup>i</sup>Pr<sub>3</sub>]MnCl (**1a**) and (B) [PhBP<sup>i</sup>Pr<sub>3</sub>]MnI (**2**). Hydrogen atoms and solvent molecules have been omitted for clarity. Selected bond distances (Å) and angles (deg) for **1a**: Mn–Cl 2.2687(7), Mn–P1 2.5300(6), Cl–Mn–P1 125.38(2), Mn–P2 2.5298(6), Mn–P3 2.5289(6), Cl–Mn–P2 120.22(2), Cl–Mn–P3 126.14(2), P1–Mn–P2 90.55(2), P2–Mn–P3 92.64(2), P1–Mn–P3 92.22(2). For **2**: Mn–I 2.6394(4), Mn–P1 2.5351(7), Mn–P2 2.5238(7), Mn–P3 2.5379(6), P1–Mn–I 127.19(2), P2–Mn–I 122.27(2), P3–Mn–I 124.09(2), P1–Mn–P2 91.39(2), P2–Mn–P3 90.68(2), P1–Mn–P3 90.95(2).

#### Scheme 1



[PhBP<sup>i</sup>Pr<sub>3</sub>]Ti with MnX<sub>2</sub> (where X = Cl or I) (Scheme 1).<sup>69</sup> This methodology has been used previously by us to prepare related compounds of divalent Fe, Ru, Co, and Ni.<sup>68,70,71</sup> The halides **1** and **2** are colorless in solution and can be crystallized from benzene/petroleum ether. Compound **1** provided light pink blocks in low yield (33%), while **2** was obtained as colorless blocks in moderate yield (62%). Because of its higher crystalline yield, compound **2** was used as the preferred starting material throughout the research described herein.

The halides **1** and **2** were characterized by various techniques including X-ray crystallography, SQUID magnetometry, and EPR spectroscopy. The solid-state structures of **1** and **2** are shown in Figures 1 and 2. The chloride complex **1** crystallizes as a mixture of its monomer (**1a**) and its dimer {[PhBP<sup>i</sup>Pr<sub>3</sub>]Mn(μ-Cl)<sub>2</sub>} (**1b**), whereas the iodide complex **2** crystallizes exclusively as a monomer. A similar monomer–dimer equilibrium was observed for [PhBP<sub>3</sub>]CoX systems in solution (where X = Cl or Br).<sup>71</sup> Solid-state structures of monomeric **1a** and **2** are pseudotetrahedral with approximate 3-fold symmetry about the axis containing B, Mn, and X (I–Mn–B = 176.6°, Cl–Mn–B = 175.5°). The Mn–P bond distances are nearly equal, as well as the P–Mn–P bond angles. For example, monomer **1a** has Mn–P distances of 2.53 Å and P–Mn–P angles that are near 90° (90.6–92.6°). Likewise, compound **2** exhibits a narrow spread in its Mn–P bond distances (2.524–2.538 Å) and P–Mn–P angles (90.7–91.4°). The Mn–P bond distances are not unusual for high-spin Mn(II)<sup>72</sup> but are



**Figure 2.** Displacement ellipsoid representations (50%) of {[PhBP<sup>i</sup>Pr<sub>3</sub>]Mn(μ-Cl)<sub>2</sub>} (**1b**) (top). Hydrogen atoms have been omitted for clarity. Selected bond distances (Å) and angles (deg) for **1b**: Mn–ClA 2.4625(9), Mn–ClA 2.5327(9), Mn–ClB 2.4630(9), Mn–ClB 2.525(1), Mn–P1 2.6162(7), Mn–P2 2.6132(7), Mn–P3A 2.552(4), Mn–P3B 2.580(4), P1–Mn–P2 86.38(2), P3A–Mn–P1 87.13(7), P3B–Mn–P1 94.03(7), P3A–Mn–P2 92.55(8), P3B–Mn–P2 86.26(7). View of the core of **1b** showing both positions of the disordered chloride and P3 atom (below). The angle between the Mn<sub>2</sub>(ClA)<sub>2</sub> and Mn<sub>2</sub>(ClB)<sub>2</sub> planes is 41°.

significantly longer than M–P distances of tris(phosphino)-borate complexes of later first-row transition metal ions. For instance, the series of structurally characterized chloride complexes, [PhBP<sup>i</sup>Pr<sub>3</sub>]MCl for M = Mn, Fe, Co, and Ni, show decreasing M–P bond distances, which reflect both their decreasing ionic radii across the series as well as their total spin, S.

The dimer **1b** crystallizes in the triclinic crystal system *P*1̄ and features a Mn<sub>2</sub>Cl<sub>2</sub> core that contains an inversion center. The chloride and one P<sup>i</sup>Pr<sub>2</sub> unit are disordered, but both can be independently refined in two positions of nearly equal populations. The solid-state structure of one conformer is shown in Figure 2 (top). A simplified structure containing all of the disordered atoms in the core is shown below it. In the dimer, the Mn center is five-coordinate and roughly square pyramidal. Two phosphorus atoms and two chlorides comprise the base, with an axial phosphine at the apex of the pyramid. The Mn–P bond distances in the dimer are even more elongated relative to the monomer **1a** (Mn–P = 2.613, 2.616, and 2.552 Å). One of these bonds (Mn–P3) is slightly shorter than the other two and corresponds to the Mn–P<sub>ax</sub> bond. The Mn–Cl bonds in the dimer are also significantly longer (2.463 and 2.533 Å) compared to the monomer (2.269 Å).

Magnetic susceptibility (SQUID) data were obtained on polycrystalline samples of **1** and **2**. The μ<sub>eff</sub> versus *T* plots are nearly identical and show very little temperature dependence (Figure 3A). From 60 to 300 K, the μ<sub>eff</sub> values average 5.94 μ<sub>B</sub> for **1** (calculated as a monomer) and 5.97 μ<sub>B</sub> for **2**. The μ<sub>eff</sub> value for **1** is identical to its solution magnetic moment of 5.9(1) in benzene, where we presume the monomeric form is favored over the dimer form (Evans method, three runs).<sup>73,74</sup> These values are consistent with five unpaired electrons per Mn, verifying the high-spin configuration of these [PhBP<sup>i</sup>Pr<sub>3</sub>]MnX species. The proximity of

(69) Shapiro, I. R.; Jenkins, D. M.; Thomas, J. C.; Day, M. W.; Peters, J. C. *Chem. Commun.* **2001**, 2152.

(70) Betley, T. A.; Peters, J. C. *Inorg. Chem.* **2003**, 42, 5074.


(71) Jenkins, D. M.; Di Bilio, A. J.; Allen, M. J.; Betley, T. A.; Peters, J. C. *J. Am. Chem. Soc.* **2002**, 124, 15336.

(72) Hebenanz, N.; Köhler, F. H.; Müller, G. *Inorg. Chem.* **1984**, 23, 3043.

(73) Sur, S. K. *J. Magn. Reson.* **1989**, 82, 169.

(74) Evans, D. F. *J. Chem. Soc.* **1959**, 2003.






**Figure 3.** (A) SQUID plot of  $\mu_{\text{eff}}$  ( $\mu_B$ ) versus  $T$  and (B)  $\chi_M T$  ( $\text{cm}^3 \text{K/mol}$ ) versus  $T$  for  $[\text{PhBP}^{\text{iPr}}_3]\text{MnCl}$ , **1**, (■), and  $[\text{PhBP}^{\text{iPr}}_3]\text{MnI}$ , **2**, (□). An expansion of the  $\chi_M T$  vs  $T$  plot at low temperatures (4–30 K) is shown in the inset.

$\mu_{\text{eff}}$  to the spin-only value ( $5.92 \mu_B$ ) further suggests that the ground states of **1** and **2** are orbitally nondegenerate as expected for a ground-state electronic configuration where all the d orbitals are singly populated ( $^6S$ ). The  $\chi_M T$  versus  $T$  plots for **1** and **2** are also similar, but distinct at very low temperatures. From 4 to 75 K,  $\chi_M T$  dips slightly for **2**, perhaps due to intermolecular paramagnetic quenching (Figure 3B).<sup>75</sup> The  $\chi_M T$  versus  $T$  plot for **1**, however, shows a gradual increase in  $\chi_M T$  in the same temperature range. The small rise in  $\chi_M T$  from 4.43 to 4.71  $\text{cm}^3 \text{K/mol}$  might be indicative of weak ferromagnetic coupling. Similarly weak coupling has been observed in another  $\text{Mn}(\mu\text{-Cl})$  dimer and might thus arise from the presence of the dimer **1b**.<sup>76</sup>

EPR spectra were also recorded on the halides in methyl-THF glass at 8 K. The iodide **2** has an axial spectrum with transitions at  $\sim 120$  and 330 mT (Figure 4A). The observed  $g$  values,  $\sim 6$  and 2, match the calculated transitions for the  $m_s = \pm 1/2$  states of a  $S = 5/2$  paramagnet with  $E/D \approx 0$ . Hyperfine structure due to the  $^{55}\text{Mn}$  nucleus ( $I = 5/2$ ) is evident in the  $g = 2$  signal, which shows a six-line splitting with  $A \approx 63$  G. The octahedral complexes  $\text{MnX}_2(\text{dmpe})_2$  (where  $X = \text{Br}, \text{I}$ ) show nearly identical EPR signatures.<sup>77</sup> Unlike **2**, the chloride **1** has an EPR spectrum with numerous transitions occurring from 0 to 1000 mT (Figure 4B). The complicated spectrum is difficult to interpret and is tentatively



**Figure 4.** EPR spectra of glassy methyl-THF solutions of (A)  $[\text{PhBP}^{\text{iPr}}_3]\text{MnI}$  (**2**) and (B)  $[\text{PhBP}^{\text{iPr}}_3]\text{MnCl}$  (**1**) at 8 K. Instrumental parameters:  $\nu = 9.38$  GHz, modulation frequency = 100 kHz, modulation amplitude = 5 G, microwave power = 0.064 mW, conversion time = 81.9 ms, time constant = 20.5 ms, sweep time = 83.9 ms, 3 scans.

attributed to a superposition of the monomer ( $\sim 130$  and 360 mT) and dimer forms of **1**. The dimer, an  $S = 5$  paramagnet, could itself give rise to multiple transitions. For both **1** and **2**, hyperfine coupling to the  $^{31}\text{P}$  nucleus ( $I = 1/2$ ) is not observed, suggesting that the spin density is tightly localized on Mn and therefore indicative of attenuated covalent character at the Mn–P bond by comparison to other  $[\text{BP}_3]\text{M}$  paramagnets we have studied.<sup>27,71,78</sup>

Compounds **1** and **2** were also studied by NMR spectroscopy and electrochemistry, but these data proved relatively uninformative. The  $^1\text{H}$  NMR spectra contained few resolvable resonances, and those that could be resolved were unusually broad by comparison to related  $[\text{PhBP}^{\text{iPr}}_3]\text{FeX}$  and  $[\text{PhBP}^{\text{iPr}}_3]\text{CoX}$  systems. The cyclic voltammograms of **1** and **2** did not reveal any reversible features (0.35 M  $[\text{tBu}_4\text{N}][\text{PF}_6]$  in THF). The lack of any reversible redox couples also contrasts the analogous halide complexes of Fe, Co, and Ni, all of which exhibit at least pseudoreversible  $\text{M}^{\text{II}}/\text{M}^{\text{I}}$  redox couples.

Surprisingly, a survey of the CCDC revealed only three structures of monomeric four-coordinate Mn halide complexes with one or two phosphine donor(s).<sup>72,79,80</sup> No such structures with three phosphine donors were found. Even

(75) Kahn, O. *Molecular Magnetism*; VCH: New York, 1993.

(76) Armentano, D.; de Munno, G.; Guerra, F.; Faus, J.; Lloret, F.; Julve, M. *Dalton Trans.* **2003**, 4626.


(77) Girolami, G. S.; Wilkinson, G.; Galas, A. M. R.; Thornton-Pett, M.; Hursthouse, M. B. *J. Chem. Soc., Dalton Trans.* **1985**, 1339.

(78) Brown, S. D.; Mehn, M. P.; Peters, J. C. *J. Am. Chem. Soc.* **2005**, *127*, 13146.

(79) Godfrey, S. M.; McAuliffe, C. A.; Ndifon, P. T.; Pritchard, R. G. *J. Chem. Soc., Dalton Trans.* **1993**, 3373.


(80) Godfrey, S. M.; McAuliffe, C. A.; Pritchard, R. G. *J. Chem. Soc., Dalton Trans.* **1993**, 371.





**Figure 5.** Displacement ellipsoid representations (50%) of [PhBP<sup>i</sup>Pr<sub>3</sub>]Mn(N<sub>3</sub>) (**3**) and [PhBP<sup>i</sup>Pr<sub>3</sub>]Mn(CH<sub>2</sub>Ph) (**4**). For clarity, hydrogen atoms have been omitted and only the phosphorus atoms of the ligand are shown. For **3**, only one of the two molecules in the asymmetric unit is shown. Selected bond distances (Å) and angles (deg) for **3**: Mn–N1 2.046(3), Mn–P1 2.534(1), Mn–P2 2.522(1), Mn–P3 2.521(2), N1–N2 1.112(5), N2–N3 1.202(6), N2–N1–Mn 128.8(3), P1–Mn–P2 92.33(4), P2–Mn–P3 93.21(5), P1–Mn–P3 91.04(5), N1–Mn–P1 123.1(1), N1–Mn–P2 125.7(1), N1–Mn–P3 122.2(1). For **4**: Mn–C28 2.138(2), Mn–P1 2.5679(4), Mn–P2 2.5507(5), Mn–P3 2.5620(5), P1–Mn–P2 91.46(1), P2–Mn–P3 89.44(1), P1–Mn–P3 89.90(1), C28–Mn–P1 117.86(5), C28–Mn–P2 126.42(5), C28–Mn–P3 130.59(5), Mn–C28–C29 111.5(1).


#### Scheme 2



monomeric five-coordinate polyphosphine-ligated Mn halides are few in number, with only three structurally characterized examples.<sup>81–83</sup> Low-coordinate Mn halides are also known for the related tridentate ligand tris(pyrazoyl)borate (Tp). A handful of monomeric four-coordinate manganese structures featuring these ligands have been reported.<sup>84–88</sup>

**Synthesis and Characterization of [PhBP<sup>i</sup>Pr<sub>3</sub>]Mn(II) Azide, Alkyls, and Amides.** A series of divalent-manganese complexes featuring other X-type ligands were readily prepared from the iodide **2** via metathesis (Scheme 2). For example, addition of excess sodium azide to **2** produced off-white [PhBP<sup>i</sup>Pr<sub>3</sub>]Mn(N<sub>3</sub>) (**3**) in good yield (73%). Compound **3** features a terminal azide ligand with a characteristic ν-(N<sub>3</sub>) stretch at 2070 cm<sup>−1</sup> (KBr, THF). Its solid-state structure is shown in Figure 5. Not surprisingly, the pseudohalide **3** exhibits a similar molecular geometry to the monomeric

#### Scheme 3



halides **1** and **2**. In a CCDC search of all structurally characterized Mn complexes containing a terminal azide, the mean Mn–N bond distance is 2.190 Å with a standard deviation of 0.096 Å. The Mn–N bond distance of 2.046(3) Å in **3** is therefore unusually short. Manganese complexes featuring similarly short Mn–N bond distances for terminal azides are typically in the oxidation state +3.<sup>53,89–92</sup>


Manganese alkyl compounds were also readily prepared from the addition of alkyl Grignards PhCH<sub>2</sub>MgBr and MeMgBr to **2** at −90° in THF. The bright yellow complex [PhBP<sup>i</sup>Pr<sub>3</sub>]Mn(CH<sub>2</sub>Ph) (**4**) and pale yellow complex [PhBP<sup>i</sup>Pr<sub>3</sub>]Mn(Me) (**5**) are soluble in hydrocarbons and can be crystallized from cyclopentane at −30 °C. The benzyl complex **4** was characterized by X-ray diffraction, and its solid-state structure is shown in Figure 5. Related examples of low-coordinate monomeric organomanganese compounds are known and include trigonal (β-diketiminato)MnR and tetrahedral (P<sub>2</sub>)MnR<sub>2</sub> species, where P<sub>2</sub> represents bidentate phosphine or two PR<sub>3</sub> ligands.<sup>93–95</sup> Despite the low valence electron count of these Mn systems, rather little has been reported concerning their reactivity patterns. One interesting exception concerns tetrahedral Mn(II) dialkyl compounds supported by diimine ligands that readily undergo alkyl migration to the diimine ligand at ambient temperatures.<sup>96</sup> We have found that the [PhBP<sup>i</sup>Pr<sub>3</sub>]-supported manganese alkyls **4** and **5** are thermally stable up to 65 °C even under an atmosphere of dihydrogen or CO. This sharply contrasts the related iron(II) alkyls [PhBP<sup>i</sup>Pr<sub>3</sub>]Fe–R, which lose alkane upon exposure to H<sub>2</sub> and bind CO readily.<sup>97</sup>

Manganese amides were generated by the addition of lithium and potassium amides to **2** at −90° in THF (Scheme 3). In the metathesis reactions with lithium amides, a

- (81) Cecconi, F.; Ghilardi, C. A.; Midollini, S.; Orlandini, A. *J. Chem. Soc., Dalton Trans.* **1992**, 33.
- (82) Beagley, B.; McAuliffe, C. A.; Minten, K.; Pritchard, R. G. *J. Chem. Soc., Dalton Trans.* **1987**, 1999.
- (83) Beagley, B.; McAuliffe, C. A.; Minten, K.; Pritchard, R. G. *J. Chem. Soc., Chem. Commun.* **1984**, 658.
- (84) Komatsuzaki, H.; Sakamoto, N.; Satoh, M.; Hikichi, S.; Akita, M.; Moro-oka, Y. *J. Am. Chem. Soc.* **1915**, 37, 6554.
- (85) Matsunaga, Y.; Fujisawa, K.; Ibi, N.; Miyashita, Y.; Okamoto, K. *Inorg. Chem.* **2005**, 44, 325.
- (86) Nabika, M.; Seki, Y.; Miyatake, T.; Ishikawa, Y.; Okamoto, K.; Fujisawa, K. *Organometallics* **2004**, 23, 4335.
- (87) Bruner, T. J.; Hascall, T.; Cowley, A. R.; Rees, L. H.; O'Hare, D. *Inorg. Chem.* **2001**, 40, 3170.
- (88) Komatsuzaki, H.; Nagasu, Y.; Suzuki, K.; Shibasaki, T.; Satoh, M.; Ebina, F.; Hikichi, S.; Akita, M.; Moro-oka, Y. *J. Chem. Soc., Dalton Trans.* **1998**, 511.

- (89) Wieghardt, K.; Bossek, U.; Nuber, B.; Weiss, J. *Inorg. Chim. Acta* **1987**, 126, 39.
- (90) Hubin, T. J.; McCormick, J. M.; Alcock, N. W.; Busch, D. H. *Inorg. Chem.* **2001**, 40, 435.
- (91) Mantel, C.; Hassan, A. K.; Pecaut, J.; Deronzier, A.; Collomb, M. N.; Duboc-Toia, C. *J. Am. Chem. Soc.* **2003**, 125, 12337.
- (92) Saha, S.; Mal, D.; Koner, S.; Bhattacharjee, A.; Gutlich, P.; Mondal, S.; Mukherjee, M.; Okamoto, K. I. *Polyhedron* **2004**, 23, 1811.
- (93) Chai, J. F.; Zhu, H. P.; Fan, H. J.; Roesky, H. W.; Magull, J. *Organometallics* **2004**, 23, 1177.
- (94) Howard, C. G.; Girolami, G. S.; Wilkinson, G.; Thornton-Pett, M.; Hursthouse, M. B. *J. Chem. Soc., Dalton Trans.* **1983**, 2631.
- (95) Chai, J. F.; Zhu, H. P.; Roesky, H. W.; He, C.; Schmidt, H. G.; Noltemeyer, M. *Organometallics* **2004**, 23, 3284.
- (96) Rioulet, V.; Copéret, C.; Basset, J. M.; Rousset, L.; Bouchu, D.; Grosvalet, L.; Perrin, M. *Angew. Chem., Int. Ed.* **2002**, 41, 3025.
- (97) Daida, E. J.; Peters, J. C. *Inorg. Chem.* **2004**, 43, 7474.






**Figure 6.** Displacement ellipsoid representations (50%) of  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}(\text{NH}(2,6\text{-}^{\text{iPr}}_2\text{-C}_6\text{H}_3))$  (**6**),  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}(\text{dbabh})$  (**7**), and  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}(1\text{-Ph(isoindolate)})$  (**8**). For clarity, hydrogen atoms (except N-H) have been omitted and only the phosphorus atoms of the ligand are shown. Selected bond distances (Å) and angles (deg) for **6**: Mn–N 1.997(2), Mn–P1 2.5877(6), Mn–P2 2.5813(6), Mn–P3 2.5637(6), P1–Mn–P2 87.96(2), P2–Mn–P3 89.95(2), P1–Mn–P3 93.11(2), N–Mn–P1 144.77(5), N–Mn–P2 105.73(5), N–Mn–P3 118.63(5). For **7**: Mn–N 1.948(1), Mn–P1 2.5357(5), Mn–P2 2.5473(5), Mn–P3 2.5503(5), P1–Mn–P2 90.46(2), P2–Mn–P3 89.51(2), P1–Mn–P3 90.71(2), N–Mn–P1 126.19(4), N–Mn–P2 126.13(4), N–Mn–P3 122.92(4). For **8**: Mn–N 2.054(2), Mn–P1 2.5308(7), Mn–P2 2.5550(7), Mn–P3 2.5718(6), P1–Mn–P2 95.13(2), P1–Mn–P3 89.54(2), P2–Mn–P3 90.09(2), N–Mn–P1 111.98(5), N–Mn–P2 136.31(5), N–Mn–P3 122.30(5).

common side-product was the free ligand  $[\text{PhBP}^{\text{iPr}}_3][\text{Li}]$ , easily identified by  $^{31}\text{P}$  NMR spectroscopy. Replacement of lithium by potassium minimizes the liberation of the  $[\text{PhBP}^{\text{iPr}}_3]$  anion. For example, 2,6-diisopropylaniline can be deprotonated with KH and added to **2** to produce yellow  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}(\text{NH}(2,6\text{-}^{\text{iPr}}_2\text{-C}_6\text{H}_3))$  (**6**) in high yield (90%).

This strategy was also employed to install 2,3:5,6-dibenzo-7-azabicyclo[2.2.1]hepta-2,5-diene (Hdbabh) as the amide functionality. The lithium amide  $\text{Li}(\text{dbabh})$  is a particularly interesting reagent to investigate because it has been successfully exploited as an N-atom transfer reagent to both Cr and Fe centers with concomitant loss of anthracene.<sup>28,98</sup> Methathesis of **2** and  $\text{Li}(\text{dbabh})$  resulted in a mixture of the amide compound  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}(\text{dbabh})$  (**7**), the bridged amide complex  $\{(\text{Hdbabh})\text{MnI}(\mu\text{-N-dbabh})\}_2$  (see Supporting Information), and the ligand  $[\text{PhBP}^{\text{iPr}}_3][\text{Li}]$ . Though the yield of the desired compound **7** was quite low, yellow crystals of **7** could be cleanly chosen from the colorless crystals of the byproducts. Attempts to improve the yield of **7** by using  $\text{K}(\text{dbabh})$  were unsuccessful. Deprotonation of Hdbabh with KH or benzyl potassium afforded a bright yellow-green solution; upon addition of **2** at  $-90^\circ\text{C}$ , the reaction solution turned orange. Red-orange crystals were grown from this solution that were clearly indicative of a species different from **7**. An X-ray diffraction study revealed the product to be a different Mn amide,  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}(1\text{-Ph(isoindolate)})$  (**8**), an isomer of **7** wherein one of the five-membered rings has opened by an overall C–C bond cleavage. This decomposition most likely occurs during the deprotonation stage of the reaction since the solution turns brilliant yellow-green upon addition of the potassium base, whereas solutions of  $\text{Li}(\text{dbabh})$  are pale pink.

The solid-state structures of the Mn(II) amides **6**, **7**, and **8** are shown in Figure 6. Collectively, these amides are similar to the other pseudotetrahedral, high-spin  $[\text{PhBP}^{\text{iPr}}_3]\text{-Mn(II)}$  species reported here in that they have typical Mn–P bond distances (ca. 2.55 Å) and P–Mn–P bond angles (ca.



**Figure 7.** Cyclic voltammogram of  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}(\text{dbabh})$  (**7**) recorded at a scan rate of 50 mV/s in THF with 0.35 M  $[\text{tBu}_4\text{N}][\text{PF}_6]$  as the supporting electrolyte. Potentials are internally referenced to  $\text{Fc}/\text{Fc}^+$ .

$90^\circ$ ). The apical N atom of **7** lies on the Mn–B axis, whereas compounds **6** and **8** exhibit a slightly canted N atom with N–Mn–B angles of  $156^\circ$  and  $167^\circ$ , respectively. The Mn–N bond distances follow the order: **7** (1.948(1) Å) < **6** (1.997(2) Å) < **8** (2.054(2) Å). This trend likely reflects the relative steric repulsion between the amide groups and the isopropyl groups of the  $[\text{PhBP}^{\text{iPr}}_3]$  ligand.

The Mn(II) alkyls and amides were studied by cyclic voltammetry in THF with 0.35 M  $[\text{tBu}_4\text{N}][\text{PF}_6]$  as the supporting electrolyte and  $\text{Ag}/\text{AgNO}_3$  as the reference electrode. Interestingly, amide **7** is distinct among this series of complexes in that it exhibits a reversible one-electron redox couple centered at  $-0.26$  V (vs  $\text{Fc}/\text{Fc}^+$ , scan rate = 50 mV/s), indicative of a  $\text{Mn}^{\text{II}}/\text{Mn}^{\text{III}}$  redox event (Figure 7). The amides **6** and **8** exhibit irreversible oxidative events near this same potential.


**Synthesis and Characterization of  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn(I)}$  Compounds.** Monovalent-manganese compounds may prove to be valuable precursors to  $\text{Mn}\equiv\text{NR}$  multiply bonded species in future studies. Indeed, oxidative nitrene transfer to Fe(I) and Co(I) precursors has proven highly effective in the synthesis of related  $\text{Fe}\equiv\text{NR}$ ,  $\text{Co}\equiv\text{NR}$ , and  $\text{Co}=\text{NR}$  species.<sup>29,34–36,99,100</sup> While monovalent-manganese complexes are numerous, they typically are octahedral 18-electron species stabilized by carbonyl ligands.<sup>101–105</sup> Low-coordinate and unsaturated Mn(I) compounds, on the other hand, are uncommon. Notable examples of this latter class include a

(98) Mindiola, D. J.; Cummins, C. C. *Angew. Chem., Int. Ed.* **1998**, *37*, 945.

(99) Betley, T. A.; Peters, J. C. *J. Am. Chem. Soc.* **2003**, *125*, 10782.

(100) Hu, X.; Meyer, K. *J. Am. Chem. Soc.* **2004**, *126*, 16322.






**Figure 8.** Displacement ellipsoid representations (50%) of [PhBP<sup>i</sup>Pr<sub>3</sub>]-Ti-MnBr(CO)<sub>4</sub> (**9**). For clarity, hydrogen atoms have been omitted. Selected bond distances (Å) and angles (deg) for **9**: Mn–Ti 2.6437(4), Mn–C28 1.913(3), Mn–C29 1.799(3), Mn–C30 1.851(3), Mn–C31 1.823(3), Mn–Br 2.5299(4), C31–Mn–Ti 172.74(8), Br–Mn–Ti 86.11(1), C29–Mn–C28 92.6(1), C29–Mn–C30 91.1(1), C28–Mn–Br 85.39(7), C29–Mn–Br 177.98(8), C30–Mn–Br 90.94(8), C31–Mn–Br 86.73(8).


three-coordinate Mn(I) dimer supported by bulky  $\beta$ -diketiminate ligands and several five-coordinate Mn compounds of the type (P $\wedge$ P)<sub>2</sub>Mn(CO)<sup>+</sup> (where P $\wedge$ P is a bidentate phosphine).<sup>106–109</sup>

To pursue a low-coordinate Mn(I) precursor, [PhBP<sup>i</sup>Pr<sub>3</sub>]-Ti and MnBr(CO)<sub>5</sub> were heated in benzene at 50 °C for several hours. The resulting yellow solution contained some TiBr precipitate, but to our surprise, the major product was a Ti–Mn adduct, [PhBP<sup>i</sup>Pr<sub>3</sub>]-Ti–MnBr(CO)<sub>4</sub> (**9**), which was isolated in 39% crystalline yield. Compound **9** is diamagnetic and exhibits a doublet in the <sup>31</sup>P NMR spectrum (45.4 ppm, *J*<sub>Ti–P</sub> = 8060 Hz) that is shifted relative to [PhBP<sup>i</sup>Pr<sub>3</sub>]-Ti (24.3 ppm, *J*<sub>Ti–P</sub> = 5913 Hz). The solid-state structure of **9** is shown in Figure 8. There are several crystal structures featuring a Ti–Mn bond, and the Ti–Mn bond of 2.6437–(4) Å is similar to those reported previously.<sup>110,111</sup>

Prolonged heating of a benzene solution of **9** resulted in its decomposition. By <sup>31</sup>P NMR spectroscopy, one major product forms upon thermolysis, characterized by a broad peak at ~50 ppm. An IR spectrum revealed two sharp stretches at 1999 and 1906 cm<sup>–1</sup> (KBr/THF). We propose this species to be the Mn(I) tricarbonyl compound, [PhBP<sup>i</sup>Pr<sub>3</sub>]-Mn(CO)<sub>3</sub> (Scheme 4). Indeed, the isoelectronic compound,




**Figure 9.** Displacement ellipsoid representations (50%) of [PhBP<sup>i</sup>Pr<sub>3</sub>]-Mn-(CN<sup>t</sup>Bu)<sub>3</sub> (**10**) from the side (A) and top (B) views. For clarity, hydrogen atoms have been omitted and only the phosphorus atoms of the ligand are shown. Selected bond distances (Å) and angles (deg) for **10**: Mn–C28 1.870(1), Mn–C33 1.864(1), Mn–C38 1.835(1), N1–C28 1.176(2), N2–C33 1.176(2), N3–C38 1.185(2), Mn–P1 2.3597(4), Mn–P2 2.3831(4), Mn–P3 2.3655(3), P1–Mn–P2 88.59(1), P2–Mn–P3 86.49(1), P1–Mn–P3 89.17(1), C33–Mn–C28 83.75(5), C38–Mn–C28 85.26(5), C38–Mn–C33 89.12(5).



**Figure 10.** Cyclic voltammogram of [PhBP<sup>i</sup>Pr<sub>3</sub>]-Mn(CN<sup>t</sup>Bu)<sub>3</sub> (**10**) recorded at a scan rate of 100 mV/s in THF with 0.35 M [nBu<sub>4</sub>][PF<sub>6</sub>] as the supporting electrolyte. Potentials are internally referenced to Fc/Fc<sup>+</sup>.

#### Scheme 4




[PhBP<sup>i</sup>Pr<sub>3</sub>]-Mn(CN<sup>t</sup>Bu)<sub>3</sub> (**10**), can be prepared by the reduction of [PhBP<sup>i</sup>Pr<sub>3</sub>]-MnI with sodium naphthalenide followed by addition of excess CN<sup>t</sup>Bu. Compound **10** is isolated in pure form but in low yield (22%) due in part to the competitive formation of [PhBP<sup>i</sup>Pr<sub>3</sub>]-Na.<sup>112</sup> Similarly, reduction of [PhBP<sup>i</sup>Pr<sub>3</sub>]-MnI with sodium naphthalenide under CO produces the presumed degradation product of **9**, [PhBP<sup>i</sup>Pr<sub>3</sub>]-Mn(CO)<sub>3</sub>. Unfortunately, this species has not been isolated in pure form. The solid-state structure of **10** confirms its assignment as a monovalent-Mn complex with three isocyanides (Figure 9). Relative to the high-spin Mn(II) species, the Mn–P bonds (2.36–2.38 Å) are more than 0.15 Å shorter. The P–Mn–P bond angles are also slightly contracted (86.5–89.2°). Compound **10** is also characterized by a reversible redox process at –0.66 V in its cyclic voltammogram, shown in Figure 10 (vs Fc/Fc<sup>+</sup>, scan rate = 100 mV/s).

(112) Attempts to minimize formation of [PhBP<sup>i</sup>Pr<sub>3</sub>]-Na by adding CN<sup>t</sup>Bu to **2** prior to the addition of sodium naphthalenide or by adding sodium naphthalenide to **2** at low temperatures gave rise to more complex reaction mixtures.

- (101) Baker, R. J.; Edwards, P. G.; Gracia-Mora, J.; Ingold, F.; Malik, K. M. A. *J. Chem. Soc., Dalton Trans.* **2002**, 3985.
- (102) Bond, A. M.; Colton, R.; van den Bergen, A.; Walter, J. N. *Inorg. Chem.* **2000**, 39, 4696.
- (103) Connolly, J.; Genge, A. R. J.; Levason, W.; Orchard, S. D.; Pope, S. J. A.; Reid, G. J. *J. Chem. Soc., Dalton Trans.* **1999**, 2343.
- (104) Connor, J. A.; Hudson, G. A. *J. Organomet. Chem.* **1974**, 73, 351.
- (105) Reimann, R. H.; Singleton, E. *J. Chem. Soc., Dalton Trans.* **1973**, 841.
- (106) Sorace, L.; Golze, C.; Gatteschi, D.; Bencini, A.; Roesky, H. W.; Chai, J.; Stückl, A. C. *Inorg. Chem.* **2006**, 45, 395.
- (107) Chai, J. F.; Zhu, H. P.; Stückl, A. C.; Roesky, H. W.; Magull, J.; Bencini, A.; Caneschi, A.; Gatteschi, D. *J. Am. Chem. Soc.* **2005**, 127, 9201.
- (108) King, W. A.; Scott, B. L.; Eckert, J.; Kubas, G. J. *Inorg. Chem.* **1999**, 38, 1069.
- (109) King, W. A.; Luo, X. L.; Scott, B. L.; Kubas, G. J.; Zilm, K. W. *J. Am. Chem. Soc.* **1996**, 118, 6782.
- (110) Schollenberger, M.; Nuber, B.; Ziegler, M. L.; Hey-Hawkins, E. *J. Organomet. Chem.* **1993**, 460, 55.
- (111) Guillard, R.; Zrineh, A.; Ferhat, M.; Tabard, A.; Mitaine, P.; Swistak, C.; Richard, P.; Lecomte, C.; Kadish, K. M. *Inorg. Chem.* **1988**, 27, 697.






**Figure 11.** DFT-optimized structures (Jaguar 5.0, B3LYP/LACVP\*\*) for (A) singlet  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}(\text{N}'\text{Bu})$  and (B) triplet  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}(\text{N}'\text{Bu})$ . Selected bond distances (Å) and angles (deg) for (A): Mn–N 1.63; Mn–P 2.28, 2.36, 2.36; Mn–N–C 144.8; P–Mn–P 96.8, 97.2, 97.7; N–Mn–P 104.7, 126.2, 127.2. For (B): Mn–N 1.66; Mn–P 2.40, 2.44, 2.45; Mn–N–C 179.2; P–Mn–P 89.5, 89.8, 91.5; N–Mn–P 121.9, 126.2, 127.0.

**DFT Models of  $\text{Mn}=\text{N}'\text{Bu}$  and  $\text{Mn}\equiv\text{N}$  Species.** DFT studies were undertaken to obtain some insight on the electronic viability of two species,  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}^{\text{III}}=\text{N}'\text{Bu}$  and  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}^{\text{IV}}\equiv\text{N}$ . We have previously used DFT methods to probe the electronic structures of  $[\text{BP}_3]\text{M}$  species with terminally bound  $\text{N}_x$  functionalities that are strongly  $\pi$ -donating, for example  $[\text{PhBP}_3]\text{Co}^{\text{III}}=\text{N}(p\text{-tolyl})$ ,  $[\text{PhBP}_3]\text{Fe}^{\text{III}}=\text{N}'\text{Bu}$ , and  $[\text{PhBP}^{\text{iPr}}_3]\text{Fe}^{\text{IV}}\equiv\text{N}$ .<sup>29,36,28</sup>

For the low-spin Fe and Co imides, a general d-orbital splitting diagram has been forwarded based upon these DFT studies. The d orbitals, which transform as  $a_1 + 2e$  under idealized 3-fold symmetry, split into two high-lying and three lower-lying orbitals. The higher-energy e set, which in simplest terms can be thought of as comprising  $d_{xz}$  and  $d_{yz}$ , is destabilized by  $\sigma^*$  and  $\pi^*$  interactions with the phosphines and imide functionality, respectively. The remaining orbitals, of predominant  $d_{z^2}$  ( $a_1$ ) and  $d_{xy}$  and  $d_{x^2-y^2}$  (e) character, lie at lower energy and are predominantly nonbonding in nature.<sup>27</sup>

If we apply this qualitative d-orbital picture to the hypothetical  $d^4$  imide,  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}=\text{N}'\text{Bu}$ , the two ground states low-spin ( $S = 0$ ) and intermediate-spin ( $S = 1$ ) need to be considered. Recently, we have reported the characterization of an unusual  $\text{Fe}^{\text{IV}}$  imide that exhibits a triplet ground state.<sup>67</sup> Because this species is isoelectronic to the hypothetical Mn(III) imide under consideration here, we might expect the latter to be a triplet as well. To test this from a theoretical standpoint, the imide  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}=\text{N}'\text{Bu}$  was optimized as a singlet and a triplet using the Jaguar computational program (B3LYP/LACVP\*\*). The input geometry was derived from the crystal structure coordinates of  $[\text{PhBP}^{\text{iPr}}_3]\text{Fe}(\text{NAd})$ , where the metal and  $\text{N}_x$  functionality were changed to Mn and N'Bu, respectively. A geometry optimization was then performed using a spin-unrestricted calculation and conducted without any symmetry constraints (see Supporting Information).

The predicted structures for the singlet and triplet states of  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}=\text{N}'\text{Bu}$  differ quite remarkably from one another (Figure 11). The optimized singlet structure has a bent Mn–N–C angle ( $145^\circ$ ) with expanded P–Mn–P angles ( $97\text{--}98^\circ$ ), whereas the optimized triplet structure features a linear Mn–N–C angle ( $179^\circ$ ) with characteristic P–Mn–P angles ( $90\text{--}92^\circ$ ). The triplet state exhibits what appears to be a more likely geometry, in accord with the



**Figure 12.** DFT-optimized structures (Jaguar 5.0, B3LYP/LACVP\*\*) for (A) doublet  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}(\text{N})$  and (B) quartet  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}(\text{N})$ . Selected bond distances (Å) and angles (deg) for (A): Mn–N 1.50; Mn–P 2.35, 2.41, 2.45; N–Mn–P 112.1, 122.3, 127.5; P–Mn–P 94.8, 95.6, 97.2. For (B): Mn–N 1.62; Mn–P 2.48, 2.49, 2.60; N–Mn–P 103.9, 109.3, 155.3; P–Mn–P 89.9, 90.5, 91.2.

fact that it is predicted by DFT to be more energetically stable by 0.74 eV than the singlet state.

On the basis of DFT studies, the known nitride  $[\text{PhBP}^{\text{iPr}}_3]\text{Fe}^{\text{IV}}\equiv\text{N}$  has a splitting diagram that differs from the structurally analogous imides in that the  $a_1$  orbital of  $d_{z^2}$  parentage is more appreciably destabilized by  $\sigma$  antibonding interactions with the nitride functionality and the phosphine donors.<sup>28</sup> This situation leaves only two orbitals at very low energy, giving rise to the singlet ground state of  $[\text{PhBP}^{\text{iPr}}_3]\text{Fe}^{\text{IV}}\equiv\text{N}$ . On the basis of this splitting scheme, the hypothetical Mn nitride  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}\equiv\text{N}$  might be expected to populate a doublet ground state. To ascertain whether this would be the case, the Mn nitride  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}(\text{N})$  was modeled as both an  $S = 1/2$  and  $3/2$  system. In the optimized quartet structure, the nitride atom is severely tilted off the Mn–B axis by  $30^\circ$  (Figure 12B) in what appears to be an unfavorable geometry. The P–Mn–P bond angles ( $90\text{--}91^\circ$ ) and Mn–P bond distances (2.48–2.60 Å) are, however, typical of the pseudotetrahedral  $[\text{PhBP}^{\text{iPr}}_3]\text{Mn}(\text{II})$  complexes described here. The Mn–N bond length of 1.62 Å seems excessively long for known  $\text{Mn}\equiv\text{N}$  species, which typically have Mn–N bond lengths in the range of 1.50–1.55 Å. These distortions in the optimized quartet structure presumably arise from population of the antibonding  $d_{z^2}$  orbital.

Relative to the quartet state, the doublet state is energetically preferred by 0.41 eV and exhibits what appears to be a more reasonable geometry (Figure 12A). The nitride atom is canted off the Mn–B axis by only  $9^\circ$  and binds tightly to Mn with a Mn–N bond length of 1.50 Å. The P–Mn–P bond angles are slightly expanded ( $95\text{--}97^\circ$ ), approaching a more tetrahedral geometry. Similar parameters were found in the DFT-optimized structure of  $[\text{PhBP}^{\text{iPr}}_3]\text{Fe}^{\text{IV}}\equiv\text{N}$ , which features expanded P–Fe–P bond angles ( $99\text{--}101^\circ$ ) and an Fe–N bond length of 1.49 Å. It is the combination of these two structural parameters that renders the  $a_1$  ( $d_{z^2}$ ) orbital more antibonding for the metal nitrides than in the case of the related metal imides.

### III. Concluding Remarks

With this report, we have extended our study of the coordination chemistry of first-row mid-to-late transition metals supported by electron-releasing  $[\text{BP}_3]$  ligands. As with Fe, Co, and Ni, the  $[\text{PhBP}^{\text{iPr}}_3]$  scaffold readily supports Mn–



(II) complexes in a pseudotetrahedral geometry. The preparative significance of the systems described herein is best underscored by noting how uncommon low-coordinate manganese complexes supported by polyphosphines are. The high donor strength and anionic charge of the [PhBP<sup>i</sup>Pr<sub>3</sub>] ligand may be advantageous over other polyphosphine ligands in general by helping to attenuate problematic phosphine dissociation. These two ligand properties are not always sufficient, however, as evident from the observed propensity to substitute manganese by an alkali metal cation or by thallium in the [PhBP<sup>i</sup>Pr<sub>3</sub>] binding pocket under certain metathetical exchange reactions. Fortunately, the halide [PhBP<sup>i</sup>Pr<sub>3</sub>]MnI does serve as a sufficiently useful reagent for most metathetical reactions, allowing the preparation of [PhBP<sup>i</sup>Pr<sub>3</sub>]Mn(II) azide, alkyl, and amide species.

DFT calculations of the hypothetical Mn(III) imide [PhBP<sup>i</sup>Pr<sub>3</sub>]Mn(N<sup>i</sup>Bu) and Mn(IV) nitride [PhBP<sup>i</sup>Pr<sub>3</sub>]Mn(N) structures suggest that their preferred ground states will be triplet and doublet, respectively. While the preparation of Mn≡N<sub>x</sub> species of these types has, at this stage, only been cursorily canvassed, it is worth noting a few experimental observations. For instance, we have explored whether the Mn(I) complex **10** might undergo facile oxidative group transfer in the presence of organic aryl and alkyl azides. We have also canvassed conditions that might have effected the oxidative deprotonation of the anilide species **6**.<sup>42</sup> Finally, attempts to generate a Mn(IV) nitride species, including thermolysis and photolysis of the amide complex **7**, as well as photolysis of the Mn azide **3**, have been explored. As yet, none of these straightforward approaches have provided a tractable Mn≡N<sub>x</sub> species. While the preparation of well-defined low-valent Mn complexes with multiply bonded linkages remains a challenge, it is one that should be surmountable. Our hope is that manganese imides and nitrides of these types will benefit from synthetic access to unsaturated Mn(I) precursors, such as a four-coordinate [PhBP<sup>i</sup>Pr<sub>3</sub>]Mn–PR<sub>3</sub> species. Oxidative nitrene or nitride group transfer might then prove synthetically more fruitful. Efforts along these lines are part of ongoing studies in our labs.

## Experimental Section

All manipulations were carried out using standard Schlenk or glovebox techniques under a dinitrogen atmosphere. Unless otherwise noted, solvents were deoxygenated and dried by thorough sparging with N<sub>2</sub> gas followed by passage through an activated alumina column. Nonhalogenated solvents were tested with a standard purple solution of benzophenone ketyl in tetrahydrofuran to confirm effective oxygen and moisture removal. Deuterated solvents were purchased from Cambridge Isotopes Laboratories, Inc. and were degassed and stored over activated 3-Å molecular sieves prior to use. The compounds [PhBP<sup>i</sup>Pr<sub>3</sub>]Tl,<sup>70</sup> benzyl potassium,<sup>113</sup> and Li(dbabh)<sup>98</sup> were prepared according to literature procedures. The reagents 2,6-diisopropyl aniline and <sup>t</sup>BuNC were dried over sieves prior to use. Sodium azide was *carefully* dried by heating at 50 °C under vacuum for 2 days. (**CAUTION: Azides are potentially explosive and should generally be stored away from**

*heat and ignition sources.*) Potassium hydride was crushed into a fine powder, washed liberally with petroleum ether, and stored at –30 °C. All other reagents were purchased from commercial vendors and used without further purification. Elemental analyses were performed by Desert Analytics, Tucson, AZ. Varian 300 MHz spectrometers were used to record the <sup>1</sup>H NMR and <sup>31</sup>P NMR spectra at ambient temperature. <sup>1</sup>H chemical shifts were referenced to residual solvent, while <sup>31</sup>P NMR chemical shifts were referenced to 85% H<sub>3</sub>PO<sub>4</sub> at 0 ppm. IR measurements were obtained with a KBr solution cell using a Bio-Rad Excalibur FTS 3000 spectrometer controlled by Bio-Rad Merlin Software (v. 2.97) set at 4 cm<sup>–1</sup> resolution. Electrochemical measurements were recorded in a glovebox under a dinitrogen atmosphere using a BAS CV 100W (Bioanalytical Systems, Inc., West Lafayette, IN). A glassy carbon working electrode, a platinum wire auxiliary-electrode, and an Ag/AgNO<sub>3</sub> nonaqueous reference electrode were used in the electrochemical studies. X-ray diffraction studies were carried out in the Beckman Institute Crystallographic Facility on a Bruker Smart 1000 CCD diffractometer. Crystals were mounted on a glass fiber with Paratone-N oil.

**EPR Measurements.** X-band EPR measurements were recorded using a Bruker EMX spectrometer (controlled by Bruker Win EPR software version 3.0) equipped with a rectangular cavity working in the TE<sub>102</sub> mode. Low-temperature measurements were conducted with an Oxford continuous-flow helium cryostat. Accurate frequency values were recorded from a frequency counter on the microwave bridge. Solution spectra were acquired in 1-methyltetrahydrofuran. Samples were prepared in a glovebox under dinitrogen in quartz EPR tubes equipped with ground-glass joints.

**Magnetic Measurements.** Magnetic measurements were recorded using a Quantum Designs SQUID magnetometer running Magnetic Property Measurement System Rev. 2 software. Data were recorded at 5000 G. The sample was suspended in the magnetometer in a plastic straw sealed under nitrogen with Lilly No. 4 gel caps. Loaded samples were centered within the magnetometer using the DC centering scan at 35 K and 5000 G. Data were acquired at 4–30 K (one data point every 2 K) and 30–300 K (one data point every 5 K). The magnetic susceptibility was adjusted for diamagnetic contributions using the constitutive corrections of Pascal's constants. The molar magnetic susceptibility (χ<sub>m</sub>) was calculated by converting the calculated magnetic susceptibility (χ) obtained from the magnetometer to a molar susceptibility (using the multiplication factor {(molecular weight)/(sample weight)(field strength)}). Effective magnetic moments were calculated using the equation below

$$\mu_{\text{eff}} = (7.997\chi_m T)^{1/2}$$

**Computational Methods.** All calculations were performed using the Jaguar 5.0 program package (Jaguar 5.0, Schrodinger, LLC, Portland, OR). The calculation employed the hybrid DFT functional B3LYP and the basis set LACVP\*\*.<sup>99</sup> Input coordinates for the geometry optimizations were derived from the solid-state structure of [PhBP<sup>i</sup>Pr<sub>3</sub>]Fe≡NAd, where Fe was replaced by Mn.<sup>99</sup> For the Mn imide, the adamantyl group was truncated to a *tert*-butyl group; and for the nitride, the adamantyl group was removed completely. Spin states and molecular charges were imposed as described in the text. The calculations were spin-unrestricted, and no symmetry constraints were used. The default values for geometry and SCF iteration cutoffs were used, and all structures converged under these criteria.

**Synthesis of [PhBP<sup>i</sup>Pr<sub>3</sub>]MnCl (**1**).** A THF solution (18 mL) of [PhBP<sup>i</sup>Pr<sub>3</sub>]Tl (250.0 mg, 36.46 mmol) was added dropwise to a

(113) Bailey, P. L.; Coxall, R. A.; Dick, C. M.; Fabre, S.; Henderson, L. C.; Herber, C.; Liddle, S. T.; Loroño-Gonzalez, D.; Parkin, A.; Parsons, S. *Chem. Eur. J.* **2003**, *9*, 4820.



**Table 1.** X-ray Diffraction Experimental Details for Compounds **1a**, **1b**, **2–4**, and **6–10**

	<b>1a</b>	<b>1b</b>	<b>2</b>	<b>3</b>	<b>4</b>
chemical formula	0.98C <sub>27</sub> H <sub>53</sub> BP <sub>3</sub> MnCl· 0.02TiC <sub>27</sub> H <sub>53</sub> BP <sub>3</sub>	C <sub>54</sub> H <sub>106</sub> B <sub>2</sub> Cl <sub>2</sub> Mn <sub>2</sub> P <sub>6</sub>	C <sub>27</sub> H <sub>53</sub> BP <sub>3</sub> MnI·C <sub>6</sub> H <sub>6</sub>	0.77C <sub>27</sub> H <sub>53</sub> BP <sub>3</sub> MnN <sub>3</sub> · 0.05TiC <sub>27</sub> H <sub>53</sub> BP <sub>3</sub> · 0.18C <sub>27</sub> H <sub>53</sub> BP <sub>3</sub> MnCl	C <sub>34</sub> H <sub>60</sub> BMnP <sub>3</sub>
fw	573.66	1143.61	741.36	1165.21	627.48
<i>T</i> (K)	100	100	100	100	100
$\lambda$ (Å)	0.71073	0.71073	0.71073	0.71073	0.71073
<i>a</i> (Å)	14.493(2)	10.950(2)	9.5424(8)	11.946(2)	9.4120(6)
<i>b</i> (Å)	12.928(2)	11.718(2)	9.9006(8)	34.586(7)	11.8746(8)
<i>c</i> (Å)	34.850(5)	13.432(3)	19.668(2)	15.423(3)	16.736(1)
$\alpha$ (°)	90	68.000(3)	89.780(1)	90	90
$\beta$ (°)	100.344(3)	76.424(4)	89.920(1)	90	105.605(1)
$\gamma$ (°)	90	78.528(3)	82.662(1)	90	90
<i>V</i> (Å <sup>3</sup> )	6424(2)	1541.5(5)	1842.9(3)	6372(2)	1801.6(2)
space group	C2/ <i>c</i> (No. 15)	<i>P</i> 1̄ (No. 2)	<i>P</i> 1̄ (No. 2)	<i>Pnc</i> 2 (No. 30)	<i>P</i> 2 <sub>1</sub> (No. 4)
<i>Z</i>	8	1	2	4	2
<i>D</i> <sub>calcd</sub> (g/cm <sup>3</sup> )	1.186	1.232	1.336	1.215	1.157
$\mu$ (cm <sup>−1</sup> )	7.3	6.9	13.5	8.3	5.2
R1, wR2 ( <i>I</i> > 2 $\sigma$ ( <i>I</i> )) <sup>a</sup>	0.0422, 0.0720	0.0356, 0.0674	0.0305, 0.0586	0.0938, 0.0783	0.0328, 0.0570

	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>
chemical formula	C <sub>39</sub> H <sub>71</sub> BMnNP <sub>3</sub>	C <sub>41</sub> H <sub>63</sub> BMnNP <sub>3</sub>	C <sub>41</sub> H <sub>63</sub> BMnNP <sub>3</sub> ·0.50C <sub>4</sub> H <sub>10</sub> O	C <sub>31</sub> H <sub>53</sub> BBrMnO <sub>4</sub> P <sub>3</sub> Tl	C <sub>42</sub> H <sub>80</sub> BMnN <sub>3</sub> P <sub>3</sub>
fw	712.63	728.58	765.64	932.67	785.75
<i>T</i> (K)	100	100	98	100	100
$\lambda$ (Å)	0.71073	0.71073	0.71073	0.71073	0.71073
<i>a</i> (Å)	10.690(1)	10.4294(9)	19.6218(9)	10.2574(5)	11.2996(7)
<i>b</i> (Å)	11.249(1)	16.812(1)	77.737(4)	18.6613(8)	17.578(1)
<i>c</i> (Å)	18.718(2)	23.393(2)	11.2629(6)	19.9243(9)	23.108(1)
$\alpha$ (°)	98.393(2)	90	90	90	90
$\beta$ (°)	101.614(2)	97.476(1)	90	100.590(1)	93.660(2)
$\gamma$ (°)	107.482(2)	90	90	90	90
<i>V</i> (Å <sup>3</sup> )	2050.8(4)	4066.7(6)	17180(2)	3748.9(3)	4580.4(5)
space group	<i>P</i> 1̄ (No. 2)	<i>P</i> 2 <sub>1</sub> / <i>c</i> (No. 14)	<i>Fdd</i> 2 (No. 43)	<i>P</i> 2 <sub>1</sub> / <i>n</i> (No. 14)	<i>P</i> 2 <sub>1</sub> / <i>n</i> (No. 14)
<i>Z</i>	2	4	16	4	4
<i>D</i> <sub>calcd</sub> (g/cm <sup>3</sup> )	1.154	1.190	1.184	1.652	1.139
$\mu$ (cm <sup>−1</sup> )	4.7	4.7	4.5	58.6	4.2
R1, wR2 ( <i>I</i> > 2 $\sigma$ ( <i>I</i> )) <sup>a</sup>	0.0419, 0.0694	0.0409, 0.0724	0.0514, 0.0731	0.0355, 0.0532	0.0437, 0.0707

$$^a \text{R1} = \sum ||F_o| - |F_c|| / \sum |F_o|, \text{wR2} = \{\sum [w(F_o^2 - F_c^2)^2] / \sum [w(F_o^2)^2]\}^{1/2}.$$

stirring THF suspension of MnCl<sub>2</sub> (46.3 mg, 36.36 mmol). The reaction mixture was stirred for 24 h and filtered through a Celite pad. The colorless solution was concentrated under reduced pressure to dryness to yield an off-white solid. After the solid was washed liberally with petroleum ether, the product was extracted into benzene. Vapor diffusion of petroleum ether into the benzene solution resulted in pale pink blocks (80 mg, 33% yield). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 300 MHz)  $\delta$  ~20 (v. br), 10.3 (br), 8.1 (br). Anal. Calcd for C<sub>27</sub>H<sub>53</sub>BClMnP<sub>3</sub>: C 56.71; H 9.34; N 0. Found: C 56.43; H 9.44; N 0.19.

**[PhBP<sup>Pr</sup><sub>3</sub>]MnI (2).** A THF solution (20 mL) of [PhBP<sup>Pr</sup><sub>3</sub>]Ti (284.7 mg, 41.52 mmol) was added dropwise to a stirring THF suspension of MnI<sub>2</sub> (129.5 mg, 41.52 mmol). The reaction mixture was stirred for 12 h and filtered through a Celite pad. The colorless solution was concentrated under reduced pressure to dryness to yield an off-white solid. After the solid was washed liberally with petroleum ether, the product was extracted into hot benzene. Vapor diffusion of petroleum ether into the benzene solution resulted in colorless crystals (172 mg, 62% yield). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 300 MHz)  $\delta$  21 (v. br), 8.8 (br), 6.6 (br). Anal. Calcd for C<sub>27</sub>H<sub>53</sub>BIMnP<sub>3</sub>: C 48.89; H 8.05; N 0. Found: C 49.12; H 7.54; N < 0.05.

**[PhBP<sup>Pr</sup><sub>3</sub>]Mn(N<sub>3</sub>) (3).** A THF solution (15 mL) of [PhBP<sup>Pr</sup><sub>3</sub>]MnI (406.0 mg, 0.612 mmol) was added to a THF suspension of NaN<sub>3</sub> (203.0 mg, 3.06 mmol). The reaction mixture was stirred for 12 h and filtered through a Celite pad. The cream-colored filtrate was concentrated under reduced pressure to dryness and extracted with benzene (258 mg, 73% yield). Single crystals suitable for X-ray diffraction were grown from vapor diffusion of petroleum ether

into the benzene solution. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 300 MHz)  $\delta$  11 (v. br), 8.3 (br). Anal. Calcd for C<sub>27</sub>H<sub>53</sub>BMnN<sub>3</sub>P<sub>3</sub>: C 56.07; H 9.24; N 7.26. Found: C 55.67; H 8.85; N 7.34.

**[PhBP<sup>Pr</sup><sub>3</sub>]Mn(CH<sub>2</sub>Ph) (4).** An Et<sub>2</sub>O solution of benzylmagnesium bromide (107  $\mu$ L, 1.41 M) was added dropwise to a THF solution of [PhBP<sup>Pr</sup><sub>3</sub>]MnI (100.0 mg, 0.151 mmol) chilled to −90 °C. The solution was allowed to warm slowly to room temperature. The yellow solution was filtered through a Celite pad, concentrated under reduced pressure to dryness, and extracted with petroleum ether (83.4 mg, 88% yield). Yellow block crystals suitable for X-ray diffraction were grown from cyclopentane at −30 °C. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 300 MHz)  $\delta$  16 (v. br), 10.3 (br), 8.3 (br). Anal. Calcd for C<sub>34</sub>H<sub>60</sub>BMnP<sub>3</sub>: C 65.08; H 9.64; N 0. Found: C 64.98; H 9.32; N < 0.05.

**[PhBP<sup>Pr</sup><sub>3</sub>]Mn(Me) (5).** An Et<sub>2</sub>O solution of methylmagnesium bromide (53.5  $\mu$ L, 2.82 M) was added dropwise to a THF solution of [PhBP<sup>Pr</sup><sub>3</sub>]MnI (100.0 mg, 0.151 mmol) chilled to −90 °C. The solution was allowed to warm slowly to room temperature. The pale yellow solution was filtered through a Celite pad, concentrated under reduced pressure to dryness, and extracted with petroleum ether (74.1 mg, 89% crude yield). Crystals were grown from cyclopentane at −30 °C (42.3 mg, 51% yield). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 300 MHz)  $\delta$  ~17 (v. br), 10.1 (br), 8.3 (br). Anal. Calcd for C<sub>28</sub>H<sub>56</sub>BMnP<sub>3</sub>: C 60.99; H 10.24; N 0. Found: C 60.85; H 9.94; N < 0.05.

**[PhBP<sup>Pr</sup><sub>3</sub>]Mn(NH(2,6-*i*-Pr<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>)) (6).** A THF solution (8 mL) of 2,6-diisopropylaniline (25.1 mg, 0.142 mmol) was added dropwise to a THF suspension of KH (5.7 mg, 0.142 mmol) chilled to −90 °C. The solution was allowed to warm slowly to room



temperature. After being stirred for 12 h, the solution was resealed to  $-90\text{ }^{\circ}\text{C}$  and a THF solution (8 mL) of [PhBP<sup>i</sup>Pr<sub>3</sub>]MnI (93.9 mg, 0.142 mmol) was added dropwise. After slowly warming to room temperature, the bright yellow reaction was concentrated under reduced pressure to dryness. The product can be extracted with petroleum ether, filtered through a Celite pad to remove salts, and concentrated under reduced pressure to afford yellow solids (90.9 mg, 90% yield). Single crystals suitable for X-ray diffraction were grown from TMS<sub>2</sub>O/petroleum ether at  $-30\text{ }^{\circ}\text{C}$ . <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 300 MHz)  $\delta$  16.3 (v. br), 10.7 (br), 8.1 (br). Anal. Calcd for C<sub>39</sub>H<sub>71</sub>BMnNP<sub>3</sub>: C 65.73; H 10.04; N 1.97. Found: C 65.81; H 9.71; N 1.88.

[PhBP<sup>i</sup>Pr<sub>3</sub>]Mn(dbabh) (7). A THF solution (5 mL) of [PhBP<sup>i</sup>Pr<sub>3</sub>]-MnI (142.2 mg, 0.214 mmol) was added dropwise to a Et<sub>2</sub>O suspension (10 mL) of Li(dbabh) (42.7 mg, 0.214 mmol) chilled to  $-90\text{ }^{\circ}\text{C}$ . The solution was allowed to warm slowly to room temperature. The yellow solution was filtered through a Celite pad, concentrated under reduced pressure to dryness, and extracted with hot Et<sub>2</sub>O. Cooling the ether solution to  $-30\text{ }^{\circ}\text{C}$  afforded yellow blocks (49.7 mg, 32% yield). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 300 MHz)  $\delta$  21 (v. br), 10.3 (br), 8.1 (br). Anal. Calcd for C<sub>41</sub>H<sub>63</sub>BMnNP<sub>3</sub>: C 67.59; H 8.72; N 1.92. Found: C 66.28; H 8.40; N 1.77.

[PhBP<sup>i</sup>Pr<sub>3</sub>]Mn(1-Ph(isoindolate)) (8). A THF solution (10 mL) of benzyl potassium (40.0 mg, 0.307 mmol) was added to a solution of Hdbabh (59.4 mg, 0.307 mmol) chilled to  $-90\text{ }^{\circ}\text{C}$ . The solution was allowed to warm slowly to room temperature. After being stirred for 24 h, the solution was resealed to  $-90\text{ }^{\circ}\text{C}$  and a THF solution (10 mL) of [PhBP<sup>i</sup>Pr<sub>3</sub>]MnI (203.9 mg, 0.307 mmol) was added dropwise. After slowly warming to room temperature, the orange solution was filtered through a Celite pad, concentrated under reduced pressure to dryness, washed with petroleum ether, and extracted with hot Et<sub>2</sub>O. Cooling the ether solution to  $-30\text{ }^{\circ}\text{C}$  afforded red-orange crystals (71.8 mg, 35% yield). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 300 MHz)  $\delta$  11 (v. br), 8.2 (br). Anal. Calcd for C<sub>41</sub>H<sub>63</sub>BMnNP<sub>3</sub>: C 67.59; H 8.72; N 1.92. Found: C 66.34; H 9.83; N 1.82.

[PhBP<sup>i</sup>Pr<sub>3</sub>]TI-MnBr(CO)<sub>4</sub> (9). [PhBP<sup>i</sup>Pr<sub>3</sub>]TI (150.0 mg, 0.219 mmol) and MnBr(CO)<sub>5</sub> (73.6 mg, 0.262 mmol) were mixed together in 10 mL of benzene. The reaction mixture was heated at  $50\text{ }^{\circ}\text{C}$  for 7 h. After cooling, the solution was concentrated under reduced pressure to dryness, extracted with benzene, filtered through a glass wool pipet, and concentrated to dryness. The yellow solids were washed liberally with petroleum ether and extracted with benzene.

Vapor diffusion of petroleum ether into the solution afforded yellow-orange crystals (79.2 mg, 39% yield). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 300 MHz)  $\delta$  7.76 (d,  $J$  = 6.9 Hz, 2H, H<sub>o</sub> of Ph), 7.53 (t,  $J$  = 6.9 Hz, 2H, H<sub>m</sub> of Ph), 7.30 (t,  $J$  = 6.9 Hz, 1H, H<sub>p</sub> of Ph), 2.00 (m, 6H, CH<sub>2</sub>P), 1.26 (m, 6H, CHMeMe'), 1.09 (d,  $J$  = 6.9 Hz, 18H, CHMeMe'), 1.01 (d,  $J$  = 6.9 Hz, 18H, CHMeMe'). <sup>31</sup>P NMR (C<sub>6</sub>D<sub>6</sub>, 121 MHz)  $\delta$  45.4 (d,  $J_{\text{TI-P}}$  = 8060 Hz). Anal. Calcd for C<sub>31</sub>H<sub>53</sub>-BBrMnO<sub>4</sub>P<sub>3</sub>TI: C 39.92; H 5.73; N 0. Found: C 40.35; H 5.83; N 0.06.

[PhBP<sup>i</sup>Pr<sub>3</sub>]Mn(CN'Bu)<sub>3</sub> (10). A THF solution (5 mL) of sodium naphthalenide (0.237 mmol, prepared by stirring excess Na and stoichiometric naphthalene in THF for 2 h, filter prior to use) was added to a stirring solution of [PhBP<sup>i</sup>Pr<sub>3</sub>]MnI (150.0 mg, 0.226 mmol). Butylisocyanide (43.7  $\mu\text{L}$ , 0.724 mmol) was then quickly added to the mixture. After 11 h, the reaction mixture was concentrated under reduced pressure to dryness, extracted with Et<sub>2</sub>O, and filtered through a glass wool pipet. The solution was again concentrated under reduced pressure to dryness, extracted with petroleum ether, and filtered through a glass wool pipet. Cooling of the solution to  $-30\text{ }^{\circ}\text{C}$  afforded pale yellow crystals (39.4 mg, 22% yield). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 500 MHz)  $\delta$  8.15 (br, 2H), 7.63 (br, 2H), 7.34 (br, 1H), 2.19 (br, 6H), 1.54 (br, 18H), 1.39 (br, 18H), 1.27 (s, 27H), 0.94 (br, 6H). <sup>31</sup>P NMR (C<sub>6</sub>D<sub>6</sub>, 121 MHz)  $\delta$  55.6 (br). Anal. Calcd for C<sub>42</sub>H<sub>80</sub>BMnN<sub>3</sub>P<sub>3</sub>: C 64.20; H 10.26; N 5.35. Found: C 63.84; H 10.10; N 5.22. IR (KBr/THF):  $\nu(\text{cm}^{-1})$  2048 (br).

**Acknowledgment.** We gratefully acknowledge the NSF (CHE-0132216) and the NIH (GM 070757) for financial support; the Beckman Institute (Caltech) for use of its crystallographic facility and SQUID magnetometer; the Materials and Process Simulation Center (Caltech, Goddard) for use of its computational facility. We thank Dr. Theodore A. Betley and Larry Henling for crystallographic assistance and Dr. David M. Jenkins for assistance with SQUID and EPR measurements.

**Supporting Information Available:** Crystallographic data and CIFs for complexes **1a**, **1b**, **2-4**, **6-10**, and {(Hdbabh)MnI( $\mu$ -N-dbabh)}<sub>2</sub>. This material is available free of charge via the Internet at <http://pubs.acs.org>.

IC060735Q