


Photoactivated in Vitro Anticancer Activity of Rhenium(I) Tricarbonyl Complexes Bearing Water-Soluble Phosphines

Sierra C. Marker,[†] Samantha N. MacMillan,[†] Warren R. Zipfel,[‡] Zhi Li,[§] Peter C. Ford,[§] and Justin J. Wilson^{*,†}[†]Department of Chemistry and Chemical Biology, Cornell University, Ithaca, New York 14853, United States[‡]Department of Biomedical Engineering, Cornell University, Ithaca, New York 14853, United States[§]Department of Chemistry and Biochemistry, University of California, Santa Barbara, Santa Barbara, California 93106-9510, United States

Supporting Information

ABSTRACT: Fifteen water-soluble rhenium compounds of the general formula $[\text{Re}(\text{CO})_3(\text{NN})(\text{PR}_3)]^+$, where NN is a diimine ligand and PR_3 is 1,3,5-triaza-7-phosphaadamantane (PTA), tris(hydroxymethyl)phosphine (THP), or 1,4-diacetyl-1,3,7-triaza-5-phosphabicyclo[3.3.1]nonane (DAPTA), were synthesized and characterized by multinuclear NMR spectroscopy, IR spectroscopy, and X-ray crystallography. The complexes bearing the THP and DAPTA ligands exhibit triplet-based luminescence in air-equilibrated aqueous solutions with quantum yields ranging from 3.4 to 11.5%. Furthermore, the THP and DAPTA complexes undergo photosubstitution of a CO ligand upon irradiation with 365 nm light with quantum yields ranging from 1.1 to 5.5% and sensitize the formation of $^1\text{O}_2$ with quantum yields as high as 70%. In contrast, all of the complexes bearing the PTA ligand are nonemissive and do not undergo photosubstitution upon irradiation with 365 nm light. These compounds were evaluated as photoactivated anticancer agents in human cervical (HeLa), ovarian (A2780), and cisplatin-resistant ovarian (A2780CP70) cancer cell lines. All of the complexes bearing THP and DAPTA exhibited a cytotoxic response upon irradiation with minimal toxicity in the absence of light. Notably, the complex with DAPTA and 1,10-phenanthroline gave rise to an IC_{50} value of 6 μM in HeLa cells upon irradiation, rendering it the most phototoxic compound in this library. The nature of the photoinduced cytotoxicity of this compound was explored in further detail. These data indicate that the phototoxic response may result from the release of both CO and the rhenium-containing photoproduct, as well as the production of $^1\text{O}_2$.



INTRODUCTION

Chemotherapy is a critical component for the treatment of nearly all types of cancer. Conventional chemotherapeutic agents, however, are nondiscriminating cytotoxic compounds that induce cell death in both cancerous and healthy tissue, a feature that is manifested in the form of undesirable patient side effects. One approach to increase selectivity and minimize toxic side effects is to employ drugs that are activated by light. This concept, known as photodynamic therapy (PDT)^{1–4} or photoactivated chemotherapy (PACT),⁵ allows for further spatiotemporal control of the cytotoxic effects to maximize cancer cell death, while minimizing healthy cell damage. An ideal candidate for a PDT or PACT agent should be nontoxic in the absence of light and therefore only give rise to toxic effects in regions of light exposure.⁶ The concept of PDT is currently employed in the clinic in the form of the FDA-approved porphyrin derivatives Photofrin, Foscan, and Verteporfin.^{7,8} A potential drawback of the currently used PDT agents is that their mechanism of action stems from their

abilities to photosensitize the formation of singlet oxygen ($^1\text{O}_2$).^{1,9–19} In hypoxic tumors, the efficacy of these PDT agents is substantially diminished due to the lower concentrations of oxygen present.^{20–23} An additional limitation of these porphyrin analogues is their poor aqueous solubility, which requires clever formulation strategies for in vivo administration.^{2,24,25} PACT is currently the subject of significant preclinical investigation.^{5,26} PACT agents generally operate under an oxygen-independent mechanism,⁵ as irradiation with light triggers release of a cytotoxic chemical compound.

Toward the development of improved PDT²⁷ and PACT⁵ agents, many research efforts have explored the potential of transition-metal complexes. In addition to functioning as $^1\text{O}_2$ photosensitizers for PDT applications, many inorganic complexes also undergo photoinduced ligand substitution

Received: October 26, 2017

reactions, which make them useful as PACT agents. Following the concept of PACT, these photosubstitution reactions can be applied to generate cytotoxic complexes from benign precursors in an oxygen-independent manner, rendering them suitable for use in hypoxic conditions.²⁸ Ruthenium PDT^{29–51} and PACT^{22,35,40,42,50,52–55} agents have received the majority of focus, because they possess ideal photophysical properties for this application. The diversity of photophysical processes accessible for other metal ion complexes have prompted researchers to expand their search for new PDT or PACT agents.


Complexes of rhenium, for example, possess rich spectroscopic and photophysical properties that can be leveraged for use in PDT or PACT.⁵⁶ This concept has been demonstrated for several rhenium(I) tricarbonyl complexes, which exhibit potent phototoxic effects in vitro.^{57–61} All of the potential rhenium-based phototoxic agents to date, however, induce cell death via production of singlet oxygen and therefore suffer from some of the same limitations as the conventional porphyrin-based PDT agents. The development of rhenium anticancer PACT agents that act via alternative mechanisms of action remains an interesting unmet research objective that could provide access to new therapeutic modalities. In this study, we describe our efforts toward this goal, in which we have developed a series of rhenium(I) tricarbonyl complexes bearing phosphine ligands that exhibit potent UVA (365 nm) light-activated toxicity against several types of cancer cell lines. Although the biological penetration depth of 365 nm light is too shallow to be useful for in vivo applications, this study provides a proof of concept for the future development of rhenium-based PDT and PACT agents.

RESULTS AND DISCUSSION

Rationale and Approach. Diimine rhenium(I) tricarbonyl complexes are usually inert to photosubstitution reactions and exhibit small luminescence quantum yields.⁶² In contrast, when a phosphine is introduced to the inner coordination sphere, the resulting complexes $[\text{Re}(\text{CO})_3(\text{NN})(\text{PR}_3)]^+$, where NN = diimine ligand and PR_3 = phosphine ligand, are brightly luminescent^{63–66} and photolabile.^{62,67} Upon irradiation with UVA light, the CO trans to the phosphine dissociates, forming the dicarbonyl complex $[\text{Re}(\text{CO})_2(\text{NN})(\text{PR}_3)(\text{S})]^+$, where S is a coordinating solvent molecule.^{62,68} Complexes such as these have been investigated as photo-CO-releasing molecules (photoCORMS),^{68–74} tools to understand the significance of CO as a biological signaling molecule. We envisioned an alternative role of these complexes as photoactivated anticancer agents. The rhenium(I) dicarbonyl photoproduct with a labile coordination site could interact with biological targets, such as DNA, in a manner similar to that reported for rhenium(I) tricarbonyl complexes.^{75–77} In this context, our laboratory has recently demonstrated that rhenium(I) tricarbonyl complexes bearing labile aqua ligands exhibit potent in vitro anticancer activity via a mechanism of action that is distinct from that of cisplatin.⁷⁸ Thus, we anticipated that the structurally similar rhenium(I) dicarbonyl phosphine photoproducts would also be active against cancer cells. Furthermore, the released CO molecule could elicit an additional cytotoxic effect (Scheme 1).^{79–84} Such “dual action” chemotherapeutic agents may be particularly effective against drug-resistant forms of cancer.

In designing our compound library, we employed the phosphines 1,3,5-triaza-7-phosphaadamantane (PTA), tris-(hydroxymethyl)phosphine (THP), and 1,4-diacetyl-1,3,7-

Scheme 1. Photoreaction of $[\text{Re}(\text{CO})_3(\text{NN})(\text{PR}_3)]^+$ Releasing 1 equiv of CO and a Labile Rhenium(I) Complex, Which May Both Give Rise to Cytotoxicity in Cancer Cells^a




^aS is a solvent molecule.

triaza-5-phosphabicyclo[3.3.1]nonane (DAPTA) because these ligands effectively confer water solubility to their coordination complexes.^{85–87} To further screen for cytotoxic effects of the rhenium photoproducts, we explored five different diimine ligands, 1,10-phenanthroline (1), 2,9-dimethyl-1,10-phenanthroline (2), 2,2'-bipyridine (3), 4,4'-dimethyl-2,2'-dipyridyl (4), and 4,4'-dimethoxy-2,2'-bipyridine (5). We hypothesized that this diverse set of compounds would possess a range of photophysical and biological properties, some of which would be valuable for use in PACT.

Synthesis and Characterization. From the 3 phosphines and 5 diimine ligands, a total of 15 rhenium tricarbonyl complexes were synthesized. The synthetic approach followed a previously reported procedure (Scheme 2).^{88,89} In the first step,

Scheme 2. Synthetic Scheme for the Preparation of the $[\text{Re}(\text{CO})_3(\text{NN})(\text{PR}_3)]^+$ Complexes.




$\text{Re}(\text{CO})_5\text{Cl}$ and a diimine ligand were heated to reflux in toluene to afford the yellow complexes *fac*- $[\text{Re}(\text{CO})_3(\text{NN})\text{Cl}]$. These complexes were treated with AgOTf to remove the axial chloride as insoluble AgCl and then allowed to react with the desired phosphine in refluxing tetrahydrofuran (THF)⁶⁸ to afford the compounds PTA-1–PTA-5, THP-1–THP-5, and DAPTA-1–DAPTA-5 (Scheme 2). The PTA and DAPTA complexes were isolated via precipitation as the triflate salts. In contrast, all of the THP complexes, except for THP-5, required additional purification by preparative high-performance liquid chromatography (HPLC). Because the HPLC mobile phase employed 0.1% trifluoroacetic acid, the THP compounds were isolated as trifluoroacetate salts, with the exception of THP-5, which was isolated as the triflate salt upon recrystallization. All of the complexes exhibited good water solubility, enabling the preparation of aqueous solutions at concentrations exceeding 1 mM. In contrast, the *fac*- $[\text{Re}(\text{CO})_3(\text{NN})\text{Cl}]$ starting materials are very poorly soluble in water.⁷⁸ Among the 15 complexes

prepared in this investigation, PTA-1 and THP-3 have previously been reported and developed as photoCORMs.^{68,83}

The complexes were characterized by ¹H (Figures S1–S15 in the Supporting Information), ³¹P{¹H} (Figures S18–S32 in the Supporting Information), and ¹⁹F (Figures S35–S49 in the Supporting Information) NMR spectroscopy, IR spectroscopy (Figures S50–S64 in the Supporting Information), and electrospray ionization mass spectrometry (ESI-MS, Figures S84–S98 in the Supporting Information). The purity of the complexes was verified with elemental analysis and HPLC (Figures S67–S81 in the Supporting Information). The IR spectra of the complexes exhibited three distinct CO stretches, as expected for *fac*-[Re(CO)₃(NN)X] (X = H₂O, Cl, Br) complexes of C_s symmetry.^{78,90–93} The energies of these stretching frequencies were invariant with respect to the coordinating diimine ligands. In contrast, a greater dependence on the nature of the axial phosphine ligand was observed. For example, PTA-1, THP-1, and DAPTA-1 exhibited pronounced CO stretches corresponding to the A'' mode at 2030, 2040, and 2038 cm⁻¹, respectively, and two less intense stretches corresponding to the two A' modes at 1940 and 1920, 1930 and 1910, and 1948 and 1914 cm⁻¹, respectively (Figures S50–S64). These values are consistent with those for other *fac*-[Re(CO)₃(NN)(PR₃)⁺] complexes.^{68,93} The presence of the axial phosphine shifts the CO stretching frequencies to higher energies in comparison to the precursor complexes *fac*-[Re(CO)₃(NN)Cl], which exhibit stretching frequencies near 2020 and 1880 cm⁻¹.⁸⁸ The CO stretching energy of the phosphine complexes is similar to those of cationic complexes such as *fac*-[Re(CO)₃(NN)(OH₂)⁺] (ν_{CO} 2030, 1930, and 1900 cm⁻¹)⁷⁸ and *fac*-[Re(CO)₃(NN)(pyridine)]⁺ (ν_{CO} 2036 and 1931 cm⁻¹).⁹⁴ The higher energy CO stretching frequencies of the phosphine complexes may therefore be a result of both the π acidity of the trans phosphine ligand and the overall cationic charge of the complex, features which decrease π back-bonding to CO π* orbitals.

The ³¹P{¹H} NMR spectra of the PTA and THP complexes show a single resonance. This resonance is shifted downfield from that of the free ligand, signifying coordination to the rhenium center.^{85,86,95,96} For the DAPTA complexes, two signals in the ³¹P NMR spectra were observed. We attribute the two signals to conformational isomers of the DAPTA ligand. As shown in Chart 1, the relative orientation of the two acetyl

Chart 1. Syn and Anti Conformational Isomers of DAPTA



groups gives rise to anti and syn isomers, which will result in distinct NMR signals if their interconversion is slow. The observations of these two isomers in other metal complexes of DAPTA was previously reported.⁹⁷ The ¹H NMR spectra of the DAPTA complexes similarly display resonances of two species that can be attributed to the two conformers. For all of the DAPTA complexes, the anti isomer was the major species in solution. This assignment is based on the ¹H NMR resonances of the methyl groups of the DAPTA ligand, which are inequivalent only for the anti isomer. Finally, the ¹⁹F NMR spectra confirm the nature of the counterions of these

complexes. As expected, the PTA and DAPTA complexes give rise to a signal at −79 ppm, which corresponds to the outer-sphere triflate ion, and the THP complexes display a signal at −76 ppm, arising from the TFA counterion, except for THP-5, which exhibits a signal corresponding to the triflate counterion at −79 ppm.

X-ray Crystallography. Single crystals of all 15 complexes were obtained, and the crystal structures (Figure 1 and Figures S99–S102 in the Supporting Information), with the exception of the previously reported THP-3,⁶⁸ were determined by X-ray crystallography. Crystal structures for PTA-1, THP-1, and DAPTA-1 are shown in Figure 1, and selected interatomic distances and angles for all 14 structures are collected in Table S2 in the Supporting Information. We note that the crystal structure of PTA-1 was recently reported, albeit for a different polymorph of this compound;⁸³ a comparison of this previously reported structure and ours reveals no significant differences with respect to the geometry of the rhenium complex.⁸³ All complexes exhibit the expected octahedral geometry for a d⁶ rhenium(I) center. All DAPTA complexes, except for DAPTA-2, crystallized as the anti isomer, which was the major species exhibited by NMR spectroscopy. No evidence for rotational disorder of the DAPTA acetyl groups was observed in the crystal structures, verifying our conformer assignments. The Re–P bond lengths were relatively invariant among complexes bearing different diimine ligands. A more pronounced dependence was observed as the phosphine ligand was altered. For example, the Re–P distances in PTA-1, THP-1, and DAPTA-1 are 2.4343(6), 2.4602(7), and 2.4449(7) Å, respectively. These values are in the range expected for [Re(CO)₃(NN)(PR₃)⁺] complexes.^{68,83,98} Among the 15 complexes, the THP complexes exhibit a slightly longer average Re–P bond distance of 2.4574 ± 0.013 Å (the error is the standard deviation over the five THP compounds) in comparison to the PTA and DAPTA complexes, for which the average Re–P distances are 2.4340 ± 0.0045 and 2.4414 ± 0.0077 Å, respectively. The longer Re–P distances found in the THP complexes are most likely a consequence of the larger cone angle of this phosphine in comparison to PTA and DAPTA.^{99,100} The Re–CO distances trans to the phosphine vary in a manner that is dependent on the nature of the phosphine as well. In general, this distance is longest for complexes of PTA and DAPTA and shortest for those of THP. For example, the Re–CO distances in PTA-1, THP-1, and DAPTA-1 are 1.962(3), 1.953(3), and 1.963(3) Å.^{68,83} The longer trans Re–CO distances in the DAPTA and PTA complexes may reflect a greater π acidity of this ligand relative to THP. For comparison, the axial Re–CO distances of *fac*-[Re(CO)₃(NN)X]⁺ complexes, where X is a solvent molecule, range from 1.882 to 1.917 Å and are therefore approximately 0.036–0.089 Å shorter than those of the phosphine derivatives.^{78,101,102}

Photophysical Properties. UV–vis and emission spectra of all 15 complexes were measured in air-equilibrated pH 7.4 phosphate-buffered saline. These data are collected in Table 1, and the UV–vis spectra of these complexes are shown in Figures S103–S107 in the Supporting Information.

Representative spectra of the phen complexes (PTA, THP, DAPTA)-1 and the bpy complexes (PTA, THP, DAPTA)-3 are given in Figure 2. The overall band shapes and structures of the absorbance profiles appear to be dependent predominantly on the nature of the diimine ligand rather than the phosphine. For example, the spectra of PTA-1, THP-1, and DAPTA-1 are nearly superimposable, as are those of PTA-3, THP-3, and




Figure 1. X-ray crystal structures of PTA-1, THP-1, and DAPTA-1. Ellipsoids are drawn at the 50% probability level. H atoms bound to C atoms and counterions are omitted for clarity.

Table 1. Photophysical Properties^a

compound	λ_{max} nm (ϵ , M ⁻¹ cm ⁻¹)	Φ_{lum} % (λ_{max} nm)	Φ_{rxn} %	τ_{air} (μs) ^b	τ_{N_2} (μs) ^c
PTA-1	226 (35900 \pm 2600), 225 (22300 \pm 600), 274 (26400 \pm 1900), 368 (3200 \pm 300)	<0.1 (n.d.)	0.06	n.d.	n.d.
THP-1	226 (36800 \pm 1300), 275 (26800 \pm 900), 322 (6400 \pm 200), 366 (3600 \pm 100)	5.1 \pm 1.1 (528)	1.4 \pm 0.4	1.5	4.8
DAPTA-1	225 (41600 \pm 7300), 275 (27200 \pm 4900), 323 (5700 \pm 900), 367 (3200 \pm 400)	10.7 \pm 0.6 (516)	1.1 \pm 0.1	1.9	5.6
PTA-2	228 (40500 \pm 1100), 253 (26100 \pm 600), 283 (26200 \pm 600), 372 (2500 \pm 300)	<0.1 (n.d.)	0.015	n.d.	n.d.
THP-2	228 (34600 \pm 1200), 286 (23400 \pm 800), 308 (11900 \pm 400), 372 (2100 \pm 70)	4.5 \pm 0.4 (518)	2.0 \pm 0.4	1.0	3.7
DAPTA-2	227 (36800 \pm 6800), 285 (23800 \pm 700), 309 (11500 \pm 300), 373 (2100 \pm 50)	7.2 \pm 0.2 (507)	1.5 \pm 0.1	2.0	7.8
PTA-3	248 (24400 \pm 800), 307 (11700 \pm 500), 319 (12900 \pm 500), 349 (3300 \pm 300)	<0.1 (538)	0.0044	n.d.	n.d.
THP-3	222 (18900 \pm 200), 249 (20100 \pm 1900), 308 (10700 \pm 400), 318 (12500 \pm 400), 343 (3600 \pm 700)	6.1 \pm 1.7 (536)	1.6 \pm 0.1	0.4	0.5
DAPTA-3	246 (22600 \pm 500), 308 (10700 \pm 200), 319 (13000 \pm 300), 345 (3400 \pm 100)	9.1 \pm 3.0 (528)	2.3 \pm 0.5	0.6	0.8
PTA-4	252 (29200 \pm 3600), 303 (13600 \pm 1800), 315 (14700 \pm 2100), 345 (4300 \pm 500)	<0.1 (530)	0.021	n.d.	n.d.
THP-4	252 (24000 \pm 3000), 304 (11800 \pm 1500), 315 (13400 \pm 1700), 338 (4600 \pm 500)	6.5 \pm 2.0 (528)	3.8 \pm 0.9	0.4	0.6
DAPTA-4	250 (8200 \pm 4300), 305 (12800 \pm 2000), 315 (14900 \pm 2300), 339 (4000 \pm 400)	11.5 \pm 3.9 (518)	5.5 \pm 0.3	0.6	1.0
PTA-5	227 (36100 \pm 2600), 249 (32900 \pm 2300), 304 (9800 \pm 700), 338 (4400 \pm 300)	<0.1 (520)	0.018	n.d.	n.d.
THP-5	223 (33900 \pm 1000), 251 (30500 \pm 900), 303 (8800 \pm 300), 337 (4500 \pm 80)	3.4 \pm 1.1 (537)	1.2 \pm 0.2	0.3	0.4
DAPTA-5	224 (41400 \pm 2000), 251 (34100 \pm 1900), 303 (9600 \pm 500), 332 (4700 \pm 300)	7.1 \pm 3.0 (527)	1.9 \pm 0.1	0.4	0.6

^an.d. = not determined due to weak or undetectable luminescence intensity. ^bLuminescence lifetime measured in air-equilibrated pH 7.4 PBS.

^cLuminescence lifetime measured in nitrogen-saturated pH 7.4 PBS.

DAPTA-3. The high-energy (<325 nm) features in these spectra most likely correspond to intraligand π – π^* transitions and are therefore expected to be only dependent on the nature of the diimine ligand. The low-energy features correspond to the metal-to-ligand charge-transfer (MLCT) transition,^{63,93,103–108} a state that dominates the photochemical properties of Re(CO)₃ complexes. Similarly, the energy of the MLCT transition is almost independent of the nature of the phosphine. This result suggests that the phosphine ligands only weakly perturb the energies of metal-based donor orbitals, so that variations in the MLCT bands can be attributed to differences in the ligand π^* orbitals. For the phen complexes (1 series), the MLCT band is centered at 367 nm; for the bpy complexes (3 series) this band occurs at 345 nm, consistent with the higher energy π^* orbitals of this less conjugated ligand.

The luminescence properties of the complexes were also evaluated (Figures S108–S112 in the Supporting Information). Representative emission spectra for complexes (PTA, THP, DAPTA)-1 are shown in Figure 3. All complexes bearing the THP and DAPTA ligands were emissive in air-equilibrated pH 7.4 phosphate-buffered saline (PBS). The quantum yields for emission under these conditions ranged from 3.4 to 6.1% for the THP complexes, depending on the nature of the diimine ligand. The DAPTA complexes were generally more emissive,

with quantum yields ranging from 7.1 to 11.5%. Notably, complexes of the PTA ligand were effectively nonemissive under these conditions. The energy of the emission, which arises from the ³MLCT state,^{109,110} varies with both the nature of the diimine ligand and the phosphine. For example, DAPTA complexes give rise to higher emission energies in comparison to the analogous THP complexes (Table 1). This result indicates that the phosphine ligand plays a larger role in the ³MLCT in comparison to the ¹MLCT state that was probed by the absorption spectra, as described above. The excited state lifetimes were also measured in both air-equilibrated (τ_{air}) and deoxygenated PBS (τ_{N_2}) (Table 1 and Figures S113–S117 in the Supporting Information), and these range from 0.2 to 2.0 μs for the air-equilibrated PBS solutions, consistent with triplet-based MLCT excited state lifetimes measured for similar rhenium complexes.^{62,93} Notably, the lifetimes of these complexes are extended under a nitrogen atmosphere, suggesting that O₂ is an effective quenching agent.

Photolysis of the axial CO ligand was investigated by monitoring the UV–vis spectral profiles of the compounds in pH 7.4 PBS as they were irradiated with 365 nm light (Figures S118–S132 in the Supporting Information). Representative spectral changes of PTA-1, THP-1, and DAPTA-1 upon irradiation are shown in Figure 4. Whereas all complexes of




Figure 2. UV-vis spectra in pH 7.4 PBS solutions for complexes (PTA, THP, DAPTA)-1 (top) and (PTA, THP, DAPTA)-3 (bottom).




Figure 3. Emission spectra in pH 7.4 PBS (excitation wavelength 350 nm, concentration 30 μ M) for complexes PTA-1 (solid blue line), THP-1 (long-dash red line), and DAPTA-1 (short-dash green line).

PTA were not photoreactive at this excitation wavelength, the THP and DAPTA complexes exhibited a marked change in their UV-vis spectra. We note that, in contrast to our results, PTA-1 was previously reported to undergo CO loss upon excitation at 365 nm.⁸³ We only observed photoreactivity of the PTA complexes when they were irradiated with higher energy UVB light (280–300 nm). Upon irradiation of the THP and DAPTA complexes, the bands near 340–360 nm decayed with the concomitant formation of a new broad feature at lower energies. In all cases, the photoconversion of these complexes

proceeded cleanly, as evidenced by the presence of well-defined isosbestic points. Furthermore, gas chromatographic analysis of three representative compounds showed that 1 equiv of CO is released per molecule (Table S3 in the Supporting Information).

Photochemical quantum yields for these compounds were measured via ferrioxalate actinometry and are given in Table 1. The nonluminescent PTA complexes were generally not photoreactive at this excitation wavelength; upper estimates for their photoreaction quantum yields are less than 0.1%. In contrast, the DAPTA and THP complexes exhibited quantum yields ranging from 1.1 to 5.5%. These values are substantially lower than those previously measured for related rhenium phosphine complexes in degassed organic solvents;^{62,68} the use of aerated aqueous solutions in this case may lead to the lower quantum yields measured for the compounds described here.

Because PDT agents act by generating singlet oxygen, we investigated the $^1\text{O}_2$ -sensitization quantum yields (Φ_Δ) for the 15 rhenium complexes in air-equilibrated pH 7.4 PBS solutions (Table 2). Upon irradiation with 365 nm light, $^1\text{O}_2$ was quantified via the *N,N*-4-dimethyl-nitrosoaniline/histidine assay following previously reported procedures.^{59,111} The quantum yields were determined in reference to those of phenanthrene ($\Phi_\Delta = 0.98$). All PTA compounds exhibited no significant production of $^1\text{O}_2$ upon irradiation. This result suggests that the excited state quenching of these complexes occurs in an O_2 -independent manner. The THP and DAPTA complexes produce $^1\text{O}_2$ with low to high quantum yields; DAPTA-2 exhibits the largest quantum yield of 70%, whereas both THP-4 and DAPTA-4 do not give rise to detectable quantities of $^1\text{O}_2$ upon irradiation.

Reinvestigation of the Photophysical Properties of THP-3. The photophysical properties of the compound THP-3 were described in a previous publication from one of our laboratories.⁶⁸ In this earlier study, a luminescence quantum yield of 18% and a photochemical quantum yield of 21% were reported. Additionally, the dicarbonyl photoproduct of THP-3 was found to exhibit photoluminescence with a maximum emission wavelength of 585 nm. Upon reanalysis of THP-3 in this investigation, we have found substantially lower luminescence and photochemical reaction quantum yields of 6.1% and 1.6%, respectively. Furthermore, the dicarbonyl photoproduct, obtained by exhaustive photolysis of THP-3, was found to be nonemissive. These contradictory results prompted further investigations of this compound in our laboratories both at Cornell and at UC Santa Barbara.

As originally reported, THP-3 was isolated as the triflate salt,⁶⁸ rather than the TFA salt. The triflate salt of THP-3 was independently prepared as previously described and characterized by elemental analysis and X-ray crystallography, which gave unit cell parameters that matched those in the previous report. The photophysical properties were investigated using conditions identical with those employed in the original study with equipment and facilities at UC Santa Barbara. The photoluminescence quantum yield was determined using rhodamine B ($\Phi_{\text{lum}} = 65\%$) as a reference standard. Quantum yields in pH 7.4 PBS in deaerated and air-equilibrated solutions were determined to be 9.7% and 8.0%, respectively (Figure S133 in the Supporting Information). The latter value is in reasonable agreement with that measured at Cornell for the TFA analogue of THP-3 under similar conditions. Furthermore, after exhaustive photolysis, no photoluminescence was detected, demonstrating that the triflate counterion also does




Figure 4. UV-vis spectra of the photoreactions of PTA-1 (110 μ M, left), THP-1 (115 μ M, middle), and DAPTA-1 (105 μ M, right) in pH 7.4 PBS over a 4 h irradiation period with 365 nm light.

Table 2. Singlet Oxygen Quantum Yields

compound	Φ_{Δ} , %
PTA-1	<1
THP-1	48.5 \pm 0.5
DAPTA-1	54.0 \pm 1
PTA-2	<1
THP-2	45.5 \pm 0.5
DAPTA-2	70.5 \pm 2.5
PTA-3	<1
THP-3	8.7 \pm 0.1
DAPTA-3	9.1 \pm 0.3
PTA-4	<1
THP-4	<1
DAPTA-4	<1
PTA-5	<1
THP-5	15.0 \pm 0
DAPTA-5	12.5 \pm 0.5

not alter this aspect of THP-3. Finally, the photochemical reaction quantum yield of this compound was reanalyzed using a FieldMaxII-TO power meter from Coherent with a PM10 V1 thermopile photodetector (calibrated) to measure the light intensities. These studies afford a quantum yield of 2.4% (Figure S134 in the Supporting Information), in reasonable agreement with the value of 1.6% obtained at Cornell. These photoreaction quantum yields are approximately a factor of 10 smaller than the value previously reported,⁶⁸ and we speculate

that a simple calculation error may account for the error in the original measurement. This also may be the case for the luminescence quantum yield, although another source of such an error would be a partially oxidized rhodamine B standard. The presence of an impurity in the original samples of THP-3 may explain the claimed photoproduct luminescence.⁶⁸ A likely candidate is *fac*-[Re(CO)₃(bpy)(OH₂)]⁺, as this complex is formed by aquation of the *fac*-[Re(CO)₃(bpy)(THF)]⁺, the starting material for THP-3. We have independently prepared and characterized *fac*-[Re(CO)₃(bpy)(OH₂)](OTf) and shown it to have an absorption spectrum close to that of THP-3 (Figure S135 in the Supporting Information) and to exhibit weak luminescence in the region of \sim 580 nm (Figure S136 in the Supporting Information). These results underline the importance of confirming the purity of all compounds prior to photophysical measurements due to the sensitivity of these measurements to small quantities of photoactive impurities.

Density Functional Theory. DFT calculations were employed to gain a deeper understanding of the photophysical properties of these compounds. The phen compounds, PTA-1, THP-1, and DAPTA-1, were explored as a representative subset of all of the complexes. Geometries were optimized at the BP86 level¹¹² of theory in the gas phase, and vertical singlet excited state energies were calculated with time-dependent DFT (TD-DFT) employing a solvent continuum dielectric model for water¹¹³ and the hybrid functional PBE0. A map of the lowest energy singlet excited states of these complexes is




Figure 5. State energy diagram of PTA-1 (left), THP-1 (middle), and DAPTA-1 (right) and the corresponding electron density difference maps for relevant excited states. Negative (white) and positive (red) electron density illustrates the transitions.




Figure 6. Single-dose (200 μM) cell viability of HeLa cells for all complexes. The compounds were tested in triplicate under both light and dark conditions. For light-irradiated samples, an irradiation time of 30 min with 365 nm light at a photon flux of $(2.38 \pm 0.31) \times 10^{-10}$ einstein/s was employed.

given in Figure 5. For THP-1 and DAPTA-1, the lowest energy singlet excited states are MLCT in character, as can be discerned from the difference density maps that show a shift of electron density from the metal to the phenanthroline ligand in the excited state. In contrast, the four lowest energy singlet excited states for PTA-1 cannot be reasonably described as MLCT states; the difference density maps of these excited states represent N 2p \rightarrow L π^* transitions or ligand-to-ligand charge-transfers where the N 2p electrons are initially localized on the PTA phosphine. The fifth excited state of PTA-1 is the definitive MLCT, as indicated by the difference density map. Because the MLCT is generally the state which mediates the luminescence and facilitates internal conversion to the photoreactive ligand field excited state, the presence of four lower energy N 2p \rightarrow L π^* states may enhance nonradiative and nonreactive decay pathways to the ground state. One can alternatively think of this as a photoinduced electron transfer (PeT) quenching mechanism, which is commonly used to design fluorescent sensors via modulation of nitrogen lone pair orbital energies.¹¹⁴ The efficient quenching of the MLCT due to the nitrogen lone pairs of the PTA complexes would, therefore, explain the lack of observed luminescence for this class of compounds. Furthermore, the lack of photosubstitution is also a consequence of the efficient quenching of the ¹MLCT state, which must occur more rapidly than intersystem crossing to states that would enable these photoreactivities.⁶² We note that the DAPTA ligand likewise possesses an N 2p lone pair that may also be available for PeT quenching. We hypothesized that the presence of the electron-withdrawing acetyl groups increase the redox potential of the nitrogen lone pair, such that PeT quenching is less thermodynamically viable. This hypothesis was further supported by examining the DFT energies of the nitrogen lone pair orbitals of PTA-1 and DAPTA-1. The energy of this orbital in PTA-1 is higher than that of DAPTA-1 by approximately 0.3 eV. As such, the nitrogen lone pair in DAPTA-1 is less susceptible to oxidation, rendering PeT quenching less thermodynamically favored.

Photoinduced Anticancer Activity. To rapidly screen for photoactivated in vitro anticancer activity, HeLa cells were

treated with the 15 compounds at a single dose of 200 μM. The cells were incubated for 4 h with the rhenium compounds and then irradiated with 365 nm light for 30 min. Another batch of cells, used as the nonirradiated control, was subjected to the same conditions in the absence of light. The viability of the cells in the presence of the 15 different compounds under the conditions of light irradiation and dark are shown in Figure 6 and Table S10 in the Supporting Information. All compounds at this concentration level, with the exception of THP-2, THP-5, and DAPTA-5, exhibited little or no toxicity in the dark (within experimental uncertainty). Upon irradiation, all of the THP and DAPTA complexes, with the exception of THP-3, induced significant cell death, indicating that these compounds are potential PDT or PACT agents. Consistent with their lack of photoreactivity, none of the PTA complexes exhibited significant cytotoxicity upon exposure to light.

On the basis of this initial screening, the most potent compounds, THP-1, THP-2, THP-5, and all DAPTA complexes, were further evaluated in full dose–response studies to determine 50% growth inhibitory concentration (IC₅₀) values in HeLa cells. These values are collected in Table 3 (Figures S140–S148 in the Supporting Information). Consistent with the single-dose studies, the compounds were all nontoxic in the absence of light. Under conditions of light irradiation at 365 nm, the compounds exhibited enhanced cytotoxicity. Notably, THP-2 and DAPTA-1 possess IC₅₀ values that are less than 10 μM. The ratio of IC₅₀ values in the dark over those determined upon irradiation give phototoxicity indices (PI) that are >34 for DAPTA-1 (Figure 7) and >21 for THP-2. The exact phototoxicity indices could not be determined because the compounds do not induce cell death in the dark at the highest concentration (200 μM) screened.

DAPTA-1 and THP-1 were further evaluated in wild-type (A2780) and cisplatin-resistant ovarian (A2780CP70) cancer cell lines (Table 4 and Figures S149–S152 in the Supporting Information). THP-1 exhibited enhanced cytotoxic effects in the presence of light in both A2780 and A2780CP70 cell lines. The IC₅₀ values of the irradiated THP-1, however, are

Table 3. IC₅₀ values in HeLa (Cervical Cancer) Cells in the Absence and Presence of 365 nm Light

compound	IC ₅₀ value (μM) HeLa		PI ^a
	dark	light	
THP-1	>200	26.4 ± 9.2	>7.6
DAPTA-1	>200	5.9 ± 1.4	>33.9
THP-2	>200	9.6 ± 4.2	>20.8
DAPTA-2	>200	19.2 ± 2.9	>10.4
DAPTA-3	>200	14.9 ± 3.2	>13.4
DAPTA-4	>200	60.3 ± 18.2	>3.3
THP-5	72.6 ± 23.2	68.0 ± 4.3	1.1
DAPTA-5	>200	24.3 ± 9.1	>8.2

^aThe phototoxic index (PI) is the dark IC₅₀ value divided by the light IC₅₀ value. For light-irradiated samples, an irradiation time 1 h with 365 nm light at a photon flux $(2.38 \pm 0.31) \times 10^{-10}$ einstein/s was employed.





Figure 7. Dose–response curves for DAPTA-1 in HeLa cells in the presence (red dashed line) and absence (black solid line) of 365 nm light irradiation. For light-irradiated samples, an irradiation time of 1 h with 365 nm light at a photon flux of $(2.38 \pm 0.31) \times 10^{-10}$ einstein/s was employed.

approximately 6-fold higher in the cisplatin-resistant cell line, indicating that this photoactivated product is susceptible to platinum resistance mechanisms. In contrast, DAPTA-1 gave rise to light-induced IC₅₀ values of 2.1 and 3.2 μM in the A2780 and A2780CP70 cell lines, indicating that this compound can circumvent cisplatin resistance.

Identifying the Cytotoxic Species. Upon photolysis of this class of compounds in water, both CO and the rhenium dicarbonyl complexes $fac-[Re(NN)(PR_3)(CO)_2(OH_2)]^+$ are released, and singlet oxygen is produced. We hypothesized that the resulting dicarbonyl rhenium photoproduct could induce

cytotoxicity by interacting covalently with biomolecules, in a similar manner to that for the corresponding tricarbonyl aqua species that we have studied.⁷⁸ To explore this possibility, we synthesized two dicarbonyl analogues, one of DAPTA-1 and the other of PTA-3, for further evaluation of their anticancer activity. The dicarbonyl analogue of DAPTA-1 was pursued because DAPTA-1 is the most potent compound in our library. In contrast, the dicarbonyl analogue of PTA-3 was investigated because this compound exhibits no phototoxicity; this compound would enable us to determine if the dicarbonyl species could elicit a cytotoxic response in the absence of CO release. The target compounds $fac-[Re(phen)(CO)_2(DAPTA)-Cl]$ (DAPTA-1A) and $fac-[Re(bpy)(CO)_2(PTA)Cl]$ (PTA-3A) were synthesized in two steps from DAPTA-1 or PTA-3 (Scheme 3). Following previously reported procedures,^{115–118}

Scheme 3. Synthetic Scheme for the Photoproducts DAPTA-1A and PTA-3A

DAPTA-1 or PTA-3 was treated with trimethylamine *N*-oxide (TMAO) and NEt₄Cl and heated to reflux in a mixture of CH₂Cl₂ and CH₃OH. This reaction induces the liberation of the axial CO ligand as CO₂, providing the open coordination site that is then occupied by the chloride ions in solution to yield DAPTA-1A or PTA-3A as red solids. The compounds DAPTA-1A (Figures S16, S33, S65, and S82 in the Supporting Information) and PTA-3A (Figures S17, S34, S66, and S83 in the Supporting Information) were characterized by conventional spectroscopic methods. The chlorido complexes were sufficiently soluble for biological studies after stirring them in water for several hours; their dissolution is hypothesized to occur with concomitant aquation of the chlorido ligand. Photolyzed solutions of DAPTA-1 give rise to a peak with the same retention time on HPLC as DAPTA-1A (Figure S153 in the Supporting Information). Furthermore, solutions of photolyzed DAPTA-1 exhibit the same UV–vis (Figure S154 in the Supporting Information) and ¹H NMR spectra (Figure S155 in the Supporting Information) as DAPTA-1A, confirming this compound to be the photoproduct.

From the photolysis of DAPTA-1 in D₂O (Figure S155 in the Supporting Information), single crystals of an aqua analogue of DAPTA-1A, $fac-[Re(phen)(DAPTA)-(CO)_2(OD_2)]^+$ (DAPTA-1A-aqua), precipitated. Crystals of PTA-3A were obtained by vapor diffusion of diethyl ether into

Table 4. IC₅₀ Values in A2780 (Ovarian) and A2780CP70 (Ovarian Cisplatin-Resistant) Cancer Cells in the Absence and Presence of 365 nm Light

compound	IC ₅₀ value (μM)					
	A2780			A2780CP70		
	dark	light	PI	dark	light	PI ^a
THP-1	>200	4.6 ± 1.4	43.5	>200	29.9 ± 7.7	6.7
DAPTA-1	>200	2.2 ± 1.1	90.9	>200	3.2 ± 0.7	62.5
cisplatin	0.18 ± 0.07			5.14 ± 1.1		

^aPhototoxic index (PI) is the dark IC₅₀ value divided by the light IC₅₀ value. For light-irradiated samples, an irradiation time 1 h with 365 nm light at a photon flux $(2.38 \pm 0.31) \times 10^{-10}$ einstein/s was employed.




Figure 8. X-ray crystal structures of **DAPTA-1A-aqua** (left) and **PTA-3A** (right). Bond lengths and angles are reported in Table S2 in the Supporting Information. Ellipsoids are drawn at the 50% probability level. H atoms bound to C atoms and counterions are omitted for clarity.

N,N-dimethylformamide (DMF). The structures of these complexes were determined by single-crystal X-ray diffraction (Figure 8). The equatorial Re–CO distances are approximately 0.06 Å shorter than those observed in **DAPTA-1** and 0.03 Å shorter than those in **PTA-3**. The decrease in distance presumably arises from the loss of a π -accepting CO ligand. **PTA-3A** also exhibits a notably smaller Cl–Re–P angle of 168.77(3)°, significantly deviating from linearity. The O–Re–P distance in **DAPTA-1A-aqua** is also considerably bent at an angle of 171.70(9)°. This axial ligand bend represents an interesting deviation from the corresponding angles found in **DAPTA-1** and **PTA-1** and other complexes of the type *fac*-[Re(CO)₃(NN)X], with C–Re–X angles between 174 and 178°. ^{102,119} A similar rhenium phosphine compound, *fac*-[Re(CO)₂(NN)(PR₃)Cl] (PR₃ = P(OEt)₃), was reported to exhibit an Cl–Re–P angle of 173°. ¹¹⁷

With purified **DAPTA-1A** and **PTA-3A** in hand, the anticancer activity of these photoproducts was analyzed. **PTA-3A** (Figure S156 in the Supporting Information) was nontoxic in HeLa cells up to 200 μ M in the dark. **DAPTA-1A** (Figure 9 and Figure S157 in the Supporting Information) exhibited a mild cytotoxic effect at concentrations >50 μ M. Additionally, solutions of **DAPTA-1** after its exhaustive photolysis gave rise to a cytotoxic response similar to that of **DAPTA-1A** (Figure S158 in the Supporting Information), supporting the toxic effects of this dicarbonyl complex. The cytotoxicity of **PTA-3A** and **DAPTA-1A** were further investigated under conditions of UVA light irradiation to probe for additional phototoxicity of these complexes. Neither complex, however, induced additional cytotoxic effects in the presence of light. Taken together, these results indicate that the phototoxicity of **DAPTA-1** could arise only in part from the dicarbonyl photoproduct and that additional cytotoxicity is elicited through a separate mechanism. The lack of cytotoxicity of **PTA-3A** is surprising because related rhenium carbonyl complexes with a labile coordination site bind covalently to biomolecules ^{75–77} and induce cell death. ^{78,120–122} Although we only investigated **DAPTA-1A** and **PTA-3A**, we hypothesize




Figure 9. Dose–response curve for **DAPTA-1A** (black solid line) in HeLa cells in the absence of light and **DAPTA-1** (red dashed line) under 365 nm light irradiation.

that the phototoxicity of the other compounds studied here could arise in part from the dicarbonyl product.

The limited toxicity of these dicarbonyl complexes could possibly be attributed to their different cellular uptakes and intracellular distributions in comparison to the tricarbonyl phosphine precursors. Upon photolysis in aqueous solution, dicarbonyl aqua complexes, such as **DAPTA-1A-aqua** (Figure 8), are formed. On coordination of water to metal centers, the pK_a decreases substantially. Depending on their pK_a values, the photogenerated dicarbonyl complexes could exist either as the charge-neutral hydroxides or as the cationic aqua complexes at physiological pH. On the basis of charge, such species would be expected to have different cellular uptake and localization. To investigate the relative abundances of these species, the pK_a values of aquated solutions of **DAPTA-1A**, **PTA-3A**, and the photoproduct of **THP-1** were determined. These solutions were titrated with base as the pH and UV–vis spectra were monitored. Upon basification, a shift in the absorbance spectra from approximately 370 to 430 nm was observed. This shift arises from the deprotonation of the aqua complex. On the basis of these data, the pK_a values for the dicarbonyl aqua

photoproducts of DAPTA-1, THP-1, and PTA-3 are 9.51, 10.07, and 8.75, respectively (Figures S159–S164 in the Supporting Information). These values are similar to those measured for related monoaqua transition-metal complexes,^{123,124} and they indicate that at physiological pH the compounds exist as the cationic aqua complexes.

The low cytotoxicity of the rhenium photoproducts, DAPTA-1A and PTA-3A, prompted us to consider the potential role of $^1\text{O}_2$ and CO in mediating cell death. Nearly all of the rhenium complexes in this study sensitize the formation of highly toxic $^1\text{O}_2$ (Table 2). No clear direct correlation between Φ_Δ and the phototoxicity exist, however, indicating that $^1\text{O}_2$ sensitization is not the only mechanism of activity of these complexes. To further test this hypothesis, the phototoxicity of DAPTA-1 was investigated under hypoxia, conditions in which $^1\text{O}_2$ sensitizers are less effective. HeLa cells were treated with DAPTA-1 with a procedure similar to that described above, except the cells were subjected to hypoxic conditions (5% CO_2 , 95% N_2) before and after exposure to 365 nm light for 1 h. Under hypoxic conditions, DAPTA-1 induced moderate phototoxicity characterized by an IC_{50} value of $20 \pm 1.5 \mu\text{M}$ (Figure S165 in the Supporting Information). This value is somewhat larger than that measured under normoxic conditions (Table 3), suggesting that $^1\text{O}_2$ sensitization does play a role in this compound's mechanism of action. The lack of a complete suppression of phototoxicity under these conditions, however, does verify that other factors, namely the dicarbonyl photoproduct and CO, may also contribute to the observed activity.

A third possible contributor to the complexes' phototoxicity is CO. Understanding the biological effects of CO is an active area of research. CO at low concentrations can have therapeutic properties, as it elicits anti-inflammatory effects and vasodilation.^{125–136} The application of CO as a cytotoxic agent for the destruction of malignant cells has also been explored.^{72,79–81,84,137–141} CO most likely kills cells by binding with high affinity to cytochrome c oxidase,¹⁴² thereby inhibiting the mitochondrial respiration pathway. Inhibition of the mitochondrial respiration pathway in this manner with CO reduces cancer cell proliferation by metabolically exhausting cells.¹⁴³ If the cytotoxic effects of these complexes are mediated solely by CO, then the photoactivated anticancer activity could potentially correlate to the quantum yield for CO release. Across the 15 compounds tested, however, we do not observe this correlation, suggesting that additional factors are in operation. One possible explanation is that the quantum yields measured in buffer are not representative of those in the complex biological media, where binding to proteins such as albumin may alter the values in an unexpected manner. For example, the photophysical properties of the organic dye Rose Bengal and TPPS (*meso*-tetra(4-sulfonatophenyl)porphyrin) are dramatically altered in the presence of albumin and the cell membrane environment.¹⁴⁴ Alternatively, differential cell uptake and intracellular localization of these compounds due to their different lipophilicities may play a large role in mediating the cytotoxic effects of the photoreleased CO and would explain the clear lack of a correlation between CO photorelease quantum yield and cytotoxicity.

CONCLUSIONS

The structural and photophysical properties of 15 rhenium tricarbonyl complexes bearing one of 3 water-soluble phosphines and 5 different diimine ligands were investigated.

THP and DAPTA complexes exhibited relatively efficient quantum yields for luminescence, photosubstitution of CO, and the production of singlet oxygen. The lack of photoreactivity of the PTA complexes was attributed to quenching from the nitrogen lone pair orbitals, via a photoinduced electron transfer mechanism. The photoreactivities of the THP and DAPTA complexes were leveraged to generate PDT or PACT agents that simultaneously released CO and a labile rhenium complex upon exposure to UV light. The complex DAPTA-1 exhibited the most potent cytotoxic effects upon light exposure in HeLa cells and in wild-type and cisplatin-resistant ovarian cancer cells. The dicarbonyl photoproduct, DAPTA-1A, was only moderately cytotoxic, suggesting that an additional mechanism of cell death may be operable, which could be attributed to $^1\text{O}_2$ and the released CO. Because no direct correlations between phototoxicity and both $^1\text{O}_2$ sensitization and CO photosubstitution reaction quantum yields could be discerned, we believe that more complicated and subtle factors are in operation for this class of compounds. Regardless, these compounds represent a new type of combined PDT and PACT agent that operates under a multi-action mechanism. They can be easily synthesized, are water-soluble, and exhibit high phototoxic indices. Although this class of compounds is activated only by UVA light, they represent an important proof of principle for the development of new phototoxic rhenium compounds. Ongoing efforts are being pursued to shift the excitation wavelength of these complexes to the red where in vivo tissue penetration would be deeper.

EXPERIMENTAL SECTION

Methods and Materials. Rhenium carbonyl was purchased from Pressure Chemicals (Pittsburgh, PA, USA). Iodobenzene dichloride was synthesized as previously reported.¹⁴⁵ The diimine ligands 2,2'-bipyridine (bpy), 4,4'-dimethyl-2,2'-dipyridyl (dmbpy), 4,4'-dimethoxy-2,2'-bipyridine (dmobpy), 1,10-phenanthroline (phen), and 2,9-dimethyl-1,10-phenanthroline (dmphen) were purchased from Sigma-Aldrich (St. Louis, MO, USA), Alfa Aesar (Heysham, England), or Beantown Chemical (Hudson, NH, USA) and were used as received. $[\text{Re}(\text{CO})_3(\text{phen})\text{Cl}]$, $[\text{Re}(\text{CO})_3(\text{dmphen})\text{Cl}]$, $[\text{Re}(\text{CO})_3(\text{bpy})\text{Cl}]$, $[\text{Re}(\text{CO})_3(\text{dmbpy})\text{Cl}]$, and $[\text{Re}(\text{CO})_3(\text{dmobpy})\text{Cl}]$ were synthesized using a previously reported procedure.^{88,89} The phosphine ligands 1,3,5-triaza-7-phosphaadamantane (PTA) and tris(hydroxymethyl)phosphine (THP) were purchased from Alfa Aesar (Heysham, England) and Acros (Geel, Belgium), respectively, and were both used as received. The compound 1,4-diacetyl-1,3,7-triaza-5-phosphabicyclo[3.3.1]nonane (DAPTA) was synthesized from PTA using a previously reported procedure.⁸⁶ All solvents were ACS grade or higher. All reactions were carried out under ambient atmospheric conditions without any effort to exclude water or oxygen.

Physical Measurements. NMR samples were prepared as solutions using $\text{MeOD-}d_4$, $\text{DMSO-}d_6$, $\text{DMF-}d_7$, or D_2O as the solvent. NMR spectra were acquired on a Varian Inova 400 MHz spectrometer. ^1H NMR chemical shifts were referenced to residual solvent peaks versus tetramethylsilane (TMS) at 0 ppm or an external standard, 1,4-dioxane, at 3.75 ppm in D_2O . ^{19}F and ^{31}P NMR spectra were referenced using an external standard of KPF_6 in D_2O (^{19}F , $\delta -72$ ppm vs CFCl_3 at 0 ppm; ^{31}P , $\delta -145$ ppm vs H_3PO_4 at 0 ppm). Samples for IR spectroscopy were prepared as KBr pellets and were analyzed on a Nicolet Avatar 370 DTGS (ThermoFisher Scientific, Waltham, MA, USA). The HPLC system (LC-20AP) used for purification of the Re-THP complexes (excluding THP-5) consisted of a SPD-20AV UV-vis detector monitored at 270 and 220 nm (Shimadzu, Japan) and a Epic Polar preparative column (120 Å, 10 μm , 25 cm \times 20 mm; ES Industries, West Berlin, NJ, USA) at a flow rate of 14 mL/min using a binary mobile phase containing 0.1% trifluoroacetic acid (TFA) and H_2O . The method for purification consisted of 0–5 min at 10%

MeOH, followed by a linear gradient of 100% MeOH for over 20 min, and then 10% MeOH for 1 min. Analytical chromatography was carried out on a LC-20AT pump with a SPD-20AV UV–vis detector monitored at 270 and 220 nm (Shimadzu, Japan) using an Ultra Aqueous C18 column (100 Å, 5 μ m, 250 mm \times 4.6 mm; Restek, Bellefonte, PA, USA) at a flow rate of 1 mL/min with a mobile phase containing 0.1% trifluoroacetic acid (TFA) in H₂O or MeOH. The method consisted of 5 min at 10% MeOH, followed by a linear gradient to 100% MeOH over 20 min. High-resolution mass spectra (HRMS) were recorded on an Exactive Orbitrap mass spectrometer in positive ESI mode (ThermoFisher Scientific, Waltham, MA, USA) with samples injected as acetonitrile/water solutions with 1% formic acid. Elemental analyses (C, H, N) were performed by Atlantic Microlab Inc. (Norcross, GA, USA). UV–visible spectra were recorded on a Cary 8454 UV–vis instrument (Agilent Technologies, Santa Clara, CA) or a Beckman Coulter DU800 UV–vis instrument using 1 cm quartz cuvettes. Lifetime measurements were collected as described below. Phototoxicity experiments were conducted using a UVA handlamp elevated 8 cm from the cells to give a photon flux of $(2.38 \pm 0.31) \times 10^{-10}$ einstein/s at 365 nm. Analytical photochemical measurements were performed using a Newport mercury/xenon arc lamp. The light output was modulated using a combination of a Newport heat-absorbing glass filter (50.8 \times 50.8 mm) with infrared cutoff (Schott KG5 filter glass), a Newport mercury line bandpass filter (25.4 mm, center wavelength 365.0 \pm 2 nm), and a Newport visible absorbing filter (50.8 \times 50.8 mm, center wavelength 340 nm) made of dark optical glass to isolate monochromatic 365 nm light. Luminescence quantum yield measurements were carried out on a Beckman Coulter DU800 UV–vis instrument and Varian Eclipse Fluorometer.

X-ray Crystallography. Single crystals were grown using vapor diffusion of diethyl ether into solutions of methanol or acetonitrile. For the compound DAPTA-1A, the crystals were isolated from a solution of D₂O, and those for PTA-3A were collected from vapor diffusion of diethyl ether into DMF. Low-temperature (223 K) X-ray diffraction data for PTA-1, THP-1, DAPTA-1, PTA-2, THP-2, DAPTA-2, PTA-3, DAPTA-3, PTA-4, THP-4, DAPTA-4, PTA-5, THP-5, DAPTA-5, DAPTA-1A, and PTA-3A were collected on a Bruker X8 Kappa diffractometer coupled to an ApexII CCD detector with graphite-monochromated Mo K α radiation (λ = 0.71073 Å). The structures were solved through intrinsic phasing using SHELXT¹⁴⁶ and refined against F^2 on all data by full-matrix least squares with SHELXL¹⁴⁷ following established refinement strategies.¹⁴⁸ All non-hydrogen atoms were refined anisotropically. Hydrogen atoms bound to carbon were included in the models at geometrically calculated positions and refined using a riding model. Hydrogen atoms bound to oxygen were located in the difference Fourier synthesis and subsequently refined semirefreely with the help of distance restraints. The isotropic displacement parameters of all hydrogen atoms were fixed to 1.2 times the U value of the atoms they are linked to (1.5 times for methyl groups). Details of the data quality and a summary of the residual values of all the refinements are given in Tables S1 and S2 in the Supporting Information.

Photochemistry. Photochemical reactions were monitored by UV–vis spectroscopy. Stock solutions of the rhenium compounds at 1–2 mM concentrations in PBS (pH 7.4) were diluted in PBS (pH 7.4) to a final volume of 3 mL at a concentration of approximately 40–150 μ M. The absorbance of these samples was between 0.1 and 0.5 at 365 nm. The solutions were stirred in a 1 cm quartz cuvette and irradiated with 365 nm light using the UVA handlamp described above. At different time points, the UV–vis spectra were acquired. The photoreaction was deemed complete when no further changes in the UV–vis spectra were observed. For DAPTA-1, the photoreaction was monitored by HPLC (Figure S153 in the Supporting Information), UV–vis (Figure S154 in the Supporting Information) and ¹H NMR spectroscopy (Figure S155 in the Supporting Information).

CO Release Measurements. These measurements were carried out as previously reported.¹⁴⁹ Millimolar stock solutions of DAPTA-1, DAPTA-4, and THP-3 were prepared by dissolving a known amount of the respective complex in 10 mL of pH 7.4 PBS. A custom designed

Schlenk cuvette⁷¹ (internal volume of 23.6 mL) was loaded with 6 mL of the stock solution and a magnetic stir bar and sealed. The UV–vis spectrum was measured using a Shimadzu UV2401 (PC) spectrophotometer. The sample was then exposed to 365 nm light using an Oriel 200–500 W/Hg arc lamp filtered by a 365 nm mercury line interference filter. The power of the incident light (50 mW) was measured using a Coherent FieldMaxII-TO Power Meter. After 1 h, the UV–vis spectrum was measured again. This was repeated until there were no further significant absorbance changes at the λ_{max} (~400 nm) of the product, indicating that the photoreaction was complete (~3 h).

A gastight syringe was used to remove a 100 μ L sample from the headspace of the cell through a special screw top fitted with a three-layer laminated silicone GC septum. This sample was analyzed by gas chromatography with thermal conductivity detection (GC-TCD) using an Agilent 6890N gas chromatograph fitted with a Carboxen 1010 PLOT fused silica capillary column ($L \times I.D.$ = 30 m \times 0.53 mm, average thickness 30 μ m). The GC inlet temperature was 230 °C. The oven was held at 35 °C for 14 min, then ramped at 20 °C/min to 245 °C, and finally held at 245 °C for 10 min. The retention time for carbon monoxide (CO) is about 11.2 min, and that for carbon dioxide (CO₂) is 21 min. Dioxygen and dinitrogen from ambient air have retention times of about 8.2 and 8.4 min, respectively. The ChemStation autointegration function was used to calculate the peak area for CO.

The amount of CO in the 100 μ L sample was calculated using a calibration curve made by injecting known amounts of CO from a Schlenk flask. The moles of CO in the headspace of the Schlenk cuvette (SC moles of CO) was calculated using the equation

$$\text{SC moles of CO} = \frac{\text{moles of CO (syringe)} \times (\text{SC headspace} + \text{injection volume})}{\text{injection volume}} \quad (1)$$

The volume of the headspace for the Schlenk cuvette (SC headspace) is calculated by subtracting the solution volume from the total volume of the Schlenk cuvette.

For CO, the partition coefficient between the gas phase and water is ~50 at ambient temperature (41 versus 0.84 mmol/(L atm)); thus, essentially all of the CO will be in the gas phase. The stoichiometric ratio (moles CO released/moles of photoreacted complex) was calculated by dividing the SC moles of CO by the moles of complex photolyzed. The average and sample standard deviation of three injections are reported. The results are shown in Table S3 in the Supporting Information.

Emission Quantum Yield. The luminescence quantum yields were measured relative to the standard quinine sulfate (Φ = 0.52, 0.05 M H₂SO₄), which was cross-referenced in our laboratory to harmaline (Φ = 0.32, 0.005 M H₂SO₄).¹⁵⁰ An excitation wavelength of 350 nm was used for the samples and standards. The compounds were measured as solutions in pH 7.4 PBS with the absorbance maintained below 0.1 to prevent inner filter effects.¹⁵⁰ At least five different concentrations of the samples and standards were measured by UV–vis and fluorescence spectroscopy, and the absorbance at 350 nm was plotted versus the integrated emission intensity. The slopes of the resulting lines were used in the equation

$$\Phi_{\text{sample}} = \Phi_{\text{ref}} \frac{S_{\text{sample}} \eta_{\text{sample}}^2}{S_{\text{ref}} \eta_{\text{ref}}^2} \quad (2)$$

where Φ_{ref} is the quantum yield of the reference, quinine sulfate, S is the slope of either the sample or the reference, and η is the refractive index.

Photochemical Reaction Quantum Yield. The photochemical reaction quantum yields were determined via potassium ferrioxalate actinometry.¹⁵¹ All solutions were prepared with minimal light exposure and were stored in the dark. To determine the photon flux of our lamp, a 1.5 mL volume of potassium ferrioxalate (6.1 mM, 0.05 M H₂SO₄) was irradiated with monochromatic 365 nm light for 30 s

with continuous stirring in a 1 cm quartz cuvette. After irradiation, a 0.1 mL aliquot of the ferrioxalate solution was added to 2.9 mL of 1,10-phenanthroline (5.7 mM, 0.5 mM H₂SO₄) containing sodium acetate buffer (1.8 M, pH 5). This procedure was repeated for an identical potassium ferrioxalate solution kept in the dark to correct for any thermal reactions. The resulting solution was shaken for 30 s and then incubated for 1 min. The absorbance was then recorded for both the irradiated solution and the dark solution. The equation used to determine photon flux is

$$q = \frac{\Delta A V_1 V_2 N_A}{1000(\Phi \epsilon V_3 t l)} \quad (3)$$

where N_A is Avogadro's number, q is photon flux (photons/s), ΔA is the change in absorbance at 510 nm for the irradiated solution minus the dark solution, V_1 (mL) is the total volume after dilution, V_2 (mL) is the volume irradiated, V_3 (mL) is the volume of irradiated sample diluted into 1,10-phenanthroline, Φ is the quantum yield at 365 nm, ϵ is the extinction coefficient (M⁻¹ cm⁻¹), l is path length (cm), and t is time (s). For potassium ferrioxalate at 365 nm, $\Phi = 1.25$ and $\epsilon = 11000$ M⁻¹ cm⁻¹.¹⁵¹ Photon flux and quantum yields were cross-referenced to the actinometer, Reinecke's salt ($\Phi = 0.38$ at 365 nm), to verify the accuracy of our measurements.¹⁵¹

The quantum yields of the rhenium compounds were determined by first measuring the photon flux of the light source with potassium ferrioxalate, as described above. The rhenium compound was irradiated in pH 7.4 PBS with the monochromatic 365 nm light. The concentrations of these samples were such that the absorbance was greater than 2 at 365 nm, ensuring that all photons were absorbed by the sample. For some samples, a correction factor ($1 - 10^{-A}$) was applied to the photon flux if the solution was below an absorbance of 2, where A is the absorbance of the sample at the irradiation wavelength, 365 nm. This correction factor accounts for the fact that in more optically dilute solutions not all photons may be absorbed. After a given time period of irradiation, the sample was diluted in 2.5 mL of PBS and the UV-vis absorbance was measured. This diluted sample was subjected to continued irradiation for 0.5–2 h until the photoreaction was complete, enabling us to determine the extinction coefficient of the photoproduct. To determine the quantum yield of the rhenium compounds, eq 3 was applied to determine Φ , using the photon flux determined from the potassium ferrioxalate actinometry. Absorbance change was monitored at 425–450 nm, depending on the sample.

Lifetime Measurements. Laser excitation for the phosphorescence lifetime measurements was provided by pulsing the 405 nm laser line from a four-line iChrome MLE laser (Toptica Photonics AG, Munich, Germany). The diode laser in the iChrome was triggered by a DG535 Digital Delay/Pulse Generator (Stanford Research, Sunnyvale, CA, USA) at 100 kHz and delivered 100 ns fwhm 405 nm excitation pulses. The 405 nm pulses were fiber-delivered to a sample-filled cuvette, and phosphorescence was collected at 90° through a second fiber for delivery to a Bialkali photomultiplier tube (HC125, Hamamatsu, Bridgewater, NJ, USA) through a 470 nm long-pass filter (HQ470lp, Chroma Technology, Bellows Falls, VT). The time-resolved photon counts were collected in 40 ns time bins using a SR430 Multichannel scaler (Stanford Research, Sunnyvale, CA). Data were transferred to a PC via the SR430 GPIB bus and fit to the standard exponential decay model using MagicPlot Pro software. Measurements were collected in PBS solutions at 1000 μ M. For deoxygenated measurements, nitrogen gas was bubbled into the PBS solutions for 15 min, and then the lifetime was determined.

Singlet Oxygen Measurements. Indirect singlet oxygen measurements were performed using the *N,N*-dimethyl-4-nitrosoaniline/histidine assay, where histidine is oxidized by singlet oxygen and the oxidized histidine reacts with *N,N*-dimethyl-4-nitrosoaniline.¹¹¹ All measurements were performed in PBS solutions containing *N,N*-dimethyl-4-nitrosoaniline (25 μ M) and histidine (10 mM).⁵⁹ The reaction mixture was irradiated with monochromatic 365 nm light in 1 cm glass cuvettes at 2 min increments, and the bleaching of *N,N*-dimethyl-4-nitrosoaniline was monitored at 440 nm. The absorbance

at 440 nm was plotted against irradiation time. The singlet oxygen quantum yield (Φ_{sample}) of the rhenium complexes was determined relative to the quantum yield of phenalenone ($\Phi_{\text{ref}} = 0.98$)¹⁵² using the formula

$$\Phi_{\text{sample}} = \Phi_{\text{ref}} \frac{S_{\text{sample}} I_{\text{ref}}}{S_{\text{ref}} I_{\text{sample}}} \quad (4)$$

where S is the slope of the plot of absorbance vs time and I is the overlap of the lamp emission spectra and the sample absorbance spectra using the formula

$$I = \int_{\lambda} I_0 [1 - 10^{A(\lambda)}] d\lambda \quad (5)$$

where I_0 is the light-flux intensity of the lamp and A is the absorbance of the compound at the irradiation wavelength. The standard phenalenone was cross-referenced to the complex *fac*-[Re(CO)₃(NNN)]⁺, where NNN = *N,N*-bis(quinolin-2-ylmethyl)-amino]-5-valeric acid (Re-COOH), reported by Leonidova et al.⁵⁹ They reported a singlet oxygen quantum yield of 20% in water. Using our experimental setup, we measured a value of 33–37%.

DFT Studies. All computational calculations were performed using version 3.0.3 of the ORCA computational package.¹⁵³ The resolution of identity (RI) approximation was used throughout for computational efficiency. Geometry optimizations were performed in the gas phase using the functional BP86,¹¹² the basis set def2-TZVP and its auxiliary def2-TZV/J,¹⁵⁴ and the zeroth-order regular approximation for relativistic effects (ZORA).^{155–157} Optimized geometries were subjected to TD-DFT calculations to determine the nature of the relevant excited states.¹¹³ For these calculations, the hybrid functional PBE0 was employed,¹⁵⁸ using the same basis sets and ZORA as described above. For the TD-DFT calculations, the SMD solvation model for water was employed.¹⁵⁹ Twenty singlet excited states were calculated for PTA-1, THP-1, and DAPTA-1. All optimized geometry coordinates are given in Tables S4–S6, the lowest energy singlet excited states are summarized in Tables S7–S9, and the frontier Kohn–Sham orbitals are located in Figures S137–S139 in the Supporting Information.

Cell Culture and Cytotoxicity. The HeLa (cervical cancer) cell line was obtained from American Type Culture Collection (ATCC) and cultured using Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% fetal bovine serum (FBS). A2780 (ovarian cancer) and A2780CP70 (cisplatin-resistant ovarian cancer) cell lines were provided by the Cell Culture Facility of Fox Chase Cancer Center¹⁶⁰ (Philadelphia, PA, USA). These cells were cultured as monolayers with Roswell Park Memorial Institute (RPMI)-1640 culture media supplemented with 10% FBS. All cell lines were grown in a humidified incubator at 37 °C with an atmosphere of 5% CO₂. Cells were passed at 80–90% confluence using trypsin/EDTA. Cells were tested monthly for mycoplasma contamination with the Plasmotest mycoplasma detection kit from InvivoGen.

The procedure for determining phototoxicity was modified from the OECD Guidelines, except MTT was used instead of Neutral Red to determine cell viability.¹⁶¹ All compounds, except for DAPTA-1A and PTA-3A, were dissolved in PBS at pH 7.4 to prepare a 1–2 mM stock solution. DAPTA-1A and PTA-3A were solubilized in water after exhaustive stirring. For single-dose studies, only HeLa cells were used. For both the single-dose and full-dose responses all cells were grown to 80–90% confluence, detached with trypsin/EDTA, seeded in 96-well plates at 2000 cells/well for HeLa cells and 4000 cells/well for A2780 and A2780CP70 cells in 100 μ L of growth media, and incubated for 24 h. The medium was removed and replaced with fresh medium (200 μ L) containing a single concentration of PTA, THP, or DAPTA-1–5. The cells were incubated for 4 h, and then the cells were irradiated with 365 nm light for 30 min (one plate under the same dosing conditions was kept outside the incubator in the dark as a control). After exposure to light, the plates were incubated for an additional 44 h, the medium was removed from the wells, and 3-(4,5-dimethylthiazol-2-yl)-2,5-tetrazolium bromide (MTT) in DMEM or RPMI (200 μ L, 1 mg/mL) was added. The additional 44 h incubation

was performed to ensure that the cells were in the logarithmic growth phase and that the cells had adequate time to regrow after exposure to the complexes. After 4 h, the MTT/DMEM or RPMI solution was removed, and the formazan crystals were dissolved in 200 μ L of an 8/1 mixture of DMSO and pH 10 glycine buffer. The absorbance at 570 nm in each well was measured using a BioTek Synergy HT plate reader. Cell viability was determined by normalizing the absorbance of the treated wells to untreated wells. The percent viability data shown are an average of three independent experiments with six replicates per concentration. The same procedure was followed as above for the full dose response, but a 1 h irradiation period was employed instead. Cell viability was again determined by normalizing the absorbance of the treated wells to untreated wells. The concentrations of the compounds versus percent viability were plotted to produce the dose–response curves, which were analyzed using a logistic sigmoid function fit with MagicPlot Pro software. The reported IC₅₀ values are the average of three independent experiments with six replicates per concentration level. For hypoxic conditions, the cells were dosed with the rhenium complex and then incubated in a hypoxia chamber (Billips-Rothenburg, Inc., Del Mar, CA, USA), which was purged with an atmosphere of 95% N₂ and 5% CO₂ for 4 h. The chamber was removed from the incubator, purged with the hypoxic gas mixture again, and then irradiated with 365 nm light for 1 h. After this irradiation period, the cells were further incubated under hypoxia at 37 °C for 16 h. The plates were then removed from the hypoxia chamber and allowed to recover under normoxic (5% CO₂, 21% O₂) conditions at 37 °C for an additional 24 h prior to analysis of cell viability with MTT, as described above. As an additional control experiment to test the effect of the UVA light on the cells, cells in the absence of rhenium complexes were subjected to the same irradiation conditions. The cell viability of the irradiated cells, relative to control cells that were not exposed to light, was always >85%, indicating that the direct light exposure alone has only a minimal effect on the cells.

pK_a Measurements. Stock solutions of DAPTA-1, THP-1, and PTA-3A were made in water at approximately 2 mM and then diluted to 84–89 μ M in water. DAPTA-1 and THP-1 were irradiated with 365 nm light until the photoreaction was complete as evidenced by UV–vis spectroscopy. PTA-3A was dissolved directly for analysis. The solutions containing the rhenium complexes were acidified to an approximate pH of 2.0 with perchloric acid. These solutions were titrated with aqueous KOH, and the pH and UV–vis absorbance spectra were measured at each stage of the titration. The titration apparatus consisted of a 100 mL water-jacketed glass vessel at 25 °C. Nitrogen gas was passed through the solution containing 30% KOH during the experiment to exclude any CO₂. The electrode was calibrated using standard buffer solutions prior to the experiment. Carbonate-free solutions of KOH were prepared in freshly boiled Milli-Q water and standardized against potassium hydrogen phthalate.¹⁶² The spectral changes resulting from the different pH values were used to determine the pK_a values using the HypSpec software package.^{162,163} Species distribution diagrams were plotted using HySS.¹⁶⁴ The titration curves were fit from a wavelength range of 360–405 nm. All absorbance spectra and titration curves are in Figures S159–S164 in the Supporting Information.

Synthesis of Re(CO)₃Cl. A mixture of rhenium carbonyl Re₂(CO)₁₀ (1.90 g, 2.91 mmol) and CH₂Cl₂ (20 mL) was stirred until Re₂(CO)₁₀ was completely dissolved. Then iodobenzene dichloride (0.711 g, 2.59 mmol) was added as a solid to the solution, and the reaction mixture was stirred for 4 h at room temperature. The desired product, a white solid, was filtered and washed with hexanes (20–30 mL). Yield: 1.268 g (68%). IR (KBr, cm^{−1}): 3435 w, 2462 w, 2157 m, 2035 s, 1958 s, 1002 w, 942 w, 912 w, 590 s, and 555 s.

Synthesis of [Re(CO)₃(phen)(PTA)]OTf (PTA-1). A mixture of [Re(CO)₃(phen)Cl] (0.224 g, 0.485 mmol) and AgOTf (0.125 g, 0.485 mmol) in THF (50 mL) was heated to reflux at 75 °C for 3 h in the dark. The white solid AgCl was removed via vacuum filtration. To the remaining yellow filtrate was added PTA (0.0915 g, 0.582 mmol), and the resulting solution was heated to reflux for an additional 15 h. The reaction mixture was cooled to room temperature, and the THF was removed using rotary evaporation. The remaining yellow solid was

dissolved in a minimal amount of methanol (5–10 mL) and filtered through Celite. The addition of diethyl ether (20–30 mL) precipitated the desired product as a yellow solid, which was isolated by filtration and dried in vacuo. Yield: 0.298 g (78%). ¹H NMR (400 MHz, MeOD-*d*₄): δ 9.53 (d, 2H, *J* = 5.2 Hz), 9.00 (d, 2H, *J* = 8.3 Hz), 8.34 (s, 2H), 8.15 (dd, 2H, *J* = 5.2, 8.3 Hz), 4.38 (d, 3H, *J* = 13.1 Hz), 4.26 (d, 3H, *J* = 13.1 Hz), 3.64 (s, 6H). ¹⁹F NMR (376 MHz, MeOD-*d*₄, external standard KPF₆): δ −79.56. ³¹P{¹H} NMR (162 MHz, MeOD-*d*₄, external standard KPF₆): δ −78.24. IR (KBr, cm^{−1}): 3460 m, 2910 w, 2360 m, 2330 m, 2030 s, 1940 s, 1920 s, 1430 m, 1265 m, 1244 m, 1153 m, 1029 m, 970 w, 950 w, 850 w, 726 w, 637 m, 440 w. ESI-MS (positive ion mode, CH₃CN/H₂O 70/30 and 1% formic acid): *m/z* 608.08745 ([M]⁺, calcd 608.08558). Anal. Calcd for PTA-1·CH₃OH (C₂₃H₂₄F₃N₅O₇PreS): C, 35.03; H, 3.07; N, 8.88. Found: C, 34.85; H, 3.02; N, 8.81.

Synthesis of [Re(CO)₃(phen)(THP)]TFA (THP-1). A mixture of [Re(CO)₃(phen)Cl] (0.100 g, 0.206 mmol) and AgOTf (0.053 g, 0.206 mmol) in THF (24 mL) was heated to reflux at 75 °C for 3 h in the dark. The white solid AgCl was removed via vacuum filtration. To the remaining yellow filtrate was added THP (0.038 g, 0.309 mmol), and the solution was heated to reflux for an additional 15 h. The reaction mixture was cooled to room temperature, and then THF was removed by rotary evaporation. The remaining yellow solid was dissolved in a minimum volume of methanol (5 mL) and filtered through Celite. The crude yellow oil (0.167 g) was purified using preparatory RP-HPLC using a CH₃OH/H₂O gradient (0–5 min, 10% CH₃OH; 5–30 min, 10–100% CH₃OH). Purified yield: 0.100 g (69%). ¹H NMR (400 MHz, MeOD-*d*₄): δ 9.56 (d, 2H, *J* = 5.2 Hz), 8.90 (d, 2H, *J* = 8.3 Hz), 8.25 (s, 2H), 8.07 (dd, 2H, *J* = 5.2, 8.3 Hz), 3.75 (t, 2H), 3.75 (s, 6H). ¹⁹F NMR (376 MHz, MeOD-*d*₄, external standard KPF₆): δ −76.37. ³¹P{¹H} NMR (162 MHz, MeOD-*d*₄, external standard KPF₆): δ 2.43. IR (KBr, cm^{−1}): 3420 m, 3210 m, 3070 m, 2830 w, 2780 w, 2650 w, 2040 s, 1930 s, 1910 w, 1670s, 1520 w, 1430 s, 1200 s, 1130 s, 1050 s, 888 m, 856 s, 800 s, 723 s, 635 w, 618 w, 538 m, 500 m, 483 w, 436 w. ESI-MS (positive ion mode, CH₃CN/H₂O 70/30 and 1% formic acid): *m/z* 575.03961 ([M]⁺, calcd 575.03763). Anal. Calcd for THP-1·H₂O (C₂₀H₁₉F₃N₅O₉Pre): C, 34.05; H, 2.71; N, 3.97. Found: C, 34.02; H, 2.61; N, 4.03.

Synthesis of [Re(CO)₃(phen)(DAPTA)]OTf (DAPTA-1). A mixture of [Re(CO)₃(phen)Cl] (0.200 g, 0.42 mmol) and AgOTf (0.106 g, 0.42 mmol) in THF (47 mL) was heated to reflux at 75 °C for 3 h in the dark. The white solid AgCl was removed via vacuum filtration. To the remaining yellow filtrate was added DAPTA (0.113 g, 0.46 mmol), and the solution was heated to reflux for an additional 15 h. The reaction mixture was cooled to room temperature, and THF was removed by rotary evaporation. The remaining yellow solid was dissolved in a minimum amount of methanol (5–10 mL) and filtered through Celite. The addition of diethyl ether (20–30 mL) afforded the desired product as a yellow solid, which was isolated by vacuum filtration and dried in vacuo. Yield: 0.255 g (73%). ¹H NMR (400 MHz, MeOD-*d*₄): DAPTA complexes show the syn and anti isomers in a 1/5 ratio; however in the aromatic region the isomers are indistinguishable, δ 9.60 (t, 2H, anti + syn), 9.02 (d, 2H, anti + syn), 8.35 (s, 2H, anti + syn), 8.18 (m, 2H, anti + syn), 5.56 (d, anti), 5.04 (d, syn), 4.75 (d, anti), 4.48 (d, syn), 4.41 (d, anti), 4.12 (m, anti), 3.93 (d, anti), 3.78 (d, anti), 3.64 (s, syn), 3.41 (d, syn), 3.34 (d, anti), 3.04 (d, syn), 2.95 (d, anti), 1.94 (s, syn), 1.91 (s, anti), 1.80 (s, anti). ¹⁹F (376 MHz, MeOD-*d*₄, external standard KPF₆): δ −79.54. ³¹P{¹H} (162 MHz, MeOD-*d*₄, external standard KPF₆): δ −44.80 ((syn), −48.75 (anti)). IR (KBr, cm^{−1}): 3452 m, 2038 s, 1948 s, 1914 s, 1643 s, 1427 m, 1330 m, 1278 m, 1258 m, 1225 w, 1148 w, 1029 w, 987 w, 888 w, 849 w, 802 w, 720 w, 638 s, 540 m, 498 w, 425 w. ESI-MS (positive ion mode, CH₃CN/H₂O 70/30 and 1% formic acid): *m/z* 680.10846 ([M]⁺, calcd 680.10671). Anal. Calcd for DAPTA-1 (C₂₃H₂₄F₃N₅O₈PreS): C, 36.23; H, 2.92; N, 8.45. Found: C, 36.35; H, 2.79; N, 8.66.

Synthesis of [Re(CO)₃(dmphen)(PTA)]OTf (PTA-2). A mixture of [Re(CO)₃(dmphen)Cl] (0.243 g, 0.473 mmol) and AgOTf (0.121 g, 0.473 mmol) in THF (54 mL) was heated to reflux at 75 °C for 3 h in the dark. The white solid AgCl was removed via vacuum filtration.

To the remaining yellow filtrate was added PTA (0.111 g, 0.71 mmol), and the solution was heated to reflux for an additional 15 h. The reaction mixture was cooled to room temperature, and then the THF was removed using rotary evaporation. The remaining yellow solid was dissolved in a minimum amount of methanol (5–10 mL) and filtered through Celite. The addition of diethyl ether (20–30 mL) afforded the desired product as a yellow solid, which was isolated by vacuum filtration and dried in vacuo. Yield: 0.290 g (78%). ^1H NMR (400 MHz, $\text{MeOD}-d_4$): δ 8.80 (d, 2H, J = 8.4 Hz), 8.23 (s, 2H), 8.13 (d, 2H, J = 8.4 Hz), 4.33 (d, 3H, J = 13.2 Hz), 4.17 (d, 3H, J = 13.2 Hz), 3.42 (s, 6H), 3.34 (s, 6H). ^{19}F NMR (376 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -79.56. $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -79.09. IR (KBr, cm^{-1}): 3448 m, 3052 w, 2913 w, 2033 s, 1958 s, 1934 s, 1445 m, 1384 w, 1266 s, 1222 w, 1150 m, 1031 m, 1013 w, 972 m, 873 m, 802 w, 743 w, 638 s, 582 w, 509 w, 440 w. ESI-MS (positive ion mode, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ 70/30 and 1% formic acid): m/z 636.11922 ($[\text{M}]^+$, calcd 636.11743). Anal. Calcd for PTA-2 ($\text{C}_{24}\text{H}_{24}\text{F}_3\text{N}_3\text{O}_8\text{PReS}$): C, 36.73; H, 3.08; N, 8.92. Found: C, 36.66; H, 3.09; N, 8.98.

Synthesis of $[\text{Re}(\text{CO})_3(\text{dmphen})(\text{THP})]\text{TFA}$ (THP-2). A mixture of $[\text{Re}(\text{CO})_3(\text{dmphen})\text{Cl}]$ (0.209 g, 0.407 mmol) and AgOTf (0.104 g, 0.407 mmol) in THF (46 mL) was heated to reflux at 75 °C for 3 h in the dark. The white solid AgCl was removed via vacuum filtration. To the remaining yellow filtrate was added THP (0.076 g, 0.612 mmol), and the solution was heated to reflux for an additional 15 h. The reaction mixture was cooled to room temperature, and then THF was removed by rotary evaporation. The crude yellow oil was purified using preparatory RP-HPLC using a $\text{CH}_3\text{OH}/\text{H}_2\text{O}$ gradient (0–5 min, 10% CH_3OH ; 5–50 min, 10–100% CH_3OH). Purified yield: 0.158 g (53%). ^1H NMR (400 MHz, $\text{MeOD}-d_4$): δ 8.70 (d, 2H, J = 8.4 Hz), 8.13 (s, 2H), 8.03 (d, 2H, J = 8.4 Hz), 3.44 (s, 6H), 3.37 (s, 6H). ^{19}F NMR (376 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -76.35. $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ 6.53. IR (KBr, cm^{-1}): 3036 m, 2837 w, 2449 w, 2044 s, 1945 s, 1909 s, 1674 s, 1594 w, 1506 w, 1442 m, 1370 w, 1202 m, 1140 m, 1043 m, 892 w, 859 m, 837 w, 800 w, 781 w, 722 w, 652 w, 619 w, 552 w, 513 w, 499 w. ESI-MS (positive ion mode, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ 70/30 and 1% formic acid): m/z 603.07053 ($[\text{M}]^+$, calcd 603.06947). Anal. Calcd for THP-2 ($\text{C}_{22}\text{H}_{23}\text{F}_3\text{N}_3\text{O}_8\text{PReS}$): C, 36.02; H, 3.16; N, 3.82. Found: C, 36.03; H, 2.90; N, 3.77.

Synthesis of $[\text{Re}(\text{CO})_3(\text{dmphen})(\text{DAPTA})]\text{OTf}$ (DAPTA-2). A mixture of $[\text{Re}(\text{CO})_3(\text{dmphen})\text{Cl}]$ (0.232 g, 0.451 mmol) and AgOTf (0.116 g, 0.51 mmol) in THF (54 mL) was heated to reflux at 75 °C for 3 h in the dark. The white solid AgCl was removed via vacuum filtration. To the remaining yellow filtrate was added DAPTA (0.144 g, 0.59 mmol), and the solution was heated to reflux for an additional 15 h. The reaction mixture was cooled to room temperature, and then the yellow precipitate was isolated by filtration and washed with a small amount of methanol (5 mL) and diethyl ether (20 mL). Yield: 0.319 g (83%). ^1H NMR (400 MHz, $\text{MeOD}-d_4$): DAPTA complexes show the syn and anti isomers in a 1/5 ratio; however in the aromatic region the isomers are indistinguishable, δ 8.85 (d, 2H, anti + syn), 8.26 (d, 2H, anti + syn), 8.16 (dd, 2H, anti + syn), 5.51 (d, anti), 4.98 (d, syn), 4.44 (d, anti), 3.93 (d, anti), 3.55 (m, anti), 3.37 (d, 3H, anti + syn), 2.98 (m, syn), 2.84 (d, anti), 1.90 (s, syn), 1.84 (s, anti), 1.46 (s, 5H, anti). ^{19}F NMR (376 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -79.57. $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -41.15 (syn), -45.58 (anti). IR (KBr, cm^{-1}): 3440 m, 3070 w, 2040 s, 1970 m, 1930 s, 1640 s, 1510 w, 1420 m, 1380 w, 1340 m, 1270 s, 1150 m, 1030 s, 989 m, 890 m, 867 m, 799 m, 705 w, 638 s, 551 m, 513 m, 469 w, 417 w. ESI-MS (positive ion mode, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ 70/30 and 1% formic acid): m/z 708.13962 ($[\text{M}]^+$, calcd 708.13801). Anal. Calcd for DAPTA-2 ($\text{C}_{27}\text{H}_{28}\text{F}_3\text{N}_3\text{O}_8\text{PReS}$): C, 37.85; H, 3.29; N, 8.17. Found: C, 38.02; H, 3.24; N, 8.09.

Synthesis of $[\text{Re}(\text{CO})_3(\text{bpy})(\text{PTA})]\text{OTf}$ (PTA-3). A mixture of $[\text{Re}(\text{CO})_3(\text{bpy})\text{Cl}]$ (0.564 g, 1.22 mmol) and AgOTf (0.314 g, 1.22 mmol) in THF (123 mL) was heated to reflux at 75 °C for 3 h in the dark. The white solid, AgCl , was removed via vacuum filtration. To the remaining yellow filtrate was added PTA (0.192 g, 1.22 mmol), and

the solution was heated to reflux for an additional 15 h. The reaction mixture was cooled to room temperature, and then the THF was removed using rotary evaporation. The remaining yellow solid was dissolved in a minimum amount of methanol (5–10 mL) and filtered through Celite. The addition of diethyl ether (20–30 mL) afforded the desired product as a yellow solid, which was isolated by vacuum filtration and dried in vacuo. Yield: 0.523 g (59%). ^1H NMR (400 MHz, $\text{MeOD}-d_4$): δ 9.13 (d, 2H, J = 5.6 Hz), 8.74 (d, 2H, J = 8.3 Hz), 8.40 (t, 2H, J = 8.3 Hz), 7.81 (t, 2H, J = 5.6 Hz), 4.47 (d, 3H, J = 13.0 Hz), 4.38 (d, 3H, J = 13.0 Hz), 3.80 (s, 6H). ^{19}F NMR (376 MHz, $\text{MeOD}-d_4$): δ -79.52. $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -77.64. IR (KBr, cm^{-1}): 3440 m, 2040 s, 1950 s, 1910 s, 1610 m, 1470 m, 1440 m, 1280 s, 1260 s, 1140 m, 1100 m, 1030 s, 1010 m, 969 m, 946 m, 903 w, 801 m, 770 s, 740 m, 631 s, 614 m, 587 m, 573 m, 512 m, 454 m. ESI-MS (positive ion mode, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ 70/30 and 1% formic acid): m/z 584.08746 ($[\text{M}]^+$, calcd 584.08558). Anal. Calcd for PTA-3 ($\text{C}_{20}\text{H}_{20}\text{F}_3\text{N}_3\text{O}_8\text{PReS}$): C, 32.79; H, 2.75; N, 9.56. Found: C, 32.51; H, 2.76; N, 9.38.

Synthesis of $[\text{Re}(\text{CO})_3(\text{bpy})(\text{THP})]\text{TFA}$ (THP-3). A mixture of $[\text{Re}(\text{CO})_3(\text{bpy})\text{Cl}]$ (0.100 g, 0.217 mmol) and AgOTf (0.056 g, 0.217 mmol) in THF (22 mL) was heated to reflux at 75 °C for 3 h in the dark. The white solid, AgCl , was removed via vacuum filtration. To the remaining yellow filtrate was added THP (0.040 g, 0.326 mmol), and the solution was heated to reflux for an additional 15 h. The reaction mixture was cooled to room temperature, and then THF was removed by rotary evaporation. The remaining yellow oil was then dissolved in a minimal amount of methanol (5 mL) and filtered through Celite. The crude yellow oil (0.170 g) was purified using preparatory RP-HPLC using a $\text{CH}_3\text{OH}/\text{H}_2\text{O}$ gradient (0–5 min, 10% CH_3OH ; 5–30 min, 10–100% CH_3OH). Purified yield: 0.070 g (45%). ^1H NMR (400 MHz, $\text{MeOD}-d_4$): δ 9.17 (d, 2H, J = 5.8 Hz), 8.64 (d, 2H, J = 8.1 Hz), 8.31 (t, 2H, J = 8.1 Hz), 7.72 (t, 2H, J = 6.4 Hz), 3.90 (s, 6H). ^{19}F NMR (376 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -79.34. $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ 2.50. IR (KBr, cm^{-1}): 3433 s, 2874 m, 2038 s, 1930 s, 1915 s, 1681 s, 1633 s, 1473 m, 1450 m, 1181 s, 1030 m, 841 w, 769 m, 731 w, 534 w, 497 w, 417 m. ESI-MS (positive ion mode, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ 70/30 and 1% formic acid): m/z 551.03927 ($[\text{M}]^+$, calcd 551.03763). Anal. Calcd for THP-3 ($\text{C}_{18}\text{H}_{19}\text{F}_3\text{N}_3\text{O}_8\text{PReS}$): C, 31.72; H, 2.81; N, 4.11. Found: C, 31.68; H, 2.61; N, 4.09.

Synthesis of $[\text{Re}(\text{CO})_3(\text{bpy})(\text{DAPTA})]\text{OTf}$ (DAPTA-3). A mixture of $[\text{Re}(\text{CO})_3(\text{bpy})\text{Cl}]$ (0.236 g, 0.51 mmol) and AgOTf (0.131 g, 0.51 mmol) in THF (55 mL) was heated to reflux at 75 °C for 3 h in the dark. The white solid precipitate, AgCl , was removed via vacuum filtration. To the remaining yellow filtrate was added DAPTA (0.150 g, 0.61 mmol), and the solution was heated to reflux for an additional 24 h. The reaction mixture was cooled to room temperature, and then the THF was removed using rotary evaporation. The remaining yellow solid was dissolved in a minimum amount of methanol (5–10 mL) and filtered through Celite. The addition of diethyl ether (20–30 mL) afforded the desired product as a yellow solid, which was isolated by vacuum filtration and dried in vacuo. Yield: 0.204 g (48%). ^1H NMR (400 MHz, $\text{MeOD}-d_4$): DAPTA complexes show the syn and anti isomers in a 1/5 ratio; however in the aromatic region the isomers are indistinguishable, δ 9.21 (t, 2H, anti + syn), 8.77 (d, 2H, anti + syn), 8.43 (t, 2H, anti + syn), 7.84 (t, 2H, anti + syn), 5.65 (d, anti), 5.12 (d, syn), 5.01 (d, anti), 4.92 (m, syn), 4.53 (d, anti), 4.22 (m, anti), 4.03 (d, anti), 3.91 (d, anti), 3.80 (s, syn), 3.58 (d, anti), 3.14 (d, anti), 2.01 (s, syn), 1.98 (s, anti), 1.90 (s, anti). ^{19}F NMR (376 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -79.56. $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -44.42 (syn), -48.33 (anti). IR (KBr, cm^{-1}): 3457 m, 2965 w, 2880 w, 2037 s, 1958 s, 1918 s, 1646 s, 1473 m, 1419 m, 1331 m, 1265 s, 1155 m, 1024 s, 987 m, 892 m, 776 m, 632 s, 515 m, 422 m. ESI-MS (positive ion mode, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ 70/30 and 1% formic acid): m/z 656.10825 ($[\text{M}]^+$, calcd 656.10671). Anal. Calcd for DAPTA-3-0.5(CH_3CH_2)₂O ($\text{C}_{25}\text{H}_{29}\text{F}_3\text{N}_3\text{O}_{8.5}\text{PReS}$): C, 35.67; H, 3.47; N, 8.32. Found: C, 35.17; H, 3.49; N, 8.38.

Synthesis of $[\text{Re}(\text{CO})_3(\text{dmbpy})(\text{PTA})]\text{OTf}$ (PTA-4). A mixture of $[\text{Re}(\text{CO})_3(\text{dmbpy})\text{Cl}]$ (0.200 g, 0.43 mmol) and AgOTf (0.111 g, 0.43 mmol) in THF (47 mL) was heated to reflux at 75 °C for 3 h in

the dark. The white solid precipitate, AgCl, was removed via vacuum filtration. To the remaining yellow filtrate was added PTA (0.102 g, 0.65 mmol), and the solution was heated to reflux for an additional 15 h. The reaction mixture was cooled to room temperature, and then the THF was removed using rotary evaporation. The remaining yellow solid was dissolved in a minimum amount of methanol (5–10 mL) and filtered through Celite. The addition of diethyl ether (20–30 mL) afforded the desired product as a yellow solid, which was isolated by vacuum filtration and dried in vacuo. Yield: 0.254 g (77%). ^1H NMR (400 MHz, $\text{MeOD}-d_4$): δ 8.91 (d, 2H, J = 5.8 Hz), 8.60 (s, 2H), 7.62 (d, 2H, J = 5.8 Hz), 4.47 (d, 3H, J = 13.2 Hz), 4.37 (d, 3H, J = 13.2 Hz), 3.78 (s, 6H), 2.67 (s, 6H). ^{19}F NMR (376 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -79.53. $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -77.22. IR (KBr, cm^{-1}): 3452 m, 2940 w, 2033 s, 1951 s, 1921 s, 1626 w, 1266 s, 1139 w, 1031 m, 972 m, 948 m, 834 m, 801 m, 740 m, 637 s, 580 m, 500 m, 436 m. ESI-MS (positive ion mode, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ 70/30 and 1% formic acid): m/z 612.11845 ($[\text{M}]^+$, calcd 612.11743). Anal. Calcd for **PTA-4** ($\text{C}_{22}\text{H}_{24}\text{F}_3\text{N}_5\text{O}_6\text{PReS}$): C, 34.74; H, 3.18; N, 9.21. Found: C, 34.97; H, 3.22; N, 9.11.

Synthesis of $[\text{Re}(\text{CO})_3(\text{dmbpy})(\text{THP})]\text{TfA}$ (THP-4). A mixture of $[\text{Re}(\text{CO})_3(\text{dmbpy})\text{Cl}]$ (0.200 g, 0.43 mmol) and AgOTf (0.111 g, 0.43 mmol) in THF (47 mL) was heated to reflux at 75 °C for 3 h in the dark. The white solid precipitate, AgCl, was removed via vacuum filtration. To the remaining yellow filtrate was added THP (0.081 g, 0.65 mmol), and the solution was heated to reflux for an additional 15 h. The reaction mixture was cooled to room temperature, and then THF was removed by rotary evaporation. The crude yellow solid (0.307 g) was purified using preparatory RP-HPLC using a $\text{CH}_3\text{OH}/\text{H}_2\text{O}$ gradient (0–5 min, 10% CH_3OH ; 5–30 min, 10–100% CH_3OH). Purified yield: 0.073 g (25%). ^1H NMR (400 MHz, $\text{MeOD}-d_4$): δ 8.96 (d, 2H, J = 5.7 Hz), 8.50 (s, 2H), 7.54 (d, 2H, J = 5.7 Hz), 3.88 (s, 6H), 2.63 (s, 6H). ^{19}F NMR (376 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -76.36. $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ 3.26. IR (KBr, cm^{-1}): 3404 m, 2900 w, 2031 s, 1935 s, 1920 s, 1675 m, 1621 w, 1440 w, 1200 m, 1130 m, 1035 m, 905 w, 830 m, 723 w, 637 w, 534 w, 508 m, 444 w. ESI-MS (positive ion mode, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ 70/30 and 1% formic acid): m/z 579.07103 ($[\text{M}]^+$, calcd 579.06893). Anal. Calcd for **THP-4** ($\text{C}_{20}\text{H}_{21}\text{F}_3\text{N}_5\text{O}_8\text{PReS}$): C, 32.79; H, 2.75; N, 9.56. Found: C, 32.51; H, 2.76; N, 9.38.

Synthesis of $[\text{Re}(\text{CO})_3(\text{dmbpy})(\text{DAPTA})]\text{OTf}$ (DAPTA-4). A mixture of $[\text{Re}(\text{CO})_3(\text{dmbpy})\text{Cl}]$ (0.267 g, 0.57 mmol) and AgOTf (0.146 g, 0.57 mmol) in THF (65 mL) was heated to reflux at 75 °C for 3 h in the dark. The white solid precipitate, AgCl, was removed via vacuum filtration. To the remaining yellow filtrate was added DAPTA (0.211 g, 0.86 mmol), and the solution was heated to reflux for an additional 15 h. The reaction mixture was cooled to room temperature, and then the THF was removed using rotary evaporation. The remaining yellow solid was dissolved in a minimum amount of methanol (5–10 mL) and filtered through Celite. The addition of diethyl ether (20–30 mL) afforded the desired product as a yellow solid, which was isolated by vacuum filtration and dried in vacuo. Yield: 0.194 g (39%). ^1H NMR (400 MHz, $\text{MeOD}-d_4$): DAPTA complexes show the syn and anti isomers in a 1/5 ratio; however in the aromatic region the isomers are indistinguishable, δ 8.99 (t, 2H, anti + syn), 8.63 (s, 2H, anti + syn), 7.65 (t, 2H, anti + syn), 5.64 (d, anti), 5.13 (d, syn), 5.00 (d, anti), 4.54 (m, anti + syn), 4.22 (m, anti), 4.02 (d, anti), 3.89 (d, anti), 3.56 (m, syn), 3.12 (d, anti), 2.68 (s, 3H, anti + syn), 2.02 (s, syn), 1.98 (s, anti), 1.91 (s, anti). ^{19}F NMR (376 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -79.53. $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -44.35 (syn), -47.93 (anti). IR (KBr, cm^{-1}): 3433 m, 2038 s, 1943 s, 1924 s, 1648 m, 1420 m, 1333 w, 1262 m, 1156 w, 1030 m, 986 w, 890 w, 826 w, 782 w, 638 m, 500 w, 421 w. ESI-MS (positive ion mode, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ 70/30 and 1% formic acid): m/z 684.13967 ($[\text{M}]^+$, calcd 684.13801). Anal. Calcd for **DAPTA-4** $\cdot \text{CH}_3\text{OH}$ ($\text{C}_{26}\text{H}_{32}\text{F}_3\text{N}_5\text{O}_9\text{PReS}$): C, 36.11; H, 3.73; N, 8.10. Found: C, 36.31; H, 3.59; N, 8.19.

Synthesis of $[\text{Re}(\text{CO})_3(\text{dmbpy})(\text{PTA})]\text{OTf}$ (PTA-5). A mixture of $[\text{Re}(\text{CO})_3(\text{dmbpy})\text{Cl}]$ (0.200 g, 0.38 mmol) and AgOTf (0.099 g, 0.38 mmol) in THF (47 mL) was heated to reflux at 75 °C for 3 h in the dark. The white solid precipitate, AgCl, was removed via vacuum filtration. To the remaining yellow filtrate was added PTA (0.079 g, 0.50 mmol), and the solution was heated to reflux for an additional 15 h. The reaction mixture was cooled to room temperature, and then the THF was removed using rotary evaporation. The remaining yellow solid was dissolved in a minimum amount of methanol (5–10 mL) and filtered through Celite. The addition of diethyl ether (20–30 mL) afforded the desired product as a yellow solid, which was isolated by vacuum filtration and dried in vacuo. Yield: 0.158 g (52%). ^1H NMR (400 MHz, $\text{MeOD}-d_4$): δ 8.84 (d, 2H, J = 6.5 Hz), 8.26 (d, 2H, J = 2.6 Hz), 7.34 (dd, 2H, J = 2.6, 6.5 Hz), 4.49 (d, 3H, J = 13.1 Hz), 4.39 (d, 3H, J = 13.1 Hz), 4.14 (s, 6H), 3.80 (s, 6H). ^{19}F NMR (376 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -79.59. $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -75.88. IR (KBr, cm^{-1}): 3443 m, 2031 s, 1942 s, 1921 s, 1615 m, 1560 w, 1497 w, 1426 w, 1350 w, 1262 m, 1242 w, 1147 w, 1029 m, 974 w, 948 w, 899 w, 828 w, 801 w, 741 w, 636 m, 581 w, 514 w, 429 w. ESI-MS (positive ion mode, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ 70/30 and 1% formic acid): m/z 644.10822 ($[\text{M}]^+$, calcd 644.10671). Anal. Calcd for **PTA-5** ($\text{C}_{22}\text{H}_{24}\text{F}_3\text{N}_5\text{O}_8\text{PReS}$): C, 33.33; H, 3.05; N, 8.84. Found: C, 33.39; H, 3.10; N, 8.80.

Synthesis of $[\text{Re}(\text{CO})_3(\text{dmbpy})(\text{THP})]\text{OTf}$ (THP-5). A mixture of $[\text{Re}(\text{CO})_3(\text{dmbpy})\text{Cl}]$ (0.297 g, 0.57 mmol) and AgOTf (0.146 g, 0.57 mmol) in THF (70 mL) was heated to reflux at 75 °C for 3 h in the dark. The white solid precipitate, AgCl, was removed via vacuum filtration. To the remaining yellow filtrate was added THP (0.092 g, 0.74 mmol), and the solution was heated to reflux for an additional 15 h. The reaction mixture was cooled to room temperature, and then THF was removed by rotary evaporation, and the residue was then dissolved in minimal methanol (5 mL) and filtered through Celite. The crude yellow oil (0.439 g) was recrystallized using vapor diffusion of diethyl ether into a solution of methanol to afford yellow crystals. Yield: 0.030 g (7%). ^1H NMR (400 MHz, $\text{MeOD}-d_4$): δ 8.88 (d, 2H, J = 6.6 Hz), 8.16 (d, 2H, J = 2.7 Hz), 7.27 (dd, 2H, J = 2.7, 6.6 Hz), 4.10 (s, 6H), 3.90 (s, 6H). ^{19}F NMR (376 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -79.57. $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ 3.16. IR (KBr, cm^{-1}): 3448 m, 2028 s, 1939 w, 1919 s, 1618 s, 1560 m, 1500 m, 1343 w, 1270 m, 1182 w, 1031 w, 905 w, 826 w, 639 w, 513 w, 434 w. ESI-MS (positive ion mode, $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ 70/30 and 1% formic acid): m/z 611.06037 ($[\text{M}]^+$, calcd 611.05875). Anal. Calcd for **THP-5** ($\text{C}_{19}\text{H}_{21}\text{F}_3\text{N}_5\text{O}_{11}\text{PReS}$): C, 30.04; H, 2.79; N, 3.69. Found: C, 30.30; H, 2.79; N, 3.89.

Synthesis of $[\text{Re}(\text{CO})_3(\text{dmbpy})(\text{DAPTA})]\text{OTf}$ (DAPTA-5). A mixture of $[\text{Re}(\text{CO})_3(\text{dmbpy})\text{Cl}]$ (0.210 g, 0.40 mmol) and AgOTf (0.103 g, 0.40 mmol) in THF (45 mL) was heated to reflux at 75 °C for 3 h in the dark. The white solid precipitate, AgCl, was removed via vacuum filtration. To the remaining yellow filtrate was added DAPTA (0.123 g, 0.52 mmol), and the solution was heated to reflux for an additional 15 h. The reaction mixture was cooled to room temperature, and then the THF was removed using rotary evaporation. The remaining yellow solid was dissolved in a minimum amount of methanol (5–10 mL) and filtered through Celite. The addition of diethyl ether (20–30 mL) afforded the desired product as a yellow solid, which was isolated by vacuum filtration and dried in vacuo. The solid material (0.343 g) was recrystallized using vapor diffusion of diethyl ether into a solution of methanol to afford pure yellow crystals. Purified yield: 0.028 g (8%). ^1H NMR (400 MHz, $\text{MeOD}-d_4$): DAPTA complexes show the syn and anti isomers in a 1/5 ratio; however in the aromatic region the isomers are indistinguishable, δ 8.91 (t, 2H, anti + syn), 8.29 (s, 2H, anti + syn), 7.37 (t, 2H, anti + syn), 5.67 (d, anti), 5.15 (d, syn), 5.02 (d, anti), 4.56 (m, anti + syn), 4.22 (m, anti), 4.15 (s, 3H, anti + syn), 4.05 (d, anti), 3.92 (d, anti), 3.54 (m, anti + syn), 3.14 (d, anti + syn), 2.03 (s, syn), 1.99 (s, anti), 1.92 (s, anti). ^{19}F NMR (376 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -79.57. $^{31}\text{P}\{^1\text{H}\}$ NMR (162 MHz, $\text{MeOD}-d_4$, external standard KPF_6): δ -43.34 (syn), -46.87 (anti). IR (KBr, cm^{-1}): 3456 m, 2040 s, 1939 s, 1919 s, 1637 s, 1500 m, 1420 m, 1340 w, 1260 m, 1150 w, 1030 s, 889 w, 796 w, 638 m, 516 w, 419 m. ESI-MS (positive

ion mode, CH₃CN/H₂O 70/30 and 1% formic acid): *m/z* 716.12966 ([M]⁺, calcd 716.12784). Anal. Calcd for DAPTA-5 (C₂₅H₂₈F₃N₅O₁₀PreS): C, 34.72; H, 3.26; N, 8.10. Found: C, 34.44; H, 3.50; N, 8.08.

Synthesis of [Re(CO)₂(phen)(DAPTA)Cl] (DAPTA-1A). Trime-thylamine *N*-oxide (TMAO, 0.040 g, 0.36 mmol) was dissolved in a mixture of CH₂Cl₂ (130 mL) and CH₃OH (5 mL). A solution of DAPTA-1 (0.262 g, 0.32 mmol) and NEt₄Cl (0.919 g, 5 mmol) in CH₂Cl₂ (105 mL) was heated to reflux at 50 °C, and the TMAO solution was added dropwise over the course of 1 h, changing the color of the solution from light yellow to dark red. The solution was heated under reflux for an additional 24 h. The CH₂Cl₂/CH₃OH was then removed via rotary evaporation, and the remaining red solid was suspended in approximately 20 mL of CH₃OH. The solid was isolated by centrifugation and then resuspended in 5–10 mL of CH₃OH. This process was repeated three times, yielding the product DAPTA-1A as a red solid. Yield: 0.150 g (94%). ¹H NMR (400 MHz, DMF-*d*₇): DAPTA complexes show the syn and anti isomers in a 1/5 ratio; however in the aromatic region the isomers are indistinguishable, δ 9.52 (m, 2H, anti + syn), 8.95 (d, 2H, anti + syn), 8.33 (s, 2H, anti + syn), 8.14 (m, 2H, anti + syn), 5.52 (d, anti), 5.06 (d, anti), 4.89 (d, syn), 4.47 (d, anti), 4.21 (m, syn), 3.93 (d, syn), 3.73 (d, anti), 3.32 (m, anti + syn), 3.12 (d, anti), 1.97 (s, syn), 1.90 (s, anti), 1.86 (s, anti). ³¹P{¹H} NMR (162 MHz, DMF-*d*₇, external standard KPF₆): δ −32.32 (syn), −34.72 (anti). IR (KBr, cm^{−1}): 3458 s, 1927s, 1839 s, 1638 s, 1427 m, 1338 m, 1240 m, 1097 w, 1050 w, 997 w, 890 w, 855 w, 795 w, 727 w, 614 w. Anal. Calcd for DAPTA-1A·2H₂O (C₂₃H₂₈ClN₅O₆Pre): C, 38.20; H, 3.90; N, 9.68. Found: C, 38.30; H, 3.94; N, 9.42.

Synthesis of [Re(CO)₂(bpy)(PTA)Cl] (PTA-3A). TMAO (0.052 g, 0.47 mmol) was dissolved in a mixture of CH₂Cl₂ (150 mL) and CH₃OH (6 mL). A mixture of PTA-3 (0.298 g, 0.41 mmol) and NEt₄Cl (1.17 g, 6.4 mmol) in CH₂Cl₂ (120 mL) was heated to reflux at 50 °C, and then the TMAO solution was added dropwise over the course of 1 h, resulting in a color change from yellow to red. The solution was heated to reflux for an additional 24 h. The CH₂Cl₂/CH₃OH was removed via rotary evaporation. To the remaining red solid was added approximately 20 mL of water. The solid was isolated by centrifugation and then resuspended in 5–10 mL of water. This process was repeated 3× to obtain the product PTA-3A as a red solid. Yield: 0.112 g (46%). ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.88 (d, 2H, *J* = 5.9 Hz), 8.67 (d, 2H, *J* = 8.3 Hz), 8.22 (t, 2H, *J* = 8.3 Hz), 7.65 (t, 2H, *J* = 5.9 Hz), 4.28 (d, 3H, *J* = 12.8 Hz), 4.15 (d, 3H, *J* = 12.8 Hz), 3.49 (s, 6H). ³¹P{¹H} NMR (162 MHz, DMSO-*d*₆, external standard KPF₆): δ −59.1. IR (KBr, cm^{−1}): 3471 m, 1912 s, 1834 s, 1605 w, 1452 w, 1239 w, 1099 w, 1014 w, 966 w, 948 w, 781 m, 594 w, 580 m, 483 w. Anal. Calcd for PTA-3A·H₂O (C₁₈H₂₂ClN₂O₃Pre): C, 35.56; H, 3.64; N, 11.50. Found: C, 35.38; H, 3.66; N, 11.36.

■ ASSOCIATED CONTENT

● Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.inorgchem.7b02747.

Complex characterization data, cell viability curves, crystal data tables, DFT-optimized geometries and excited states, UV–vis and emission spectra, transient emission decay profiles, and species distribution diagrams (PDF)

Accession Codes

CCDC 1581545–1581560 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail for J.J.W.: jjw275@cornell.edu.

ORCID

Zhi Li: 0000-0002-8799-0283

Peter C. Ford: 0000-0002-5509-9912

Justin J. Wilson: 0000-0002-4086-7982

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This research was supported by Cornell University and by the Office of the Assistant Secretary of Defense for Health Affairs through the Ovarian Cancer Research Program under award no. W81XWH-17-1-0097. This work made use of the NMR facility at Cornell University, which is supported, in part, by the NSF under award no. CHE-1531632. Studies at UCSB were supported by the National Science Foundation under award no. CHE-1565702. Dr. Nikki Thiele (Cornell University) and Mr. Jacob Barrett (UCSB) are thanked for assistance with the p*K*_a measurements and the CO release quantification, respectively.

■ REFERENCES

- (1) Dolmans, D. E. J. G. J.; Fukumura, D.; Jain, R. K. Photodynamic Therapy for Cancer. *Nat. Rev. Cancer* **2003**, *3*, 380–387.
- (2) Castano, A. P.; Demidova, T. N.; Hamblin, M. R. Mechanisms in Photodynamic Therapy: Part One—Photosensitizers, Photochemistry and Cellular Localization. *Photodiagn. Photodyn. Ther.* **2004**, *1*, 279–293.
- (3) Castano, A. P.; Mroz, P.; Hamblin, M. R. Photodynamic Therapy and Anti-Tumour Immunity. *Nat. Rev. Cancer* **2006**, *6*, 535–545.
- (4) Agostinis, P.; Berg, K.; Cengel, K. A.; Foster, T. H.; Girotti, A. W.; Gollnick, S. O.; Hahn, S. M.; Hamblin, M. R.; Juzeniene, A.; Kessel, D.; Korbelik, M.; Moan, J.; Mroz, P.; Nowis, D.; Piette, J.; Wilson, B. C.; Golab, J. Photodynamic Therapy of Cancer: An Update. *CA-Cancer J. Clin.* **2011**, *61*, 250–281.
- (5) Farrer, N. J.; Salassa, L.; Sadler, P. J. Photoactivated Chemotherapy (PACT): The Potential of Excited-State d-Block Metals in Medicine. *Dalton Trans.* **2009**, 10690–10701.
- (6) Taub, A. F. Photodynamic Therapy: Other Uses. *Dermatol. Clin.* **2007**, *25*, 101–109.
- (7) DeRosa, M. C.; Crutchley, R. J. Photosensitized Singlet Oxygen and Its Applications. *Coord. Chem. Rev.* **2002**, *233–234*, 351–371.
- (8) O'Connor, A. E.; Gallagher, W. M.; Byrne, A. T. Porphyrin and Nonporphyrin Photosensitizers in Oncology: Preclinical and Clinical Advances in Photodynamic Therapy. *Photochem. Photobiol.* **2009**, *85*, 1053–1074.
- (9) Yang, J. Z.; Van Vugt, D. A.; Kennedy, J. C.; Reid, R. L. Intrauterine 5-Aminolevulinic Acid Induces Selective Fluorescence and Photodynamic Ablation of the Rat Endometrium. *Photochem. Photobiol.* **1993**, *57*, 803–807.
- (10) Bonnett, R. Photosensitizers of the Porphyrin and Phthalocyanine Series for Photodynamic Therapy. *Chem. Soc. Rev.* **1995**, *24*, 19–33.
- (11) Ochsner, M. Photophysical and Photobiological Processes in the Photodynamic Therapy of Tumours. *J. Photochem. Photobiol., B* **1997**, *39*, 1–18.
- (12) Dougherty, T. J.; Gomer, C. J.; Henderson, B. W.; Jori, G.; Kessel, D.; Korbelik, M.; Moan, J.; Peng, Q. Photodynamic Therapy. *J. Natl. Cancer Inst.* **1998**, *90*, 889–905.
- (13) Sharman, W. M.; Allen, C. M.; van Lier, J. E. Photodynamic Therapeutics: Basic Principles and Clinical Applications. *Drug Discovery Today* **1999**, *4*, 507–517.

- (14) MacDonald, I. J.; Dougherty, T. J. Basic Principles of Photodynamic Therapy. *J. Porphyrins Phthalocyanines* **2001**, *5*, 105–129.
- (15) Pervaiz, S.; Olivo, M. Art and Science of Photodynamic Therapy. *Clin. Exp. Pharmacol. Physiol.* **2006**, *33*, 551–556.
- (16) Palumbo, G. Photodynamic Therapy and Cancer: A Brief Sightseeing Tour. *Expert Opin. Drug Delivery* **2007**, *4*, 131–148.
- (17) Juarranz, Á.; Jaén, P.; Sanz-Rodríguez, F.; Cuevas, J.; González, S. Photodynamic Therapy of Cancer. Basic Principles and Applications. *Clin. Transl. Oncol.* **2008**, *10*, 148–154.
- (18) Bhuvaneswari, R.; Gan, Y. Y.; Soo, K. C.; Olivo, M. The Effect of Photodynamic Therapy on Tumor Angiogenesis. *Cell. Mol. Life Sci.* **2009**, *66*, 2275–2283.
- (19) Plaetzer, K.; Krammer, B.; Berlanda, J.; Berr, F.; Kiesslich, T. Photophysics and Photochemistry of Photodynamic Therapy: Fundamental Aspects. *Lasers Med. Sci.* **2009**, *24*, 259–268.
- (20) Harris, A. L. Hypoxia—A Key Regulatory Factor in Tumour Growth. *Nat. Rev. Cancer* **2002**, *2*, 38–47.
- (21) Brown, J. M. Tumor Hypoxia in Cancer Therapy. *Methods Enzymol.* **2007**, *435*, 296–321.
- (22) Howerton, B. S.; Heidary, D. K.; Glazer, E. C. Strained Ruthenium Complexes Are Potent Light-Activated Anticancer Agents. *J. Am. Chem. Soc.* **2012**, *134*, 8324–8327.
- (23) Bozzini, G.; Colin, P.; Betrouni, N.; Maurage, C. A.; Leroy, X.; Simonin, S.; Martin-Schmitt, C.; Villers, A.; Mordon, S. Efficiency of 5-ALA Mediated Photodynamic Therapy on Hypoxic Prostate Cancer: A Preclinical Study on the Dunning R3327-AT2 Rat Tumor Model. *Photodiagn. Photodyn. Ther.* **2013**, *10*, 296–303.
- (24) Kinsella, T. J.; Colussi, V. C.; Oleinick, N. L.; Sibata, C. H. Photodynamic Therapy in Oncology. *Expert Opin. Pharmacother.* **2001**, *2*, 917–927.
- (25) Baas, P.; van Mansom, I.; van Tinteren, H.; Stewart, F. A.; van Zandwijk, N. Effect of N-Acetylcysteine on Photofrin-Induced Skin Photosensitivity in Patients. *Lasers Surg. Med.* **1995**, *16*, 359–367.
- (26) Farrer, N. J.; Sadler, P. J. Photochemotherapy: Targeted Activation of Metal Anticancer Complexes. *Aust. J. Chem.* **2008**, *61*, 669–674.
- (27) Josefsen, L. B.; Boyle, R. W. Photodynamic Therapy and the Development of Metal-Based Photosensitizers. *Met.-Based Drugs* **2008**, *2008*, 1–23.
- (28) Lameijer, L. N.; Ernst, D.; Hopkins, S. L.; Meijer, M. S.; Askes, S. H. C.; Le Dévédec, S. E.; Bonnet, S. A Red-Light-Activated Ruthenium-Caged NAMPT Inhibitor Remains Phototoxic in Hypoxic Cancer Cells. *Angew. Chem., Int. Ed.* **2017**, *56*, 11549–11553.
- (29) Charlesworth, P.; Truscott, T. G.; Brooks, R. C.; Wilson, B. C. The Photophysical Properties of a Ruthenium-Substituted Phthalocyanine. *J. Photochem. Photobiol., B* **1994**, *26*, 277–282.
- (30) Zhou, J.; Liu, J.; Feng, Y.; Wei, S.; Gu, X.; Wang, X.; Zhang, B. Synthesis and Characterization of the Monomer Ruthenium Complex of Hypocrellin B. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 3067–3070.
- (31) Davia, K.; King, D.; Hong, Y.; Swavey, S. A Porphyrin-Ruthenium Photosensitizer as a Potential Photodynamic Therapy Agent. *Inorg. Chem. Commun.* **2008**, *11*, 584–586.
- (32) Schmitt, F.; Govindaswamy, P.; Süß-Fink, G.; Ang, W. H.; Dyson, P. J.; Juillerat-Jeanneret, L.; Therrien, B. Ruthenium Porphyrin Compounds for Photodynamic Therapy of Cancer. *J. Med. Chem.* **2008**, *51*, 1811–1816.
- (33) Gianferrara, T.; Bergamo, A.; Bratsos, I.; Milani, B.; Spagnul, C.; Sava, G.; Alessio, E. Ruthenium-Porphyrin Conjugates with Cytotoxic and Phototoxic Antitumor Activity. *J. Med. Chem.* **2010**, *53*, 4678–4690.
- (34) Sun, Y.; Joyce, L. E.; Dickson, N. M.; Turro, C. Efficient DNA Photocleavage by $[\text{Ru}(\text{bpy})_2(\text{dppn})]^{2+}$ with Visible Light. *Chem. Commun.* **2010**, *46*, 2426–2428.
- (35) Carneiro, Z. A.; de Moraes, J. C. B.; Rodrigues, F. P.; de Lima, R. G.; Curti, C.; da Rocha, Z. N.; Paulo, M.; Bendhack, L. M.; Tedesco, A. C.; Formiga, A. L. B.; da Silva, R. S. Photocytotoxic Activity of a Nitrosyl Phthalocyanine Ruthenium Complex – A System Capable of Producing Nitric Oxide and Singlet Oxygen. *J. Inorg. Biochem.* **2011**, *105*, 1035–1043.
- (36) Johnpeter, J. P.; Schmitt, F.; Denoyelle-Di-Muro, E.; Wagnières, G.; Juillerat-Jeanneret, L.; Therrien, B. Photoactive Sawhorse-Type Diruthenium Tetracarbonyl Complexes. *Inorg. Chim. Acta* **2012**, *393*, 246–251.
- (37) Pernot, M.; Bastogne, T.; Barry, N. P. E.; Therrien, B.; Koellensperger, G.; Hann, S.; Reshetov, V.; Barberi-Heyob, M. Systems Biology Approach for *in vivo* Photodynamic Therapy Optimization of Ruthenium-Porphyrin Compounds. *J. Photochem. Photobiol., B* **2012**, *117*, 80–89.
- (38) Zhang, J.-X.; Zhou, J.-W.; Chan, C.-F.; Lau, T. C.-K.; Kwong, D. W. J.; Tam, H.-L.; Mak, N.-K.; Wong, K.-L.; Wong, W.-K. Comparative Studies of the Cellular Uptake, Subcellular Localization, and Cytotoxic and Phototoxic Antitumor Properties of Ruthenium-(II)–Porphyrin Conjugates with Different Linkers. *Bioconjugate Chem.* **2012**, *23*, 1623–1638.
- (39) Lincoln, R.; Kohler, L.; Monro, S.; Yin, H.; Stephenson, M.; Zong, R.; Chouai, A.; Dorsey, C.; Hennigar, R.; Thummel, R. P.; McFarland, S. A. Exploitation of Long-Lived ^3IL Excited States for Metal–Organic Photodynamic Therapy: Verification in a Metastatic Melanoma Model. *J. Am. Chem. Soc.* **2013**, *135*, 17161–17175.
- (40) Albani, B. A.; Peña, B.; Leed, N. A.; de Paula, N. A. B. G.; Pavani, C.; Baptista, M. S.; Dunbar, K. R.; Turro, C. Marked Improvement in Photoinduced Cell Death by a New Tris-Heteroleptic Complex with Dual Action: Singlet Oxygen Sensitization and Ligand Dissociation. *J. Am. Chem. Soc.* **2014**, *136*, 17095–17101.
- (41) Frei, A.; Rubbiani, R.; Tubafard, S.; Blaque, O.; Anstaett, P.; Felgenträger, A.; Maisch, T.; Spiccia, L.; Gasser, G. Synthesis, Characterization, and Biological Evaluation of New Ru(II) Polypyridyl Photosensitizers for Photodynamic Therapy. *J. Med. Chem.* **2014**, *57*, 7280–7292.
- (42) Joshi, T.; Pierroz, V.; Mari, C.; Gemperle, L.; Ferrari, S.; Gasser, G. A Bis(dipyridophenazine)(2-(2-pyridyl)pyrimidine-4-carboxylic acid)ruthenium(II) Complex with Anticancer Action upon Photo-deprotection. *Angew. Chem., Int. Ed.* **2014**, *53*, 2960–2963.
- (43) Mari, C.; Pierroz, V.; Rubbiani, R.; Patra, M.; Hess, J.; Spingler, B.; Oehninger, L.; Schur, J.; Ott, I.; Salassa, L.; Ferrari, S.; Gasser, G. DNA Intercalating Ru^{II} Polypyridyl Complexes as Effective Photosensitizers in Photodynamic Therapy. *Chem. - Eur. J.* **2014**, *20*, 14421–14436.
- (44) Fong, J.; Kasimova, K.; Arenas, Y.; Kaspler, P.; Lazic, S.; Mandel, A.; Lilge, L. A Novel Class of Ruthenium-Based Photosensitizers Effectively Kills *in vitro* Cancer Cells and *in vivo* Tumors. *Photochem. Photobiol. Sci.* **2015**, *14*, 2014–2023.
- (45) Huang, H.; Yu, B.; Zhang, P.; Huang, J.; Chen, Y.; Gasser, G.; Ji, L.; Chao, H. Highly Charged Ruthenium(II) Polypyridyl Complexes as Lysosome-Localized Photosensitizers for Two-Photon Photodynamic Therapy. *Angew. Chem., Int. Ed.* **2015**, *54*, 14049–14052.
- (46) Shi, G.; Monro, S.; Hennigar, R.; Colpitts, J.; Fong, J.; Kasimova, K.; Yin, H.; DeCoste, R.; Spencer, C.; Chamberlain, L.; Mandel, A.; Lilge, L.; McFarland, S. A. Ru(II) Dyads Derived from α -Oligothiophenes: A New Class of Potent and Versatile Photosensitizers for PDT. *Coord. Chem. Rev.* **2015**, *282*–283, 127–138.
- (47) Kaspler, P.; Lazic, S.; Forward, S.; Arenas, Y.; Mandel, A.; Lilge, L. A Ruthenium(II) Based Photosensitizer and Transferrin Complexes Enhance Photo-Physical Properties, Cell Uptake, and Photodynamic Therapy Safety and Efficacy. *Photochem. Photobiol. Sci.* **2016**, *15*, 481–495.
- (48) Mari, C.; Huang, H.; Rubbiani, R.; Schulze, M.; Würthner, F.; Chao, H.; Gasser, G. Evaluation of Perylene Bisimide-Based Ru^{II} and Ir^{III} Complexes as Photosensitizers for Photodynamic Therapy. *Eur. J. Inorg. Chem.* **2017**, *2017*, 1745–1752.
- (49) Pierroz, V.; Rubbiani, R.; Gentili, C.; Patra, M.; Mari, C.; Gasser, G.; Ferrari, S. Dual Mode of Cell Death upon Photo-Irradiation of a Ru^{II} Polypyridyl Complex in Interphase or Mitosis. *Chem. Sci.* **2016**, *7*, 6115–6124.
- (50) Havrylyuk, D.; Heidary, D. K.; Nease, L.; Parkin, S.; Glazer, E. C. Photochemical Properties and Structure-Activity Relationships of

- Ru^{II} Complexes with Pyridylbenzazole Ligands as Promising Anticancer Agents. *Eur. J. Inorg. Chem.* **2017**, 2017, 1687–1694.
- (51) Mari, C.; Rubbiani, R.; Gasser, G. Biological Evaluation of Nitrile Containing Ru(II) Polypyridyl Complexes as Potential Photodynamic Therapy Agents. *Inorg. Chim. Acta* **2017**, 454, 21–26.
- (52) Wachter, E.; Heidary, D. K.; Howerton, B. S.; Parkin, S.; Glazer, E. C. Light-Activated Ruthenium Complexes Photobind DNA and are Cytotoxic in the Photodynamic Therapy Window. *Chem. Commun.* **2012**, 48, 9649–9651.
- (53) Scambellone, M. A.; David, A.; Garner, R. N.; Dunbar, K. R.; Turro, C. Cellular Toxicity Induced by the Photorelease of a Caged Bioactive Molecule: Design of a Potential Dual-Action Ru(II) Complex. *J. Am. Chem. Soc.* **2013**, 135, 11274–11282.
- (54) Hidayatullah, A. N.; Wachter, E.; Heidary, D. K.; Parkin, S.; Glazer, E. C. Photoactive Ru(II) Complexes with Dioxinophenanthroline Ligands Are Potent Cytotoxic Agents. *Inorg. Chem.* **2014**, 53, 10030–10032.
- (55) Mari, C.; Pierroz, V.; Leonidova, A.; Ferrari, S.; Gasser, G. Towards Selective Light-Activated Ru^{II}-Based Prodrug Candidates. *Eur. J. Inorg. Chem.* **2015**, 2015, 3879–3891.
- (56) Kumar, A.; Sun, S.-S.; Lees, A. J. Photophysics and Photochemistry of Organometallic Rhenium Diimine Complexes. In *Photophysics of Organometallics*; Lees, A. J., Ed.; Springer Berlin Heidelberg: Berlin, Heidelberg, 2010; pp 1–36.
- (57) Kastl, A.; Dieckmann, S.; Wähler, K.; Völker, T.; Kastl, L.; Merkel, A. L.; Vultur, A.; Shannan, B.; Harms, K.; Ocker, M.; Parak, W. J.; Herlyn, M.; Meggers, E. Rhenium Complexes with Visible-Light-Induced Anticancer Activity. *ChemMedChem* **2013**, 8, 924–927.
- (58) Leonidova, A.; Gasser, G. Underestimated Potential of Organometallic Rhenium Complexes as Anticancer Agents. *ACS Chem. Biol.* **2014**, 9, 2180–2193.
- (59) Leonidova, A.; Pierroz, V.; Rubbiani, R.; Heier, J.; Ferrari, S.; Gasser, G. Towards Cancer Cell-Specific Phototoxic Organometallic Rhenium(I) Complexes. *Dalton Trans.* **2014**, 43, 4287–4294.
- (60) Wähler, K.; Ludewig, A.; Szabo, P.; Harms, K.; Meggers, E. Rhenium Complexes with Red-Light-Induced Anticancer Activity. *Eur. J. Inorg. Chem.* **2014**, 2014, 807–811.
- (61) Quntal, L.; Raposinho, P.; Mendes, F.; Santos, I.; Navarro-Ranninger, C.; Alvarez-Valdes, A.; Huang, H.; Chao, H.; Rubbiani, R.; Gasser, G.; Quiroga, A. G.; Paulo, A. Combining Imaging and Anticancer Properties with New Heterobimetallic Pt(II)/M(I) (M = Re ^{99m}Tc) Complexes. *Dalton Trans.* **2017**, 46, 14523–14536.
- (62) Koike, K.; Okoshi, N.; Hori, H.; Takeuchi, K.; Ishitani, O.; Tsubaki, H.; Clark, I. P.; George, M. W.; Johnson, F. P. A.; Turner, J. J. Mechanism of the Photochemical Ligand Substitution Reactions of *fac*-[Re(bpy)(CO)₃(PR₃)]⁺ Complexes and the Properties of Their Triplet Ligand-Field Excited States. *J. Am. Chem. Soc.* **2002**, 124, 11448–11455.
- (63) Wrighton, M.; Morse, D. L. The Nature of the Lowest Excited State in Tricarbonylchloro-1,10-phenanthrolinerhenium(I) and Related Complexes. *J. Am. Chem. Soc.* **1974**, 96, 998–1003.
- (64) Caspar, J. V.; Meyer, T. J. Application of the Energy Gap Law to Nonradiative, Excited-State Decay. *J. Phys. Chem.* **1983**, 87, 952–957.
- (65) Kalyanasundaram, K. Luminescence and Redox Reactions of the Metal-to-Ligand Charge-Transfer Excited State of Tricarbonylchloro-(polypyridyl)rhenium(I) Complexes. *J. Chem. Soc., Faraday Trans. 2* **1986**, 82, 2401–2415.
- (66) Kalyanasundaram, K. Mononuclear Complexes of Re(I). In *Photochemistry of Polypyridine and Porphyrin Complexes*; Academic Press: London, 1992; pp 321–329.
- (67) Koike, K.; Tanabe, J.; Toyama, S.; Tsubaki, H.; Sakamoto, K.; Westwell, J. R.; Johnson, F. P. A.; Hori, H.; Saitoh, H.; Ishitani, O. New Synthetic Routes to Biscarbonylbipyridinerhenium(I) Complexes *cis,trans*-[Re(X₂bpy)(CO)₂(PR₃)(Y)]ⁿ⁺ (X₂bpy = 4,4'-X₂-2,2'-bipyridine) via Photochemical Ligand Substitution Reactions, and Their Photophysical and Electrochemical Properties. *Inorg. Chem.* **2000**, 39, 2777–2783.
- (68) Pierri, A. E.; Pallaoro, A.; Wu, G.; Ford, P. C. A Luminescent and Biocompatible PhotoCORM. *J. Am. Chem. Soc.* **2012**, 134, 18197–18200.
- (69) Kunz, P. C.; Huber, W.; Rojas, A.; Schatzschneider, U.; Spingler, B. Tricarbonylmanganese(I) and -Rhenium(I) Complexes of Imidazol-Based Phosphane Ligands: Influence of the Substitution Pattern on the CO Release Properties. *Eur. J. Inorg. Chem.* **2009**, 2009, 5358–5366.
- (70) Pfeiffer, H.; Rojas, A.; Niesel, J.; Schatzschneider, U. Sonogashira and “Click” Reactions for the N-Terminal and Side-Chain Functionalization of Peptides with [Mn(CO)₃(tpm)]⁺-Based CO Releasing Molecules (tpm = tris(pyrazolyl)methane). *Dalton Trans.* **2009**, 0, 4292–4298.
- (71) Rimmer, R. D.; Richter, H.; Ford, P. C. A Photochemical Precursor for Carbon Monoxide Release in Aerated Aqueous Media. *Inorg. Chem.* **2010**, 49, 1180–1185.
- (72) Brückmann, N. E.; Wahl, M.; Reiß, G. J.; Kohns, M.; Wätjen, W.; Kunz, P. C. Polymer Conjugates of Photoinducible CO-Releasing Molecules. *Eur. J. Inorg. Chem.* **2011**, 2011, 4571–4577.
- (73) Kretschmer, R.; Gessner, G.; Görls, H.; Heinemann, S. H.; Westerhausen, M. Dicarbonyl-bis(cysteamine)iron(II): A Light Induced Carbon Monoxide Releasing Molecule Based on Iron (CORM-S1). *J. Inorg. Biochem.* **2011**, 105, 6–9.
- (74) Schatzschneider, U. PhotoCORMs: Light-Triggered Release of Carbon Monoxide from the Coordination Sphere of Transition Metal Complexes for Biological Applications. *Inorg. Chim. Acta* **2011**, 374, 19–23.
- (75) Zobi, F.; Spingler, B.; Fox, T.; Alberto, R. Toward Novel DNA Binding Metal Complexes: Structure and Basic Kinetic Data of [M(9MeG)₂(CH₃OH)(CO)₃]⁺ (M = ⁹⁹Tc, Re). *Inorg. Chem.* **2003**, 42, 2818–2820.
- (76) Zobi, F.; Blacque, O.; Schmalte, H. W.; Spingler, B.; Alberto, R. Head-to-Head (HH) and Head-to-Tail (HT) Conformers of *cis*-Bis Guanine Ligands Bound to the [Re(CO)₃]⁺ Core. *Inorg. Chem.* **2004**, 43, 2087–2096.
- (77) Zobi, F.; Spingler, B.; Alberto, R. Guanine and Plasmid DNA Binding of Mono- and Trinuclear *fac*-[Re(CO)₃]⁺ Complexes with Amino Acid Ligands. *ChemBioChem* **2005**, 6, 1397–1405.
- (78) Knopf, K. M.; Murphy, B. L.; MacMillan, S. N.; Baskin, J. M.; Barr, M. P.; Boros, E.; Wilson, J. J. In Vitro Anticancer Activity and in Vivo Biodistribution of Rhenium(I) Tricarbonyl Aqua Complexes. *J. Am. Chem. Soc.* **2017**, 139, 14302–14314.
- (79) Niesel, J.; Pinto, A.; Peindy N'Dongo, H. W.; Merz, K.; Ott, I.; Gust, R.; Schatzschneider, U. Photoinduced CO Release, Cellular Uptake and Cytotoxicity of a Tris(pyrazolyl)methane (tpm) Manganese Tricarbonyl Complex. *Chem. Commun.* **2008**, 1798–1800.
- (80) Carrington, S. J.; Chakraborty, I.; Mascharak, P. K. Rapid CO Release from a Mn(I) Carbonyl Complex Derived from Azopyridine upon Exposure to Visible Light and Its Phototoxicity toward Malignant Cells. *Chem. Commun.* **2013**, 49, 11254–11256.
- (81) Carrington, S. J.; Chakraborty, I.; Bernard, J. M. L.; Mascharak, P. K. Synthesis and Characterization of a “Turn-on” photoCORM for Trackable CO Delivery to Biological Targets. *ACS Med. Chem. Lett.* **2014**, 5, 1324–1328.
- (82) Carrington, S. J.; Chakraborty, I.; Bernard, J. M. L.; Mascharak, P. K. A Theranostic Two-Tone Luminescent PhotoCORM Derived from Re(I) and (2-Pyridyl)-benzothiazole: Trackable CO Delivery to Malignant Cells. *Inorg. Chem.* **2016**, 55, 7852–7858.
- (83) Chakraborty, I.; Carrington, S. J.; Roseman, G.; Mascharak, P. K. Synthesis, Structures, and CO Release Capacity of a Family of Water-Soluble PhotoCORMs: Assessment of the Biocompatibility and Their Phototoxicity toward Human Breast Cancer Cells. *Inorg. Chem.* **2017**, 56, 1534–1545.
- (84) Kawahara, B.; Moller, T.; Hu-Moore, K.; Carrington, S.; Faull, K. F.; Sen, S.; Mascharak, P. K. Attenuation of Antioxidant Capacity in Human Breast Cancer Cells by Carbon Monoxide through Inhibition of Cystathionine β-Synthase Activity: Implications in Chemotherapeutic Drug Sensitivity. *J. Med. Chem.* **2017**, 60, 8000–8010.

- (85) Fisher, K. J.; Alyea, E. C.; Shahnazarian, N. A. ^{31}P NMR Study of the Water Soluble Derivatives of 1,3,5-Triaza-7-Phosphaadamantane (PTA). *Phosphorus, Sulfur Silicon Relat. Elem.* **1990**, *48*, 37–40.
- (86) Darensbourg, D. J.; Ortiz, C. G.; Kamplain, J. W. A New Water-Soluble Phosphine Derived from 1,3,5-Triaza-7-Phosphaadamantane (PTA), 3,7-Diacetyl-1,3,7-Triaza-5-phosphabicyclo[3.3.1]nonane. Structural, Bonding, and Solubility Properties. *Organometallics* **2004**, *23*, 1747–1754.
- (87) Gonsalvi, L.; Peruzzini, M. 1,3,5-Triaza-7-Phosphaadamantane (PTA). In *Encyclopedia of Reagents for Organic Synthesis*; Wiley: Chichester, U.K., 2010; pp 3–7.
- (88) Kurz, P.; Probst, B.; Spingler, B.; Alberto, R. Ligand Variations in $[\text{ReX}(\text{diimine})(\text{CO})_3]$ Complexes: Effects on Photocatalytic CO_2 Reduction. *Eur. J. Inorg. Chem.* **2006**, *2006*, 2966–2974.
- (89) Smieja, J. M.; Kubiak, C. P. $\text{Re}(\text{bipy-tBu})(\text{CO})_3\text{Cl}$ —improved Catalytic Activity for Reduction of Carbon Dioxide: IR-Spectroelectrochemical and Mechanistic Studies. *Inorg. Chem.* **2010**, *49*, 9283–9289.
- (90) Giordano, P. J.; Wrighton, M. S. The Nature of the Lowest Excited State in *fac*-Tricarbonylhalobis(4-phenylpyridine)rhenium(I) and *fac*-tricarbonylhalobis(4,4'-bipyridine)rhenium(I): Emissive Organometallic Complexes in Fluid Solution. *J. Am. Chem. Soc.* **1979**, *101*, 2888–2897.
- (91) Dattelbaum, D. M.; Martin, R. L.; Schoonover, J. R.; Meyer, T. J. Molecular and Electronic Structure in the Metal-to-Ligand Charge Transfer Excited States of *fac*- $[\text{Re}(4,4'\text{-X}_2\text{bpy})(\text{CO})_3(4\text{-Etpy})]^+$ ($\text{X} = \text{CH}_3, \text{H}, \text{CO}_2\text{Et}$). Application of Density Functional Theory and Time-Resolved Infrared Spectroscopy. *J. Phys. Chem. A* **2004**, *108*, 3518–3526.
- (92) Ramos, L. D.; Sampaio, R. N.; de Assis, F. F.; de Oliveira, K. T.; Homem-de-Mello, P.; Patrocínio, A. O. T.; Frin, K. P. M. Contrasting Photophysical Properties of Rhenium(I) Tricarbonyl Complexes Having Carbazole Groups Attached to the Polypyridine Ligand. *Dalton Trans.* **2016**, *45*, 11688–11698.
- (93) Gonçalves, M. R.; Frin, K. P. M. Synthesis, Characterization, Photophysical and Electrochemical Properties of Rhenium(I) Tricarbonyl Diimine Complexes with Triphenylphosphine Ligand. *Polyhedron* **2017**, *132*, 20–27.
- (94) Fredericks, S. M.; Luong, J. C.; Wrighton, M. S. Multiple Emissions from Rhenium(I) Complexes: Intraligand and Charge-Transfer Emission from Substituted Metal Carbonyl Cations. *J. Am. Chem. Soc.* **1979**, *101*, 7415–7417.
- (95) Darensbourg, D. J.; Robertson, J. B.; Larkins, D. L.; Reibenspies, J. H. Water-Soluble Organometallic Compounds. 7. Further Studies of 1,3,5-Triaza-7-Phosphaadamantane Derivatives of Group 10 Metals, Including Metal Carbonyls and Hydrides. *Inorg. Chem.* **1999**, *38*, 2473–2481.
- (96) DuBois, D. L.; Miedaner, A. Mediated Electrochemical Reduction of CO_2 . Preparation and Comparison of an Isoelectronic Series of Complexes. *J. Am. Chem. Soc.* **1987**, *109*, 113–117.
- (97) Mohr, F.; Sanz, S.; Tiekink, E. R. T.; Laguna, M. Water-Soluble and Water-Stable Organometallic Gold(II) Complexes. *Organometallics* **2006**, *25*, 3084–3087.
- (98) Tsubaki, H.; Tohyama, S.; Koike, K.; Saitoh, H.; Ishitani, O. Effect of Intramolecular π – π and CH – π Interactions between Ligands on Structure, Electrochemical and Spectroscopic Properties of *fac*- $[\text{Re}(\text{bpy})(\text{CO})_3(\text{PR}_3)]^+$ ($\text{bpy} = 2,2'$ -bipyridine; $\text{PR}_3 =$ trialkyl or triarylphosphines). *Dalton Trans.* **2005**, 385–395.
- (99) Stahl, L.; Ernst, R. D. Equilibria Studies Involving Ligand Coordination to “Open Titanocenes”: Phosphine and Pentadienyl Cone Angle Influences and the Existence of These Electron-Deficient Molecules. *J. Am. Chem. Soc.* **1987**, *109*, 5673–5680.
- (100) Tolman, C. A. Steric Effects of Phosphorus Ligands in Organometallic Chemistry and Homogeneous Catalysis. *Chem. Rev.* **1977**, *77*, 313–348.
- (101) Connick, W. B.; Di Bilio, A. J.; Schaeffer, W. P.; Gray, H. B. The Red Form of $[\text{Re}(\text{phen})(\text{CO})_3(\text{H}_2\text{O})]\text{-CF}_3\text{SO}_3\text{-H}_2\text{O}$. *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.* **1999**, *55*, 913–916.
- (102) Salignac, B.; Grundler, P. V.; Cayemittes, S.; Frey, U.; Scopelliti, R.; Merbach, A. E.; Hedinger, R.; Hegetschweiler, K.; Alberto, R.; Prinz, U.; Raabe, G.; Kölle, U.; Hall, S. Reactivity of the Organometallic *fac*- $[(\text{CO})_3\text{Re}^{\text{I}}(\text{H}_2\text{O})_3]^+$ Aquaion. Kinetic and Thermodynamic Properties of H_2O Substitution. *Inorg. Chem.* **2003**, *42*, 3516–3526.
- (103) Sacksteder, L.; Zipp, A. P.; Brown, E. A.; Streich, J.; Demas, J. N.; DeGraff, B. A. Luminescence Studies of Pyridine α -Diimine Rhenium(I) Tricarbonyl Complexes. *Inorg. Chem.* **1990**, *29*, 4335–4340.
- (104) Tapolsky, G.; Duesing, R.; Meyer, T. J. Synthetic Control of Excited-State Properties in Ligand-Bridged Complexes of Rhenium(I). Intramolecular Energy Transfer by an Electron-Transfer/Energy-Transfer Cascade. *Inorg. Chem.* **1990**, *29*, 2285–2297.
- (105) Worl, L. A.; Duesing, R.; Chen, P.; Della Ciana, L.; Meyer, T. J. Photophysical Properties of Polypyridyl Carbonyl Complexes of Rhenium(I). *J. Chem. Soc., Dalton Trans.* **1991**, 849–858.
- (106) Wallace, L.; Rillema, D. P. Photophysical Properties of Rhenium(I) Tricarbonyl Complexes Containing Alkyl- and Aryl-Substituted Phenanthrolines as Ligands. *Inorg. Chem.* **1993**, *32*, 3836–3843.
- (107) Itokazu, M. K.; Polo, A. S.; de Faria, D. L. A.; Bignozzi, C. A.; Iha, N. Y. M. Syntheses and Spectroscopic Characterization of *fac*- $[\text{Re}(\text{CO})_3(\text{phen})(\text{L})]\text{PF}_6$, $\text{L} = \text{trans- and cis-1,2-bis(4-pyridyl)-ethylene}$. *Inorg. Chim. Acta* **2001**, *313*, 149–155.
- (108) Polo, A. S.; Itokazu, M. K.; Frin, K. M.; de Toledo Patrocínio, A. O.; Iha, N. Y. M. Light Driven *Trans*-to-*Cis* Isomerization of Stilbene-like Ligands in *fac*- $[\text{Re}(\text{CO})_3(\text{NN})(\text{trans-L})]^+$ and Luminescence of Their Photoproducts. *Coord. Chem. Rev.* **2006**, *250*, 1669–1680.
- (109) Kalyanasundaram, K.; Gratzel, M. Photosensitization and Photocatalysis Using Inorganic and Organometallic Compounds. *Catalysis by Metal Complexes*; Kalyanasundaram, K., Grätzel, M., Eds.; Springer Netherlands: Dordrecht, The Netherlands, 1993; Vol. 14.
- (110) Balzani, V.; Juris, A.; Venturi, M.; Campagna, S.; Serroni, S. Luminescent and Redox-Active Polynuclear Transition Metal Complexes. *Chem. Rev.* **1996**, *96*, 759–833.
- (111) Kraljić, I.; Mohsni, S. El. A New Method for the Detection of Singlet Oxygen in Aqueous Solutions. *Photochem. Photobiol.* **1978**, *28*, 577–581.
- (112) Becke, A. D. Density-Functional Exchange-Energy Approximation with Correct Asymptotic Behavior. *Phys. Rev. A: At., Mol., Opt. Phys.* **1988**, *38*, 3098–3100.
- (113) Adamo, C.; Jacquemin, D. The Calculations of Excited-State Properties with Time-Dependent Density Functional Theory. *Chem. Soc. Rev.* **2013**, *42*, 845–856.
- (114) de Silva, A. P.; Gunaratne, H. Q. N.; Gunlaugsson, T.; Huxley, A. J. M.; McCoy, C. P.; Rademacher, J. T.; Rice, T. E. Signaling Recognition Events with Fluorescent Sensors and Switches. *Chem. Rev.* **1997**, *97*, 1515–1566.
- (115) Basolo, F. Early Kinetic Studies on CO Substitution Reactions of Metal Carbonyls. *J. Organomet. Chem.* **1990**, *383*, 579–586.
- (116) Basolo, F. Associative Substitution Reactions of Metal Carbonyls. *Inorg. Chim. Acta* **1985**, *100*, 33–39.
- (117) Kurtz, D. A.; Dhakal, B.; Donovan, E. S.; Nichol, G. S.; Felton, G. A. N. Non-Photochemical Synthesis of $\text{Re}(\text{diimine})(\text{CO})_2(\text{L})\text{Cl}$ ($\text{L} =$ Phosphine or Phosphite) Compounds. *Inorg. Chem. Commun.* **2015**, *59*, 80–83.
- (118) Shen, J.-K.; Gao, Y.-C.; Shi, Q.-Z.; Basolo, F. Kinetic Studies of CO Substitution of Metal Carbonyls in the Presence of O-Atom Transfer Reagents. *Coord. Chem. Rev.* **1993**, *128*, 69–88.
- (119) Schutte, M.; Kemp, G.; Visser, H. G.; Roodt, A. Tuning the Reactivity in Classic Low-Spin d^6 Rhenium(I) Tricarbonyl Radiopharmaceutical Synthon by Selective Bidentate Ligand Variation ($\text{L}_1\text{L}'$ -bid; $\text{L}_1\text{L}' = \text{N}, \text{N}', \text{N}, \text{O}$, and O, O' Donor Atom Sets) in *fac*- $[\text{Re}(\text{CO})_3(\text{L}_1\text{L}'\text{-bid})(\text{MeOH})]^n$ Complexes. *Inorg. Chem.* **2011**, *50*, 12486–12498.

- (120) Kaplanis, M.; Stamatakis, G.; Papakonstantinou, V. D.; Paravatou-Petsotas, M.; Demopoulos, C. A.; Mitsopoulou, C. A. Re(I) Tricarbonyl Complex of 1,10-Phenanthroline-5,6-Dione: DNA Binding, Cytotoxicity, Anti-Inflammatory and Anti-Coagulant Effects towards Platelet Activating Factor. *J. Inorg. Biochem.* **2014**, *135*, 1–9.
- (121) Balakrishnan, G.; Rajendran, T.; Murugan, K. S.; Kumar, M. S.; Sivasubramanian, V. K.; Ganesan, M.; Mahesh, A.; Thirunalasundari, T.; Rajagopal, S. Interaction of Rhenium(I) Complex Carrying Long Alkyl Chain with Calf Thymus DNA: Cytotoxic and Cell Imaging Studies. *Inorg. Chim. Acta* **2015**, *434*, 51–59.
- (122) Medley, J.; Payne, G.; Banerjee, H. N.; Giri, D.; Winstead, A.; Wachira, J. M.; Krause, J. A.; Shaw, R.; Pramanik, S. K.; Mandal, S. K. DNA-Binding and Cytotoxic Efficacy Studies of Organorhenium Pentylcarbonate Compounds. *Mol. Cell. Biochem.* **2015**, *398*, 21–30.
- (123) Llobet, A.; Doppelt, P.; Meyer, T. J. Redox Properties of Aqua Complexes of Ruthenium(II) Containing the Tridentate Ligands 2,2′:6′,2″-Terpyridine and Tris(1-pyrazolyl)methane. *Inorg. Chem.* **1988**, *27*, 514–520.
- (124) Dadci, L.; Elias, H.; Frey, U.; Hörnig, A.; Koelle, U.; Merbach, A. E.; Paulus, H.; Schneider, J. S. π -Arene Aqua Complexes of Cobalt, Rhodium, Iridium, and Ruthenium: Preparation, Structure, and Kinetics of Water Exchange and Water Substitution. *Inorg. Chem.* **1995**, *34*, 306–315.
- (125) Motterlini, R.; Gonzales, A.; Foresti, R.; Clark, J. E.; Green, C. J.; Winslow, R. M. Heme Oxygenase-1 Derived Carbon Monoxide Contributes to the Suppression of Acute Hypertensive Responses In Vivo. *Circ. Res.* **1998**, *83*, 568–577.
- (126) Otterbein, L. E. Carbon Monoxide: Innovative Anti-Inflammatory Properties of an Age-Old Gas Molecule. *Antioxid. Redox Signaling* **2002**, *4*, 309–319.
- (127) Otterbein, L. E.; Zuckerbraun, B. S.; Haga, M.; Liu, F.; Song, R.; Usheva, A.; Stachulak, C.; Bodyak, N.; Smith, R. N.; Csizmadia, E.; Tyagi, S.; Akamatsu, Y.; Flavell, R. J.; Billiar, T. R.; Tzeng, E.; Bach, F. H.; Choi, A. M. K.; Soares, M. P. Carbon Monoxide Suppresses Arteriosclerotic Lesions Associated with Chronic Graft Rejection and with Balloon Injury. *Nat. Med.* **2003**, *9*, 183–190.
- (128) Neto, J. S.; Nakao, A.; Kimizuka, K.; Romanosky, A. J.; Stolz, D. B.; Uchiyama, T.; Nalesnik, M. A.; Otterbein, L. E.; Murase, N. Protection of Transplant-Induced Renal Ischemia-Reperfusion Injury with Carbon Monoxide. *Am. J. Physiol. Renal, Fluid Electrolyte Physiol.* **2004**, *287*, F979–F989.
- (129) Kim, H. P.; Rytter, S. W.; Choi, A. M. K. CO as a Cellular Signaling Molecule. *Annu. Rev. Pharmacol. Toxicol.* **2006**, *46*, 411–449.
- (130) Bilban, M.; Haschemi, A.; Wegiel, B.; Chin, B. Y.; Wagner, O.; Otterbein, L. E. Heme Oxygenase and Carbon Monoxide Initiate Homeostatic Signaling. *J. Mol. Med.* **2008**, *86*, 267–279.
- (131) Nakao, A.; Kaczorowski, D. J.; Sugimoto, R.; Billiar, T. R.; McCurry, K. R. Application of Heme Oxygenase-1, Carbon Monoxide and Biliverdin for the Prevention of Intestinal Ischemia/Reperfusion Injury. *J. Clin. Biochem. Nutr.* **2008**, *42*, 78–88.
- (132) Halilovic, A.; Patil, K. A.; Bellner, L.; Marrazzo, G.; Castellano, K.; Cullaro, G.; Dunn, M. W.; Schwartzman, M. L. Knockdown of Heme Oxygenase-2 Impairs Corneal Epithelial Cell Wound Healing. *J. Cell. Physiol.* **2011**, *226*, 1732–1740.
- (133) Tavares, A. F. N.; Teixeira, M.; Romão, C. C.; Seixas, J. D.; Nobre, L. S.; Saraiva, L. M. Reactive Oxygen Species Mediate Bactericidal Killing Elicited by Carbon Monoxide-Releasing Molecules. *J. Biol. Chem.* **2011**, *286*, 26708–26717.
- (134) Wu, M.-L.; Ho, Y.-C.; Yet, S.-F. A Central Role of Heme Oxygenase-1 in Cardiovascular Protection. *Antioxid. Redox Signaling* **2011**, *15*, 1835–1846.
- (135) Gullotta, F.; di Masi, A.; Ascenzi, P. Carbon Monoxide: An Unusual Drug. *IUBMB Life* **2012**, *64*, 378–386.
- (136) Ling, K.; Men, F.; Wang, W.-C.; Zhou, Y.-Q.; Zhang, H.-W.; Ye, D.-W. Carbon Monoxide and Its Controlled Release: Therapeutic Application, Detection, and Development of Carbon Monoxide Releasing Molecules (CORMs). *J. Med. Chem.* **2017**, DOI: 10.1021/acs.jmedchem.6b01153.
- (137) Jackson, C. S.; Schmitt, S.; Dou, Q. P.; Kodanko, J. J. Synthesis, Characterization, and Reactivity of the Stable Iron Carbonyl Complex [Fe(CO)(N4Py)](ClO₄)₂: Photoactivated Carbon Monoxide Release, Growth Inhibitory Activity, and Peptide Ligation. *Inorg. Chem.* **2011**, *50*, 5336–5338.
- (138) Peng, P.; Wang, C.; Shi, Z.; Johns, V. K.; Ma, L.; Oyer, J.; Copik, A.; Igarashi, R.; Liao, Y. Visible-Light Activatable Organic CO-Releasing Molecules (PhotoCORMs) That Simultaneously Generate Fluorophores. *Org. Biomol. Chem.* **2013**, *11*, 6671–6674.
- (139) Romão, C. C.; Vieira, H. L. A. Metal Carbonyl Prodrugs: CO Delivery and Beyond. In *Bioorganometallic Chemistry*; Wiley-VCH: Weinheim, Germany, 2014; pp 165–202.
- (140) Loboda, A.; Jozkowicz, A.; Dulak, J. HO-1/CO System in Tumor Growth, Angiogenesis and Metabolism – Targeting HO-1 as an Anti-Tumor Therapy. *Vasc. Pharmacol.* **2015**, *74*, 11–22.
- (141) Üstün, E.; Özgür, A.; Coşkun, K. A.; Demir, S.; Özdemir, İ.; Tutar, Y. CO-Releasing Properties and Anticancer Activities of Manganese Complexes with Imidazole/benzimidazole Ligands. *J. Coord. Chem.* **2016**, *69*, 3384–3394.
- (142) Boczkowski, J.; Poderoso, J. J.; Motterlini, R. CO–Metal Interaction: Vital Signaling from a Lethal Gas. *Trends Biochem. Sci.* **2006**, *31*, 614–621.
- (143) Wegiel, B.; Gallo, D.; Csizmadia, E.; Harris, C.; Belcher, J.; Vercellotti, G. M.; Penacho, N.; Seth, P.; Sukhatme, V.; Ahmed, A.; Pandolfi, P. P.; Helczynski, L.; Bjartell, A.; Persson, J. L.; Otterbein, L. E. Carbon Monoxide Expedites Metabolic Exhaustion to Inhibit Tumor Growth. *Cancer Res.* **2013**, *73*, 7009–7021.
- (144) Rabek, J. F. Photophysics in the Membrane Environment. In *Photophysics and Photochemistry*; CRC Press: Boca Raton, FL, 1991; Vol. 3, pp 157–161.
- (145) Hepples, C.; Murphy, G. K. Synthesis of 3,3-Dichloro-2-Oxindoles from Isatin-3-*p*-Tosylhydrazones and (Dichloroiodo)-benzene. *Tetrahedron Lett.* **2015**, *56*, 4971–4974.
- (146) Sheldrick, G. M. Crystal Structure Refinement with SHELXL. *Acta Crystallogr., Sect. C: Struct. Chem.* **2015**, *71*, 3–8.
- (147) Sheldrick, G. M. A Short History of SHELX. *Acta Crystallogr., Sect. A: Found. Crystallogr.* **2008**, *64*, 112–122.
- (148) Müller, P. Practical Suggestions for Better Crystal Structures. *Crystallogr. Rev.* **2009**, *15*, 57–83.
- (149) Li, Z.; Pierri, A. E.; Huang, P. J.; Wu, G.; Iretskii, A. V.; Ford, P. C. Dinuclear PhotoCORMs: Dioxygen-Assisted Carbon Monoxide Uncaging from Long-Wavelength-Absorbing Metal–Metal-Bonded Carbonyl Complexes. *Inorg. Chem.* **2017**, *56*, 6094–6104.
- (150) Brouwer, A. M. Standards for Photoluminescence Quantum Yield Measurements in Solution (IUPAC Technical Report). *Pure Appl. Chem.* **2011**, *83*, 2213–2228.
- (151) Montalti, M.; Credi, A.; Luca, P.; Gandolfi, M. T. Chemical Actinometry. In *Handbook of Photochemistry*, 3rd ed.; CRC Press: Boca Raton, FL, 2006; pp 601–616.
- (152) Schmidt, R.; Tanielian, C.; Dunsbach, R.; Wolff, C. Phenalenone, a Universal Reference Compound for the Determination of Quantum Yields of Singlet Oxygen O₂ (¹Δ_g) Sensitization. *J. Photochem. Photobiol., A* **1994**, *79*, 11–17.
- (153) Neese, F. The ORCA Program System. *WIREs Comput. Mol. Sci.* **2012**, *2*, 73–78.
- (154) Pantazis, D. A.; Chen, X.-Y.; Landis, C. R.; Neese, F. All-Electron Scalar Relativistic Basis Sets for Third-Row Transition Metal Atoms. *J. Chem. Theory Comput.* **2008**, *4*, 908–919.
- (155) van Lenthe, E.; van der Avoird, A.; Wormer, P. E. S. Density Functional Calculations of Molecular Hyperfine Interactions in the Zero Order Regular Approximation for Relativistic Effects. *J. Chem. Phys.* **1998**, *108*, 4783–4796.
- (156) van Willen, C. Molecular Density Functional Calculations in the Regular Relativistic Approximation: Method, Application to Coinage Metal Diatomics, Hydrides, Fluorides and Chlorides, and Comparison with First-Order Relativistic Calculations. *J. Chem. Phys.* **1998**, *109*, 392–399.
- (157) Weigend, F.; Ahlrichs, R. Balanced Basis Sets of Split Valence, Triple Zeta Valence and Quadruple Zeta Valence Quality for H to Rn:

Design and Assessment of Accuracy. *Phys. Chem. Chem. Phys.* **2005**, *7*, 3297–3305.

(158) Adamo, C.; Barone, V. Toward Reliable Density Functional Methods without Adjustable Parameters: The PBE0 Model. *J. Chem. Phys.* **1999**, *110*, 6158–6170.

(159) Marenich, A. V.; Cramer, C. J.; Truhlar, D. G. Universal Solvation Model Based on Solute Electron Density and on a Continuum Model of the Solvent Defined by the Bulk Dielectric Constant and Atomic Surface Tensions. *J. Phys. Chem. B* **2009**, *113*, 6378–6396.

(160) Godwin, A. K.; Meister, A.; O'Dwyer, P. J.; Huang, C. S.; Hamilton, T. C.; Anderson, M. E. High Resistance to Cisplatin in Human Ovarian Cancer Cell Lines is Associated with Marked Increase of Glutathione Synthesis. *Proc. Natl. Acad. Sci. U. S. A.* **1992**, *89*, 3070–3074.

(161) OECD Guideline For Testing of Chemicals: In Vitro 3T3 NRU Phototoxicity Test: <https://ntp.niehs.nih.gov/iccvam/suppdocs/feddocs/oecd/oecd432-508.pdf> (accessed Dec 7, 2017).

(162) Weekes, D. M.; Ramogida, C. F.; de Guadalupe Jaraquemada-Peláez, M.; Patrick, B. O.; Apte, C.; Kostelnik, T. I.; Cawthray, J. F.; Murphy, L.; Orvig, C. Dipicolinate Complexes of Gallium(III) and Lanthanum(III). *Inorg. Chem.* **2016**, *55*, 12544–12558.

(163) Equilibrium Constants from Solution Spectrophotometric Data: <http://www.hyperquad.co.uk/HypSpec.htm> (accessed Dec 12, 2017).

(164) Alderighi, L.; Gans, P.; Ienco, A.; Peters, D.; Sabatini, A.; Vacca, A. Hyperquad Simulation and Speciation (HySS): A Utility Program for the Investigation of Equilibria Involving Soluble and Partially Soluble Species. *Coord. Chem. Rev.* **1999**, *184*, 311–318.