


# Diphenylphosphidoboratabenzene: An Anionic Analogue of Triphenylphosphine

Diego A. Hoic, William M. Davis, and Gregory C. Fu\*


Department of Chemistry  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Received May 10, 1996

Stimulated by its relationship to the ubiquitous triphenylphosphine ligand (**1**),<sup>1</sup> we have prepared<sup>2</sup> and begun to explore the coordination chemistry of the diphenylphosphidoboratabenzene anion (DPB; **2**). DPB may be viewed as a negatively charged, essentially isosteric, variant of PPh<sub>3</sub>; within this context, comparative studies of PPh<sub>3</sub> and DPB complexes, both for a given metal and for metals which are adjacent in the periodic table, could lead to useful new insights into reactivity.<sup>3</sup> In this communication, we report the first stage of our investigation into the chemistry of DPB, specifically, synthetic and structural work which establishes the viability of DPB as a ligand for an array of transition metals.



A wide range of DPB adducts can be generated through treatment of transition metal halides with potassium diphenylphosphidoboratabenzene (K-DPB). For example, reaction of Cp<sub>2</sub>ZrHCl with K-DPB in the presence of PMe<sub>3</sub> leads to displacement of the chloride ligand and formation of Cp<sub>2</sub>ZrH(DPB)(PMe<sub>3</sub>) (**3**; eq 1). A single-crystal X-ray diffraction study of this complex (Figure 1a; Table 1a)<sup>4</sup> reveals a structure very similar to that found for Cp<sub>2</sub>ZrH(SiPh<sub>3</sub>)(PMe<sub>3</sub>).<sup>5</sup> The P–C bonds of the DPB ligand are ~0.12 Å shorter than the P–B bond, and the ligand adopts a slightly distorted tetrahedral geometry.



We have also established that the iodide of CpFe(CO)<sub>2</sub>I is readily substituted by DPB, producing CpFe(CO)<sub>2</sub>(DPB) (**4**, eq

(1) For a review of phosphine complexes of transition metals, see: Dias, P. B.; de Piedade, M. E. M.; Simoes, J. A. M. *Coord. Chem. Rev.* **1994**, 135, 737–807.

(2) Qiao, S.; Hoic, D. A.; Fu, G. C. *J. Am. Chem. Soc.* **1996**, 118, 6329–6330.

(3) This general approach has proved to be extremely interesting in early transition metal metallocene chemistry. For examples and leading references, see: (a) Crowther, D. J.; Baenziger, N. C.; Jordan, R. F. *J. Am. Chem. Soc.* **1991**, 113, 1455–1457. (b) Quan, R. W.; Bazan, G. C.; Kiely, A. F.; Schaefer, W. P.; Bercaw, J. E. *J. Am. Chem. Soc.* **1994**, 116, 4489–4490.


(4) Data for compound **3**: A wheat-colored plate (0.08 × 0.09 × 0.39 mm; grown from toluene/reaction-mixture at –35 °C for three weeks) was mounted with a glass fiber on a Siemens SMART/CCD three circle diffractometer ( $\chi$  fixed at 54.78°). Data collection was done at –120 °C using Mo K $\alpha$  radiation. The crystal was found to be monoclinic, belonging to the space group P2<sub>1</sub>/c. The cell constants are  $a = 9.1017(6)$  Å,  $b = 17.6952(12)$  Å,  $c = 17.8481(12)$  Å,  $\beta = 104.2750(10)^\circ$ ,  $V = 2785.8(3)$  Å<sup>3</sup>,  $Z = 4$ ,  $\rho_{\text{calc}} = 1.334$  g/cm<sup>3</sup>. 10 886 reflections were collected, of which 3977 were unique ( $R_{\text{int}} = 0.0550$ ). Solution was done by direct methods. The final refinement by full-matrix least-squares was done on  $F^2$  (3971 data, 312 parameters) and yielded  $R_1 = 0.0641$  and  $wR_2 = 0.1104$  (for data with  $I > 2\sigma(I)$ ); GOF = 1.411. All computations were handled by the Siemens software package (SMART, SAINT, SHELXTL).

(5) Kreutzer, K. A.; Fisher, R. A.; Davis, W. M.; Spaltenstein, E.; Buchwald, S. L. *Organometallics* **1991**, 10, 4031–4035.

**Table 1.** Selected Bond Distances (Å) and Angles (deg) for (a) Cp<sub>2</sub>ZrH(DPB)(PMe<sub>3</sub>), (b) CpFe(CO)<sub>2</sub>(DPB), and (c) Rh(PMe<sub>3</sub>)<sub>3</sub>(DPB)

| (a) Cp <sub>2</sub> ZrH(DPB)(PMe <sub>3</sub> ) |           |                  |           |
|-------------------------------------------------|-----------|------------------|-----------|
| Zr–Cp<br>(average Zr–C)                         | 2.506     | P(1)–Zr–H        | 58(2)     |
| Zr–P(1)                                         | 2.738(1)  | P(1)–Zr–P(2)     | 115.09(5) |
| Zr–P(2)                                         | 2.657(2)  | P(2)–Zr–H        | 57(2)     |
| Zr–H                                            | 1.81(5)   | C(17)–P(1)–C(11) | 100.1(2)  |
| P(1)–C(17)                                      | 1.841(5)  | C(17)–P(1)–Zr    | 111.8(2)  |
| P(1)–C(11)                                      | 1.848(6)  | C(17)–P(1)–B     | 106.6(2)  |
| P(1)–B                                          | 1.966(6)  | C(11)–P(1)–Zr    | 118.5(2)  |
|                                                 |           | C(11)–P(1)–B     | 104.4(3)  |
|                                                 |           | Zr–P(1)–B        | 114.0(2)  |
| (b) CpFe(CO) <sub>2</sub> (DPB)                 |           |                  |           |
| Fe(1)–Cp<br>(average Fe–C)                      | 2.097     | B(1)–P(1)–Fe(1)  | 115.3(3)  |
| Fe(1)–P(1)                                      | 2.276(2)  | B(1)–P(1)–C(21)  | 109.0(4)  |
| Fe(1)–C(6)                                      | 1.741(10) | B(1)–P(1)–C(31)  | 106.8(3)  |
| Fe(1)–C(7)                                      | 1.743(10) | C(31)–P(1)–C(21) | 104.0(3)  |
| P(1)–B(1)                                       | 1.967(9)  | C(31)–P(1)–Fe(1) | 114.7(3)  |
| P(1)–C(21)                                      | 1.840(8)  | C(21)–P(1)–Fe(1) | 106.4(2)  |
| P(1)–C(31)                                      | 1.838(8)  | P(1)–Fe(1)–C(6)  | 95.4(3)   |
| C(6)–O(2)                                       | 1.170(9)  | P(1)–Fe(1)–C(7)  | 92.3(3)   |
| C(7)–O(1)                                       | 1.167(9)  | C(6)–Fe(1)–C(7)  | 94.8(4)   |
| (c) Rh(PMe <sub>3</sub> ) <sub>3</sub> (DPB)    |           |                  |           |
| Rh–P(1)                                         | 2.299(2)  | P(1)–Rh–P(2)     | 96.13(7)  |
| Rh–P(2)                                         | 2.285(2)  | P(1)–Rh–P(3)     | 146.36(8) |
| Rh–P(3)                                         | 2.307(2)  | P(1)–Rh–P(4)     | 92.59(7)  |
| Rh–P(4)                                         | 2.306(2)  | P(2)–Rh–P(3)     | 93.60(8)  |
| P(1)–B                                          | 1.927(8)  | P(2)–Rh–P(4)     | 148.03(8) |
| P(1)–C(1)                                       | 1.856(8)  | P(3)–Rh–P(4)     | 95.96(8)  |
| P(1)–C(7)                                       | 1.871(7)  | B–P(1)–Rh        | 104.2(3)  |
|                                                 |           | B–P(1)–C(1)      | 109.3(3)  |
|                                                 |           | B–P(1)–C(7)      | 105.4(3)  |
|                                                 |           | C(1)–P(1)–Rh     | 123.5(3)  |
|                                                 |           | C(1)–P(1)–C(7)   | 102.8(3)  |
|                                                 |           | C(7)–P(1)–Rh     | 110.5(3)  |

2). The X-ray crystal structure of **4** (Figure 1b; Table 1b)<sup>6</sup> displays a three-legged piano-stool geometry typical of CpFeL<sub>2</sub>X complexes, with a nearly staggered conformation about the Fe–P bond (dihedral angle [Cp centroid–Fe(1)–P(1)–C(31)] = –168°).<sup>7</sup> As in the case of zirconium complex **3**, the P–C bonds of the DPB ligand of **4** are shorter than the P–B bond (by ~0.13 Å).




[CpFe(CO)<sub>2</sub>(PPh<sub>3</sub>)]<sup>+</sup> and CpFe(CO)<sub>2</sub>(PPh<sub>2</sub>) are isoelectronic with complex **4**.<sup>8,9</sup> Comparison of the C–O stretching frequencies (Table 2) suggests that the iron atom of [CpFe(CO)<sub>2</sub>(PPh<sub>3</sub>)]<sup>+</sup> is the least electron-rich and that of CpFe(CO)<sub>2</sub>(PPh<sub>2</sub>) is the most electron-rich. It is important to note that the diphenylphosphido group is unique among the three phosphorus ligands in that it bears a “lone pair” which can contribute electron density to the metal. The IR data as well as the Fe–

(6) Data for compound **4**: Orange plate (0.28 × 0.12 × 0.07 mm; from pentane/toluene/THF at –35 °C). Monoclinic, C2/c,  $a = 17.1939(14)$  Å,  $b = 15.0376(13)$  Å,  $c = 15.9249(13)$  Å,  $\beta = 92.5420(10)^\circ$ ,  $V = 4113.4(6)$  Å<sup>3</sup>,  $Z = 8$ ,  $\rho_{\text{calc}} = 1.415$  g/cm<sup>3</sup>. 5705 reflections, 1914 unique ( $R_{\text{int}} = 0.0713$ ). The final refinement (1898 data, 262 parameters) yielded  $R_1 = 0.0649$  and  $wR_2 = 0.1389$  for data with  $I > 2\sigma(I)$ ; GOF = 1.290.

(7) For a discussion, see: Brunner, H.; Hammer, B.; Kruger, C.; Angermund, K.; Bernal, I. *Organometallics* **1985**, 4, 1063–1068.

(8) (a) [CpFe(CO)<sub>2</sub>PPh<sub>3</sub>]Cl·3H<sub>2</sub>O: Riley, P. E.; Davis, R. E. *Organometallics* **1983**, 2, 286–292. See also: Davison, A.; Green, M. L. H.; Wilkinson, G. *J. Chem. Soc.* **1961**, 3172–3177. (b) [CpFe(CO)<sub>2</sub>PPh<sub>3</sub>]<sup>+</sup>[(NC)<sub>2</sub>CC(CN)C(CN)<sub>2</sub>]<sup>–</sup>: Sim, G. A.; Woodhouse, D. I.; Knox, G. R. *J. Chem. Soc., Dalton Trans.* **1979**, 629–635. (c) [CpFe(CO)<sub>2</sub>PPh<sub>3</sub>]PF<sub>6</sub>: Janik, T. S.; Krajowski, L. M.; Churchill, M. R. *J. Chem. Cryst.* **1995**, 25, 751–754. [CpFe(CO)<sub>2</sub>PPh<sub>3</sub>]Cl·3H<sub>2</sub>O, [CpFe(CO)<sub>2</sub>PPh<sub>3</sub>]<sup>+</sup>[(NC)<sub>2</sub>CC(CN)C(CN)<sub>2</sub>]<sup>–</sup>, and [CpFe(CO)<sub>2</sub>PPh<sub>3</sub>]PF<sub>6</sub> differ by less than 0.01 Å for all of the bond distances reported in Table 2.

(9) CpFe(CO)<sub>2</sub>(PPh<sub>2</sub>): Burckett-St. Laurent, J. C. T. R.; Haines, R. J.; Nolte, C. R.; Steen, N. D. C. T. *Inorg. Chem.* **1980**, 19, 577–587.



**Figure 1.** ORTEP illustrations, with thermal ellipsoids drawn at the 35% probability level, of transition metal complexes of DPB: (a)  $\text{Cp}_2\text{ZrH}(\text{DPB})(\text{PMe}_3)$ , (b)  $\text{CpFe}(\text{CO})_2(\text{DPB})$ , and (c)  $\text{Rh}(\text{PMe}_3)_3(\text{DPB})$ .

(CO) and C–O bond distances (Table 2) indicate that the DPB ligand donates significantly more electron density to iron than does the  $\text{PPh}_3$  ligand.


Finally, we have determined that treatment of  $\text{Rh}(\text{PMe}_3)_3\text{Cl}$  with K-DPB generates  $\text{Rh}(\text{PMe}_3)_3(\text{DPB})$  (**5**; eq 3). A single-crystal X-ray diffraction study of complex **5** (Figure 1c; Table 1c)<sup>10</sup> reveals a distorted square planar geometry, with angles

(10) Data for compound **5**: Dark red prisms ( $0.22 \times 0.22 \times 0.13$  mm; from pentane/THF at room temperature). Monoclinic,  $P2_1/n$ ,  $a = 9.4031(8)$  Å,  $b = 17.692(2)$  Å,  $c = 17.672(2)$  Å,  $\beta = 95.6350(10)$ ,  $V = 2925.8(4)$  Å<sup>3</sup>,  $Z = 4$ ,  $\rho_{\text{calc}} = 1.344$  g/cm<sup>3</sup>. 7705 reflections, 2710 unique ( $R_{\text{int}} = 0.1446$ ). The final refinement (2640 data, 290 parameters) yielded  $R_1 = 0.0572$  and  $wR_2 = 0.1000$  for data with  $I > 2\sigma(I)$ ; GOF = 1.166.

**Table 2.** Comparison of  $[\text{CpFe}(\text{CO})_2(\text{PPh}_3)]\text{Cl} \cdot 3\text{H}_2\text{O}$ ,<sup>8</sup>  $\text{CpFe}(\text{CO})_2(\text{DPB})$ , and  $\text{CpFe}(\text{CO})_2(\text{PPh}_2)$ <sup>9</sup>

|                                                                               | $\nu(\text{CO})$ (cm <sup>-1</sup> )          | av bond distances (Å) |       |
|-------------------------------------------------------------------------------|-----------------------------------------------|-----------------------|-------|
|                                                                               |                                               | Fe–(CO)               | C–O   |
| $[\text{CpFe}(\text{CO})_2(\text{PPh}_3)]\text{Cl} \cdot 3\text{H}_2\text{O}$ | 2025, 2070 (Nujol)                            | 1.771                 | 1.139 |
| $\text{CpFe}(\text{CO})_2(\text{DPB})$                                        | 1982, 2024 (KBr)                              | 1.742                 | 1.168 |
|                                                                               | 1989, 2035 (CH <sub>2</sub> Cl <sub>2</sub> ) |                       |       |
| $\text{CpFe}(\text{CO})_2(\text{PPh}_2)$                                      | 1966, 2015 (cyclohexane)                      | not available         |       |

between cis phosphorus ligands ranging from 92 to 96° and angles between trans phosphorus ligands of 146° and 148° (P–Rh–P).<sup>11</sup> The Rh–P bond distance for the  $\text{PMe}_3$  group trans to DPB is 2.307(2) Å, and the Rh–P bond distances for the  $\text{PMe}_3$  groups cis to DPB are 2.285(2) and 2.306(2) Å. The P–C bonds of the DPB ligand of  $\text{Rh}(\text{PMe}_3)_3(\text{DPB})$  are only ~0.06 Å shorter than the P–B bond, about half the difference found for **3** and **4**, possibly reflecting the lower Lewis acidity of the rhodium center.<sup>12</sup>



In conclusion, we have established that a range of complexes containing DPB, a new anionic ligand, can be synthesized via substitution reactions of transition metal halides. It is worth noting that the chemistry of K-DPB differs from that of its nitrogen analogue, potassium diphenylamidoboratabenzene, which does not afford an  $\eta^1$  adduct in any of the reactions described above.<sup>13,14</sup> Given the structural similarity of  $\text{PPh}_3$  (**1**) and DPB (**2**), we anticipate that studies comparing the reactivity of complexes which bear these ligands may prove to be particularly interesting. The development of a chiral variant of DPB is also underway.<sup>15</sup>

**Acknowledgment.** Support has been provided by the American Cancer Society, the Camille and Henry Dreyfus Foundation, and the National Science Foundation (Young Investigator Award, with funding from DuPont, Hoechst Celanese, Merck, Pfizer, Procter & Gamble, Rohm & Haas, and Upjohn). Acknowledgment is made to the donors of the Petroleum Research Fund, administered by the ACS, for partial support of this research.

**Supporting Information Available:** A listing of experimental procedures and compound characterization data (31 pages). See any current masthead page for ordering and Internet access instructions.

JA9615740

(11) For a closely related structure, see: Thorn, D. L.; Harlow, R. L. *Inorg. Chem.* **1990**, 29, 2017–2019.

(12) A preliminary reactivity study indicates that treatment of complex **5** with H<sub>2</sub> results in the formation of two rhodium hydride species. Further investigations are underway.

(13) DPB adducts **3–5** are the first boratabenzene complexes in which the ligand is  $\sigma$ -bound, rather than  $\pi$ -bound, to the metal. For leading references to  $\pi$ -bound boratabenzene complexes, see: (a) Herberich, G. E. In *Comprehensive Organometallic Chemistry II*; Abel, E. W.; Stone, F. G. A.; Wilkinson, G., Eds.; Pergamon: New York, 1995; Vol. 1, Chapter 5. (b) Herberich, G. E.; Ohst, H. *Adv. Organomet. Chem.* **1986**, 25, 199–236. (c) Bazan, G. C.; Rodriguez, G.; Ashe, A. J., III; Al-Ahmad, S.; Muller, C. *J. Am. Chem. Soc.* **1996**, 118, 2291–2292.

(14)  $(\eta^6\text{-C}_5\text{H}_5\text{BNPh}_2)\text{Rh}(\text{PMe}_3)_2$  is formed cleanly upon treatment of  $\text{Rh}(\text{PMe}_3)_3\text{Cl}$  with potassium diphenylamidoboratabenzene.

(15) Examples of anionic  $\eta^1$  ligands in which the ligating atom is both stereogenic and configurationally stable are rare. See: Brunner, H.; Zettlmeier, W. *Handbook of Enantioselective Catalysis*; VCH: New York, 1993; Vol. 2.