

ORIGINAL ARTICLE

Proline-based phosphoramidite reagents for the reductive ligation of S-nitrosothiols

Chung-Min Park¹, Tyler D Biggs¹ and Ming Xian

S-Nitrosothiols (RSNOs) have many biological implications but are rarely used in organic synthesis. In this work we report the development of proline-based phosphoramidite substrates that can effectively convert RSNOs to proline-based sulfenamides through a reductive ligation process. A unique property of this method is that the phosphine oxide moiety on the ligation products can be readily removed under acidic conditions. In conjugation with the facile preparation of RSNOs from the corresponding thiols (RSHs), this method provides a new way to prepare proline-based sulfenamides from simple thiol starting materials.

The Journal of Antibiotics advance online publication, 13 January 2016; doi:10.1038/ja.2015.144

INTRODUCTION

Sulfur-nitrogen (-S-N-) linkages are unique chemical moieties. These structures often show interesting bioactivities (Figure 1). For example, sulfonylureas are a class of important herbicides.^{1,2} Sulfanilamides have been used as common drugs for infections.^{3,4} N-Thiolated β -lactams are found to have antibacterial, antifungal and anticancer effects.^{5–12} Sulfenamides exhibit antimicrobial activities against various infectious pathogens.¹³ Proline-based sulfenamides have been recognized as potential antiproliferative and anti-infective agents.^{14–16} Because of these activities, the preparation of these molecules, especially the formation of the -S-N- linkages, has become an active area in organic synthesis. So far, many methods have been developed for the construction of the S-N bonds, including: (1) sulfonylation between sulfonyl chlorides and amines,^{17,18} (2) amination of thiols with N-halo compounds or amines in the presence of oxidizing reagents to give unsubstituted and N-substituted sulfenamides,¹⁹ (3) the treatment of disulfides with ammonia or amines in the presence of silver or mercuric salts in alkaline medium,^{20,21} (4) sulfonylation of primary and secondary amines with the esters of sulfenic acids,²² (5) derivations of N-chlorothio-compounds,^{23,24} (6) the [2,3]-sigmatropic rearrangement of S-allylsulfynimines,²⁵ (7) reductive cleavage of sulfinimide acids in the presence of thiophenols,²⁶ (8) electrolysis of 2-mercaptobenzothiazole-amine or bisbenzothiazol-2-yl disulfide-amine^{27,28} and (9) the preparation of sulfin- or sulfon-amides by the oxidation of sulfenamides in the presence of the oxidants.^{29,30}

S-Nitrosothiols (RSNOs) are a group of -S-N- bond containing molecules. The formation of S-nitrosothiols in biological systems is an important post-translational modification elicited by nitric oxide.^{31–33} RSNOs include protein-based adducts (through cysteine residues) and small molecules (such as S-nitrosoglutathione and S-nitrosocysteine).

Small organic thiols (RSHs) can also be easily converted to the corresponding RSNOs under mild nitrosation conditions. Normally, RSNOs are unstable species that makes their detection in biological systems very challenging. Moreover, the instability of RSNOs makes these compounds unattractive for synthetic chemists. As a result, the application of RSNOs in synthesis has been rarely reported. Interestingly, in our recent work on RSNO bioorthogonal reactions (aiming at the development of novel detection methods for protein RSNO formation),^{34–41} we have discovered some unusual reactivity and properties of RSNOs. In our opinion, RSNOs are powerful synthons that can be used to introduce S, N and/or O atoms into molecular structures. Herein, we wish to report a method to prepare proline-based sulfenamides from readily available RSH substrates via RSNO intermediates. The novel reactions and synthetic strategies introduced in this article could be applied for an effective synthesis of bioactive molecules including a proline moiety.

RESULTS AND DISCUSSION

In our previous work, we discovered that RSNOs and triarylphosphines (2 equivalents) can rapidly react to generate reactive thioazaylide intermediates in high yields under mild conditions.³⁴ Thioazaylides are potent nucleophilic species. Upon manipulating the electrophilic groups attached to the phosphine reagents, thioazaylides can be trapped as stable products. For example, as shown in Scheme 1, when an *ortho*-ester group was attached to the phosphine reagent (compound 3), the reactive thioazaylide 4 could be trapped via an intramolecular acyl transfer to afford sulfenamide 5.^{35–37} This reductive ligation process provides a way for capturing unstable biological RSNO as stable and detectable conjugates. In our opinion, this reaction, in conjunction with easy RSNO formation from RSH, would be also useful for the synthesis of sulfenamide derivatives from

Department of Chemistry, Washington State University, Pullman, WA, USA

¹These authors contributed equally to this work.

Correspondence: Professor M Xian, Department of Chemistry, Washington State University, 468 Fulmer Hall, Pullman, WA 99164, USA.

E-mail: mxian@wsu.edu

Received 9 November 2015; revised 7 December 2015; accepted 13 December 2015






Figure 1 Representative molecules having sulfur-nitrogen (-S-N-) linkages.



Scheme 1 Reductive ligations of *S*-nitrosothiols.



Scheme 2 Reactions between proline-based phosphoramidites and TrSNO.



















simple RSH starting materials. However, the use of phosphine substrates like **3** would lead to products like **5** that contain an unnecessary triphenylphosphine oxide moiety. The removal of this bulky group from the final products would require harsh conditions. Therefore, seeking phosphine substrates that can undergo the reductive ligation while leading to a readily removable phosphine oxide moiety from the products is desirable. With this idea in mind, we proposed that proline-based phosphoramidites like **6** would be suitable substrates. As shown in Scheme 1, the reaction between RSNO and **6** should still follow the reductive ligation process. The resultant phosphoramidate moiety can be considered as a NH-protecting group and removable under acidic condition. As such, proline-linked sulfenamides could be prepared from simple RSH and phosphoramidite **6**.

To test this idea, three proline-based phosphoramidites (Scheme 2, **9–11**) were prepared. These substrates were treated with TrSNO (**12a**) to explore their reactivity. The selection of TrSNO as the RSNO model compound was because of its remarkable stability and ease of synthesis. We found the characteristic green color of TrSNO disappeared immediately when treated with all three phosphoramidites,

suggesting the formation of thioazaylides was fast. For **9** and **10**, the reactions stopped at this stage as no desired ligation product was obtained even when the reaction time was extended to 24 h. With **11**, however, we obtained the desired ligation product **16a**. Apparently, this can be explained by the factor that phenyl ester is a better leaving group than the methyl and *t*-butyl esters. We tested a series of different solvents for this reaction. The best was found to be a mixture of THF/phosphate buffer (pH 7.4, 20 mM, 3:1) that gave **16a** in 84% yield.

As phosphoramidite **11** was found to be the most reactive substrate for this ligation, it was applied to other RSNO substrates to test the generality of the reaction. A series of primary, secondary and tertiary RSNOs were freshly prepared from RSH and then used in this study without any purification. The results were summarized in Table 1. In all cases, the desired ligation products were obtained. For relatively stable tertiary RSNO substrates, the corresponding ligation products were isolated in modest to high yields (entries 1–4). Secondary and primary RSNO substrates gave slightly lower yields (entries 5–9). Interestingly, the purification of secondary RSNO products (entries 5 and 6) was found to be very difficult because they overlapped with the

Table 1 Summary of the reactions between 11 and S-nitrosothiols (RSNOs)

$\text{R-SH} \xrightarrow{\text{nitrosation}} \text{R-SNO} \xrightarrow[\text{THF/buffer}]{\text{11}} \text{16}$			
entry	RSNO	product	yield (%)
1	 12a	 16a	84
2	 12b	 16b	92
3	 12c	 16c	54
4	 12d	 16d	57
5	 12e	 16e	82 ^a
6	 12f	 16f	40 ^a
7	 12g	 16g	62
8	 12h	 16h	42
9	 12i	 16i	40




^aYields determined by NMR analysis.

phosphine oxide by-product. Nevertheless, the formation of the products was clearly confirmed by NMR and mass spectroscopy analysis. These results demonstrated that RSH can be readily converted to proline-based sulfenamides by this two-step SNO-ligation process.

Although phosphoramidite **9** did not show good reactivity to convert RSNO to the desired sulfenamides, we did observe some unique reactivity of **9**. As shown in Scheme 3, the treatment of a cysteine SNO derivative **12g** with **9** gave dehydroalanine **18** in a modest yield (37%) together with the disulfide product (46%).

These results indicated that the reaction indeed proceeded to form the thioazaylide intermediate **17**. Presumably, the methyl ester was not reactive enough. Hence, the acyl transfer was slow and nonproductive. Instead, the azaylide underwent an intramolecular β -elimination on the cysteine substrate to form dehydroalanine **18**.

In the proline-based sulfenamide products, the diphenylphosphoryl moiety could be considered as the protecting group of proline. We expected it could be removed under acidic

Scheme 3 The reaction between **9** and **12g**.Scheme 4 Deprotection of **16g**.Scheme 5 The reaction between **11** and HNO.

conditions.⁴² We then tested the deprotection of **16g** (Scheme 4). Indeed, a cleavage cocktail of 10% water in TFA provided sulfenamide **19** almost quantitatively. This result revealed that the -S-N- bond on the sulfenamide compounds was quite stable under acidic conditions. Given the efficiency of this protocol, it can find applications in making water-soluble proline-based sulfenamides.

HNO is the one-electron reduced/protonated form of nitric oxide. It shows distinct physiology and pharmacology from nitric oxide.⁴³ The reductive ligation was also found to be effective for HNO.⁴⁴ Several specific fluorescent probes for HNO detection have been developed based on the triarylphosphine template.^{45–47} We next wondered whether the proline-based phosphoramidite would work for HNO and tested the reaction between **11** and HNO (generated from Angeli's salt). As shown in Scheme 5, the reaction proved to be fast and efficient. The desired ligation product **21** was obtained in a comparable yield (50%) as the one obtained from triarylphosphines,⁴⁴ suggesting the proline-based phosphoramidite may be useful for the design of novel HNO sensors.

CONCLUSIONS

In summary, we have developed a reductive ligation of *S*-nitrosothiols using *N*-diphenylphosphine proline ester substrates. This reaction was found to be effective for all small-molecule *S*-nitrosothiols (primary, secondary and tertiary), as well as for HNO. The ligation products bear a removable phosphine oxide moiety on the proline residue. In conjugation with the facile preparation of RSNOs from the corresponding RSHs, this novel method provides a unique way to prepare proline-based bioactive sulfenamides from simple thiol starting materials.

METHODS

Instrumentation

¹H NMR spectra, ¹³C NMR and ³¹P NMR were recorded at 300 MHz (VX 300, Varian, Palo Alto, CA, USA) and are reported in p.p.m. on the δ scale relative to residual CHCl₃ (δ 7.25 for ¹H and δ 77.0 for ¹³C). These experiments were performed at room temperature. Mass spectra were recorded using an electrospray ionization mass spectrometry (ESI, Thermo Finnigan LCQ Advantage, San Jose, CA, USA) or MALDI-TOF (matrix-assisted laser desorption ionization time-of-flight) mass spectrometry. Mass data were reported in units of *m/z* for [M+H]⁺ or [M+Na]⁺.

Preparation of proline-based phosphoramidites

To a solution of the proline ester HCl salt in CH₂Cl₂ (total *c* = 0.30 M) was added freshly distilled 2.5 equivalent of triethylamine followed by PPh₂Cl (1 equivalent) at 0 °C. The mixture was slowly warmed to room temperature and stirred for 3–4 h. Upon completion (monitored by TLC), the reaction mixture was filtered to remove the triethylammonium salt. The filtrate was concentrated and diluted with EtOAc, and washed with satd NaHCO₃, water and brine. The organic layer was dried with anhydrous MgSO₄, filtered and concentrated. The crude product was purified by flash column chromatography (with preneutralized silica gel by 3% triethylamine in hexanes).

Compound 9: ¹H NMR (300 MHz, CDCl₃) δ 7.59–7.27 (m, 10H), 4.22 (ddd, *J* = 8.8, 6.0, 3.5 Hz, 1H), 3.63 (s, 3H), 3.17–3.06 (m, 1H), 2.88 (m, 1H), 2.22–1.70 (m, 4H); ³¹P NMR (122 MHz, CDCl₃) δ 49.31; ¹³C NMR (75 MHz, CDCl₃) δ 176.00, 138.93, 132.88, 132.61, 132.23, 131.98, 128.90, 128.40, 128.33, 128.27, 128.19, 65.28, 64.87, 51.96, 47.64, 47.57, 31.80, 31.71, 25.90; MS (ESI) *m/z* calcd for C₁₈H₂₁NO₂P [M+H]⁺ 314.1, found 314.1.

Compound 10: ¹H NMR (300 MHz, CDCl₃) δ 7.49 (ddd, *J* = 8.1, 5.4, 1.5 Hz, 2H), 7.39–7.28 (m, 8H), 4.07 (ddd, *J* = 8.6, 6.6, 3.4 Hz, 1H), 3.17–3.06 (m, 1H), 2.83 (dtd, *J* = 9.2, 6.8, 2.7 Hz, 1H), 2.07 (dq, *J* = 12.0, 8.3 Hz, 1H), 1.99–1.82 (m, 2H), 1.76–1.64 (m, 1H), 1.40 (s, 9H). ¹³C NMR (75 MHz, CDCl₃) δ 174.76, 139.27, 132.93, 132.66, 132.26, 132.01, 128.81, 128.38, 128.29, 128.19,

128.12, 80.76, 66.37, 65.95, 47.72, 47.65, 31.83, 31.74, 28.24, 25.87. ^{31}P NMR (122 MHz, CDCl_3) δ 49.64.

Compound 11: ^1H NMR (300 MHz, CDCl_3) δ 7.59–7.15 (m, 13H), 7.07–6.95 (m, 2H), 4.47 (ddd, $J=8.3, 6.3, 3.8$ Hz, 1H), 3.21 (dddd, $J=8.7, 7.5, 5.1, 1.0$ Hz, 1H), 2.95 (dtd, $J=9.4, 7.0, 2.4$ Hz, 1H), 2.36–2.14 (m, 2H), 2.09–1.93 (m, 1H), 1.83 (m, 1H); ^{31}P NMR (122 MHz, CDCl_3) δ 49.99; ^{13}C NMR (75 MHz, CDCl_3) δ 174.0, 150.9, 138.8, 138.7, 138.6, 132.9, 132.7, 132.2, 131.9, 129.5, 129.0, 128.5, 128.3, 128.2, 125.9, 121.6, 65.6, 65.1, 47.8, 47.7, 32.0, 31.9, 25.9; MS (ESI) m/z calcd for $\text{C}_{23}\text{H}_{23}\text{NO}_2\text{P}$ $[\text{M}+\text{H}]^+$ 376.1, found 376.2.

Preparation of S-nitrosothiols

The thiol starting material (RSH, 0.2 mmol) was dissolved in 1 ml of MeOH followed by the addition of 1 N HCl (1 ml) at room temperature. To this solution was then added freshly prepared 1 N NaNO_2 (1 ml) in water in dark (total $c=0.07$ M). The color of the reaction was immediately turned to red (for primary and secondary RSNO) or green (for tertiary RSNO). The mixture was stirred for 10–15 min at room temperature. Upon completion (monitored by TLC), the RSNO product was directly extracted with cold diethyl ether (1 ml \times 3) in dark. The organic layers were collected and dried. The solvent was removed to provide the RSNO product that was then used for the ligation reaction without further purification.

General reductive ligation procedure

To the freshly prepared RSNO product was added a solution of 2 equivalent of **11** in 3:1 THF-aqueous buffer (pH 7.4, degassed by bubbling with argon). The final concentration was ~ 0.1 M. The reaction was monitored by TLC and it was usually completed within 15–30 min at room temperature. The reaction mixture was extracted with ethyl acetate. The combined organic layers were washed with water and brine, dried by anhydrous Na_2SO_4 and concentrated. The crude product was purified by flash column chromatography.

Thioazaylide 15: ^1H NMR (300 MHz, CDCl_3) δ 7.81–7.55 (m, 4H), 7.47–7.17 (m, 23H), 6.94–6.80 (m, 3H), 4.35 (td, $J=8.8, 3.5$ Hz, 1H), 3.14 (qd, $J=6.5, 2.8$ Hz, 2H), 2.30 (m, 1H), 2.13–1.99 (m, 1H), 1.99–1.78 (m, 2H); ^{31}P NMR (122 MHz, CDCl_3) δ 18.3; FT-IR (thin film) 3025.4, 2985.3, 1728.8 (strong, C=O, carbonyl group), 1601.3, 1450.0, 1372.3, 1268.0, 1247.1, 1070.2 cm^{-1} ; MS (ESI) m/z calcd for $\text{C}_{42}\text{H}_{37}\text{N}_2\text{NaO}_5\text{PS}$ $[\text{M}+\text{Na}]^+$ 687.2, found 687.1.

Compound 16a: ^1H NMR (300 MHz, CDCl_3) δ 7.81 (s, 1H), 7.76–7.24 (m, 25H), 3.95 (dt, $J=9.2, 5.5$ Hz, 1H), 3.01 (p, $J=8.1$ Hz, 1H), 2.66–2.51 (m, 1H), 1.92 (m, 2H), 1.68–1.61 (m, 1H), 1.34–1.26 (m, 1H); ^{31}P (122 MHz, CDCl_3) δ 30.2; ^{13}C NMR (75 MHz, CDCl_3) δ 178.99, 150.71, 145.25, 133.24, 133.12, 132.87, 132.74, 132.01, 131.59, 130.54, 130.04, 129.81, 129.51, 128.59, 128.41, 128.22, 128.15, 127.56, 126.42, 126.00, 121.61, 120.73, 115.57, 63.10, 60.11, 49.03, 32.18, 25.76; MS (ESI) m/z calcd for $\text{C}_{36}\text{H}_{33}\text{N}_2\text{NaO}_2\text{PS}$ $[\text{M}+\text{Na}]^+$ 611.2, found 611.1.

Compound 16b: ^1H NMR (300 MHz, CDCl_3) δ 9.55 (t, 5.6 Hz, 1H, NH), 9.12 (d, $J=9.5$ Hz, 1H, NH), 8.88 (s, 1H, NH), 8.12–6.82 (m, 20H), 5.18 (d, $J=9.0$ Hz, 1H), 4.46–4.33 (m, 2H), 4.20–4.02 (m, 1H), 3.44 (m, 1H), 3.31 (m, 1H), 2.39 (m, 1H), 2.17 (m, 3H), 1.36 (d, $J=10.8$ Hz, 6H); ^{31}P (122 MHz, CDCl_3) δ 30.2; ^{13}C NMR (75 MHz, CDCl_3) δ 173.13, 169.58, 168.43, 157.53, 150.52, 133.19, 133.06, 133.03, 132.99, 132.53, 132.39, 132.21, 132.06, 131.05, 130.90, 130.49, 129.75, 129.68, 129.62, 129.33, 129.16, 129.01, 128.93, 128.85, 128.61, 128.23, 128.12, 127.42, 126.29, 121.54, 121.42, 119.59, 115.85, 62.20, 60.24, 55.76, 47.78, 43.79, 32.61, 25.91; MS (ESI) m/z calcd for $\text{C}_{36}\text{H}_{39}\text{N}_4\text{NaO}_4\text{PS}$ $[\text{M}+\text{Na}]^+$ 677.2, found 677.1.

Compound 16c: ^1H NMR (400 MHz, CDCl_3) δ 8.72 (s, 1H), 7.82 (m, 4H), 7.58–7.40 (m, 6H), 4.18 (t, $J=7.5$ Hz, 1H), 3.19 (dd, $J=12.2, 6.0$ Hz, 2H), 2.65–2.25 (m, 5H), 2.03 (t, $J=11.8$ Hz, 2H), 1.90 (dd, $J=13.6, 10.1$ Hz, 5H), 1.42 (d, $J=7.3$ Hz, 3H), 1.25 (d, $J=8.4$ Hz, 3H), 1.00 (d, $J=6.0$ Hz, 2.5H), 0.93 (d, $J=7.1$ Hz, 0.5H); ^{31}P (122 MHz, CDCl_3) δ 28.70; ^{13}C NMR (101 MHz, CDCl_3) δ 211.43, 175.80, 132.34, 132.25, 132.15, 131.96, 131.86, 128.89, 128.79, 128.66, 77.32, 77.00, 76.68, 62.25, 57.80, 52.67, 52.24, 48.20, 36.73, 34.47, 31.11, 31.05, 30.90, 29.74, 25.64, 25.16, 25.10, 22.28, 22.19; MS (Maldi) m/z calcd for $\text{C}_{27}\text{H}_{36}\text{N}_2\text{O}_3\text{PS}$ $[\text{M}+\text{H}]^+$ 499.2178, found 499.2201.

Compound 16d: ^1H NMR (300 MHz, CDCl_3) δ 8.64 (br-s, 1H, NH), 7.92–7.72 (m, 4H), 7.52 (dddd, $J=12.8, 9.3, 7.3, 2.3$ Hz, 6H), 4.21 (ddd,

$J=8.5, 6.5, 2.2$ Hz, 1H), 3.20 (m, 2H), 2.43 (dd, $J=12.7, 5.9$ Hz, 1H), 2.13–2.02 (m, 1H), 1.91 (m, 2H), 1.28 (s, 9H); ^{31}P (122 MHz, CDCl_3) δ 30.0; ^{13}C NMR (75 MHz, CDCl_3) δ 175.98, 132.67, 132.52, 132.39, 132.32, 132.19, 129.65, 129.25, 129.13, 129.08, 128.97, 121.40, 62.58, 48.91, 48.35, 31.49, 28.92, 25.42, 25.33; MS (Maldi) m/z calcd for $\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_2\text{PS}$ $[\text{M}+\text{H}]^+$ 403.1609, found 403.1604.

Compound 16g: ^1H NMR (300 MHz, CDCl_3) δ 8.97 (s, 1H), 8.30 (d, $J=8.3$ Hz, 1H), 8.09–7.97 (m, 2H), 7.90–7.67 (m, 4H), 7.62–7.36 (m, 9H), 5.02 (ddd, $J=8.3, 6.3, 4.3$ Hz, 1H), 4.27–4.02 (m, 1H), 3.74 (s, 3H), 3.32 (dd, $J=14.8, 6.4$ Hz, 1H), 3.04 (m, 3H), 2.35–2.17 (m, 1H), 2.01–1.84 (m, 1H), 1.72 (m, 2H); ^{31}P (122 MHz, CDCl_3) δ 30.1; ^{13}C NMR (75 MHz, CDCl_3) δ 176.84, 171.32, 167.73, 133.84, 132.72, 132.50, 132.37, 132.21, 132.07, 131.97, 129.21, 129.10, 128.93, 128.65, 127.91, 61.90, 52.96, 51.28, 48.72, 42.00, 30.95, 25.32; MS (ESI) m/z calcd for $\text{C}_{28}\text{H}_{30}\text{N}_3\text{NaO}_5\text{PS}$ $[\text{M}+\text{Na}]^+$ 574.3, found 574.3.

Compound 16h: ^1H NMR (300 MHz, CDCl_3) δ 8.92 (d, $J=7.6$ Hz, 1H), 8.44 (s, 1H), 7.75 (tdd, $J=12.0, 8.3, 1.4$ Hz, 4H), 7.67–7.38 (m, 6H), 7.01 (d, $J=7.2$ Hz, 3H), 6.97–6.85 (m, 3H), 5.38 (td, $J=9.5, 3.2$ Hz, 1H), 4.63 (td, $J=8.3, 5.0$ Hz, 1H), 4.07 (td, $J=7.0, 5.0$ Hz, 1H), 3.67 (s, 3H), 3.42 (m, 1H), 3.27–2.99 (m, 3H), 2.84 (dd, $J=13.9, 8.9$ Hz, 1H), 2.49 (dd, $J=14.1, 9.9$ Hz, 1H), 2.23–2.10 (m, 2H), 2.08 (s, 3H), 2.02–1.76 (m, 3H); ^{31}P (122 MHz, CDCl_3) δ 30.4; ^{13}C NMR (75 MHz, CDCl_3) δ 177.22, 171.75, 171.46, 171.04, 136.73, 132.65, 132.52, 132.33, 132.19, 129.31, 129.20, 129.10, 129.04, 128.93, 128.41, 126.81, 62.41, 54.36, 52.55, 49.93, 43.67, 37.95, 25.70, 23.40; MS (ESI) m/z calcd for $\text{C}_{32}\text{H}_{37}\text{N}_4\text{NaO}_6\text{PS}$ $[\text{M}+\text{Na}]^+$ 659.2, found 659.3.

Compound 16i: ^1H NMR (300 MHz, CDCl_3) δ 8.21 (d, $J=7.4$ Hz, 1H), 7.80 (m, 5H), 7.45 (m, 6H), 7.19 (d, $J=8.1$ Hz, 1H), 4.93 (m, 1H), 4.75 (m, 1H), 4.50 (m, 1H), 4.01 (ddd, $J=8.2, 5.9, 2.3$ Hz, 2H), 3.75–3.57 (m, 3H), 3.29–3.11 (m, 2H), 2.40–2.24 (m, 1H), 2.10–2.03 (m, 1H), 1.98 (m, 3H), 1.92–1.81 (m, 4H), 1.47–1.29 (m, 3H); ^{31}P (122 MHz, CDCl_3) δ 30.4; ^{13}C NMR (75 MHz, CDCl_3) δ 171.22, 170.56, 170.56, 170.08, 132.61, 132.57, 132.54, 132.51, 132.42, 132.37, 132.30, 132.24, 131.76, 131.52, 130.05, 129.84, 129.19, 129.10, 129.03, 128.93, 62.06, 62.02, 53.70, 52.63, 48.47, 48.06, 48.01, 42.04, 31.27, 31.19, 25.26, 25.17, 23.24, 17.97, 17.57; MS (ESI) m/z calcd for $\text{C}_{26}\text{H}_{35}\text{N}_4\text{NaO}_5\text{PS}$ $[\text{M}+\text{Na}]^+$ 569.2, found 569.3.

Dehydroalanine (Dha) 18: ^1H NMR (300 MHz, CDCl_3) δ 8.54 (br, 1H), 7.85–7.81 (m, 2H), 7.57–7.43 (m, 3H), 6.79 (s, 1H), 5.98 (d, 1H, $J=1.4$ Hz), 3.88 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 166.0, 165.0, 134.4, 132.3, 131.2, 129.0, 127.1, 109.1, 53.3; MS (ESI) m/z calcd for $\text{C}_{11}\text{H}_{12}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 206.1, found 206.2.

Deprotection of the diphenylphosphoryl group

A sulfenamide product (0.05 mmol) was treated with 1 ml of cold 10% water in TFA at 0°C (total $c=0.05$ M). The resulted solution was stirred for 30 min at 0°C . Upon completion, the excess TFA was removed with hexanes as the co-solvent under vacuum. To the remaining mixture was added cold diethyl ether to solidify the proline-sulfenamide TFA salt. The solid product was further washed with cold diethyl ether (2 ml \times 5) and dried to provide the final product.

Compound 19: ^1H NMR (300 MHz, CDCl_3) δ 10.31 (br-s, 1H), 9.70 (br-s, 1H), 9.09 (s, 1H), 7.57–7.35 (m, 5H), 4.91 (m, 1H), 4.64 (m, 1H), 3.74 (s, 3H), 3.38 (m, 2H), 3.23 (m, 2H), 2.32 (m, 1H), 2.01 (m, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 172.40, 171.21, 168.61, 132.96, 128.96, 128.70, 127.61, 60.54, 53.21, 51.71, 47.05, 40.52, 29.64, 24.59; MS (Maldi) m/z calcd for $\text{C}_{16}\text{H}_{22}\text{N}_3\text{O}_4\text{S}$ $[\text{M}+\text{H}]^+$ 352.1331, found 352.1320.

The reaction between 11 and HNO

To an argon sparged mixture of acetonitrile and water was added **11** (0.21 mmol), and to this stirring mixture was added freshly prepared Angeli's salt (0.1 mmol, $\text{Na}_2\text{N}_2\text{O}_3$). The resulting solution was allowed to stir until the reaction was completed (by TLC, or 20 min). The product **21** (50% yield) was isolated by extraction and flash column chromatography.

Compound 21: ^1H NMR (300 MHz, CDCl_3) δ 10.04 (br-s, 1H, NH), 7.96–7.68 (m, 5H), 7.64–7.39 (m, 5H), 5.85 (br-s, 1H, NH), 4.03 (ddd, $J=8.1, 5.7, 2.3$ Hz, 1H), 3.18 (m, 2H), 2.46–2.18 (m, 1H), 2.18–1.99 (m, 1H), 1.89 (ddt, $J=13.5, 8.9, 4.2$ Hz, 2H); ^{31}P (122 MHz, CDCl_3) δ 28.6; ^{13}C NMR

(75 MHz, CDCl₃) δ 176.32, 132.66, 132.62, 132.59, 132.55, 132.44, 132.41, 132.31, 132.28, 131.65, 131.39, 129.92, 129.71, 129.22, 129.13, 129.05, 128.96, 62.12, 62.08, 60.45, 48.07, 48.02, 31.29, 31.21, 25.27, 25.18; MS (ESI) m/z calcd for C₁₇H₁₉N₂NaO₂P [M+Na]⁺ 337.1, found 337.1.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

This work is supported by the NSF (0844931) and the NIH (R01GM088226).

- Chaleff, R. S. & Mauvais, C. J. Acetolactate synthase is the site of action of two sulfonylurea herbicides in higher plants. *Science* **224**, 1443–1445 (1984).
- Wang, J. G. *et al.* Structure-activity relationships for a new family of sulfonylurea herbicides. *J. Comput. Aided Mol. Des.* **19**, 801–820 (2005).
- Cooper, L. Z., Madoff, M. A. & Weinstein, L. Absorption and excretion of sulfamethomidine, a "long-acting" sulfonamide. *Antibiot. Chemother.* **12**, 618–629 (1962).
- Shah, S. S., Rivera, G. & Ashfaq, M. Recent advances in medicinal chemistry of sulfonamides. Rational design as anti-tumoral, anti-bacterial and anti-inflammatory agents. *Mini. Rev. Med. Chem.* **13**, 70–86 (2013).
- Turos, E. *et al.* N-thiolated bicyclic and monocyclic β -lactams. *Tetrahedron* **56**, 5571–5578 (2000).
- O'Driscoll, M. *et al.* Studies on the antifungal properties of N-thiolated β -lactams. *Bioorg. Med. Chem.* **16**, 7832–7837 (2008).
- Prosen, K. R. *et al.* The impact of fatty acids on the antibacterial properties of N-thiolated β -lactams. *Bioorg. Med. Chem. Lett.* **21**, 5293–5295 (2011).
- Revell, K. D., Heldreth, B., Long, T. E., Jang, S. & Turos, E. N-thiolated β -lactams: studies on the mode of action and identification of a primary cellular target in *Staphylococcus aureus*. *Bioorg. Med. Chem.* **15**, 2453–2467 (2007).
- Heldreth, B. *et al.* N-Thiolated β -lactam antibacterials: effects of the N-organothio substituent on anti-MRSA activity. *Bioorg. Med. Chem.* **14**, 3775–3784 (2006).
- Turos, E. *et al.* N-Thiolated β -lactams: a new family of anti-Bacillus agents. *Bioorg. Med. Chem. Lett.* **16**, 2084–2090 (2006).
- Turos, E. *et al.* N-thiolated β -lactams: novel antibacterial agents for methicillin-resistant *Staphylococcus aureus*. *Bioorg. Med. Chem. Lett.* **12**, 2229–2231 (2002).
- Mishra, R. K. *et al.* N-thiolated 2-oxazolidinones: a new family of antibacterial agents for methicillin-resistant *Staphylococcus aureus* and *Bacillus anthracis*. *Bioorg. Med. Chem. Lett.* **16**, 2081–2083 (2006).
- Shang, J. L. *et al.* Synthesis and evaluation of novel sulfenamides as novel anti methicillin-resistant *Staphylococcus aureus* agents. *Bioorg. Med. Chem. Lett.* **23**, 724–727 (2013).
- Crasto, C. F. *et al.* Synthesis and activity of analogues of the isoleucyl tRNA synthetase inhibitor SB-203207. *Bioorg. Med. Chem.* **11**, 2687–2694 (2003).
- Stefanska, A. L., Cassels, R., Ready, S. J. & Warr, S. R. SB-203207 and SB-203208, two novel isoleucyl tRNA synthetase inhibitors from a *Streptomyces* sp. I. Fermentation, isolation and properties. *J. Antibiot. (Tokyo)* **53**, 357–363 (2000).
- Huang, S. *et al.* 1,2,4-Triazolylaminoarylsulfonamide derivatives as antiproliferative agents, their preparation, pharmaceutical compositions, and use in therapy, WO2006042215A1 (2006).
- Koval, I. V. Progress in the chemistry of perhalomethanesulphenyl halides. *Russ. Chem. Rev.* **60**, 830–849 (1991).
- Koval, I. V. Sylphenyl chlorides in organic synthesis. *Russ. Chem. Rev.* **64**, 731–751 (1995).
- Koval, I. V. Thiols as synthons. *Russ. Chem. Rev.* **62**, 769–786 (1993).
- Davis, F. A. *et al.* Chemistry of the sulfur-nitrogen bond. VI. Convenient one-step synthesis of sulfenimines (S-aryl thiooximes). *J. Org. Chem.* **38**, 2809–2813 (1973).
- Davis, F. A. *et al.* Chemistry of the sulfur-nitrogen bond. 12. Metal-assisted synthesis of sulfenamide derivatives from aliphatic and aromatic disulfides. *J. Org. Chem.* **42**, 967–972 (1977).
- Petrov, K. A., Rudnev, G. V. & Sorokin, V. D. Sulphenamides and their derivatives. *Russ. Chem. Rev.* **59**, 832–843 (1990).
- Mueller, W. H. & Butler, P. E. β -Chloroalkylsulfenamides. Addition of dimethylamino-sulphenyl chloride to unsaturated hydrocarbons. *J. Org. Chem.* **33**, 2111–2113 (1968).
- Kühle, E. One hundred years of sulfenic acid chemistry IIb. Substitution and cyclization reactions of sulphenyl halides. *Synthesis* **1971**, 617–638 (1971).
- Koval, I. V. Progress in the chemistry of sulphilimines. *Russ. Chem. Rev.* **59**, 819–831 (1990).
- Pel'kis, N. P. & Levchenko, E. S. Interaction of areniminosulfinic acid-derivatives with thiophenols. *Zh. Org. Khim.* **22**, 387–391 (1986).
- Torii, S., Tanaka, H. & Ukida, M. Electrosynthesis of hetero-hetero atom bonds. 2. An efficient preparation of (2-benzothiazolyl)- and thiocarbamoylsulfenamides by electrolytic cross-coupling reaction of 2-mercaptobenzothiazole, bis(2-benzothiazolyl) disulfide, and/or bis(dialkylthiocarbamoyl) disulfides with various amines. *J. Org. Chem.* **43**, 3223–3227 (1978).
- Torii, S., Tanaka, H. & Ukida, M. Electrosynthesis of hetero-hetero atom bonds. 3. Sodium bromide promoted electrolytic cross-coupling reaction of imides with disulfides. Convenient synthesis of N-(cyclohexylthio)phthalimide, an important prevulcanization inhibitor. *J. Org. Chem.* **44**, 1554–1557 (1979).
- Davis, F. A., Friedman, A. J. & Nadir, U. K. Chemistry of the sulfur-nitrogen bond. 14. Arenesulfenic acids from N-alkyldienearenesulfenimines (sulfenimines). *J. Am. Chem. Soc.* **100**, 2844–2852 (1978).
- Haake, M., Gebbing, H. & Benack, H. A simple method for the oxidation of sulfenamides to sulfenimines. *Synthesis* **1979**, 97–97 (1979).
- Zhang, Y. & Hogg, N. S-Nitrosothiols: cellular formation and transport. *Free Radical Biol. Med.* **38**, 831–838 (2005).
- Lancaster, J. R. Jr. Protein cysteine thiol nitrosation: maker or marker of reactive nitrogen species-induced nonerythroid cellular signaling? *Nitric Oxide* **19**, 68–72 (2008).
- Smith, B. C. & Marletta, M. A. Mechanisms of S-nitrosothiol formation and selectivity in nitric oxide signaling. *Curr. Opin. Chem. Biol.* **16**, 498–506 (2012).
- Wang, H. & Xian, M. Fast reductive ligation of S-nitrosothiols. *Angew. Chem. Int. Ed.* **47**, 6598–6601 (2008).
- Zhang, J., Wang, H. & Xian, M. Exploration of the "traceless" reductive ligation of S-nitrosothiols. *Org. Lett.* **11**, 477–480 (2009).
- Zhang, J. *et al.* Reductive ligation mediated one-step disulfide formation of S-nitrosothiols. *Org. Lett.* **12**, 4208–4211 (2010).
- Zhang, J., Wang, H. & Xian, M. An unexpected Bis-ligation of S-nitrosothiols. *J. Am. Chem. Soc.* **131**, 3854–3855 (2009).
- Pan, J. & Xian, M. Disulfide formation via sulfenamides. *Chem. Commun.* **47**, 352–354 (2011).
- Wang, H., Zhang, J. & Xian, M. Facile formation of dehydroalanine from S-nitrosocysteines. *J. Am. Chem. Soc.* **131**, 13238–13239 (2009).
- Pan, J., Devarie-Baez, N. O. & Xian, M. Facile amide formation via S-nitrosothioacids. *Org. Lett.* **13**, 1092–1094 (2011).
- Devarie-Baez, N. O., Zhang, D., Li, S., Whorton, A. R. & Xian, M. Direct methods for detection of protein S-nitrosylation. *Methods* **62**, 171–176 (2013).
- Ramage, R., Hopton, D., Parrott, M. J., Kenner, G. W. & Moore, G. A. Phosphinamides: a new class of amino protecting groups in peptide synthesis. *J. Chem. Soc. Perkin Trans. 1*, 1357–1370 (1984).
- Fukuto, J. M. *et al.* The physiological chemistry and biological activity of nitroxyl (HNO): the neglected, misunderstood, and enigmatic nitrogen oxide. *Chem. Res. Toxicol.* **18**, 790–801 (2005).
- Reisz, J. A., Klorig, E. B., Wright, M. W. & King, S. B. Reductive phosphine-mediated ligation of nitroxyl (HNO). *Org. Lett.* **11**, 2719–2721 (2009).
- Reisz, J. A., Zink, C. N. & King, S. B. Rapid and selective nitroxyl (HNO) trapping by phosphines: kinetics and new aqueous ligations for HNO detection and quantitation. *J. Am. Chem. Soc.* **133**, 11675–11685 (2011).
- Miao, Z. *et al.* A selective phosphine-based fluorescent probe for nitroxyl in living cells. *Bioorg. Med. Chem. Lett.* **25**, 16–19 (2015).
- Kawai, K. *et al.* A reductant-resistant and metal-free fluorescent probe for nitroxyl applicable to living cells. *J. Am. Chem. Soc.* **135**, 12690–12696 (2013).