ORIGINAL RESEARCH





Synthesis and anticancer activity of multisubstituted purines and xanthines with one or two propynylthio and aminobutynylthio groups

Alicja Kowalska¹ · Krystian Pluta¹ · Małgorzata Latocha²

Received: 3 November 2017 / Accepted: 10 February 2018 © The Author(s) 2018. This article is an open access publication

Abstract

A synthesis of new 2,6-disubstituted and 2,6,8-trisubstituted 7-methylpurines as well as 8-substituted 3,7-dimethylxanthines containing a triple bond chain have been worked out. Purinethiones and xanthinethiones were converted into propynylthio derivatives, which were then further transformed via a Mannich reaction into aminobutynylthio derivatives (amine = pyrrolidine, piperidine, morpholine, and diethylamine). The products thus obtained represent various types of the purine and xanthine structure: 8-mono-, 2,6- and 6,8-dipropynylthio, 6- and 8-monoaminobutynylthio, 2,6- and 6,8-diaminobutynylthio derivatives. All of these compounds were tested for their anticancer activity against human glioblastoma SNB-19, human adenocarcinoma MDA-MB-231, and melanoma C-32 cell lines. The anticancer activity depends on the nature of the substituent and its localization in the purine and xanthine framework. Generally, compounds possessing two alkynylthio groups (propynylthio or aminobutynylthio) were more active than those possessing only one group. Some compounds exhibited stronger or similar anticancer activity to cisplatin. All compounds were found to be 2,6-dipropynylthio-7-methylpurine **4**, 2-chloro-6,8-dipropynylthio-7-methylpurine **14**, and 2-chloro-6,8-di(N-morpholinylbutynylthio)-7-methylpurine **15c** acting selectively on glioblastoma SNB-19, melanoma C-32, and adenocarcinoma MDA-MB-231 with the IC₅₀ = $0.07-4.08 \mu g/mL$.

Keywords Propynylthiopurines · Aminobutynylthiopurines · Aminobutynylthioxanthines · Anticancer activity · Structure–activity relationship

Introduction

Purine is a privileged heterocyclic nucleus existing in the chemical architecture of various bioactive compounds. It is an important pharmacophore interacting with the synthesis

Electronic supplementary material The online version of this article (https://doi.org/10.1007/s00044-018-2155-3) contains supplementary material, which is available to authorized users.

Alicja Kowalska kowalska@sum.edu.pl

² Department of Cell Biology, School of Pharmacy with the Division of Laboratory Medicine The Medical University of Silesia Jedności 8, 41-200 Sosnowiec, Poland and function of nucleic acids and enzymes. Synthetically modified purines are inhibitors of protein kinase, fructose biphosphatase, adenylation enzyme, and adenosine receptor modulators (Bliman et al. 2014; Sharma et al. 2016).

The physicochemical properties of many substituted purine derivatives provide the potential for tight binding to deep hydrophobic folds of physiologically disease relevant proteins, such as thyrosine kinases, sulfatases, and phosphorylases. Purines could be the selectively target of these protein hydrophobic folds, influencing the progression of a disease (Riva-Toniolo et al. 2003).

Conventional thiopurines (6-mercaptopurine, 6-thioguanine, and azathioprine) are incorporated into the structure of natural biomolecules—they modify nucleic acid synthesis and are commonly used as effective anticancer and immunosuppressive drugs in the treatment of acute leukemia and autoimmune hepatitis (Prima et al. 2013; Deswal and Srivastava 2017). Thiopurines remain the first line of therapy

¹ Department of Organic Chemistry, School of Pharmacy with the Division of Laboratory Medicine The Medical University of Silesia Jagiellońska 4, 41-200 Sosnowiec, Poland

in inflammatory bowel disease, Crohn's disease, ulcerative colitis, and autoimmune hepatitis (Herreras and Iborra 2017; Ward et al. 2017; Warner et al. 2016).

6-Thiopurine derivatives containing the 1,2,3-triazole moiety present higher values regarding the inhibition of parasite multiplication than chloroquine and show antimalarial and antileishmanial activities (Corrales et al. 2011).

The pharmacologically active xanthines are commonly used for their effects as stimulants, bronchodilators, phosphodiesterase inhibitors, CFTR chloride-channel activators, and adenosine receptor antagonists. The selectivity of caffeine analogs for A2A adenosine receptors was increased by the replacement of one or two methyl groups of caffeine with a propyl or propynyl substituent. 3,7-Dimethyl-1-propynylxanthine has been used as a selective A2 antagonist. 1-Propynyltheobromine and 1,3-dipropynyl-7-methylxanthine are approximately fourfold more potent than caffeine at the ryanodine sensitive calcium release channels, and fivefold more potent than caffeine as GABA_A receptor modulators. The range of clinical applications for xanthines has continued to widen and currently includes their use as anticonvulsants, nootropics, and therapeutics for the treatment of migraine (Daly et al. 2007; Thomas et al. 2013; Zagorska et al. 2015).

The introduction of substituents in positions 2, 6, and 9 of the purine ring enhances both binding affinity and selectivity regarding kinases. This enzyme family plays a critical role in the proliferation of human tumor cells, as well as in the survival and migration of neoplasia. There was an observed selectivity of 2-acetylenyl-6,9-disubstituted purine analogs regarding the inhibition of cyclindependent kinase (CDK1-cyclin B) (Legraverend et al. 2000; Sharma et al. 2016). The inhibitors of this enzyme play an important role in cell cycle regulation in a number of disease conditions, in particular cancer, and have been characterized as tumor suppressors. It is highly notable that the 2-nitrogen atom, which is characteristic of all other purine-based cyclin-dependent kinase inhibitors (CKIs), can be replaced by an sp-hybridized carbon atom, which effectively occupies the ribose binding pocket (Sharma et al. 2016).

Some substituents in positions 2 and 6 of the purine modulate both the affinity and selectivity of adenosine agonists for the different receptor subtypes. The 2-alkynyl adenosine and their N-ethylcarboxamide analogs, such as 2hexynyl (HENECA) and 2-(R,S)-phenylhydroxypropynyl (R,S-PHPNECA), have proven to be highly potent A_{2A} adenosine receptor agonists (Volpini et al. 2002, 2005; Endo et al. 2014). The type of alkynyl chain in the 2 position seems to be very important for the potency at the A_{2A} receptor. Selective A_{2A} receptor subtype antagonists have emerged as an attractive target for Parkinson's disease therapy as well as other neurodegenerative disorders such as schizophrenia and epilepsy. They are also effective as hypoglycemic agents (Baraldi et al. 1998; Endo et al. 2014). Apadenoson (ATL-146e) has entered phase III clinical trials as a pharmacologic stress agent for use in myocardial perfusion imaging (Foitzik et al. 2009).

Alkynyl chains in the 8 position of adenosine comprise a very selective ligand for the A_3 receptor subtype. Thus, it is possible to modulate the activity at the adenosine receptor subtype by introduction of alkynyl chains in either the 2 or 8 positions of 9-ethyladenine (Volpini et al. 2001, 2005). 8-Substituted 2-alkynyl-N9-propynyladenine derivatives were found to antagonize the receptor and may be effective in the treatment of Parkinson's disease (Endo et al. 2014; Ivachtchenko et al. 2013). 2-Propynyladenosinyl phenyl ether compositions as A_{2A} receptor agonists stimulate mammalian coronary vasodilatation and can be useful as therapeutics for coronary artery disease (Zablocki et al. 2001).

The presence of the alkynyl chain in 2-substituted adenosines is also very important for purinergic receptor pharmacology. Purine ionotropic receptors could serve as attractive targets for pharmacological interventions in various diseases. 2-Alkynyl derivatives of adenine, adenosine, N-alkylcarboxamidoadenosine, and adenine nucleotide are selective ligands for P2X ionotropic receptors (Koles et al. 2007; Dal Ben et al. 2011). 2-Hexynyl and 2-phenylethynyladenosine mono-, di-, and triphosphates are P2Y₁ and P2Y₁₂ receptor agonists and adenylyl cyclase modulators. Through interaction with P2Y₁ and P2Y₁₂ receptors, they effectively inhibit platelet aggregation and may be used in the prevention and treatment of arterial thrombosis as highly potent antithrombotic agents (Cristalli et al. 2005; Dal Ben et al. 2011).

Most 6-alkynyl-2-oxopurines and 6-phenylalkynylpurine nucleoside analogs exhibit cytotoxicity against a human chronic myelogenous leukemic cell line, K-562, that is comparable to or better than the known anticancer drugs, such as 6-mercaptopurine and fludarabine (Andersen et al. 2002; Brathe et al. 2003). The alkynyl-modified nucleosides substituted specifically in the 8 position, such as 8alkynyladenosines and 2'-deoxyadenosines, proved to be very selective ligands for the A3 adenosine receptor subtype, and showed moderate anti-HCV and anti HIV-1 activity in cell culture (Kuchar et al. 2008; Dimopoulou et al. 2015). Oligonucleotides containing the alkynyl group in the 1 position of 6-thiopurine or 6-thioguanine moiety have demonstrated potent antiviral activity in several assays, including the human immunodeficiency virus (Broom et al. 1997). Some of the 2- or 6-alkynylpurine derivatives were prepared for use in antibacterial prevention and therapy (Atamanyuk et al. 2012). 6-Alkynylpurines also show profound inhibitory activity against human 15lipoxygenase. The inhibitors of these enzymes have been

implicated in the oxidation of low density lipoproteins, and could have potential as drugs against artherosclerosis and other major diseases, such as for instance cancer, Parkinson's, Altzheimer's disease, heart infraction, and rheumatoid arthritis, which are also linked to free radicals (Brathe et al. 2002; Berg et al. 2005). 6-Alkynylpurines as cytokine analogs also possess a profound plant growth stimulation effect (Brathe et al. 2003).

Experimental

All commercially available organic solvents and reagents were from Sigma-Aldrich and Chempur and were used without further purification.

The melting points were determined in open capillary tubes on a Boetius melting point apparatus and were uncorrected. The ¹H NMR and ¹³C NMR spectra were recorded on a Bruker AscendTM 600 spectrometer operating at 600 MHz and 150 MHz for ¹H and ¹³C nuclei, respectively, in deuterochloroform and dimethyl sulphoxide- d_6 with tetramethylsilane as internal standard. Shifts were given in ppm, coupling constant (J) values was presented in hertz (Hz), and the abbreviations were as follows: s (singlet), d (doublet), t (triplet), and m (multiplet). High resolution mass spectral analysis (HR ESIMS) was performed on a Bruker Impact II spectrometer (in the acetonitrile: 1% formic acid, 8:2 v/v). The reactions were monitored by thin layer chromatography (TLC) using sheets coated with silica gel 60_{F254} (Merck) and chloroform, ethanol (9:1 and 5:1) as the solvents. Purity of the synthesized compounds was confirmed by TLC on the same way. Spots were detected by their absorption under UV light ($\lambda = 254$ nm) and the chromatograms were further visualized by iodine vapor. Column chromatography separations were carried out with Merck Kieselgel 60 or aluminum oxide 90 (Merck) using a mixture of chloroform-ethanol (99:1, v/v) as an eluent.

2-Chloro-6-propynylthio-7-methylpurine **1**, (Kowalska et al. 2015), 7-methylpurine-2,6-dithione **3** (Kowalska 2007), and 2,6-dichloro-7-methylpurine **11** (Kowalska et al. 1985) were obtained according to the previously described method. Theobromine was commercially available from Sigma-Aldrich.

Synthesis of 2-chloro-6-aminobut-2-ynyltio-7methylpurines 2a-c

2-Chloro-6-alkylaminobut-2-ynylthio-7-methylpurines **2a-c** were prepared from 6-propynylthio derivative **1** (0.12 g, 0.5 mmol) and amine (piperidine, morpholine, and diethy-lamine, 0.7 mmol) according to the previously described procedure (Kowalska et al. 2015).

2-Chloro-6-(4-N-piperidinylbut-2-ynylthio)-7methylpurine 2a

It was obtained as a pale yellow solid (0.14 g, 82%); mp 122–124 °C (EtOH); ¹H NMR (CDCl₃), δ : 1.44 (m, 2H, CH₂), 1.65 (m, 4H, 2CH₂), 2.54 (m, 4H, 2NCH₂), 3.33 (s, 2H, SCH₂), 4.13 (s, 3H, NCH₃), 4.24 (t, *J* = 2.1 Hz, 2H, CCH₂), 8.03 (s, 1H, H-8); (DMSO-d₆), δ : 1.23 (m, 2H, CH₂), 1.41 (m, 4H, 2CH₂), 2.28 (m, 4H, 2NCH₂), 3.20 (s, 2H, SCH₂), 4.05 (s, 3H, NCH₃), 4.25 (t, *J* = 2.4 Hz, 2H, CCH₂), 8.60 (s, 1H, H-8), ¹³C NMR (DMSO-d₆), δ : 18.86 (CH₂, SCH₂CC), 23.88 (CH₂), 25.81 (2CH₂), 34.87 (CH₃, NCH₃), 47.44 (CH₂, CCH₂), 52.68 (2CH₂, NCH₂), 79.22 (C, SCH₂CC), 80.34 (C, SCH₂CC), 123.21 (C, C-5), 151.09 (CH, C-8), 152.38 (C, C-2), 154.27 (C, C-6), 160.55 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₁₅H₁₉ClN₅S 336.1050 found 336.1051.

2-Chloro-6-(4-N-morpholinylbut-2-ynylthio)-7methylpurine 2b

It was obtained as a pale yellow solid (0.16 g, 94%); mp 168–169 °C (EtOH); ¹H NMR (CDCl₃), δ : 2.62 (m, 4H, 2NCH₂), 3.36 (s, 2H, SCH₂), 3.78 (m, 4H, 2OCH₂), 4.13 (s, 3H, NCH₃), 4.25 (s, 2H, CCH₂), 8.03 (s, 1H, H-8); (DMSO-d₆), δ : 2.36 (m, 4H, 2NCH₂), 3.24 (t, J = 2.4 Hz, 2H, SCH₂), 3.52 (m, 4H, 2OCH₂), 4.05 (s, 3H, NCH₃), 4.27 (t, J = 2.4 Hz, 2H, CCH₂), 8.59 (s, 1H, H-8); ¹³C NMR (DMSO-d₆), δ : 18.74 (CH₂, SCH₂CC), 34.86 (CH₃, NCH₃), 46.96 (CH₂, CCH₂), 51.95 (2CH₂, NCH₂), 66.45 (2CH₂, OCH₂), 78.82 (C, SCH₂CC), 80.58 (C, SCH₂CC), 123.12 (C, C-5), 151.09 (CH, C-8), 152.35 (C, C-2), 154.24 (C, C-6), 160.53 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₁₄H₁₇ClN₅OS 338.0842 found 338.0840.

2-Chloro-6-(4-diethylaminobut-2-ynylthio)-7methylpurine 2c

It was obtained as a pale yellow solid (0.14 g, 86%); mp 117–118 °C (EtOH); ¹H NMR (CDCl₃), δ : 1.06 (t, J = 7.2 Hz, 6H, 2CH₃), 2.54 (q, J = 7.2 Hz, 4H, 2NCH₂), 3.44 (s, 2H, SCH₂), 4.13 (s, 3H, NCH₃), 4.24 (t, J = 2.1 Hz, 2H, CCH₂), 8.02 (s, 1H, H-8); (DMSO-d₆), δ : 0.87 (t, J = 7.2 Hz, 6H, 2CH₃), 2.32 (q, J = 7.2 Hz, 4H, 2NCH₂), 3.33 (s, 2H, SCH₂), 4.05 (s, 3H, NCH₃), 4.24 (s, 2H, CCH₂), 8.59 (s, 1H, H-8), ¹³C NMR (DMSO-d₆), δ : 12.84 (2CH₃, CH₂CH₃), 18.93 (CH₂, SCH₂CC), 34.86 (CH₃, NCH₃), 40.74 (CH₂, NCH₂), 46.85 (2CH₂, CCH₂), 78.66 (C, SCH₂CC), 80.10 (C, SCH₂CC), 123.22 (C, C-5), 151.09 (CH, C-8), 152.38 (C, C-2), 154.26 (C, C-6), 160.55 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₁₄H₁₉ClN₅S 324.1050 found 324.1053.

Synthesis of 2,6-di(prop-2-ynylthio)-7-methylpurine 4

The mixture of 7-methylpurine-2,6-dithione **3** (0.10 g, 0.5 mmol) and potassium *tert*-butoxide (0.16 g, 1.44 mmol) in 10 mL of DMF was stirred at room temperature for 0.5 h. Then, 80% solution of propargyl bromide (0.17 g, 1.44 mmol) in dry toluene (0.25 mL) was added. The reaction mixture was stirred for an additional 24 h at room temperature and then poured into 25 mL of water. The resulted solid was filtered off and washed with water to give compound **4**.

It was obtained as a pale yellow solid (0.11 g, 79%); mp 162–164 °C (EtOH); ¹H NMR (CDCl₃), δ : 2.21 (t, J = 2.7 Hz, 1H, CH), 2.28 (t, J = 2.7 Hz, 1H, CH), 4.04 (d, J = 2.7 Hz, 2H, SCH₂), 4.11 (s, 3H, NCH₃), 4.22 (d, J = 2.7 Hz, 2H, SCH₂), 8.09 (s, 1H, H-8); (DMSO-d₆), δ : 3.14 (t, J = 2.4 Hz, 1H, CH), 3.26 (t, J = 2.4 Hz, 1H, CH), 4.02 (s, 3H, NCH₃), 4.06 (d, J = 2.4 Hz, 2H, SCH₂), 4.27 (d, J = 2.4 Hz, 2H, SCH₂), 8.47 (s, 1H, H-8); ¹³C NMR (DMSO-d₆), δ : 17.82 (CH₂, SCH₂CCH), 19.50 (CH₂, SCH₂CCH), 34.62 (CH₃, NCH₃), 73.56 (CH, SCH₂CCH), 74.52 (CH, SCH₂CCH), 80.09 (C, SCH₂CCH), 81.24 (C, SCH₂CCH), 121.42 (C, C-5), 149.92 (CH, C-8), 152.16 (C, C-2), 159.87 (C, C-6), 161.61 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₁₂H₁₁N₄S₂ 275.0425 found 275.0417.

General procedure for synthesis of 2,6-di(aminobut-2-ynylthio)-7-methylpurines 5a-d

To a mixture of propynylthio derivative **2** (0.137 g, 0.5 mmol) and paraformaldehyde (0.067 g, 1.4 mmol) in 10 mL of dry dioxane, appropriate amines (pyrrolidine, piperidine, morpholine, or diethylamine, 1.4 mmol) and CuCl (0.01 g) were added. The reaction mixture was stirred at temperature 80 °C for 4 h. In the case of dimethylamine, the reaction was performed at 50 °C for 24 h. After cooling, the small amount of solid was filtered off and residue was extracted with chloroform (3×5 mL), dried with anhydrous Na₂SO₄, and evaporated in vacuo. The dried residue was dissolved in CHCl₃ and purified by thin layer preparative chromatography (silica gel, CHCl₃ – EtOH, 5:1 v/v) to give compound **5a-d**.

2,6-Di(4-N-pyrrolidinylbut-2-ynylthio)-7methylpurine 5a

It was obtained as a dark yellow solid (0.175 g, 79.5%); mp 97–99 °C (EtOH); ¹H NMR (CDCl₃), δ : 1.85 (m, 8H, 4CH₂), 2.64 (m, 8H, 4NCH₂), 3.44 (t, J = 2.1 Hz, 2H, SCH₂), 3.47 (t, J = 2.1 Hz, 2H, SCH₂), 3.98 (s, 3H, NCH₃), 4.06 (m, 2H, CCH₂), 4.21 (m, 2H, CCH₂), 7.92 (s, 1H, H-8); (DMSO-d₆), δ : 1.61 (m, 8H, 4CH₂), 2.43 (m, 8H,

4NCH₂), 3.36 (s, 4H, SCH₂), 4.01 (s, 3H, NCH₃), 4.10 (t, J = 1.8 Hz, 2H, CCH₂), 4.30 (t, J = 1.8 Hz, 2H, CCH₂), 8.46 (s, 1H, H-8); ¹³C NMR (DMSO-d₆), δ : 18.41 (CH₂, SCH₂CC), 19.97(CH₂, SCH₂CC), 23.72 (4CH₂), 34.64 (CH₃, NCH₃), 42.70 (CH₂, CCH₂), 42.81 (CH₂, CCH₂), 52.01 (2CH₂, NCH₂), 52.07 (2CH₂, NCH₂), 78.36 (C, SCH₂CC), 79.30 (C, SCH₂CC), 80.23 (C, SCH₂CC), 81.23 (C, SCH₂CC), 121.42 (C, C-5), 149.85 (CH, C-8), 152.28 (C, C-2), 159.86 (C, C-6), 161.88 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₂₂H₂₉N₆S₂ 441.1895 found 441.1886.

2,6-Di(4-N-piperidinylbut-2-ynylthio)-7methylpurine 5b

It was obtained as a dark yellow solid (0.18 g, 77%); mp 130–132 °C (EtOH); ¹H NMR; (CDCl₃), δ: 1.40 (m, 4H, 2CH₂), 1.60 (m, 8H, 4CH₂), 2.47 (m, 8H, 4NCH₂), 3.28 (m, 4H, 2SCH₂), 4.07 (s, 3H, NCH₃), 4.07 (t, J = 2.1 Hz, 2H, CCH₂), 4.23 (t, *J* = 2.1 Hz, 2H, CCH₂), 7.92 (s, 1H, H-8); (DMSO-d₆), *δ*: 1.23 (m, 4H, 2CH₂), 1.43 (m, 8H, 4CH₂), 2.30 (m, 8H, 4NCH₂), 3.21 (m, 4H, 2SCH₂), 4.01 (s, 3H, NCH₃), 4.10 (s, 2H, CCH₂), 4.32 (s, 2H, CCH₂), 8.46 (s, 1H, H-8); ¹³C NMR (DMSO-d₆), δ: 18.49 (CH₂, SCH₂CC), 20.05 (CH₂, SCH₂CC), 23.89 (CH₂), 23.90 (CH₂), 25.78 (2CH₂), 25.80 (2CH₂), 34.62 (CH₃ NCH₃), 47.48 (CH₂, CCH₂), 47.53 (CH₂, CCH₂), 52.80 (4CH₂, NCH₂), 78.22 (C, SCH₂CC), 79.15 (C, SCH₂CC), 80.72 (C, SCH₂CC), 81.72 (C, SCH₂CC), 121.42 (C, C-5), 149.82 (CH, C-8), 152.31 (C, C-2), 159.86 (C, C-6), 161.90 (C, C-4); HR ESIMS m/z $[M + H]^+$ calcd. for $C_{24}H_{33}N_6S_2$ 469.2208 found 469.2205.

2,6-Di(4-N-morpholinylbut-2-ynylthio)-7methylpurine 5c

It was obtained as a dark yellow solid (0.175 g, 74%); mp 121–123 °C (EtOH); ¹H NMR (CDCl₃), δ: 2.55 (m, 8H, 4NCH₂), 3.30 (m, 4H, 2SCH₂), 3.73 (m, 8H, 4OCH₂), 4.08 (s, 3H, NCH₃), 4.08 (t, J = 2.1 Hz, 2H, CCH₂), 4.23 (t, J =2.1 Hz, 2H, CCH₂), 7.93 (s, 1H, H-8); (DMSO-d₆), δ: 2.35 (m, 8H, 4NCH₂), 3.23 (s, 2H, SCH₂), 3.25 (s, 2H, SCH₂), 3.52 (m, 8H, 4OCH₂), 4.01 (s, 3H, NCH₃), 4.11 (s, 2H, CCH₂), 4.32 (s, 2H, CCH₂), 8.46 (s, 1H, H-8); ¹³C NMR (DMSO-d₆), *δ*: 18.39 (CH₂, SCH₂CC), 19.99 (CH₂, SCH₂CC), 34.63 (CH₃ NCH₃), 47.03 (CH₂, CCH₂), 47.06 (CH₂, CCH₂), 52.00 (2CH₂, NCH₂), 52.09 (2CH₂, NCH₂), 66.47 (4CH₂, OCH₂), 77.77 (C, SCH₂CC), 78.76 (C, SCH₂CC), 80.95 (C, SCH₂CC), 82.08 (C, SCH₂CC), 121.43 (C, C-5), 149.83 (CH, C-8), 152.32 (C, C-2), 159.86 (C, C-6), 161.82 (C, C-4); HR ESIMS $m/z [M + H]^+$ calcd. for C₂₂H₂₉N₆O₂S₂ 473.1793 found 473.1795.

2,6-Di(4-diethylaminobut-2-ynylthio)-7methylpurine 5d

It was obtained as a dark yellow solid (0.173 g, 78%); mp 77–79 °C (EtOH); ¹H NMR (CDCl₃), δ : 1.06 (m, 12H, 4CH₃), 2.54 (m, 8H, 4NCH₂), 3.45 (m, 4H, 2SCH₂), 4.07 (t, J = 2.1 Hz, 2H, CCH₂), 4.08 (s, 3H, NCH₃), 4.22 (t, J = 2.1 Hz, 2H, CCH₂), 7.92 (s, 1H, H-8); (DMSO-d₆), δ : 0.91 (m, 12H, 4CH₃), 2.37 (m, 8H, 4NCH₂), 3.35 (m, 4H, SCH₂), 4.01 (s, 3H, NCH₃), 4.09 (m, 2H, CCH₂), 4.29 (m, 2H, CCH₂), 8.46 (s, 1H, H-8); ¹³C NMR (DMSO-d₆), δ : 12.80 (4CH₃, CH₂CH₃), 17.80 (CH₂, SCH₂CC), 20.01 (CH₂, SCH₂CC), 34.62 (CH₃, NCH₃), 40.84 (4CH₂, NCH₂), 46.92 (2CH₂, CCH₂), 74.52 (2C, SCH₂CC), 80.03 (2C, SCH₂CC), 121.42 (C, C-5), 149.91 (CH, C-8), 152.02 (C, C-2), 159.89 (C, C-6), 161.86 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₂₂H₃₃N₆S₂ 445.2208 found 445.2204.

Synthesis of 8-bromo-3,7-dimethylxanthine 7

To a stirred suspension of theobromine **6** (2.00 g, 10 mmol) in 4 mL of carbon tetrachloride and 12 mL of nitrobenzene, bromine (2.90 g, 0.9 mL, 18 mmol) dissolved in 1.5 mL of nitrobenzene was added dropwise. When the addition was completed, the reaction mixture was refluxed with stirring for 5 h. The suspension was then carefully poured with stirring into 40 mL of acetone. The resulting white precipitate was filtered, washed with acetone and ether, and dried in vacuo (Koppel et al. 1962). The crude product was purified by column chromatography (silica gel, CHCl₃ - EtOH, 99:1, v/v) to give compound **7**.

It was obtained as a pale yellow solid (2.98 g, 69%), mp 298–300 °C, dec (EtOH), 298 °C dec (Koppel et al. 1962) ¹H NMR (DMSO-d₆), δ : 3.31 (s, 3H, N3CH₃), 3.82 (s, 3H, N7CH₃), 11.28 (s, 1H, N1H); ¹³C NMR (DMSO-d₆), δ : 28.96 (CH₃, N3CH₃), 34.10 (CH₃, N7CH₃), 109.58 (C, C-5), 128.84 (CH, C-8), 149.48 (C, C-2), 151.04 (C, C-6), 154.58 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₇H₈BrN₄O₂ 258.9831 found 258.9834.

Synthesis of 3,7-dimethylxanthine-8-thione 8

To the suspension of 8-bromo derivative 7 (0.26 g, 1 mmol) in 40 mL of anhydrous ethanol, sodium hydrosulfide (NaSH \times 2 H₂O, 0.92 g, 10 mmol) was added and the mixture was stirred and refluxed for 2 h. After cooling the alcohol was evaporated in vacuo and dry residue was dissolved in 5% NaOH solution. The reaction product was precipitated with 15% hydrochloric acid to give compound **8**.

It was obtained as a dark yellow solid (0.20 g, 95%), mp 322-324 °C (EtOH), 319-320 °C (Carson et al. 1998); ¹H NMR (DMSO-d₆), δ : 3.30 (s, 3H, N3CH₃), 3.64 (s, 3H, N7CH₃), 11.34 (s, 1H, N1H), 13.64 (s, 1H, N9H); ¹³C

NMR (DMSO-d₆), δ : 30.39 (CH₃, N3CH₃), 32.29 (CH₃, N7CH₃), 104.66 (C, C-5), 140.91 (CH, C-8), 150.12 (C, C-2), 153.11 (C, C-6), 164.19 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₇H₉N₄O₂S 213.0446 found 213.0438.

Synthesis of 8-(prop-2-ynylthio)-3,7dimethylxanthine 9

8-(Prop-2-ynylthio)-3,7-dimethylxanthine 9 was obtained from xanthine-8-thione 8 (0.11 g, 0.5 mmol) with propargyl bromide in DMF according to the previously described procedure (Kowalska et al. 2015).

It was obtained as a pale yellow solid (0.100 g, 80%); mp 252–253 °C (EtOH); ¹H NMR (DMSO-d₆), δ : 3.29 (t, J = 2.4 Hz, 1H, CH), 3.36 (s, 3H, N3CH₃), 3.79 (s, 3H, N7CH₃), 4.07 (d, J = 2.4 Hz, 2H, SCH₂), 11.17 (s, 1H, NH); ¹³C NMR (DMSO-d₆), δ : 21.64 (CH₂, SCH₂CCH), 28.99 (CH₃, N3CH₃), 32.79 (CH₃, N7CH₃), 75.26 (C, SCH₂CCH), 79.93 (C, SCH₂CCH), 109.21 (C, C-5), 148.19 (CH, C-8), 149.80 (C, C-2), 151.12 (C, C-6), 154.69 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₁₀H₁₁N₄O₂S 251.0603 found 251.0600.

Synthesis of 8-(aminobut-2-ynylthio)-3,7dimethylxanthine 10a-d

8-(Aminobut-2-ynylthio)-3,7-dimethylxanthines **10a-d** were obtained from 8-propynylthio derivative **9** (0.125 g, 0.5 mmol) and appropriate amines (pyrrolidine, piperidine, morpholine, and diethylamine, 0.7 mmol) in dry dioxane according to the described procedure (Kowalska et al. 2015).

8-(4-N-pyrrolidinylbut-2-ynylthio)-3,7dimethylxanthine 10a

It was obtained as a pale yellow solid (0.139 g, 84%); mp 195–197 °C (EtOH); ¹H NMR (DMSO-d₆), δ : 1.56 (m, 4H, 2CH₂), 2.32 (m, 4H, 2NCH₂), 3.33 (s, 2H, SCH₂), 3.33 (s, 3H, N3CH₃), 3.80 (s, 3H, N7CH₃), 4.07 (t, *J* = 2.4 Hz, 2H, CCH₂), 11.19 (s, 1H, NH); ¹³C NMR (DMSO-d₆), δ : 22.36 (CH₂, SCH₂CC), 23.65 (2CH₂), 28.95 (CH₃, N3CH₃), 32.83 (CH₃, N7CH₃), 42.48 (CH₂, CCH₂), 51.73 (2CH₂, NCH₂), 80.10 (C, SCH₂CC), 80.15 (C, SCH₂CC), 109.11 (C, C-5), 148.28 (CH, C-8), 149.82 (C, C-2), 151.11 (C, C-6), 154.68 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₁₅H₂₀N₅O₂S 334.1338 found 334.1343.

8-(4-N-piperidinylbut-2-ynylthio)-3,7dimethylxanthine 10b

It was obtained as a pale yellow solid (0.120 g, 69%); mp 223–225 °C (EtOH); ¹H NMR (DMSO- d_6), δ : 1.23 (m, 2H,

CH₂), 1.39 (m, 4H, 2CH₂), 2.19 (m, 4H, 2NCH₂), 3.18 (t, J = 1.8 Hz, 2H, SCH₂), 3.32 (s, 3H, N3CH₃), 3.81 (s, 3H, N7CH₃), 4.08 (t, J = 1.8 Hz, 2H, CCH₂), 11.19 (s, 1H, NH); ¹³C NMR (DMSO-d₆), δ : 22.42 (CH₂, SCH₂CC), 23.90 (CH₂), 25.81 (2CH₂), 28.95 (CH₃, N3CH₃), 32.87 (CH₃, N7CH₃), 47.37 (CH₂, CCH₂), 52.57 (2CH₂, NCH₂), 79.92 (C, SCH₂CC), 80.62 (C, SCH₂CC), 109.12 (C, C-5), 148.25 (CH, C-8), 149.84 (C, C-2), 151.11 (C, C-6), 154.69 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₁₆H₂₂N₅O₂S 348.1494 found 348.1489.

8-(4-N-morpholinylbut-2-ynylthio)-3,7dimethylxanthine 10c

It was obtained as a pale yellow solid (0.126 g, 72%); mp 174–176 °C (EtOH); ¹H NMR (DMSO-d₆), δ : 2.24 (m, 4H, 2NCH₂), 3.24 (t, J = 1.8 Hz, 2H, SCH₂), 3.33 (s, 3H, N3CH₃), 3.81 (s, 3H, N7CH₃), 4.09 (t, J = 1.8 Hz, 2H, CCH₂), 4.47 (t, J = 1.8 Hz, 4H, 2OCH₂), 11.15 (s, 1H, NH); ¹³C NMR (DMSO-d₆), δ : 22.30 (CH₂, SCH₂CC), 28.96 (CH₃, N3CH₃), 32.86 (CH₃, N7CH₃), 46.80 (CH₂, CCH₂), 51.53 (2CH₂, NCH₂), 66.43 (2CH₂, 2OCH₂), 79.28 (C, SCH₂CC), 81.15 (C, SCH₂CC), 109.18 (C, C-5), 148.13 (CH, C-8), 149.86 (C, C-2), 151.15 (C, C-6), 154.73 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₁₅H₂₀N₅O₃S 350.1287 found 350.1279.

8-(4-diethylaminobut-2-ynylthio)-3,7dimethylxanthine 10d

It was obtained as a pale yellow solid (0.131 g, 78%); mp 186–188 °C (EtOH); ¹H NMR (DMSO-d₆), δ : 0.86 (t, J = 7.2 Hz, 6H, 2CH₃), 2.24 (q, J = 7.2 Hz, 4H, 2NCH₂), 3.30 (s, 2H, SCH₂), 3.30 (s, 3H, N3CH₃), 3.80 (s, 3H, N7CH₃), 4.07 (s, 2H, CCH₂), 11.18 (s, 1H, NH); ¹³C NMR (DMSO-d₆), δ : 12.82 (2CH₃), 22.43 (SCH₂CC), 28.94, (CH₃, N3CH₃), 32.85 (CH₃, N7CH₃), 40.69 (CH₂, CCH₂), 46.71 (2CH₂, NCH₂), 79.33 (C, SCH₂CC), 80.36 (C, SCH₂CC), 109.10 (C, C-5), 148.26 (CH, C-8), 149.85 (C, C-2), 151.12 (C, C-6), 154.67 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₁₅H₂₂N₅O₂S 336.1494 found 336.1491.

Synthesis of 2-chloro-6,8-dibromo-7-methylpurine 12

To a stirred suspension of 2,6-dichloro-7-methylpurine **11** (2.00 g, 9.8 mmol) in 4 mL of carbon tetrachloride and 12 mL of nitrobenzene, bromine (4.90 g, 1.6 mL, 31 mmol) dissolved in 3.0 mL of nitrobenzene was added dropwise. When the addition was completed, the reaction mixture was refluxed with stirring for 5 h. The suspension was then carefully poured with stirring into 40 mL of acetone. The resulting white precipitate was filtered, washed with acetone

and ether, and dried in vacuo. Crude product was purified by column chromatography (silica gel, $CHCl_3$ - ETOH, 99:1, v/v) to give compound **12**.

It was obtained as a pale yellow solid (2.15 g, 67%), mp 218–220 °C, (EtOH); ¹H NMR (DMSO-d₆), δ : 4.02 (s, 3H, N7CH₃), ¹³C NMR (DMSO-d₆), δ : 35.22 (CH₃, NCH₃), 127.07 (C, C-5), 133.88 (CH, C-8), 142.90 (C, C-2), 151.46 (C, C-6), 161.03 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₆H₄Br₂ClN₄ 324.8491 found 324.8489.

Synthesis of 2-chloro-7-methylpurine-6,8-dithione 13

A solution of 6,8-dibromo derivative **12** (0.326 g, 1 mmol) and sodium hydrosulfide (NaSH \times 2H₂O, 0.92 g, 10 mmol) in 40 mL ethanol was refluxed for 2 h. The alcohol was evaporated in vacuo and dry residue was dissolved in 5% NaOH solution. The reaction product was precipitated with 15% hydrochloric acid to give compound **13**.

It was obtained as a dark yellow solid (0.225 g, 97%), mp 296–298 °C (EtOH); ¹H NMR (DMSO-d₆), δ : 4.03 (s, 3H, N7CH₃), 13.88 (s, 1H, N1H), 13.91 (s, 1H, N9H); ¹³C NMR (DMSO-d₆), δ : 34.62 (CH₃, NCH₃), 121.42 (C, C-5), 149.92 (CH, C-8), 152.16 (C, C-2), 159.87 (C, C-6), 161.61 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₆H₆ClN₄S₂ 232.9722 found 232.9720.

Synthesis of 2-chloro-6,8-di(prop-2-ynylthio)-7methylpurine 14

2-Chloro-6,8-di(prop-2-ynylthio)-7-methylpurine 14 was obtained by alkylation of purine-6,8-dithione 13 (0.12 g, 0.5 mmol) with propargyl bromine in DMF according to the procedure described for the synthesis compound 4.

It was obtained as a pale yellow solid (0.105 g, 68%); mp 161–163 °C (EtOH), ¹H NMR (DMSO-d₆), δ : 3.26 (t, J = 2.4 Hz, 1H, CH), 3.31 (t, J = 2.4 Hz, 1H, CH), 3.88 (s, 3H, NCH₃), 4.22 (d, J = 2.4 Hz, 2H, SCH₂), 4.26 (d, J = 2.4 Hz, 2H, SCH₂); ¹³C NMR (DMSO-d₆), δ : 18.22 (CH₂, SCH₂CCH), 21.02 (CH₂, SCH₂CCH), 33.90 (CH₃, NCH₃), 74.68 (CH, SCH₂CCH), 75.05 (CH, SCH₂CCH), 79.61 (C, SCH₂CCH), 79.68 (C, SCH₂CCH), 125.50 (C, C-5), 150.98 (CH, C-8), 152.08 (C, C-2), 159.62 (C, C-6), 159.97 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₁₂H₁₀ClN₄S₂ 309.0035 found 309.0032.

Synthesis of 2-chloro-6,8-di(aminobut-2-ynylthio)-7methylpurines 15a-d

2-Chloro-6,8-di(aminobut-2-ynylthio)-7-methylpurines **15a-d** were obtained from 6,8-dipropynylthio derivative **14** (0.15 g, 0.5 mmol) and appropriate amines ((pyrrolidine, piperidine, morpholine, and diethylamine, 0.7 mmol) in dry dioxane according to the previously described procedure for the synthesis of 2,6-di(aminobut-2-ynylthio)-7-methylpurines **5a-d**.

2-Chloro-6,8-di(4-N-pyrrolidinylbut-2-ynylthio)-7methylpurine 15a

It was obtained as a pale yellow solid (0.183 g, 77%); mp 133–135 °C (EtOH); ¹H NMR (DMSO-d₆), δ : 1.58 (m, 8H, 4CH₂), 2.38 (m, 8H, 4NCH₂), 3.34 (t, J = 2.4 Hz, 2H, SCH₂), 3.36 (t, J = 2.4 Hz, 2H, SCH₂), 3.88 (s, 3H, NCH₃), 4.22 (t, J = 2.4 Hz, 2H, CCH₂), 4.27 (t, J = 2.4 Hz, 2H, CCH₂); ¹³C NMR (DMSO-d₆), δ : 18.93 (CH₂, SCH₂CC), 18.96 (CH₂, SCH₂CC), 21.70 (CH₃, NCH₃), 23.66 (2CH₂), 23.71 (2CH₂), 33.91 (CH₂, CCH₂), 33.92 (CH₂, CCH₂), 42.57 (CH₂, NCH₂), 42.63 (CH₂, NCH₂), 51.83 (CH₂, NCH₂), 51.95 (CH₂, NCH₂), 79.32 (C, SCH₂CC), 79.38 (C, SCH₂CC), 79.89 (C, SCH₂CC), 80.03 (C, SCH₂CC), 125.52 (C, C-5), 143.07 (CH, C-8), 151.20 (C, C-2), 152.19 (C, C-6), 159.84 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₂₂H₂₈ClN₆S₂ 475.1505 found 475.1491.

2-Chloro-6,8-di(4-N-piperidinylbut-2-ynylthio)-7methylpurine 15b

It was obtained as a pale yellow solid (0.165 g, 74%); mp 108–110 °C (EtOH); ¹H NMR (DMSO-d₆), δ : 1.23 (m, 4H, 2CH₂), 1.40 (m, 8H, 4CH₂), 2.26 (m, 8H, 4NCH₂), 3.19 (s, 4H, 2SCH₂), 3.89 (s, 3H, NCH₃), 4.23 (t, J = 2.4 Hz, 2H, CCH₂), 4.28 (t, J = 2.4 Hz, 2H, CCH₂); ¹³C NMR (DMSO-d₆), δ : 18.97 (2CH₂, SCH₂CC), 21.77 (CH₃, NCH₃), 23.85 (CH₂), 23.88 (CH₂), 25.76 (2CH₂), 25.81 (2CH₂), 33.94 (CH₂, CCH₂), 33.95 (CH₂, CCH₂), 47.41 (CH₂, NCH₂), 47.43 (CH₂, NCH₂), 52.67 (CH₂, NCH₂), 52.71 (CH₂, NCH₂), 79.14 (C, SCH₂CC), 79.75 (C, SCH₂CC), 80.38 (C, SCH₂CC), 80.43 (C, SCH₂CC), 125.54 (C, C-5), 143.10 (CH, C-8), 151.24 (C, C-2), 152.22 (C, C-6), 159.81 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₂₄H₃₂ClN₆S₂ 503.1818 found 503.1814.

2-Chloro-6,8-di(4-N-morpholinylbut-2-ynylthio)-7methylpurine 15c

It was obtained as a pale yellow solid (0.172 g, 70%); mp 112–113 °C (EtOH); ¹H NMR (DMSO-d₆), δ : 2.30 (m, 4H, 2NCH₂), 2.35 (m, 4H, 2NCH₂), 3.23 (m, 4H, SCH₂), 3.47 (m, 4H, 2OCH₂), 3.51 (m, 4H, 2OCH₂), 3.89 (s, 3H, NCH₃), 4.25 (t, J = 2.4 Hz, 2H, CCH₂), 4.29 (t, J = 2.4 Hz, 2H, CCH₂); ¹³C NMR (DMSO-d₆), δ : 18.89 (CH₂, SCH₂CC), 18.91 (CH₂, SCH₂CC), 21.73 (CH₃, NCH₃), 33.93 (CH₂, CCH₂), 33.94 (CH₂, CCH₂), 46.90 (CH₂, NCH₂), 46.96 (CH₂, NCH₂), 51.85 (CH₂, 2NCH₂), 66.46 (CH₂, 4OCH₂), 78.83 (C, SCH₂CC), 79.33 (C, SCH₂CC),

 $\begin{array}{l} 80.62 \ (C, \ SCH_2CC), \ 80.74 \ (C, \ SCH_2CC), \ 125.51 \ (C, \ C-5), \\ 143.05 \ (CH, \ C-8), \ 151.23 \ (C, \ C-2), \ 152.18 \ (C, \ C-6), \ 159.95 \\ (C, \ C-4); \ HR \ ESIMS \ m/z \ [M+H]^+ \ calcd. \ for \\ C_{22}H_{28}ClN_6O_2S_2 \ 507.1404 \ found \ 507.1398. \end{array}$

2-Chloro-6,8-di(diethylaminobut-2-ynylthio)-7methylpurine 15d

It was obtained as a pale yellow solid (0.194 g, 81%); mp 88–90 °C (EtOH); ¹H NMR (DMSO-d₆), δ : 0.87 (m, 12H, 4CH₃), 2.32 (m, 8H, 4NCH₂), 3.33 (s, 4H, 2SCH₂), 3.88 (s, 3H, NCH₃), 4.22 (t, J = 2.4 Hz, 2H, CCH₂), 4.27 (t, J = 2.4 Hz, 2H, CCH₂); ¹³C NMR (DMSO-d₆), δ : 12.76 (4CH₃,), 19.06 (CH₂, SCH₂CC), 19.09 (CH₂, SCH₂CC), 21.82 (CH₃, NCH₃), 33.93 (CH₂, CCH₂), 33.94 (CH₂, CCH₂), 46.83 (CH₂, 2NCH₂), 46.85 (CH₂, 2NCH₂), 78.41 (C, SCH₂CC), 79.17 (C, SCH₂CC), 80.19 (C, SCH₂CC), 80.36 (C, SCH₂CC), 125.97 (C, C-5), 143.09 (CH, C-8), 151.22 (C, C-2), 152.22 (C, C-6), 159.81 (C, C-4); HR ESIMS m/z [M + H]⁺ calcd. for C₂₂H₃₂ClN₆S₂ 479.1818 found 479.1813.

Antiproliferative assay in vitro

Cell culture

Compounds were evaluated for their anticancer activity using three cultured cell lines: SNB-19 (human glioblastoma, DSMZ - German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany), C-32 (human amelanotic melanoma, ATCC-American Type Culture Collection, Manassas, VA, USA), MDA-MB-231 (human adenocarcinoma mammary gland, ATCC, Manassas, VA, USA), and HFF-1 (human fibroblast cell line, ATCC, Manassas, VA, USA). The cultured cells were kept at 37 °C and 5% CO₂. The cells were seeded $(1 \times 10^4 \text{ cells/well/100})$ µL DMEM supplemented with 10% FCS and streptomycin and penicillin) using 96-well plates (Corning). The cells were counted in a hemocytometer (Burker's chamber) using a phase contrast Olympus IX50 microscope equipped with Sony SSC-DC58 AP camera and Olympus DP10 digital camera.

Proliferation assay

The antiproliferative effect of the compounds obtained from both the cancer and the normal cells was determined using the Cell Proliferation Reagent WST-1 assay (Roche Diagnostics, Mannheim, Germany). This colorimetric assay is based on the viable cell's ability to cause the bright redcolored stable tetrazolium salt (2-(4-iodophenyl)-3-(4nitrophenyl)-5-(2,4-disulfophenyl)-2H-tetrazolium, monosodium salt) to cleave to the dark red soluble formazan by cellular enzymes. An expansion in the number of viable cells results in an increase in the overall activity of mitochondrial dehydrogenases in the sample. An increase in the amount of formazan dye formed correlates to the number of metabolically active cells in the culture. The formazan dye produced by metabolically active cells is quantified by a scanning ELISA reader that measures the absorbance of the dve solution at appropriate wavelengths. The examined cells were exposed to the tested compounds for 72 h at various concentrations between 0.1 and 100 µg/mL (prepared initially at a concentration of 1 mg/mL in DMSO). The control was performed in order to check that DMSO has no effect on the cells at the concentration used. The cells were incubated with WST-1 (10 µL) for 1 h and the absorbance of the samples was measured against a background control at 450 nm using a microplate reader with a reference wavelength at 600 nm. The results are expressed as the means of at least two independent experiments performed in triplicate. The antiproliferative activity of the tested compound was compared to cisplatin. The IC₅₀ values (a concentration of a compound that is required for 50% inhibition) were calculated from the dose-response relationship with respect to control.

Result and discussion

Chemistry

2-Chloro-6-(prop-2-ynylthio)-7-methylpurine 1 was obtained via S-alkylation of the appropriate 6-purinethione with propargyl bromide (as solution in toluene) in DMF at room temperature in the presence of potassium tert-butoxide (Kowalska et al. 2015). Using this procedure, 2,6dipropynylthio-7-methylpurine 4 was obtained from 2,6purinethione 3. 2-Chloro-6-propynylthio-7-methylpurine 1 and 2.6-dipropynylthio-7-methylpurine 4, which were transformed via a Mannich reaction with paraformaldehyde and secondary amine (pyrrolidine, piperidine, morpholine, and diethylamine) in dry dioxane in the presence of catalytic amounts of CuCl into 2-chloro-6-aminobut-2-ynylthio-7-methylpurines 2a-c, as well as 2,6-di(aminobut-2-ynylthio)-7-methylpurines 5a-d in yields of 82-94% and 74-79.5%, respectively. Whereas 2- and 6-chloropurines are easily obtained from oxopurines, the introduction of the halogen atom in the 8 position requires halogenation with bromine (Koppel et al. 1962). Compounds 6 and 11 were brominated with bromine (as a solution in nitrobenzene) followed by a refluxing of the reaction mixture for 5 h. In the case of compound 11, the bromination proceeded not only in the 8 position but also in ipso in the 6 position. 8-Xanthinethione 8 and 8-purinethione 13 were obtained in the sodium hydrosulfide reaction. In order to obtain 8aminobutynylthioxanthines 10a-d and 6,8-diaminobutynylthiopurines 15a-d, the same sequence of propynylation and a Mannich reaction was used.

Scheme 1 Scheme 2 Scheme 3 Scheme 4

Anticancer activity

For the biological tests, 18 propynylthio- and aminobutynylthiopurines and xanthines of 4 types were selected:

a. 2-chloro-6-aminobutynylthio derivatives containing the piperidinyl, morpholinyl, and diethylamino groups (**2a**-**c**),

b. 2,6-dipropynylthio (4) and 2,6-diaminobutynylthiopurines containing the pyrrolidinyl, piperidinyl, morpholinyl, and diethylamino groups (5a-d),

c. 8-propynylthio (9) and 8-aminobutynylthioxanthines derivatives containing the pyrrolidinyl, piperidinyl, morpholinyl, and diethylamino groups (10a-d),

d. the 2-chloro-6,8-dipropynylthio (14) and 2-chloro-6,8diaminobutynylthiopurines containing pyrrolidinyl, piperidinyl, morpholinyl, and diethylamino groups (15a-d),

The anticancer activity of the compounds obtained was investigated in vitro using cultured human glioblastoma SNB-19, human adenocarcinoma MDA-MB-231, and melanoma C-32 cell lines. Normal human fibroblasts (HFF-1) were used as a control and cisplatin was used as a reference drug. Table 1 shows the activity of alkynylthiopurines and xanthines **2a-c**, **4**, **5a-d**, **9**, **10a-d**, **14**, and **15a-d** as the IC₅₀ values. The tested compounds exhibited different activities against three cancer cell lines. The SNB-19 and C-32 lines were more sensitive than the MDA-MB-231 line for most compounds.

As shown in Table 1, 6,8-disubstituted derivatives 14 and 15a-d were the most active group of the tested





Scheme 3 Synthesis of 8propynylthio- and 8aminobutynylthio-3,7dimethylxanthines 9 and 10a-d





compounds and in some cases were more or at least as active as cisplatin. 2-Chloro-6,8-dipropynylthiopurine **14** and its pyrrolidinylmethyl derivative **15a** exhibited strong antiproliferative activity (IC₅₀ values of <1 µg/mL) against the SBN-19 line. The last compound was also active against two other lines (IC₅₀ = 3.8–4.5 µg/mL). Compounds **15c** and **15d** (containing two morpholinylbutynylthio and die-thylaminobutynylthio groups) showed selective activity against the SNB-19 and C-32 lines (IC₅₀ = 2.9–5.3 µg/mL).

The activity of 2,6-disubstituted compounds (4 and its aminomethyl derivatives **5a-c**) were also selective. 2,6-Dipropynylthiopurine 4 was more active than cisplatin

against the MDA-MB-231 and C-32 cell lines (IC₅₀ \leq 1.77 µg/mL). The introduction of the pyrrolidinylmethyl and piperidinylmethyl moieties to the propynylthio chain (compounds **5a** and **5b**) slightly decreased activity against selected lines. On the other hand, derivative **5c** (the morpholinylmethyl moiety that was introduced) was very active (more so than cisplatin) with the IC₅₀ value of 0.35 µg/mL against the C-32 cell line.

2-Chloro-6-aminobutynyltiopurines **2a-c** were less active against all cell lines with IC_{50} values over 11.88 µg/mL.

The other monoalkynylthiosubstituted derivatives **9** and **10a-d** containing the 3,7-dimethylxanthine framework showed

Scheme 4 Synthesis of 2chloro-6,8-dipropynylthio- and 2-chloro-6,8diaminobutynylthio-7methylpurines 14 and 15a-d



 Table 1
 Antiproliferative activity of alkynylthiopurines and xanthines

Compounds	Antiproliferative activity IC ₅₀ (µg/mL)			
	SNB-19	C-32	MDA-MB-231	HFF-1
2a	28.80	21.01	33.40	35.42
2b	35.40	125.00	66.28	20.09
2c	17.40	11.88	48.06	8.65
4	15.01	1.77	1.00	10.00
5a	19.41	6.81	19.95	10.00
5b	4.64	6.39	7.35	3.98
5c	21.54	0.35	50.80	0.37
5d	48.93	23.86	38.49	5.95
9	191.10	103.46	169.34	153.65
10a	176.62	51.08	471.50	34.26
10b	166.17	355.00	363.50	30.39
10c	1.69	222.72	6.54	5.87
10d	1.48	26.36	68.12	3.68
14	0,07	270.35	10.00	20.09
15a	0.52	4.55	3.83	3.63
15b	11.62	19.08	38.07	5.50
15c	26.10	4.08	74.05	36.38
15d	2.91	5.33	10.00	1.00
Cisplatin	1.46	5.71	11.85	1.21

very different activity. In contrast to the very active 2,6- and 6,8-dipropynylthio derivatives **4** and **14**, 8-propynylthio derivative **9** was found inactive ($IC_{50} > 100 \mu g/mL$). Surprisingly,

its derivatives **10d** and **10c** (containing the diethylaminobutynylthio and morpholinylbutynylthio groups) exhibited strong activity against the SBN-19 line (IC₅₀ < $1.7 \mu g/mL$). Two other derivatives (**10a** and **10b**) were found rather inactive.

All the compounds were tested for their cytotoxicity on the normal human fibroblasts (HFF-1). Some compounds (**5b-d**, **10c-d**, **15a-b**, and **15d**) as well as cisplatin also turned out to be cytotoxic for fibroblasts. Very active compounds **4** (against adenocarcinoma MDA-MB-231 and melanoma C-32), **14** (against glioblastoma SNB-19), and **15c** (against melanoma C-32) were found to have relatively weak cytotoxicity.

Conclusion

While searching for alkynylthiopurines and aminoalkynylthiopurines and xanthines with the anticancer activity, we worked out a synthesis of 18 new multisubstituted 7methylpurines and 3,7-dimethylxanthines containing one or two propynylthio groups (in 2, 6, and 8 positions), which were further transformed via a Mannich reaction with cyclic and open chain amines and formaldehyde into aminobutynylthio derivatives. The products obtained represent various types of the purine and xanthine structure containing the triple bond: 8-mono-, 2,6-, and 6,8-dipropynylthio derivatives **4**, **9**, and **14**, 6- and 8-monoaminobutynylthio derivatives **2a-c** and **10a-d**, 2,6- and 6,8-diaminobutynylthio derivatives **5a-d** and **15a-d**. All tested compounds exhibited different anticancer activity against human glioblastoma SNB-19, human adenocarcinoma MDA-MB-231, and melanoma C-32 cell lines depending on the nature of the substituent and its localization in the purine framework. Some compounds exhibited stronger or similar anticancer activity to cisplatin. A small number of compounds were also cytotoxic against the normal human fibroblasts (HFF-1). The most promising compounds with strong anticancer activity and relatively low cytotoxicity turned out to be 2,6-dipropynylthio-7-methylpurine **4**, 2-chloro-6,8-dipropynylthio-7-methylpurine **14**, and 2-chloro-6,8-di(N-morpho-linylbutynylthio)-7-methylpurine **15c**.

Acknowledgements The synthesis and the structure elucidation are supported by the Medical University of Silesia (grant KNW-1-006/K/ 6/O.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://crea tivecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Andersen G, Gundersen LL, Nissen-Meyer J, Rise F, Spilberg B (2002) Cytotoxic and antibacterial activity of 2-oxopurine derivatives. Bioorg Med Chem Lett 12:567–569
- Atamanyuk D, Denis A, Gerusz V, Ledoussal B, Bonvin Y, Desroy N, Gold J, Moreau F, Oxoby M (2012) Purine derivatives and their use as pharmaceuticals for prevention or treatment of bacterial infections. European Patent WO2012172043 (A1)
- Baraldi PG, Cacciari B, de Las Infantas MJP, Romagnoli R, Spalluto G, Volpini R, Costanzi S, Vittori S, Cristalli G, Melman N, Park KS, Ji X, Jacobson KA (1998) Synthesis and biological activity of a new series of N⁶-arylcarbamoyl, 2-(Ar)alkynyl- N⁶-arylcarbamoyl, and N⁶-carboxamido derivatives of adenosine-5'-Nethyluronamide as A₁ and A₃ adenosine receptor agonists. J Med Chem 41:3174–3185
- Berg TC, Gundersen LL, Eriksen AB, Malterud KE (2005) Synthesis of optically active 6-alkynyl- and 6-alkylpurines as cytokinin analogs and inhibitors of 15-lipoxygenase; studes of intramolecular cyclization of 6-(hydroxyalkyn-1-yl)purines. J Org Chem 23:4988–4994
- Bliman D, Pettersson M, Bood M, Grotli M (2014) 8-Bromination of 2,6,9-trisubstituted purines with pyridinium tribromide. Tetrahedron Lett 55:2929–2931
- Brathe A, Andersen G, Gundersen LL, Malterud KE, Rise F (2002) Antioxidant activity of synthetic cytokinin analogues: 6-alkynyland 6-alkenylpurines as novel 15-lipoxygenase inhibitors. Bioorg Med Chem 10:1581–1586
- Brathe A, Gundersen LL, Nissen-Meyer J, Rise F, Spilsberg B (2003) Cytotoxic activity of 6-alkynyl- and 6-alkenylpurines. Bioorg Med Chem Lett 13:877–880

- Broom AD, Gall AA, Meyer RB (1997) Oligonucleotides containing n-methyl thiolated bases having antiviral activity. US Patent 5652359A
- Carson DA, Cottam HB, Wasson DB (1998) Compounds for inhibition of ceramide-mediated signal transduction. US Patent 5843943
- Corrales RCNR, de Souza NB, Pinheiro LS, Abramo C, Coimbra ES, da Silva AD (2011) Thiopurine derivatives containing triazole and steroid: synthesis, antimalarial and antileishmanial activities. Biomed Pharmacother 65:198–203
- Cristalli G, Podda GM, Costanzi S, Lambertucci C, Lecchi A, Vittori S, Volpini R, Zighetti ML, Cattaneo M (2005) Effects of 5'phosphate derivatives of 2-hexynyl adenosine and 2phenylethynyl adenosine on responses of human platelets mediated by P2Y receptors. J Med Chem 48:2763–2766
- Daly JW (2007) Caffeine analogs: biomedical impact. Cell Mol Life Sci 64:2153–2169
- Dal Ben D, Buccioni M, Lambertucci C, Marucci G, Volpini R, Cristalli G (2011) The importance of alkynyl chain presence for the activity of adenine nucleosides/nucleotides on purinergic receptors. Curr Med Chem 18:1444–1463
- Deswal S, Srivastava A (2017) Role of allopurinol in optimizing thiopurine therapy in patients with autoimmune hepatitis. J Clin Exp Hepatol 7:55–62
- Dimopoulou A, Manta S, Parmenopoulou V, Kollatos N, Christidou O, Triantakonstanti VV, Schols D, Komiotis D (2015) An easy microwave-assisted synthesis of C8-alkynyl adenine pyranonucleosides as novel cytotoxic antitumor agents. Front Chem 3:21
- Endo K, Deguchi K, Matsunaga H, Tomaya K, Yamada K (2014) 8-Substituted 2-alkynyl-N⁹-propargyladenines as A_{2A} adenosine receptor antagonists. Bioorg Med Chem 22:3072–3082
- Foitzik RC, Devine SM, Hausler NE, Scammells PJ (2009) Linear and convergent approaches to 2-substituted adenosine-5'-N-alkylcarboxamides. Tetrahedron 65:8851–8857
- Herreras J, Iborra M (2017) When should thiopurine therapy during sustained remission in inflammatory bowel disease be stopped? Adv Res Gastroentero Hepatol 4:555647
- Ivachtchenko AV, Mitkin OD, Kadieva MG, Okun IM (2013) Substituted phenoxyacetic acids and esters and amides thereof, comprising a 2,6-dioxo-2,3,6,7-tetrahydro-1H-purine-8-yl fragment and constituting A_{2A} adenosine receptor antagonists, and the use thereof. European Patent WO2013058681
- Koles L, Furst S, Illes P (2007) Purine ionotropic (P2X) receptors. Curr Pharm Des 13:2368–2384
- Koppel HC, Springer RH, Robins RK, Schneider FH, Cheng CC (1962) Synthesis of some 8-substituted bis(β-chloroethyl)amino derivatives of naturally occurring N-methylated purines. J Org Chem 27:2173–2177
- Kowalska A (2007) Synthesis of bis(7-methyl-2- or 6-purinyl) disulfides. Phosphorus Sulfur Silicon 182:2881–2890
- Kowalska A, Latocha M, Pluta K (2015) Synthesis and anticancer activity of thiosubstituted purines. Med Chem Res 24:3107–3116
- Kowalska A, Maślankiewicz A, Syrek B, Ciepliński P (1985) 2,6-Dialkoxy-7-methylpurines. Mon Chem 116:341–351
- Kuchar M, Hocek M, Pohl R, Votruba I, Shih I, Mabery E, Mackman R (2008) Synthesis, cytostatic, and antiviral activity of novel 6-[2-(dialkylamino)ethyl]-, 6-(2-alkoxyethyl)-, 6-[2-(alkylsulfanyl) ethyl]-, and 6-[2-(dialkylamino)vinyl]purine nucleosides. Bioorg Med Chem 16:1400–1424
- Legraverend M, Tunnah P, Noble M, Ducrot P, Ludwig O, Grierson DS, Leost M, Meijer L, Endicott J (2000) Cyclin-dependent kinase inhibition by new C-2 alkynylated purine derivatives and molecular structure of a CDK2- inhibitor compex. J Med Chem 43:1282–1292

- Prima FD, Shenoy A, Moses SR, Shabaraya AR (2013) Antitumor activity of mercaptopurine in combination with trikatu and gomutra on 20-methylcholantrene induced carcinogenesis. J Appl Pharm Sci 3:20–23
- Riva-Toniolo C, Muller S, Schaub J, Brill WKD (2003) Catalysis of nucleophilic aromatic substitutions in the 2,6,8-trisubstituted purines and application in the synthesis of combinatorial libraries. Mol Div 6:43–53
- Sharma S, Singh J, Ojha R, Singh H, Kaur M, Bedi PMS, Nepali K (2016) Design strategies, structure activity relationship and mechanistic insights for purines as kinase inhibitors. Eur J Med Chem 112:298–346
- Thomas R, Lee J, Chevalier V, Sadler S, Selesniemi K, Hatfield S, Sitkovsky M, Ondrehen NJ, Jones GB (2013) Design and evaluation of xanthine based adenosine receptor antagonists: potential hypoxia targeted immunotherapies. Bioorg Med Chem 21:7453–7464
- Ward MG, Patel KV, Kariyawasam VC, Goel R, Warner B, Elliott TR, Blaker PA, Irving PM, Marinaki AM, Sanderson JD (2017) Thioguanine in inflammatory bowel disease: long-term efficacy and safety. United Eur. Gastroenterol J 5:563–570
- Warner B, Johnston E, Arenas-Hernandez M, Marinaki A, Irving P, Sanderson J (2016) A practical guide to thiopurine

prescribing and monitoring in IBD. Frontline Gastroenterol 0: 1-6

- Volpini R, Costanzi S, Lambertucci C, Vittori S, Klotz KN, Lorenzen A, Cristalli G (2001) Introduction of alkynyl chains on C-8 of adenosine led to very selective antagonists of the A₃ adenosine receptor. Bioorg Med Chem Lett 11:1931–1934
- Volpini R, Costanzi S, Lambertucci C, Vittori S, Cristalli G (2002) Purine nucleosides bearing 1-alkynyl chains as adenosine receptor agonists. Curr Pharm Design 8:99–110
- Volpini R, Costanzi S, Lambertucci C, Vittori S, Martini C, Trincavelli ML, Klotz KN, Cristalli G (2005) 2-And 8-alkynyl-9-ethyladenines: synthesis and biological activity at human and rat adenosine receptors. Purinergic Signal 1:173–181
- Zablocki AJ, Palle VP, Elzein EO, Nudelman G (2001) Propargyl phenyl ether A_{2A} receptor agonists. US Patent 6180615 B1
- Zagorska A, Kołaczkowski M, Bucki A, Siwek A, Kazek G, Satała G, Bojarski AJ, Partyka A, Wesołowska A, Pawłowski M (2015) Structure-activity relationships and molecular studies of novel arylpiperazinylalkyl purine-2,4-diones and purine-2,4,8-triones with antidepressant and anxiolytic-like activity. Eur J Med Chem 97:142–154