

A new homobimetallic arenediyl diplatinum(II) unit as building block for macromolecular synthesis. X-ray crystal structure of $[C_6H_2\{CH_2NMe_2\}_2-1,5-\{PtCl(PPh_3)\}_2-2,4]$

Michel D. Meijer^a, Arjan W. Kleij^a, Martin Lutz^b, Anthony L. Spek^{b,*}, Gerard van Koten^{a,*}

^a Debye Institute, Department of Metal-Mediated Synthesis, Utrecht University, Padualaan 8, 3584 CH, Utrecht, The Netherlands

^b Department of Crystal and Structural Chemistry, Bijvoet Center for Biomolecular Research, Utrecht University, Padualaan 8, 3584 CH Utrecht The Netherlands

Received 11 June 2001; accepted 28 July 2001

Abstract

Treatment of the diaminobenzene $[C_6H_4\{CH_2NMe_2\}_2-1,3]$ (NCN-H, **1**) with one or two equivalents of *cis*-PtCl₂(DMSO)₂ leads to exclusive formation of the doubly cycloplatinated species $[C_6H_4\{CH_2NMe_2\}_2-1,5-\{PtCl(DMSO)\}_2-2,4]$ (**3**), which upon addition of triphenylphosphine yields the bisphosphine adduct $[C_6H_4\{CH_2NMe_2\}_2-1,5-\{PtCl(PPh_3)\}_2-2,4]$ (**4**). The X-ray molecular structure of **4** revealed the presence of highly distorted square planar Pt(II) centers which is caused by close proximity of the two phosphine donor ligands. Complexes of type **3** can be regarded as suitable starting materials for the directional build-up of larger macromolecular structures. © 2001 Published by Elsevier Science B.V.

Keywords: Arylamine ligands; Building blocks; C–H activation; Platinum

1. Introduction

Macromolecular synthesis with organometallic complexes is a field of research that is increasingly investigated with the ultimate goal of creating novel materials with unprecedented properties [1]. Prerequisites for the build up of large metal-containing structures seem to be the synthetic availability and stability of such metalated synthons [2]. Additionally, these organometallic complexes should include weak donor ligands which upon exchange with polydentate Lewis bases can give rise to self-assembled structures. Recently, the preparation of metalated (*N*),*C*,*N*- [3,4] or *S*,*C*,*S*-chelating [5] aryl ligands suitable as fragments of larger synthetic supramolecules has been reported. Simultaneously, the use of platinum(II) dimethylsulfoxides as mild and versatile metalating reagents for introduction of Pt(II)

centers onto aromatic ligands has been developed [4,6]. Herein, we report our first results on the synthesis of a doubly cycloplatinated diaminoarene ligand by treatment of $C_6H_4\{CH_2NMe_2\}_2-1,3$ (**1**, NCN-H) with *cis*-PtCl₂(DMSO)₂, and the subsequent exchange of the pendant dimethylsulfoxide ligands for triphenylphosphine donor ligands.

2. Results and discussion

2.1. Synthesis and characterization of diplatinated species

Direct cycloplatination of the NCN-H ligand (**1**) with *cis*-PtCl₂(DMSO)₂ in a 1:1 molar ratio was initially aimed at the activation of the 1-position (H_a, Scheme 1), which would afford known **2**. To this end, equimolar amounts of *cis*-PtCl₂(DMSO)₂ and **1** (Fig. 1) were mixed in refluxing MeOH in the presence of NaOAc as an external base. After initial dissolution of the platinum salt, a white precipitate was formed, which was isolated (48% yield) and characterized by NMR spec-

* Corresponding authors. Tel.: +31-30-2533120; fax: +31-30-2523615 (G. van Koten).

E-mail addresses: a.l.spek@chem.uu.nl (A.L. Spek), g.vankoten@chem.uu.nl (G. van Koten).


troscopy. The ^1H -NMR spectrum displayed the presence of a single organometallic product, which to our surprise was identified as the doubly platinated NCN derivative $[\text{C}_6\text{H}_2\{\text{CH}_2\text{NMe}_2\}_{2-1,5-\{\text{PtCl}(\text{DMSO})\}_2-2,4}]$ (**3**). Illustrative in the ^1H -NMR spectrum of **3** is the presence of a low-field, singlet resonance at 8.10 ppm with platinum couplings ($^3J(\text{Pt}-\text{H}) = 51.8$ Hz), which was attributed to H_b (Scheme 1). Additionally, H_a was observed as a single resonance at 6.78 ppm.

Remarkably, no *mono*-metalated species were detected. Earlier results in analogous palladation reactions of the NCN-H ligand with $\text{Pd}(\text{OAc})_2$ and $\text{Li}_2[\text{PdCl}_4]$ also yielded biscyclometalated complexes [3d], [7]. These results suggest strongly that the intermediate monometalated species is more susceptible to electrophilic metalation than NCN-H itself, affording exclusively the bismetalated product. Moreover, the observed regioselectivity suggests that the 1-position of the NCN ligand is more sterically hindered with respect to the 3- and 5-position [8]. We also carried out the analogous cycloplatination reaction of NCN-H (**1**) in the presence of two equivalents of the platinum(II) salt. As expected, **3** was formed selectively (78% isolated yield) and no other platinum species could be detected.

Complex **3** proved to be rather insoluble in common organic solvents. Therefore, a PPh_3 -for-DMSO ligand exchange in CH_2Cl_2 was carried out. This procedure afforded the bisphosphine complex **4** in almost quantitative yield as white needles. Complex **4** was analyzed using multinuclear NMR spectroscopy, elemental analyses and also by a single crystal structure determination (see Section 2.2). Of particular note is the high field shift of $\text{Ar}-\text{H}_b$ positioned at 5.77 ppm (cf., 8.10 ppm in **3**). It seems likely that this high field shift arises from specific shielding caused by the π -electron density of the phenyl rings of each of the PPh_3 ligands in a close proximity to H_b . The proposed *trans*-position of both phosphine ligands (with respect to the NMe_2 donor fragment) in **4** is supported by ^{31}P -NMR spectroscopy, which shows a single singlet flanked by platinum satellites at 15.8 ppm, $^1J(\text{Pt}-\text{P}) = 4377$ Hz. This coupling constant closely relates to previously reported data on similar complexes [4].

2.2. X-ray crystal structure of **4**

Compound **4** was subjected to an X-ray crystal structure determination. The molecular structure is shown in



Scheme 1. Cycloplatination of the NCN-H ligand (**1**). Reagents and conditions: (i) $\text{PtCl}_2(\text{DMSO})_2$, NaOAc, MeOH, 65 °C; (ii) PPh_3 , CH_2Cl_2 .




Fig. 1. Displacement ellipsoid plot (50% probability level) of the molecular structure of **4**, with the adopted numbering scheme. All hydrogen atoms have been omitted for clarity. Selected bond lengths (Å) and angles (°): Pt(1)–Cl(1) 2.4007(8), Pt(1)–P(1) 2.2242(7), Pt(1)–N(1) 2.141(3), Pt(1)–C(1) 2.016(3), Pt(2)–Cl(2) 2.4007(6), Pt(2)–P(2) 2.2268(7), Pt(2)–N(2) 2.159(2), Pt(2)–C(3) 2.009(2), C(1)–Pt(1)–N(1) 80.02(10), N(1)–Pt(1)–Cl(1) 90.70, Cl(1)–Pt(1)–P(1) 90.68(3), P(1)–Pt(1)–C(1) 98.75(8), C(1)–Pt(1)–Cl(1) 169.58(8), N(1)–Pt(1)–P(1) 177.68(6), C(3)–Pt(2)–P(2) 96.83(7), P(2)–Pt(2)–Cl(2) 94.38(2), Cl(2)–Pt(2)–N(2) 90.40(6), N(2)–Pt(2)–C(3) 79.99(9), C(3)–Pt(2)–Cl(2) 165.12(7), N(2)–Pt(2)–P(2) 170.54(6).

Fig. 1. Relevant bond angles and distances have been deposited as Section 5.

The two platinum centers are positioned opposite to the central arene plane, 0.221(1) Å above the plane for Pt1 and 0.253 Å below the plane for Pt2, respectively. They are in distorted planar environments with dihedral angles of 5.15(9) and 14.06(11)° between the N–Pt–C and P–Pt–Cl planes for Pt1 and Pt2, respectively. This distortion is caused by the strain of the two five membered chelate rings. Interestingly, these chelate rings have different conformations: the first ring is best described as a twist conformation with N1 0.774(2) Å above the plane of the central arene ring; the second ring is in an envelope conformation with N2 1.212(2) Å below the central arene plane. The N–Pt–C bite angles of 80.03(9) and 79.98(9)° are significantly smaller than the ideal values. The two PPh₃ ligands are located *trans* to the amino groups. The conformations of the PPh₃ groups slightly differ: the first ligand is in a *gauche* conformation with respect to the Pt1–Cl1 bond, while the second ligand is in a *staggered* conformation to Pt2–Cl2. We assume that the PPh₃ ligands behave like ‘cogwheels’ and that this interaction between them determines the conformation. Moreover, in this conformation, a close proximity between the aromatic hydrogen H2 and the two phenyl groups of the PPh₃ ligands is present, as supported by the NMR spectroscopic studies. The distance between H2 and C31 is 2.79 Å, which is significantly shorter than the sum of the covalent radii (C + H = 2.90 Å). Moreover, the H2–C30 (3.03 Å) and H2–C25 (2.23 Å) distances are indicative for the close proximity of H2 and the phenyl groups.

3. Conclusions

The presented results show that bisplatinum(II) derivatives of the diaminoarene ligand NCN-H are easily accessible via a direct and selective cycloplatination route. Moreover, recent experiments have unambiguously demonstrated the chemical stability of complexes of type **4** in subsequent synthesis [4c,d]. Future research programmes are focused on application of the presented building blocks in macromolecular synthesis by using suitable ligands with a predefined directional character.

4. Experimental

4.1. General

All chemicals were purchased from Acros/Aldrich and used without further purification. Solvents were distilled from appropriate drying agents *prior* to use.

The starting materials *cis*-PtCl₂(DMSO)₂ [6] and NCN-H (**1**) [9] were prepared according to literature. NMR spectroscopic measurements were carried out on a Varian Inova 300 MHz spectrometer at room temperature (r.t.). Chemical shifts are in ppm and referenced to residual solvent resonances. Elemental analyses were performed by Dornis and Kolbe, Mikroanalytisch Labor, Mülheim a/d Ruhr, Germany.

4.2. [C₆H₂{CH₂NMe₂}₂-1,5-{PtCl(DMSO)}₂-2,4] (**3**)

To a solution of C₆H₄(CH₂NMe₂)₂-1,3 (0.32 g, 1.70 mmol) in MeOH (40 ml) was added solid *cis*-PtCl₂(DMSO)₂ (1.45 g, 3.43 mmol) and NaOAc (0.29 g, 3.45 mmol). The reaction mixture was stirred at 65 °C and within 15 min a clear yellow solution was obtained. The solution was stirred at 65 °C for 3 h whereupon a white solid precipitated from solution. The product was collected by decantation, washed with MeOH (50 ml), Et₂O (2 × 50 ml) and dried in vacuo to yield a white solid (1.07 g, 78%). ¹H-NMR (CDCl₃): δ 8.10 (s, ³J(Pt–H) = 51.8 Hz, 1H, Ar-*H*_{ortho}), 6.78 (s, 1H, Ar-*H*), 3.89 (s, ³J(Pt–H) = 40.6 Hz, 4H, CH₂N), 3.54 (s, ³J(Pt–H) = 23.4 Hz, 12H, SCH₃), 2.90 (s, ³J(Pt–H) = 33.2 Hz, 12H, NCH₃). ¹³C{¹H} NMR (CDCl₃): δ 142.5, 139.6 (²J(Pt–C) = 65.9 Hz), 134.9, 115.2 (³J(Pt–C) = 40.2 Hz), 74.8 (²J(Pt–C) = 54.8 Hz, CH₂N), 52.0 (NCH₃), 46.4 (²J(Pt–C) = 65.5 Hz, SCH₃). The compound proved to be insoluble for a decent ¹³C{¹H} NMR spectroscopic analysis. Anal. Calcd. for C₁₆H₃₀Cl₂N₂O₂Pt₂S₂: C, 23.79; H, 3.74; N, 3.47. Found: C, 23.81; H, 3.72; N, 3.39%.

4.3. [C₆H₂{CH₂NMe₂}₂-1,5-{PtCl(PPh₃)}₂-2,4] (**4**)

To a suspension of **1** (0.18 g, 0.19 mmol) in CH₂Cl₂ (10 ml) was added PPh₃ (0.16 g, 0.61 mmol) dissolved in CH₂Cl₂ (5 ml). The suspension gradually turned into a clear, yellow solution. After 4 h, Et₂O (60 ml) was added to the mixture whereupon white crystals separated from the solution, which were dried in vacuo (0.22 g, 95%). ¹H-NMR (CDCl₃): δ 7.39–7.24 (m, 20H, PAr-*H*), 7.14–7.06 (m, 10H, PAr-*H*), 6.82 (s, 1H, Ar-*H*), 5.77 (s, ⁴J(P–H) = 5.20 Hz, ³J(Pt–H) = 55.0 Hz, 1H, Ar-*H*_{ortho}), 3.94 (s, 4H, ³J(Pt–H) = 30.0 Hz, CH₂N), 2.70 (d, ³J(Pt–H) = 31.4 Hz, ³J(P–H) = 1.8 Hz, 12H, NCH₃). ¹³C{¹H} NMR (CDCl₃): δ 146.7 (²J(P–C) = 4.8 Hz), 140.2, 137.8 (¹J(P–C) = 7.0 Hz, PAr-*C*_{ipso}), 134.9 (²J(P–C) = 11.8 Hz, PAr-*C*_{ortho}), 131.3, 130.5, 129.8 (⁴J(P–C) = 2.5 Hz, PAr-*C*_{para}), 127.4 (³J(P–C) = 10.9 Hz, PAr-*C*_{meta}), 115.1 (³J(Pt–C) = 34.8 Hz), 73.9 (²J(Pt–C) = 50.7 Hz, CH₂N), 50.1 (NMe₂). ³¹P{¹H}-NMR (CDCl₃, 81 MHz): δ 15.8 (¹J(Pt–P) = 4377 Hz). Anal. Calcd. for C₄₈H₄₈Cl₂N₂P₂Pt₂: C, 49.03; H, 4.11; N 2.38. Found: C, 48.88; H, 4.18; N, 2.31%.

4.4. Crystal structure determination of **4**

Complex 4: $\text{C}_{48}\text{H}_{48}\text{Cl}_2\text{N}_2\text{P}_2\text{Pt}_2\cdot\text{C}_4\text{H}_{10}\text{O}$, $F_w = 1250.02$, colourless needle, $0.48 \times 0.09 \times 0.06 \text{ mm}^3$, monoclinic, $P2_1/c$ (No. 14), $a = 13.2504(1)$, $b = 23.3587(2)$, $c = 17.2594(2) \text{ \AA}$, $\beta = 116.5033(6)^\circ$, $V = 4780.59(8) \text{ \AA}^3$, $Z = 4$, $\rho = 1.737 \text{ g cm}^{-3}$, 76701 measured reflections, 10999 unique reflections ($R_{\text{int}} = 0.051$). Intensities were measured on a Nonius Kappa CCD diffractometer with rotating anode (Mo– $\text{K}\alpha$, $\lambda = 0.71073 \text{ \AA}$) at a temperature of 150 K. Analytical absorption correction with the program PLATON [10] (routine TOMPA, $\mu = 6.06 \text{ mm}^{-1}$, 0.19–0.82 transmission). The structure was solved with automated Patterson methods with the program DIRDIF97 [11] and refined with the program SHELXL97 [12] against F^2 of all reflections up to a resolution of $(\sin \theta/\lambda)_{\text{max}} = 0.65 \text{ \AA}^{-1}$. Non-hydrogen atoms were refined freely with anisotropic displacement parameters, hydrogen atoms were refined as rigid groups. The crystal structure contains disordered diethyl ether solvent molecules. Their contribution to the structure factors was secured by back-Fourier transformation (program PLATON [10], CALC SQUEEZE, 172 e^- /unit cell in voids of $635.7 \text{ \AA}^3$ per unit cell). R ($I > 2\sigma(I)$): $R1 = 0.0209$, $wR2 = 0.0433$. R (all data): $R1 = 0.0269$, $wR2 = 0.0446$. $S = 1.036$. The drawings, structure calculations, and checking for higher symmetry were performed with the program PLATON [10].

5. Supplementary material

Crystallographic data for the structural analysis have been deposited with the Cambridge Crystallographic Data Centre, CCDC no. 163055 for compound **4**. Copies of this information may be obtained free of charge from The Director, CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (Fax: +44-1223-336033; e-mail: deposit@ccdc.cam.ac.uk or www: http://www.ccdc.cam.ac.uk)

Acknowledgements

The authors thank Dr P. Steenwinkel for valuable experimental assistance and stimulating discussions and The Council for Chemical Sciences (CW) of the Dutch

Organization for Scientific Research (NWO) for financial contribution.

References

- [1] (a) M. Fujita, Chem. Soc. Rev. 27 (1998) 417;
(b) P.J. Stang, B. Olenyuk, Acc. Chem. Res. 30 (1997) 502;
(c) S. Leininger, B. Olenyuk, P.J. Stang, Chem. Rev. 100 (2000) 853.
- [2] M. Albrecht, R.A. Gossage, M. Lutz, A.L. Spek, G. van Koten, Chem. Eur. J. 6 (2000) 1431. Recently, we reported on the use of highly stable organoplatinum complexes in the construction of larger (dendritic) macromolecules.
- [3] (a) S. Trofimenko, Inorg. Chem. 12 (1973) 1215;
(b) S. Chakladar, P. Paul, A.K. Mukherjee, S.K. Dutta, K.K. Nanda, D. Podder, K. Nag, J. Chem. Soc., Dalton Trans. (1992) 3119;
(c) M. López-Torres, A. Fernández, J.J. Fernández, A. Suárez, S. Castro-Juiz, J.M. Vila, M.T. Pereira, Organometallics 20 (2001) 1350;
(d) P. Steenwinkel, R.A. Gossage, T. Maunala, D.M. Grove, G. van Koten, Chem. Eur. J. 4 (1998) 763.
- [4] (a) A.W. Kleij, R.J.M. Klein Gebbink, P.A.J. van den Nieuwenhuijzen, H. Kooijman, M. Lutz, A.L. Spek, G. van Koten, Organometallics 20 (2001) 634;
(b) A.W. Kleij, R.J.M. Klein Gebbink, M. Lutz, A.L. Spek, G. van Koten, J. Organomet. Chem. 621 (2001) 190;
(c) M.D. Meijer, E. de Wolf, M. Lutz, A.L. Spek, G.P.M. van Klink, G. van Koten, G. Organometallics 20 (2001) 4198;
(d) M.D. Meijer, A.W. Kleij, B.S. Williams, D. Ellis, M. Lutz, A.L. Spek, G.P.M. van Klink, G. van Koten, Organometallics, in press.
- [5] (a) J.R. Hall, S.J. Loeb, G.K.H. Shimizu, G.P.A. Yap, Angew. Chem. Int. Ed. 37 (1997) 121;
(b) S.J. Loeb, G.K.H. Shimizu, J. Chem. Soc., Chem. Commun. (1993) 1395.
- [6] P.R.R. Ranatunge-Bandarage, B.H. Robinson, J. Simpson, Organometallics 13 (1994) 500.
- [7] Note that reaction of NCN-H with $\text{Pd}(\text{OAc})_2$ yielded mixtures of mono- and bispalladated products, dependent on the experimental conditions, see [3d].
- [8] For selective metalation on the 1-position, this position should be activated prior to the transmetalation reaction, for example, by introducing 1-SiMe₃ groups or by C–H activation at the 1-position by lithium reagents (see, for example, [2] and [3d]).
- [9] P. Steenwinkel, Multinuclear Organometallics Based on Aryldiamine Ligands, PhD thesis Utrecht University, Utrecht (1998).
- [10] A.L. Spek, PLATON. A multipurpose crystallographic tool. Utrecht University, The Netherlands, 2000.
- [11] P.T. Beurskens, G. Admiraal, G. Beurskens, W.P. Bosman, S. Garcia-Granda, R.O. Gould, J.M.M. Smits, C. Smykalla, The DIRDIF97 program system, Technical Report of the Crystallography Laboratory, University of Nijmegen, The Netherlands, 1997.
- [12] G.M. Sheldrick, SHELXL-97. Program for crystal structure refinement. University of Göttingen, Germany, 1997.