


This article was downloaded by: [University of New Mexico]

On: 22 November 2014, At: 16:15

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

### REACTIONS OF CYCLIC AZA-YLIDES WITH $\beta$ -DICARBONYL COMPOUNDS

Tetsuya Uchiyama<sup>a</sup>, Masayuki Inoue<sup>a</sup>, Hideki Masuda<sup>a</sup>, Tetsuya Fujimoto<sup>a</sup> & Iwao Yamamoto<sup>a</sup>

<sup>a</sup> Department of Functional Polymer Science, Faculty of Textile Science and Technology, Shinshu University, Ueda, Nagano, 386-8567, Japan  
Published online: 04 Oct 2006.

To cite this article: Tetsuya Uchiyama, Masayuki Inoue, Hideki Masuda, Tetsuya Fujimoto & Iwao Yamamoto (2001) REACTIONS OF CYCLIC AZA-YLIDES WITH  $\beta$ -DICARBONYL COMPOUNDS, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 175:1, 99-108, DOI: [10.1080/10426500108040259](https://doi.org/10.1080/10426500108040259)

To link to this article: <http://dx.doi.org/10.1080/10426500108040259>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or

indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

# REACTIONS OF CYCLIC AZA-YLIDES WITH $\beta$ -DICARBONYL COMPOUNDS

TETSUYA UCHİYAMA, MASAYUKI INOUE, HIDEKI MASUDA,  
TETSUYA FUJIMOTO and IWAO YAMAMOTO\*

*Department of Functional Polymer Science, Faculty of Textile Science and Technology, Shinshu University, Ueda, Nagano 386-8567 Japan*

*(Received November 10, 2000.; In final form January 25, 2001)*

The reactions of five- and six-membered cyclic iminophosphoranes, **3** and **4**, with  $\beta$ -dicarbonyl compounds gave enamino phosphine oxide derivatives **5a-h** in good yields. Compounds **5b** and **5e** were stereoselectively converted to 1,6- and 1,7-diene **9a** and **9b**.

**Keywords:** cyclic iminophosphoranes;  $\beta$ -dicarbonyl compounds; aza-Wittig reaction; Horner-Wittig Reaction

## INTRODUCTION


Cyclic iminophosphoranes **3** and **4**<sup>[1]</sup> have two different kinds of nucleophilic atoms, i.e., carbon and nitrogen on the  $\alpha$  position of the phosphorus atom. Recently, we reported tandem reactions with alkylating agents<sup>[2]</sup> using these two nucleophilic sites. Furthermore, the reactions of cyclic iminophosphoranes at the nitrogen nucleophilic site with  $\alpha$ -diketones gave imino phosphine oxides in good yields. Also, the imino phosphine oxides were treated with LDA in THF to produce the cyclic imines<sup>[3]</sup>.

In this paper, we describe the reactions of cyclic iminophosphoranes **3** and **4** with various  $\beta$ -dicarbonyl derivatives having an active methylene group.

\* Corresponding Author.

## RESULTS AND DISCUSSION

A reaction of cyclic aza-ylide **3**, generated from aminophosphonium salt **1** in the presence of sodium hydride, with 2,4-pentanedione gave enamino-phosphine oxide derivatives **5a** in 68% yields. Similarly, enamino-phosphine oxide derivatives **5b-5f** were obtained in good yields. (TABLE I)

TABLE I Reaction of Cyclic Iminophosphoranes with  $\beta$ -dicarbonyl Derivatives

| Entry | Ylides | $R_1$           | $R_2$   | Base   | Products | Yields <sup>a</sup> | ratio <sup>b</sup> |
|-------|--------|-----------------|---|--------|----------|---------------------|--------------------|
| 1     | 3      | CH <sub>3</sub> | CH <sub>3</sub>   | NaH    | 5a       | 68                  | >99:1              |
| 2     | 3      | CH <sub>3</sub> | Ph  | NaH    | 5b, 6b   | 60                  | 75:25              |
| 3     | 3      | CH <sub>3</sub> | <i>p</i> -CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub> | NaH    | 5c       | 61                  | >99:1              |
| 4     | 4      | CH <sub>3</sub> | CH <sub>3</sub>   | NaH    | 5d       | 66                  | >99:1              |
| 5     | 4      | CH <sub>3</sub> | Ph  | NaH    | 5e       | 89                  | >99:1              |
| 6     | 4      | CH <sub>3</sub> | <i>p</i> -CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub> | NaH    | 5f       | 87                  | >99:1              |
| 7     | 3      | CH <sub>3</sub> | OEt   | NaHMDS | 5g       | 7                   | >99:1              |
| 8     | 3      | CF <sub>3</sub> | OEt   | NaHMDS | 5h       | 61                  | >99:1              |

a. Isolated Yields.

b. Determined by <sup>1</sup>H-NMR.

The structures of the products were determined from spectral data. Especially, the vinylic carbons of **5a** were observed at 96.02 and 163.3ppm in the <sup>13</sup>C-NMR spectrum. In the <sup>1</sup>H-NMR spectrum of the products **5a**, the amine proton appeared at about 10.87 ppm as a broad triplet signal. Furthermore, nuclear Overhauser effect was observed between methyl proton ( $R_1$ ) and olefin proton. These results and literature data<sup>[5]</sup> suggest that the compound **5a** has an *Z*-enamine structure.

In the case of a reaction of **3** with benzoyl acetone, a mixture of isomers, **5b** and **6b**, was obtained in a ratio of 75:25. The major isomer **5b** showed an NH resonance at 11.49ppm in the  $^1\text{H}$ -NMR spectrum, and the minor isomer did so at 10.79ppm. The downfield shift of the NH proton should be the result of a hydrogen bond between the amino hydrogen and the carbonyl oxygen<sup>[6]</sup>. The NOESY spectrum of a mixture of **5b** and **6b** indicated between methyl proton ( $R_1$ ) and olefin proton of **5b**.

Furthermore, the carbonyl resonance of **5a** was observed at 195.4ppm in the  $^{13}\text{C}$ -NMR spectra. On the other hand, that of **5b** was observed at 187.9ppm. The higher field signal of the carbonyl carbon is attributable the result of a phenyl-carbonyl resonance effect<sup>[7]</sup>. These results suggest the compound **5b** has a benzoyl group, not an acetyl group. The regiochemistry of the reactions might be the result of a steric effect of the phenyl group.

The reaction of **3** with ethyl acetoacetate provided **5g** in very low yield. (7%) The yield was improved to 61% when ethyl 4,4,4-trifluoroacetoacetate was used as a substrate. It was known that the great electrophilicity of fluoroalkyl carboxylic acid derivatives enables phosphorus ylides to add to the carbonyl group of esters<sup>[4]</sup> and even amides<sup>[8]</sup>.

Treatment of a mixture of **5b** and **6b** with two equimolar amounts of LDA at 0°C gave a dark orange solution, which reacted with benzaldehyde to give the erythro alcohol **7a** in 8% yield. However, when using five equimolar amounts of LDA, the reaction progressed to form the product **7a** in 68% yield exclusively, after purification by column chromatography.

The structure and stereochemistry of **7a-c** were determined by  $^1\text{H}$ -NMR and  $^{13}\text{C}$ -NMR spectra. In the  $^1\text{H}$ -NMR spectrum, the coupling constant was observed at  $J_{AB} < 1\text{Hz}$ . The threo isomer ( $J_{AB} = 6-9\text{Hz}$ ) was not detected. Warren<sup>[9]</sup> and we<sup>[10]</sup> also have reported that reactions of alkyl or aryl (diphenyl) phosphine oxides with aldehydes gave preferentially erythro adducts.

Conversions of erythro adducts **7a**, **8a** and **7c** into the 1,6- and 1,7-diene **9a** and **9b** with elimination of the diphenylphosphinate ion were performed by treatment with sodium hydride in *N,N*-dimethylformamide (SCHEME 1) The results indicated that elimination of the diphenylphosphinate ion proceeded with retention of the configuration in the alcohols **7a+8a** and **7c**.

In summary, cyclic 2-azaphosphonium salts can react with  $\beta$ -dicarbonyl compounds to give enamino phosphine oxide derivatives in good yields, and we showed that the two kinds of nucleophilic sites could be used for formation of a carbon-nitrogen bond and a stereoselective carbon-carbon double bond.





TABLE II Reactions of Enamine Derivatives with Aldehydes

| enamine | Base                  | Ar  | Products | Yield <sup>a</sup> | ratio <sup>b</sup> |
|---------|-----------------------|---|----------|--------------------|--------------------|
| 5b+6b   | LDA(2eq.)             | Ph  | 7a       | 8                  | n.d                |
| 5b+6b   | LDA(5eq.)             | Ph  | 7a, 8a   | 68                 | 80:20              |
| 5b+6b   | LDA(5eq.)             | <i>p</i> -ClC <sub>6</sub> H <sub>4</sub> | 7b, 8b   | 51                 | 93:7               |
| 5b+6b   | <i>n</i> -BuLi (5eq.) | Ph  | 7a, 8a   | 13                 | 80:20              |
| 5e      | LDA(5eq.)             | Ph  | 7c       | 70                 | >99:1              |
| 5e      | <i>n</i> -BuLi        | Ph  | 7c       | 14                 | >99:1              |

a. Isolated Yields.

b. Determined by <sup>1</sup>H-NMR.

SCHEME I

## EXPERIMENTAL SECTION

## 4-[3-(Diphenyl-phosphinoyl)-propylamino]-pent-3-en-2-one 5a

A mixture of sodium hydride (60% dispersion in mineral oil, 0.09 g, 2.25 mmol) and 1,1-diphenyl-2-azaphosphorinanium perchlorate **1** (0.68 g, 2

mmol) in dry THF (10 ml) was stirred for 30min at room temperature and 10min at reflux temperature. After cooling to room temperature, to this mixture was added dropwise a solution of 2,4-pentanedione (0.30 g, 3 mmol) in dry THF (5 ml) at room temperature and stirred for 20hr at reflux temperature. The reaction mixture was allowed to cool to room temperature. Water (10 ml) and dichloromethane (50 ml) were then added dropwise, and the organic layer was separated. The aqueous layer was extracted with dichloromethane. The combined organic extracts were washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give yellow crystals. (0.45 g, 66%): m.p. 139–140°C; IR (KBr) 3060, 2935, 1600, 1545, 1440, 1180, 1115  $\text{cm}^{-1}$ ;  $^1\text{H-NMR}$  (400.13MHz,  $\text{CDCl}_3$ )  $\delta$  1.79–2.05 (m, 2H), 1.85 (s, 3H), 2.00 (s, 3H), 2.31–2.38 (m, 2H), 3.31–3.36 (m, 2H), 4.96 (s, 1H), 7.27–7.70 (m, 10H), 10.87 (brt, 1H);  $^{13}\text{C-NMR}$  (100.61MHz,  $\text{CDCl}_3$ )  $\delta$  18.96 (s), 23.05 (d,  $^2J_{\text{PC}}=3.67\text{Hz}$ ), 27.14 (d,  $^1J_{\text{PC}}=72.02\text{Hz}$ ), 29.15 (s), 43.57 (d,  $^3J_{\text{PC}}=12.81\text{Hz}$ ), 96.02 (s), 128.85–135.46 (m) 163.31 (s), 195.38 (s); MS  $m/z$  342 ( $\text{M}^++1$ ); HRMS calcd. for  $\text{C}_{20}\text{H}_{25}\text{NO}_2\text{P}$  ( $\text{M}^++1$ ) 342.1623, found 342.1617.

### Compound 5b and 6b

Prepared as above. Yellow crystals (89% yield): m.p. 135–140°C; IR (KBr) 3050, 2930, 1590, 1575, 1540, 1430  $\text{cm}^{-1}$ ;  $^1\text{H-NMR}$  (400.13MHz,  $\text{CDCl}_3$ )  $\delta$  1.75–2.02 (m, 2H), 1.97 (s, 2.25H), 2.08 (s, 0.75H), 2.24–2.29 (m, 0.5H), 2.35–2.42 (m, 1.5H), 3.18–3.21 (m, 0.5H), 3.39–3.44 (m, 1.5H), 5.05 (s, 0.25H), 5.66 (s, 0.75H), 7.43–7.85 (m, 15H), 10.79 (brt, 0.25H) 11.49 (brt, 0.75H);  $^{13}\text{C-NMR}$  5b (100.61MHz,  $\text{CDCl}_3$ )  $\delta$  19.23 (s), 22.70 (d,  $^2J_{\text{PC}}=3.66\text{Hz}$ ), 26.76 (d,  $^1J_{\text{PC}}=72.02\text{Hz}$ ), 43.46 (d,  $^3J_{\text{PC}}=12.81\text{Hz}$ ), 92.50 (s), 126.88–140.47 (m), 164.83 (s), 187.99 (s); 6b  $\delta$  23.08 (s), 26.70 (d,  $^1J_{\text{PC}}=71.64\text{Hz}$ ), 44.64 (d,  $^3J_{\text{PC}}=15.24\text{Hz}$ ), 97.15 (s), 165.01(s), 195.95 (s); MS  $m/z$  403 ( $\text{M}^+$ ); HRMS calcd. for  $\text{C}_{25}\text{H}_{26}\text{NO}_2\text{P}$  ( $\text{M}^+$ ) 403.1701, found 403.1731.

### Compound 5c

Prepared as above. Yellow crystals (87% yield): m.p. 162–164°C; IR (KBr) 3060, 2920, 1575, 1515, 1435, 1290, 1160, 1110  $\text{cm}^{-1}$ ;  $^1\text{H-NMR}$  (89.56MHz,  $\text{CDCl}_3$ )  $\delta$  2.10–3.53 (m, 6H), 1.97 (s, 3H), 2.37 (s, 3H), 5.64

(s, 1H), 7.14–7.87 (m, 14H), 11.44 (brt, 1H);  $^{13}\text{C}$ -NMR (22.39MHz,  $\text{CDCl}_3$ )  $\delta$  19.26 (s), 22.01 (d,  $^2J_{\text{PC}}=3.66\text{Hz}$ ), 26.75 (d,  $^1J_{\text{PC}}=72.63\text{Hz}$ ), 43.42 (d,  $^3J_{\text{PC}}=12.20\text{Hz}$ ), 92.29 (s), 126.90–137.68 (m), 164.53 (s), 187.88 (s); MS  $m/z$  418 ( $\text{M}^++1$ ); HRMS calcd. for  $\text{C}_{26}\text{H}_{29}\text{NO}_2\text{P}(\text{M}^++1)$  418.1921, found 418.1947.

### Compound 5d

Prepared as above. Yellow crystals (68% yield): m.p. 96–98°C; IR (KBr) 3420, 3050, 2940, 1610, 1570, 1440, 1300, 1180, 1120  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (89.56MHz,  $\text{CDCl}_3$ )  $\delta$  1.68–3.23 (m, 6H), 1.86 (s, 3H), 1.98 (s, 3H), 4.93 (s, 1H), 7.29–7.86 (m, 10H), 10.82 (brt, 1H);  $^{13}\text{C}$ -NMR (22.39MHz,  $\text{CDCl}_3$ )  $\delta$  18.71 (s), 19.45 (d,  $^2J_{\text{PC}}=3.66\text{Hz}$ ), 29.66 (d,  $^1J_{\text{PC}}=72.02\text{Hz}$ ), 31.62 (d,  $^3J_{\text{PC}}=12.82\text{Hz}$ ), 42.61 (s), 95.29 (s), 128.45–135.19 (m), 164.53 (s), 194.76 (s); MS  $m/z$  356 ( $\text{M}^++1$ ); HRMS calcd. for  $\text{C}_{21}\text{H}_{27}\text{NO}_2\text{P}(\text{M}^++1)$  356.1779, found 356.1765.

### Compound 5e

Prepared as above. A yellow syrup (89% yield): IR (neat) 3380, 3050, 2940, 1610, 1570, 1440, 1300, 1180, 1120  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400.13MHz,  $\text{CDCl}_3$ )  $\delta$  1.71–1.78 (m, 4H), 2.01 (s, 3H), 2.03–2.35 (m, 2H), 3.27–3.31 (m, 2H), 5.65 (s, 1H), 7.28–7.90 (m, 15H), 11.44 (brt, 1H);  $^{13}\text{C}$ -NMR (100.61MHz,  $\text{CDCl}_3$ )  $\delta$  19.22 (s), 19.33 (s), 29.38 (d,  $^1J_{\text{PC}}=71.64\text{Hz}$ ), 31.18 (d,  $^3J_{\text{PC}}=12.95\text{Hz}$ ), 42.85 (s), 92.11 (s), 126.84–140.39 (m), 164.80 (s), 187.61 (s); MS  $m/z$  418 ( $\text{M}^++1$ ); HRMS calcd. for  $\text{C}_{26}\text{H}_{29}\text{NO}_2\text{P}(\text{M}^++1)$  418.1936, found 418.1938.

### Compound 5f

Prepared as above. A yellow syrup (61% yield): IR (neat) 3420, 3070, 2950, 1600, 1590, 1440, 1180, 1120  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (89.56MHz,  $\text{CDCl}_3$ )  $\delta$  1.74–3.39 (m, 8H), 2.00 (s, 3H), 2.37 (s, 3H), 5.62 (s, 1H), 7.14–7.85 (m, 10H), 11.39 (brt, 1H);  $^{13}\text{C}$ -NMR (22.39MHz,  $\text{CDCl}_3$ )  $\delta$  19.31 (s), 19.39 (s), 29.55 (d,  $^1J_{\text{PC}}=72.02\text{Hz}$ ), 31.34 (d,  $^3J_{\text{PC}}=13.43\text{Hz}$ ), 42.93 (s), 92.04 (s), 127.01–140.75 (m), 164.50 (s), 187.71 (s); MS  $m/z$  432 ( $\text{M}^++1$ ).



**Compound 5g**

To a suspension of 1,1-diphenyl-2-azaphospholanium perchlorate **1** (0.68 g, 2 mmol) in dry THF (10 ml) was added a 1 M solution of NaH-MDS in THF (2.0 ml, 2.2 mmol) at room temperature with stirring. After 15 min, a solution of ethyl acetoacetate (0.26 g, 2 mmol) in THF (10 ml) was added to the mixture and stirred for 1 hr. Ten of water was then added and extracted with dichloromethane. The organic layer was added over anhydrous sodium sulfate and concentrated in vacuo to give a yellow syrup **5** g(0.05 g, 7%): IR (neat) 3400, 2940, 1640, 1600, 1440, 1260, 1175, 1120  $\text{cm}^{-1}$ ;  $^1\text{H-NMR}$  (400.13MHz,  $\text{CDCl}_3$ )  $\delta$  1.24 (t, 3H,  $J=7.0\text{Hz}$ ), 1.83 (s, 3H), 1.89–2.42 (m, 4H), 3.14–3.19 (m, 2H), 4.08 (q, 2H,  $J=7.0\text{Hz}$ ), 4.44 (s), 7.29–7.86 (m, 10H), 8.55 (m, 1H);  $^{13}\text{C-NMR}$  (100.61MHz,  $\text{CDCl}_3$ )  $\delta$  15.20 (s), 19.67 (s), 23.70 (d,  $^2J_{\text{PC}}=3.66\text{Hz}$ ), 27.39 (d,  $^1J_{\text{PC}}=72.02\text{Hz}$ ), 43.88 (d,  $^3J_{\text{PC}}=13.3\text{Hz}$ ), 58.83 (s), 83.73 (s), 127.47–132.54 (m), 162.14 (s), 171.19 (s); MS 385  $m/z$  ( $\text{M}^+$ ); HRMS calcd. for  $\text{C}_{22}\text{H}_{28}\text{O}_3\text{NP}$  ( $\text{M}^+$ ) 385.1806, found 385.1780.

**Compound 5h**

Prepared as above. Pale yellow crystals (61% yield): m.p. 78–79°C; IR (KBr) 3275, 2980, 1675, 1620, 1440, 1280, 1195, 1120  $\text{cm}^{-1}$ ;  $^1\text{H-NMR}$  (400.13MHz,  $\text{CDCl}_3$ )  $\delta$  1.26 (t, 3H,  $J=7.4\text{Hz}$ ), 1.86–1.93 (m, 2H), 2.28–2.35 (m, 2H), 3.34 (q, 2H,  $J_{\text{HH}}=6.60\text{Hz}$ ,  $J_{\text{HH}}=6.56\text{Hz}$ ), 4.15 (q, 2H,  $J=7.4\text{Hz}$ ), 5.08 (s, 1H), 7.27–7.77 (m, 10H), 8.16 (brt, 1H);  $^{13}\text{C-NMR}$  (100.61MHz,  $\text{CDCl}_3$ )  $\delta$  14.27 (s), 22.86 (d,  $^2J_{\text{PC}}=3.66\text{Hz}$ ), 27.15 (d,  $^1J_{\text{PC}}=72.63\text{Hz}$ ), 44.40 (d,  $^3J_{\text{PC}}=15.24\text{Hz}$ ), 59.78 (s), 85.11 (d,  $^3J_{\text{CF}}=5.34\text{Hz}$ ), 120.77 (q,  $^1J_{\text{CF}}=275.12\text{Hz}$ ), 148.20 (q,  $^2J_{\text{CF}}=31.25\text{Hz}$ ), 169.81 (s); MS 425  $m/z$  ( $\text{M}^+$ ); HRMS calcd. for  $\text{C}_{21}\text{H}_{23}\text{F}_3\text{NO}_3\text{P}$  ( $\text{M}^+$ ) 425.1367, found 425.1344.

**3-[3-(Diphenyl-phosphinoyl)-4-hydroxy-4-phenyl-butylamino]-1-phenyl-but-2-en-1-one 7a**

To an LDA solution (5.65 ml, 11.3 mmol) was added acyclic enamine phosphine oxides **5b**, **6b** (0.910 g, 2.26 mmol) in 10 ml of dry THF solution at  $-78^\circ\text{C}$ , and stirred for 10min at this temperature. A solution of benzaldehyde (0.288 g, 2.71 mmol) in THF was then added dropwise at same

temperature. The mixture was stirred for 1 hr at  $-78^{\circ}\text{C}$ . After aqueous workup at room temperature, the mixture was extracted with ether, dried over anhydrous sodium sulfate, and concentrated to give the crude products. This crude product was purified by column chromatography of silica gel using ethyl acetate as the eluent to give **7a** as a yellow syrup. (0.770 g, 67%): IR (neat) 3350, 3050, 1595, 1540, 1435, 1320, 1285  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400.13MHz,  $\text{CDCl}_3$ )  $\delta$  1.41. (s, 2.4H), 1.71–2.87 (m, 5H), 2.13 (s, 0.6H), 4.73 (brs, 0.2H), 4.86 (brs, 0.8H), 4.90 (s, 0.2H), 5.27 (d, 0.2H,  $^3J_{\text{PH}}=9.36\text{Hz}$ ) 5.31 (d, 0.8H,  $^3J_{\text{PH}}=8.84\text{Hz}$ ), 5.50 (s, 0.8H), 7.19–8.04 (m, 20H), 10.52 (brt, 0.2H), 11.94 (brt, 0.8H);  $^{13}\text{C}$ -NMR (100.61MHz,  $\text{CDCl}_3$ ) **7a**  $\delta$  18.32 (s), 21.88 (s), 40.97 (d,  $^1J_{\text{PC}}=67.07\text{Hz}$ ), 41.42 (d,  $^3J_{\text{PC}}=8.38\text{Hz}$ ), 69.89 (s), 92.55 (s), 125.04–143.16 (m), 164.79 (s), 187.63 (s); **8a**  $\delta$  97.50 (s), 165.06 (s), 195.77 (s); MS 509  $m/z$  ( $\text{M}^+$ ); HRMS calcd. for  $\text{C}_{32}\text{H}_{32}\text{NO}_3\text{P}$  ( $\text{M}^+$ ) 509.2120, found 509.2090.

### Compound 7b

Prepared as above. A yellow syrup (51% yield): IR (neat) 3380, 3050, 1595, 1580, 1540, 1435, 1320, 1260  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400.13MHz,  $\text{CDCl}_3$ )  $\delta$  1.46 (s, 3H), 1.73–2.84 (m, 5H), 4.93 (m, 1H), 5.20 (d, 0.07H,  $^3J_{\text{PH}}=9.32\text{Hz}$ ), 5.26 (d, 0.93H,  $^3J_{\text{PH}}=8.84\text{Hz}$ ), 5.27 (s, 0.07H), 5.53 (s, 0.93H), 7.23–8.04 (m, 19H), 10.50 (brt, 0.07H), 11.22 (brt, 0.93H);  $^{13}\text{C}$ -NMR (100.61MHz,  $\text{CDCl}_3$ ) **7b**  $\delta$  18.34 (s), 21.87 (s), 40.80 (d,  $^1J_{\text{PC}}=67.06\text{Hz}$ ), 41.32 (d,  $^3J_{\text{PC}}=7.62\text{Hz}$ ), 69.71 (s), 92.75 (s), 126.65–140.55 (m), 164.77 (s), 187.84 (s); **7b**  $\delta$  97.55 (s), 165.00 (s), 195.91 (s); MS  $m/z$  558 ( $\text{M}^++1$ ); HRMS calcd. for  $\text{C}_{33}\text{H}_{34}\text{NO}_3\text{PCl}$  ( $\text{M}^++1$ ) 558.1962, found 558.1969.

### Compound 7c

Prepared as above. A yellow syrup (70% yield): IR (neat) 3380, 3050, 1595, 1580, 1540, 1435, 1320, 1260  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400.13MHz,  $\text{CDCl}_3$ )  $\delta$  0.84–0.93 (m, 1H), 1.07–1.13 (m, 1H), 1.58–1.71 (m, 1H), 1.69 (s, 3H), 1.88–1.98 (m, 1H), 2.51–2.55 (m, 1H), 2.72 (dt, 1H,  $J_{\text{HH}_\text{C}}=7.04\text{Hz}$ ,  $J_{\text{HH}_\text{N}}=6.80\text{Hz}$ ), 4.81 (brs, 1H), 5.28 (d, 1H,  $^3J_{\text{PH}}=9.12\text{Hz}$ ), 5.53 (s, 1H), 7.20–8.04 (m, 20H), 11.22 (brt, 1H);  $^{13}\text{C}$ -NMR (100.61MHz,  $\text{CDCl}_3$ )  $\delta$  19.02 (s), 19.04 (s), 30.19 (d,  $^3J_{\text{PC}}=6.10\text{Hz}$ ), 42.99 (s), 44.41 (d,  $^1J_{\text{PC}}=66.30\text{Hz}$ ), 70.61 (d,

$^2J_{\text{PC}}=2.29\text{Hz}$ ), 91.94 (s), 125.37–142.03 (m), 164.59 (s), 187.28 (s); MS  $m/z$  523 ( $M^+$ ); HRMS calcd. for  $\text{C}_{33}\text{H}_{34}\text{NO}_3\text{P}$  ( $M^+$ ) 523.2276, found 523.2248.

### (Z)-1-Phenyl-3-(4-phenyl-3-butenylamino)-2-butene-1-one 9a

A solution of the alcohol 9a (0.588 g, 1.12 mmol) in dry dimethylformamide (DMF) was added dropwise to a suspension of sodium hydride (60% dispersion in mineral oil, 0.225 g, 4.60 mmol) at  $0^\circ\text{C}$ , and the mixture was stirred at r.t. for 8h. After aqueous workup, the mixture was extracted with ether, and the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, and concentrated to give the crude product. The crude product was purified by column chromatography on silica gel with AcOEt as eluent to give 9a as a yellow syrup (0.226 g, 66%): IR (neat) 3060, 3030, 1605, 1545, 1325, 1295  $\text{cm}^{-1}$ ;  $^1\text{H-NMR}$  (400.13MHz,  $\text{CDCl}_3$ )  $\delta$  1.99 (s, 3H), 2.61–2.67 (m, 2H), 3.33–3.41 (m, 2H), 5.63–5.70 (m, 2H), 6.58 (d, 1H,  $J_{\text{HH}}=11.64\text{Hz}$ ), 7.16–7.86 (m, 10H), 11.49 (brt, 1H);  $^{13}\text{C-NMR}$  (100.61MHz,  $\text{CDCl}_3$ )  $\delta$  19.52 (s), 29.38 (s), 43.43 (s), 92.43 (s), 126.06–140.65 (m), 164.75 (s), 187.90 (s); MS  $m/z$  291 ( $M^+$ ); HRMS calcd. for  $\text{C}_{20}\text{H}_{21}\text{NO}$  ( $M^+$ ) 291.1623, found 291.1626.

### Compound 9b

Prepared as above. A yellow syrup (58% yield): IR (neat) 3060, 3020, 2940, 1605, 1590, 1440, 1325, 1290  $\text{cm}^{-1}$ ;  $^1\text{H-NMR}$  (400.13MHz,  $\text{CDCl}_3$ )  $\delta$  1.73–1.80 (m, 2H), 1.96 (s, 3H), 2.40–2.46 (m, 2H), 3.23–3.28 (m, 2H), 5.48–5.65 (m, 2H), 6.46 (d, 1H,  $J_{\text{HH}}=11.64\text{Hz}$ ), 7.16–7.95 (m, 10H), 11.45 (brt, 1H);  $^{13}\text{C-NMR}$  (100.61MHz,  $\text{CDCl}_3$ )  $\delta$  19.73 (s), 26.11 (s), 30.77 (s), 42.95 (s), 92.47 (s), 121.49–140.60 (m), 164.64 (s), 187.35 (s); MS  $m/z$  306 ( $M^++1$ ); HRMS calcd. for  $\text{C}_{21}\text{H}_{24}\text{NO}$  ( $M^++1$ ) 306.1858, found 306.1896.

### Acknowledgements

This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, and Culture of Japan (No. 11650889). We are grateful to the Ihara Chemical Co., Ltd., for the gift of triphenylphosphine. We thank Welfide Co., Ltd., for the MS and HRMS measurement.

### References

- [1] Sakai, T., Kodama, T., Fujimoto, T., Ohta, K., Yamamoto, I. *J. Org. Chem.*, **1997**, *62*, 6627.
- [2] Yamamoto, I., Tashiro, K., Uchiyama, T., Fujimoto, T. *Phosphorus, Sulfur and Silicon*, **1999**, Vol. 148, 11.
- [3] Sakai, T., Ida, K., Uchiyama, T., Fujimoto, T., Ohta, K., Yamamoto, I. *Heterocyclic Commun.*, **1996**, *2*, 233.
- [4] (a) Bestmann, H. J., Dornauer, H., Rostok, K. *Chem. Ber.* **1970**, *103*, 2011.  
(b) Bégué, J.P., Mesureur, D. *J. Fluorine Chem.* **1988**, *39*, 271.
- [5] Stefani, H. A., Costa, I. M., Silva, D. O. *Synthesis*, **2000**, *11*, 1526.
- [6] Emsley, J. W., Feeney, J., Sutcliffe, L. H. "High Resolution Nuclear Magnetic Resonance Spectroscopy", Pergamon Press, Oxford, London, Edinburgh, New York, Paris, Frankfurt, pp. 818–819.
- [7] G. C. Levy, G. L. Nelson, "Carbon-13 Nuclear Magnetic Resonance for Organic Chemists", Wiley-Interscience, New York, London, Sydney, Toronto, **1972**, pp. 112–114.
- [8] Bégué, J.P., Mesureur, D. *Synthesis* **1989**, 309.
- [9] (a) Buss, A.D., Warren S. *J. Chem. Soc., Chem. Commun.*, **1981**, 100.  
(b) Buss, A. D., Warren, S. *Tetrahedron Lett.*, **1983**, *24*, 3931.
- [10] Yamamoto, I., Fujimoto, T., Ohta, K., Matsuzaki, K. *J. Chem. Soc., Perkin Trans.* **1987**, *1*, 1537.