

Manganese-Catalyzed α -Alkylation of Secondary Alcohols with Primary Alcohols under Phosphine-Free Conditions

Tingting Liu, Liandi Wang, Kaikai Wu, and Zhengkun Yu

ACS Catal., Just Accepted Manuscript • DOI: 10.1021/acscatal.8b01960 • Publication Date (Web): 05 Jul 2018

Downloaded from <http://pubs.acs.org> on July 5, 2018

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



Manganese-Catalyzed β -Alkylation of Secondary Alcohols with Primary Alcohols under Phosphine-Free Conditions


Tingting Liu,^{†,‡} Liandi Wang,[†] Kaikai Wu,[†] and Zhengkun Yu^{*,†,§}

[†] Dalian Institute of Chemical Physics, Chinese Academy of Sciences, 457 Zhongshan Road, Dalian 116023, P. R. China

[‡] University of Chinese Academy of Sciences, Beijing 100049, P. R. China

[§] State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 354 Fenglin Road, Shanghai 200032, P. R. China

ABSTRACT: Manganese(I) complexes bearing a pyridyl-supported pyrazolyl-imidazolyl ligand efficiently catalyzed the direct β -alkylation of secondary alcohols with primary alcohols under phosphine-free conditions. The β -alkylated secondary alcohols were obtained in moderate to good yields with water formed as the by-product through a borrowing hydrogen pathway. β -Alkylation of cholesterol was also effectively achieved. The present protocol provides a concise atom-economical method for C–C bond formation from primary and secondary alcohols.




KEYWORDS: manganese • alcohols • alkylation • borrowing hydrogen • cholesterol

Construction of carbon–carbon bonds is of great importance in organic synthesis.¹ More and more concern on the consequence of climate change and dwindling crude oil reserves results in the search for alternative carbon resources for the formation of C–C bonds.² Thus, readily available alcohols have recently been paid much attention to be utilized as the electrophiles or alkylating agents to establish C–C bonds.³ The traditional method to prepare β -alkylated alcohols from secondary alcohols usually requires a multistep process which involves oxidation of the secondary alcohols, alkylation with alkyl halides, and reduction of the β -alkylated ketones. The drawbacks of such a protocol include waste generation, expensive and toxic noble-metal catalyst, limited availability of the starting materials. β -Alkylation of secondary alcohols can also be realized by means of hydrogen autotransfer or borrowing hydrogen which has become an important synthetic strategy in organic chemistry due to its high efficiency, low cost, and versatility.⁴ The operational simplicity, ready availability of the starting materials from renewable resources, and generation of water as the only stoichiometric by-product make the borrowing hydrogen strategy be atom-economical, sustainable, and environmentally benign.⁵ Direct β -alkylation of secondary alcohols with primary alcohols have been documented under ruthenium,⁶ iridium,⁷ palladium,⁸ and copper⁹ catalysis or under metal-free conditions¹⁰ through a borrowing hydrogen strategy.

It has been desired to apply non-noble metal catalysis in organic synthesis due to the economic and ecological benefits as well as different reactivities of the metals. In this regard, base metals such as iron, cobalt, and nickel have been


achieved much progress in hydrogenation and dehydrogenation reactions.^{11–13} Manganese is the third most abundant metal in the earth's crust which offers an attractive alternative to noble metals. In particular, it is capable of existing in several oxidation states. Milstein and co-workers reported α -olefination of nitriles,^{14a} dehydrogenative cross-coupling of alcohols with amines,^{14b} *N*-formylation of amines with methanol,¹⁵ and deoxygenation of alcohols¹⁶ catalyzed by pincer-type Mn–PNP complexes. With a manganese complex as the catalyst, diverse alcohols underwent dehydrogenation-based processes such as *N*-alkylation of amines,¹⁷ C-alkylation of ketones,¹⁸ dehydrogenation of alcohols to esters,¹⁹ and methanol to H₂ and CO₂,²⁰ synthesis of pyrroles, quinolines,



Scheme 1. Selected Known Manganese Complex Catalysts


and pyrimidines.²¹ Hydrogenations of ketones, nitriles, esters, CO₂, and amides have also been realized with manganese complex catalysts.^{22,23} Very recently, Kempe, et al. reported manganese-catalyzed dehydrogenative alkylation or α -olefination of alkyl-substituted *N*-heteroarenes with alcohols.²⁴ The typical phosphine-bearing amine-based PNP–MnH,^{14a} triazine-based PNP–MnH,^{21c} pyridine-based PNP–MnBr,^{21d} and amine-based PNP–MnBr^{23c} complex catalysts have been documented (Scheme 1).

Recently, we have reported versatile pyridyl-based NNN, CNN and NNP ligands for the construction of ruthenium complex catalysts for transfer hydrogenation of ketones, dehydrogenation of *N*-heterocyclic compounds, synthesis of pyrrole derivatives, and Oppenauer-type oxidation of secondary alcohols.²⁵ Among these ruthenium complex catalysts, Ru(III) complex **A** exhibited a high catalytic activity in the β -alkylation of secondary alcohols.^{6b} During our continuous investigation of transition-metal complex catalysts bearing a unsymmetrical tridentate ligand, we found that the transition-metal complexes bearing a phosphine-free pyridyl-supported 2,6-(mixed *N*-heterocycles) ligand can usually exhibit enhanced catalytic activity and selectivity. Encouraged by these findings, we envisioned that these ligands might also be employed to construct manganese complex catalysts. Herein, we disclose the synthesis of Mn(I)-NN complexes **1** and their catalytic activities in β -alkylation of secondary alcohols with primary alcohols (Scheme 2).



Scheme 2. Relevant Ligands and Their Ru(III) and Mn(I) Complexes

Treatment of our previously reported pyridyl-supported pyrazolyl-imidazolyl ligand **L1**,^{25c} that is, 2-(3,5-dimethylpyrazol-1-yl)-6-(benzimidazol-2-yl)pyridine, with $\text{Mn}(\text{CO})_3\text{Br}$ (1 equiv) in THF at 25 °C under a nitrogen atmosphere led to complex **1a** in 83% yield (Scheme 3). In a similar fashion, complex **1b** was obtained in 87% yield from the same reaction in methanol. The Mn–Br bond in **1a** was transformed to Mn–OMe in **1b**. Reacting $\text{Mn}(\text{CO})_3\text{Br}$ with the *N*-methylated ligand **L2** in methanol only afforded complex **1c** (80%), and the reaction exhibited no solvent effect. For the comparison study, bidentate ligand **L3** without bearing the 3,5-dimethylpyrazol-1-yl (Me_2pz) moiety was also applied in the reaction with $\text{Mn}(\text{CO})_3\text{Br}$ in methanol, giving complex **1d** in 85% yield. It is noteworthy that stirring a solution of complex **1a** in methanol at ambient temperature for 24 h quantitatively gave complex **1b**. The Mn–Br bond in **1a** was transformed to



Conditions: **L1-L3** (0.5 mmol), $\text{Mn}(\text{CO})_3\text{Br}$ (0.5 mmol), solvent (25 mL), 0.1 MPa N_2 , 25 °C, 24 h. (i) Synthesis of **1a**: **L1**, THF, 83% yield; (ii) Synthesis of **1b**: **L1**, MeOH, 87% yield; (iii) Synthesis of **1c**: **L2**, MeOH, 80% yield; (iv) Synthesis of **1d**: **L3**, MeOH, 85% yield.

Scheme 3. Synthesis of Manganese Complexes

Mn–OMe in **1b** by reacting with methanol and release of hydrogen bromide under the stated conditions, while

complexes **1c** and **1d** could not undergo the same type of reactions.

Complexes **1** were fully characterized by NMR, FT-IR, and elemental analyses, which are consistent with their compositions. The proton NMR signals of complex **1a** was shifted 0.1–2.1 ppm downfield as compared with those of the free ligand **L1**, suggesting that coordination of the ligand to manganese diminishes the electron density on the ligand. Three coordinating carbonyl groups in $\text{Mn}(\text{CO})_3$ moiety of complex **1a** were confirmed by the $^{13}\text{C}\{^1\text{H}\}$ NMR signals at 225.4, 220.2, and 215.2 ppm. The proton NMR signals of complex **1b** were broadened and its $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum could not be obtained due to the possibility of a dynamic process.²⁶ The proton NMR spectrum of complex **1c** exhibited one singlet at 4.40 ppm, corresponding to the proton resonance of *N*-methyl group, and other proton resonance signals were shifted 0.1–0.5 ppm downfield as compared with those of the free ligand **L2** due to its coordination with the manganese center. In the $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of **1c**, the singlets at 226.6, 221.0, 216.6, and 33.1 ppm revealed the presence of three coordinating carbonyl groups and *N*-methyl, respectively. The NMR spectra of complex **1d** were in accordance with its composition, and the carbonyl resonance signals appeared at 224.2, 221.3, and 220.8 ppm, respectively. As compared with the infrared spectrum of ligand **L1**, the stretching vibrations of the carbonyls in $\text{Mn}(\text{CO})_3$ of complex **1a** were assigned at 2025 and 1933 cm^{-1} , respectively, corresponding to two types of coordinating carbonyls. In a similar manner, two types of carbonyl stretching vibrations were observed at 2027 and 1909 cm^{-1} , 2024 and 1935 cm^{-1} , and 2025 and 1933 cm^{-1} for complexes **1b-1d**, respectively.





Figure 1. Perspective view of complex **1b** with thermal ellipsoids at the 30% probability level.

Single crystals of complex **1b** suitable for X-ray diffraction were obtained by slow vapor diffusion of *n*-pentane into a saturated solution of the complex in $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (v/v, 1/2) at 4 °C. The molecular structure of complex **1b** adopts an octahedral geometry with the metal center coordinated by the pyridyl and imidazolyl nitrogen atoms of ligand **L1** and three carbonyls, and bonded to the methoxy oxygen atom (Figure 1). The potentially tridentate ligand **L1** actually acts as a bidentate ligand in complex **1b**, forming two coordination bonds of Mn(1)–N(3) (2.13 Å) and Mn(1)–N(4) (2.02 Å). The unsymmetrical coordinating pyridyl-imidazolyl moiety occupies the two meridional sites with two carbonyls in a quasiplanar disposition, and the manganese atom is nearly situated in such a plane. The methoxy oxygen atom is

positioned *trans* to the third carbonyl with a C(18)-Mn(1)-O(4) angle of 176°, and bond lengths of Mn(1)-O(4) (2.08 Å) and Mn(1)-C(18) (1.79 Å), suggesting that the axial carbonyl and methoxy oxygen atom are almost linearly positioned at the two sides of the quasiplane. The two coordinating nitrogen atoms are also nearly linearly arranged with the two equatorial carbonyls, exhibiting angles of N(3)-Mn(1)-C(19) (175°) and N(4)-Mn(1)-C(20) (176°), respectively.

The catalytic activities of complexes **1** were evaluated in the C-alkylation of 1-phenylethanol (**2a**) with benzyl alcohol (**3a**) (Table 1). With 2.1 mol % loading, complexes **1a–1d** were applied as the catalysts for the reaction in the presence of 30 mol % *t*BuOK as the base in refluxing toluene. In the case of using complex **1a**, 86% conversion was achieved for **2a** with a 67/33 selectivity for the target C-alkylation product **4a** and C-alkylation/dehydrogenation product **4a'** (Table 1, entry 2), while the reaction proceeded less efficiently without a catalyst (Table 1, entry 1). The reaction efficiency was remarkably improved with complex **1b** as the catalyst, reaching 96% conversion for **2a** and a 94/6 selectivity for **4a/4a'**, and the corresponding β -alkylation product **4a** was isolated in 90% yield (Table 1, entry 3). Under the same conditions, both complexes **1c** and **1d** behaved less efficiently to the reaction than complex **1b** as the catalyst (Table 1, entries 4 and 5). Presence of both the N-H functionality and 3,5-dimethylpyrazol-1-yl moiety in the ligand (**L1**) seems to be

Table 1. Screening of Reaction Conditions^a




| Entry | Catalyst | Base | Conversion of 2a ^b (%) | 4a : 4a' ^c (molar ratio) ^b |
|-----------------|------------------------|----------------------------|------------------------------------------|----------------------------------------------------------------|
| 1 | | <i>t</i> BuOK | 83 | 40:60 |
| 2 | 1a | <i>t</i> BuOK | 86 | 67:33 |
| 3 | 1b | <i>t</i> BuOK | 96 | 94:6 (90) ^c |
| 4 | 1c | <i>t</i> BuOK | 84 | 53:47 |
| 5 | 1d | <i>t</i> BuOK | 92 | 78:22 |
| 6 | 1b | <i>t</i> BuONa | 87 | 92:8 |
| 7 | 1b | KOH | 80 | 88:12 |
| 8 | 1b | NaOH | 74 | 86:14 |
| 9 | 1b | | <1 | |
| 10 | 1b ^d | <i>t</i> BuOK | 91 | 65:35 |
| 11 | 1b | <i>t</i> BuOK ^e | 90 | 90:10 |
| 12 | 1b | <i>t</i> BuOK ^f | 84 | 91:9 |
| 13 ^g | 1b | <i>t</i> BuOK | 78 | 88:12 |
| 14 ^h | 1b | <i>t</i> BuOK | 66 | 89:11 |

^aConditions: **2a** (2 mmol), **3a** (2 mmol), catalyst **1** (2.1 mol %), base (30 mol %), toluene (1 mL), 0.1 MPa N₂, 110 °C, 24 h. ^bDetermined by GC analysis. ^cIsolated yield of **4a** given in parentheses. ^d1.0 mol %. ^e20 mol %. ^f40 mol %. ^g100 °C. ^hSolvent-free conditions.

crucial to render the formation of complex **1b**. The order of the catalytic activities for these complexes was thus obtained

as **1b** > **1d** > **1a** > **1c** (Table 1, entries 2–5). With **1b** as the catalyst, the reaction conditions were further optimized. An obvious base effect was observed on the alkylation efficiency with the order *t*BuOK > *t*BuONa > KOH > NaOH >> no base (Table 1, entries 3 and 6–9). Lowering the catalyst loading to 1.0 mol %, varying the base loading to 20–40 mol %, or lowering the reaction temperature to 100 °C reduced the conversion of **2a** with deteriorated selectivities of **4a/4a'** (Table 1, entries 10–13). Equivalent base was needed in the hidden alkylation of alcohol reported by Kempe and coworkers.^{21b} Under the solventless conditions, the reaction could not efficiently occur (Table 1, entry 14). The high catalytic activity of complex **1b** is presumably attributed to the presence of an NH functionality in the ligand and the OMe ligand attached on the metal center. Methanol may be released to form a coordinatively unsaturated (16-electron) Mn(I) species which thus enhances the interaction between the substrate and the catalytically active Mn(I) metal center, accelerating the reaction.

Table 2. Scope of Primary Alcohols (3)^a




^aConditions: **2a** (2 mmol), **3** (2 mmol), 2.1 mol % **1b**, 30 mol % *t*BuOK, 0.1 MPa N₂, toluene (1 mL), 110 °C, 24 h. ^b4.2 mol % **1b**, *o*-xylene (1 mL), 140 °C, 48 h.

Under the optimized conditions, the protocol generality was explored by applying a variety of primary alcohols (Table 2). Benzyl alcohols (**3**) bearing an electron-donating methyl, methoxyl, isopropyl, or 3,4-methylenedioxy substituent reacted with **2a** to form the target products **4b–4h** in good

yields (78–88%). Methyl group exhibited no obvious steric impact on the yields of **4b–4d** (84–85%), while methoxyl, isopropyl, and 3,4-methylenedioxy demonstrated various electronic/steric effects on the product yields of **4e–4h** (78–88%). The *ortho*-substituents such as 2-F and 2-Cl on the aryl group of the benzyl alcohol substrate showed negative electronic/steric effects, resulting in products **4i** (53%) and **4j** (68%), whereas the *m*- and *p*-Cl, and *p*-Br-substituted benzyl alcohols reacted well with **2a** to afford the target products **4k–4m** in 78–82% yields. An obvious substituent effect was revealed on the reactivity of the chloro-substituted benzyl alcohols. 2-Naphthylmethanol efficiently underwent the reaction with **2a**, forming **4n** in 90% yield without showing a steric effect. With a higher catalyst loading (4.2 mol %) and longer reaction time (48 h) at a higher temperature (140 °C), 2- and 4-pyridylmethanols also reacted with **2a** to give **4o** and **4p** in 54–60% yields. In these cases, the catalyst might be partially deactivated by the coordination of the pyridyl atom to the metal center of the catalyst. Middle-chain aliphatic primary alcohols exhibited reactivities lower than those of the benzyl alcohols, and their reactions with **2a** were conducted

Table 3. Scope of Secondary Alcohols (2)^a




^aConditions: **2a** (2 mmol), **3** (2 mmol), 2.1 mol % **1b**, 30 mol % *t*BuOK, 0.1 MPa N₂, toluene (1 mL), 110 °C, 24 h. ^b4.2 mol % **1b**, *o*-xylene (1 mL), 140 °C, 48 h.

under the relatively harsh conditions, forming the target products **4q–4s** in 53–61% yields. These results have revealed that benzyl alcohols are more reactive than aliphatic primary alcohols in the β -alkylation of secondary alcohols.


Then, the reactions of benzyl alcohol (**3a**) with structurally diverse secondary alcohols were investigated (Table 3). Various functional groups such as methyl, methoxyl, chloro,

bromo, and trifluoromethyl could be tolerant on the aryl group of 1-arylethanol (**2**). *o*-, *m*-, and *p*-methyls did not exhibit an obvious impact on the formation of **5a–5c** (81–84%). 2-OMe and 4-OMe diminished the yields of **5d** (62%) and **5e** (74%). 3-Chloro also led to a relatively low yield for **5f** (70%), while 4-chloro facilitated the production of **5g** (80%). Unexpectedly, 3- and 4-Br groups promoted the reactions to yield the target products **5h** and **5i** in good yields (83–93%). The strong electron-withdrawing trifluoromethyl did not favor the reactions to produce **5j** (60%) and **5k** (70%). In a similar fashion, 1-(2-naphthyl)ethanol efficiently reacted with **3a** to afford **5l** (82%), exhibiting no steric effect. 1-(2-Pyridyl)ethanol and aliphatic secondary alcohols reacted with **3a** less efficiently, giving **5m–5p** in 54–61% yields by means of 4.2 mol % **1b** as the catalyst under relatively harsh conditions.

Next, the substrate scope was extended to cyclic secondary alcohols. Thus, the reactions of cyclopentanol **6** with benzyl alcohols were explored (Scheme 4). In order to reach double *C*-alkylation, 2 equiv of benzyl alcohol was applied in the presence of 4.2 mol % catalyst. To our delight, cyclopentanol **6** efficiently reacted with benzyl alcohols, affording the di(β -alkylated) secondary alcohol products, that is, 2,5-dibenzylcyclopentanol **7**, in 77–83% yields. However, cyclohexanol, 1,2,3,4-tetrahydro-naphthalen-1-ol, and 2,3-dihydro-1*H*-inden-1-ol could not effectively react with benzyl alcohols under the stated conditions. The substrate scope was further extended to potentially useful cyclic secondary alcohols. In a similar manner, cholesterol and its derivative




Scheme 4. Reactions of Cyclopentanol with Benzylic Alcohols



were employed to react with benzyl alcohols to access *C*-alkylated cholesterol derivatives (eqs 1 and 2). Unexpectedly, cholesterol (**8a**) reacted with excess of benzyl alcohol (3 equiv) afforded the monoalkylation product **9a** (57%) with the olefinic C=C bond hydrogenated. The cholesterol derivative, that is, 5 α -cholestan-3 β -ol (**8b**), also reacted well with benzyl alcohol and 1-(2-methylphenyl)ethanol to form the corresponding *C*-monoalkylation products **9a** (60%) and **9b** (56%), respectively. It should be noted that dialkylation did not occur to **8a** and **8b** under the stated conditions. The reactivity of the secondary carbon atoms of non-cyclic alcohols was also investigated. 1-Phenylpropan-1-ol was reacted with benzyl alcohol to give no target product even under harsh reaction conditions (160 °C, 48 h). This result revealed that the reactivity of the secondary carbon atoms of non-cyclic alcohols was much lower than that of cyclic alcohols.

This methodology was further applied for one-pot synthesis of flavan derivatives from simple alcohols.²⁷ Under the standard conditions, secondary alcohol **2a** was reacted with 2-bromobenzyl alcohol (**3t**). The intermediate alcohol was not isolated, and then directly subjected to the copper(I) catalysis, resulting in 2-phenylchroman (**10**) in 30% yield (Scheme 5).



Scheme 5. One-Pot Synthesis of Flavan Derivative 10

In summary, we have developed efficient manganese-catalyzed direct β -alkylation of secondary alcohols with primary alcohols under phosphine-free conditions. The present protocol offers a concise strategy to construct transition-metal complex catalysts from readily available phosphine-free ligands, and provides a useful alkylation method for secondary alcohols and cholesterol derivatives.

ASSOCIATED CONTENT

Supporting Information.

The Supporting Information is available free of charge on the Internet <http://pubs.acs.org>.

Experimental materials and procedures

Analytical data and NMR spectra of compounds and complexes

X-ray crystallographic analysis for **1b**

AUTHOR INFORMATION

Corresponding Author

* E-mail: zkvu@dicp.ac.cn.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

We are grateful to the National Natural Science Foundation of China (21672209) for support of this research.

REFERENCES

- (1) Rygus, J. P. G.; Crudden, C. M. Enantiospecific and Iterative Suzuki-Miyaura Cross-Couplings. *J. Am. Chem. Soc.* **2017**, *139*, 18124-18347.
- (2) Poliakoff, M.; Fitzpatrick, J. M.; Farren, T. R.; Anastas, P. T. Green Chemistry: Science and Politics of Change. *Science* **2002**, *297*, 807-810.
- (3) (a) Kim, S. W.; Zhang, W.; Krische, M. J. Catalytic Enantioselective Carbonyl Allylation and Propargylation via Alcohol-Mediated Hydrogen Transfer: Merging the Chemistry of Grignard and Sabatier. *Acc. Chem. Res.* **2017**, *50*, 2371-2380. (b) Chelucci, G. Ruthenium and Osmium Complexes in C-C Bond-Forming Reactions by Borrowing Hydrogen Catalysis. *Coord. Chem. Rev.* **2017**, *331*, 1-36. (c) Obara, Y. Recent Advances in α -Alkylation Reactions using Alcohols with Hydrogen Borrowing Methodologies. *ACS Catal.* **2014**, *4*, 3972-3981.
- (4) (a) Corma, A.; Navas, J.; Sabater, M. J. Advances in One-Pot Synthesis through Borrowing Hydrogen Catalysis. *Chem. Rev.* **2018**, *118*, 1410-1459. (b) Crabtree, R. H. Homogeneous Transition Metal Catalysis of Acceptorless Dehydrogenative Alcohol Oxidation: Applications in Hydrogen Storage and to Heterocycle Synthesis. *Chem. Rev.* **2017**, *117*, 9228-9246. (c) Huang, F.; Liu, Z. Q.; Yu, Z. K. C-Alkylation of Ketones and Related Compounds by Alcohols: Transition-Metal-Catalyzed Dehydrogenation. *Angew. Chem. Int. Ed.* **2016**, *55*, 862-875. (d) Yang, Q.; Wang, Q. F.; Yu, Z. K. Substitution of Alcohols by N-nucleophiles via Transition Metal-Catalyzed Dehydrogenation. *Chem. Soc. Rev.* **2015**, *44*, 2305-2329.
- (5) Schleppehorst, C.; Maji, B.; Glorius, F. Ruthenium-NHC Catalyzed α -Alkylation of Methylene Ketones Provides Branched Products through Borrowing Hydrogen Strategy. *ACS Catal.* **2016**, *6*, 4184-4188.
- (6) (a) Roy, B. C.; Debnath, S.; Chakrabarti, K.; Paul, B.; Maji, M.; Kundu, S. *ortho*-Amino Group Functionalized 2,2'-Bipyridine based Ru(II) Complex Catalyzed Alkylation of Secondary Alcohols, Nitriles and Amines using Alcohols. *Org. Chem. Front.* **2018**, *5*, 1008-1018. (b) Wang, Q. F.; Wu, K. K.; Yu, Z. K. Ruthenium(III)-Catalyzed β -Alkylation of Secondary Alcohols with Primary Alcohols. *Organometallics* **2016**, *35*, 1251-1256.
- (7) (a) Ruiz-Botella, S.; Peris, E. Unveiling the Importance of π -Stacking in Borrowing-Hydrogen Processes Catalyzed by Iridium Complexes with Pyrene Tags. *Chem. Eur. J.* **2015**, *21*, 15263-15271. (b) Jiménez, M. V.; Fernández-Tornos, J.; Modrego, F. J.; Pérez-Torrente, J. J.; Oro, L. A. Oxidation and β -Alkylation of Alcohols Catalyzed by Iridium(I) Complexes with Functionalised N-Heterocyclic Carbene Ligands. *Chem. Eur. J.* **2015**, *21*, 17877-17889.
- (8) Kose, O.; Saito, S. Cross-Coupling Reaction of Alcohols for Carbon-Carbon Bond Formation using Pincer-Type NHC/Palladium Catalysts. *Org. Biomol. Chem.* **2010**, *8*, 896-900.
- (9) (a) Tan, D.-W.; Li, H.-X.; Zhu, D.-L.; Li, H.-Y.; Young, D. J.; Yao, J.-L.; Lang, J.-P. Ligand-Controlled Copper(I)-Catalyzed Cross-Coupling of Secondary and Primary Alcohols to α -Alkylated Ketones, Pyridines, and Quinolines. *Org. Lett.* **2018**, *20*, 608-611. (b) Liao, S.; Yu, K.; Li, Q.; Tian, H.; Zhang, Z.; Yu, X.; Xu, Q. Copper-Catalyzed C-Alkylation of Secondary Alcohols and Methyl Ketones with Alcohols Employing the Aerobic Relay Race Methodology. *Org. Biomol. Chem.* **2012**, *10*, 2973-2978.
- (10) (a) Xu, Q.; Chen, J.; Tian, H.; Yuan, X.; Li, S.; Zhou, C.; Liu, J. Catalyst-Free Dehydrative α -Alkylation of Ketones with Alcohols: Green and Selective Autocatalyzed Synthesis of Alcohols and Ketones. *Angew. Chem. Int. Ed.* **2014**, *53*, 225-229. (b) Xu, Q.; Chen, J.; Liu, Q. Aldehyde-Catalyzed Transition Metal-Free Dehydrative β -Alkylation of Methyl Carbinols with Alcohols. *Adv. Synth. Catal.* **2013**, *355*, 697-704. (c) Allen, L. J.; Crabtree, R. H. Green Alcohol Couplings without Transition Metal Catalysts: Base-Mediated β -Alkylation of Alcohols in Aerobic Conditions. *Green Chem.* **2010**, *12*, 1362-1364.
- (11) (a) Balaraman, E.; Nandakumar, A.; Jaiswal, G.; Sahoo, M. K. Iron-Catalyzed Dehydrogenation Reactions and Their Applications in

Sustainable Energy and Catalysis. *Catal. Sci. Technol.* **2017**, *7*, 3177-3195. (b) Zell, T.; Milstein, D. Hydrogenation and Dehydrogenation Iron Pincer Catalysts Capable of Metal-Ligand Cooperation by Aromatization/De aromatization. *Acc. Chem. Res.* **2015**, *48*, 1979-1994.

(12) (a) Freitag, F.; Irrgang, T.; Kempe, R. Cobalt-Catalyzed Alkylation of Secondary Alcohols with Primary Alcohols via Borrowing Hydrogen/Hydrogen Autotransfer. *Chem. Eur. J.* **2017**, *23*, 12110-12113. (b) Zhang, G.; Wu, J.; Zeng, H.; Zhang, S.; Yin, Z.; Zheng, S. Cobalt-Catalyzed α -Alkylation of Ketones with Primary Alcohols. *Org. Lett.* **2017**, *19*, 1080-1083.

(13) Vellakkaran, M.; Singh, K.; Banerjee, D. An Efficient and Selective Nickel-Catalyzed Direct N-Alkylation of Anilines with Alcohols. *ACS Catal.* **2017**, *7*, 8152-8158.

(14) (a) Chakraborty, S.; Das, U. K.; Ben-David, Y.; Milstein, D. Manganese Catalyzed α -Olefination of Nitriles by Primary Alcohols. *J. Am. Chem. Soc.* **2017**, *139*, 11710-11713. (b) Mukherjee, A.; Nerush, A.; Leitus, G.; Shimon, L. J. W.; Ben-David, Y.; Espinosa Jalapa, N. A.; Milstein, D. Manganese-Catalyzed Environmentally Benign Dehydrogenative Coupling of Alcohols and Amines to Form Aldimines and H₂: A Catalytic and Mechanistic Study. *J. Am. Chem. Soc.* **2016**, *138*, 4298-4301.

(15) Chakraborty, S.; Gellrich, U.; Diskin-Posner, Y.; Leitus, G.; Avram, L.; Milstein, D. Manganese-Catalyzed N-Formylation of Amines by Methanol Liberating H₂: A Catalytic and Mechanistic Study. *Angew. Chem. Int. Ed.* **2017**, *56*, 4229-4233.

(16) Bauer, J. O.; Chakraborty, S.; Milstein, D. Manganese-Catalyzed Direct Deoxygenation of Primary Alcohols. *ACS Catal.* **2017**, *7*, 4462-4466.

(17) (a) Das, U. K.; Ben-David, Y.; Diskin-Posner, Y.; Milstein, D. N-Substituted Hydrazones by Manganese-Catalyzed Coupling of Alcohols with Hydrazine: 'Borrowing Hydrogen' and Acceptorless Dehydrogenation in One System. *Angew. Chem. Int. Ed.* **2017**, *51*, 2179-2182. (b) Garbe, M.; Junge, K.; Beller, M. Homogeneous Catalysis by Manganese-Based Pincer Complexes. *Eur. J. Org. Chem.* **2017**, 4344-4362. (c) Elangovan, S.; Neumann, J.; Sortais, J.-B.; Junge, K.; Darcel, C.; Beller, M. Efficient and Selective N-Alkylation of Amines with Alcohols Catalysed by Manganese Pincer Complexes. *Nat. Commun.* **2016**, *7*, 12641.

(18) Peña-López, M.; Piehl, P.; Elangovan, S.; Neumann, H.; Beller, M. Manganese-Catalyzed Hydrogen-Autotransfer C-C Bond Formation: α -Alkylation of Ketones with Primary Alcohols. *Angew. Chem. Int. Ed.* **2016**, *55*, 14967-14971.

(19) (a) Shao, Z.; Wang, Y.; Liu, Y.; Wang, Q.; Fu, X.; Liu, Q. A General and Efficient Mn-Catalyzed Acceptorless Dehydrogenative Coupling of Alcohols with Hydroxides into Carboxylates. *Org. Chem. Front.* **2018**, *5*, 1248-1256. (b) Nguyen, D. H.; Trivelli, X.; Capet, F.; Paul, J.-F.; Dumeignil, F.; Gauvin, R. M. Manganese Pincer Complexes for the Base-Free, Acceptorless Dehydrogenative Coupling of Alcohols to Esters: Development, Scope, and Understanding. *ACS Catal.* **2017**, *7*, 2022-2032.

(20) Andérez-Fernández, M.; Vogt, L. K.; Fischer, S.; Zhou, W.; Jiao, H.; Garbe, M.; Elangovan, S.; Junge, K.; Junge, H.; Ludwig, R.; Beller, M. A Stable Manganese Pincer Catalyst for the Selective Dehydrogenation of Methanol. *Angew. Chem. Int. Ed.* **2017**, *56*, 559-562.

(21) (a) Kallmeier, F.; Kempe, R. Manganese Complexes for (De)Hydrogenation Catalysis: A Comparison to Cobalt and Iron Catalysts. *Angew. Chem. Int. Ed.* **2018**, *57*, 46-60. (b) Deibl, N.; Kempe, R. Manganese-Catalyzed Multicomponent Synthesis of Pyrimidines from Alcohols and Amidines. *Angew. Chem. Int. Ed.* **2017**, *56*, 1663-1666. (c) Kallmeier, F.; Dudzic, B.; Irrgang, T.; Kempe, R. Manganese-Catalyzed Sustainable Synthesis of Pyrroles from Alcohols and Amino Alcohols. *Angew. Chem. Int. Ed.* **2017**, *56*, 7261-7265. (d) Mastalir, M.; Glatz, M.; Pittenauer, E.; Allmaier, G.; Kirchner, K. Sustainable Synthesis of Quinolines and Pyrimidines Catalyzed by Manganese PNP Pincer Complexes. *J. Am. Chem. Soc.* **2016**, *138*, 15543-15546.

(22) (a) Filonenko, G. A.; van Putten, R.; Hensen, E. J. M.; Pidko, E. A. Catalytic (De)Hydrogenation Promoted by Non-Precious Metals-Co, Fe and Mn: Recent Advances in an Emerging Field. *Chem. Soc. Rev.* **2018**, *47*, 1459-1483. (b) Bertini, F.; Glatz, M.; Gorgas, N.; Stöger, B.; Peruzzini, M.; Veiros, L. F.; Kirchner, K.; Gonsalvi, L. Carbon Dioxide Hydrogenation Catalysed by Well-Defined Mn(I) PNP Pincer Hydride Complexes. *Chem. Sci.* **2017**, *8*, 5024-5029. (c) Dubey, A.; Nencini, L.; Fayzullin, R. R.; Nervi, C.; Khusnutdinova, J. R. Bio-Inspired Mn(I) Complexes for the Hydrogenation of CO₂ to Formate and Formamide. *ACS Catal.* **2017**, *7*, 3864-3868. (d) Bruneau-Voisine, A.; Wang, D.; Dorcet, V.; Roisnel, T.; Darcel, C.; Sortais, J.-B. Transfer Hydrogenation of Carbonyl Derivatives Catalyzed by an Inexpensive Phosphine-Free Manganese Precatalyst. *Org. Lett.* **2017**, *19*, 3656-3659.

(23) (a) Wei, D.; Bruneau-Voisine, A.; Valyaev, D. A.; Lugan, N.; Sortais, J.-B. Manganese Catalyzed Reductive Amination of Aldehydes using Hydrogen as a Reductant. *Chem. Commun.* **2018**, *54*, 4302-4305. (b) Espinosa-Jalapa, N. A.; Nerush, A.; Shimon, L. J. W.; Leitus, G.; Avram, L.; Ben-David, Y.; Milstein, D. Manganese-Catalyzed Hydrogenation of Esters to Alcohols. *Chem. Eur. J.* **2017**, *23*, 5934-5938. (c) Elangovan, S.; Topf, C.; Fischer, S.; Jiao, H.; Spannenberg, A.; Baumann, W.; Ludwig, R.; Junge, K.; Beller, M. Selective Catalytic Hydrogenations of Nitriles, Ketones, and Aldehydes by Well-Defined Manganese Pincer Complexes. *J. Am. Chem. Soc.* **2016**, *138*, 8809-8814. (d) Kallmeier, F.; Irrgang, T.; Dietel, T.; Kempe, R. Highly Active and Selective Manganese C=O Bond Hydrogenation Catalysts: The Importance of the Multidentate Ligand, the Ancillary Ligands, and the Oxidation State. *Angew. Chem. Int. Ed.* **2016**, *55*, 11806-11809.

(24) Zhang, G.; Irrgang, T.; Dietel, T.; Kallmeier, F.; Kempe, R. Manganese-Catalyzed Dehydrogenative Alkylation or α -Olefination of Alkyl-Substituted N-Heteroarenes with Alcohols. *Angew. Chem. Int. Ed.* **2018**, *57*, 10102/anie.201801573.

(25) (a) Wang, Q. F.; Chai, H. N.; Yu, Z. K. Acceptorless Dehydrogenation of N-Heterocycles and Secondary Alcohols by Ru(II)-NNC Complexes Bearing a Pyrazoyl-indolyl-pyridine Ligand. *Organometallics* **2018**, *37*, 584-591. (b) Chai, H. N.; Wang, L. D.; Liu, T. T.; Yu, Z. K. A Versatile Ru(II)-NNP Complex Catalyst for the Synthesis of Multisubstituted Pyrroles and Pyridines. *Organometallics*, **2017**, *36*, 4936-4932. (c) Chai, H. N.; Liu, T. T.; Yu, Z. K. Cooperative N-H and CH₂ Skeleton Effects on the Catalytic Activities of Bimetallic Ru(II)-NNN Complexes: Experimental and Theoretical Study. *Organometallics* **2017**, *36*, 4136-4144. (d) Liu, T. T.; Chai, H. N.; Wang, L. D.; Yu, Z. K. Exceptionally Active Assembled Dinuclear Ruthenium(II)-NNN Complex Catalysts for Transfer Hydrogenation of Ketones. *Organometallics* **2017**, *36*, 2914-2921. (e) Zeng, F. L.; Yu, Z. K. Construction of Highly Active Ruthenium(II) NNN Complex Catalysts Bearing a Pyridyl-Supported Pyrazolyl-Imidazolyl Ligand for Transfer Hydrogenation of Ketones. *Organometallics* **2009**, *28*, 1855-1862.

(26) (a) Faller, J. W. In *Comprehensive Organometallic Chemistry III*; Crabtree, R. H., Mingos, D. M. P., Eds.; Elsevier: Oxford, U.K., 2007; pp 407-427. (b) Viviente, E. M.; Pregosin, P. S.; Schott, D. In *Mechanisms in Homogeneous Catalysis: A Spectroscopic Approach*; Heaton, B., Ed.; Wiley: Germany, 2005; pp 1-80.

(27) Shee, S.; Paul, B.; Panja, D.; Roy, B. C.; Chakrabarti, K.; Ganguli, K.; Das, A.; Das, G. K.; Kundu, S. Tandem Cross Coupling Reaction of Alcohols for Sustainable Synthesis of β -Alkylated Secondary Alcohols and Flavan Derivatives. *Adv. Synth. Catal.* **2017**, *359*, 3888-3893.

Table of Contents (TOC) Graphic:

