

Electrophile-Driven Regioselective Synthesis of Functionalized Quinolines


Shaukat Ali, Hai-Tao Zhu, Xiao-Feng Xia, Ke-Gong Ji, Yan-Fang Yang, Xian-Rong Song, and Yong-Min Liang*

State Key Laboratory of Applied Organic Chemistry, Lanzhou University, and State Key Laboratory of Solid Lubrication, Lanzhou Institute of Chemical Physics, Chinese Academy of Science, Lanzhou 730000, People's Republic of China

liangym@lzu.edu.cn

Received March 16, 2011

ABSTRACT



Highly substituted 3-iodoquinolines bearing different alkyl and aryl moieties can be synthesized in moderate to excellent yields (up to 99%) by 6-endo-dig iodocyclization of 2-tosylaminophenylprop-1-yn-3-ols with molecular iodine (I_2) under mild conditions. The cyclization is highly regioselective and the resulting 3-iodoquinolines can be further functionalized by various coupling reactions.

Quinoline derivatives are a major class of heterocycles which occur in various natural products, especially in alkaloids. The quinoline skeleton is often used for the construction of many synthetic compounds with diverse pharmacological properties.¹ In particular, halogen-containing quinolines are of special interest because the halogen atom sometimes plays an important role in the compound's bioactivity, and such compounds provide a further avenue for functionalization.² As a result, a number of attempts have been made to prepare and functionalize them since the late 1800s. The structural core of quinoline can be synthesized by various conventional methods.^{3–8} However, these classical syntheses often make use of elevated temperatures and prolonged reaction times and/or use of strong acids and bases, which cannot be applied

to sensitive substrates. There are also many new metal-catalyzed protocols for preparing quinolines.^{9–12} Thus, a metal free, mild, and environmentally benign protocol for preparing quinolines from readily available and simple substrates is still of high demand because of their extreme significance. In recent years, the electrophilic cyclization of heteroatom nucleophiles, such as oxygen, nitrogen, and sulfur, with tethered alkynes has proven to be an effective

(1) Kouznetsov, V. V.; Vargas Méndez, L. Y.; Meléndez Gómez, C. M. *Curr. Org. Chem.* **2005**, *9*, 141–161.

(2) (a) Newhouse, B. J.; Bordner, J.; Augeri, D. J.; Litts, C. S.; Kleinman, E. F. *J. Org. Chem.* **1992**, *57*, 6991. (b) Torii, S.; Xu, L. H.; Sadakane, M.; Okumoto, H. *Synlett* **1992**, 513. (c) Nobuhide, M.; Yoshinobu, Y.; Hiroshi, I.; Yoshio, O.; Tamejiro, H. *Tetrahedron Lett.* **1993**, *24*, 8263. (d) Croisey-Delcey, M.; Croisy, A.; Carrez, D.; Huel, C.; Chiaroni, A.; Ducrot, P.; Bisagni, E.; Jin, L.; Leclercq, G. *Bioorg. Med. Chem.* **2000**, *8*, 2629.

(3) Manske, R. H. F.; Kukla, M. *Org. React.* **1953**, *7*, 59.

(4) Bergstrom, F. W. *Chem. Rev.* **1944**, *35*, 153.

(5) Cheng, C.-C.; Yan, S.-J. *Org. React.* **1982**, *28*, 37.

(6) Jones, G. In *The Chemistry of Heterocyclic Compounds*; Weissberger, A., Taylor, E. C., Eds.; John Wiley and Sons: Chichester, UK, 1977; Vol. 32, Part I, pp 93–318.

(7) Reitsema, R. H. *Chem. Rev.* **1948**, *43*, 47.

(8) Bergstrom, F. W. *Chem. Rev.* **1944**, *35*, 156.

(9) For selected papers, see: (a) Berman, A. M.; Lewis, J. C.; Bergman, R. G.; Ellman, J. A. *J. Am. Chem. Soc.* **2008**, *130*, 14926. (b) Horn, J.; Marsden, S. P.; Nelson, A.; House, D.; Weingarten, G. G. *Org. Lett.* **2008**, *10*, 4117.

(10) (a) Arisawa, M.; Nishida, A.; Nakagawa, M. *J. Organomet. Chem.* **2006**, *691*, 5109. (b) Movassaghi, M.; Hill, M. D. *J. Am. Chem. Soc.* **2006**, *128*, 4592.

(11) Liu, X.-Y.; Che, C.-M. *Angew. Chem., Int. Ed.* **2008**, *47*, 3805.


(12) For selected papers, see: (a) Zhang, Z.; Tan, J.; Wang, Z. *Org. Lett.* **2008**, *10*, 173. (b) Sakai, N.; Annaka, K.; Fujita, A.; Sato, A.; Konakahara, T. *J. Org. Chem.* **2008**, *73*, 4160. (c) Arcadi, A.; Aschi, M.; Marinelli, F.; Verdecchia, M. *Tetrahedron* **2008**, *64*, 5354. (d) Li, L.; Jones, W. D. *J. Am. Chem. Soc.* **2007**, *129*, 10707. (e) Gabriele, B.; Mancuso, R.; Salerno, G.; Ruffolo, G.; Plastina, P. *J. Org. Chem.* **2007**, *72*, 6873. (f) Sakai, N.; Annaka, K.; Konakahara, T. *J. Org. Chem.* **2006**, *71*, 3653. (g) Godet, T.; Belmont, P. *Synlett* **2008**, 2513.

(13) (a) Bi, H.-P.; Guo, L.-N.; Duan, X.-H.; Gou, F.-R.; Huang, S.-H.; Liu, X.-Y.; Liang, Y.-M. *Org. Lett.* **2007**, *9*, 397. (b) Xie, Y.-X.; Liu, X.-Y.; Wu, L.-Y.; Han, Y.; Zhao, L.-B.; Fan, M.-J.; Liang, Y.-M. *Eur. J. Org. Chem.* **2008**, 1013. (c) Wen, S.-G.; Liu, W.-M.; Liang, Y.-M. *J. Org. Chem.* **2008**, *73*, 4342. (d) Xie, Y.-X.; Yan, Z.-Y.; Qian, B.; Deng, W.-Y.; Wu, L.-Y.; Liu, X.-Y.; Liang, Y.-M. *Chem. Commun.* **2009**, 5451. (e) Xie, Y.-X.; Yan, Z.-Y.; Wang, D.-Z.; Wu, L.-Y.; Qian, B.; Liu, X.-Y.; Liang, Y.-M. *Eur. J. Org. Chem.* **2009**, 2283. (f) Ji, K.-G.; Zhu, H.-T.; Yang, F.; Shu, X.-Z.; Zhao, S.-C.; Shaukat, A.; Liu, X.-Y.; Liang, Y.-M. *Chem.—Eur. J.* **2010**, *16*, 1615.

method for the synthesis of heterocyclic compounds.^{13–31} We¹³ and others have reported that electrophilic cyclization¹⁴ of alkynes can be a very powerful tool for the preparation of a wide variety of important heterocyclic compounds due to the mild, efficient, and clean reactions. Many important heterocycles, such as benzo[*b*]thiophenes,¹⁵ benzofurans,¹⁶ 2,3-dihydropyrroles and pyrroles,¹⁷ furans,^{13b,18} dihydropyrans,^{13f,19} indoles,²⁰ isochromenes,²¹ isocoumarins and R-pyrones,²² isoquinolines and quinolines,²³ isoxazoles²⁴ and oxazoles,²⁵ furanones,²⁶ furopyridines,²⁷ spiro[4,5]trienones,²⁸ coumestans and coumestrols,²⁹ naphthols,³⁰ and naphthalenes,³¹ have been reported based on this strategy. Also very recently, Wai's group reported the synthesis of *N*-tosylated indoles using 2-tosylaminophenylprop-1-yn-3-ols as starting materials in the presence of gold catalyst.³² We envisioned that the treatment of 2-tosylaminophenylprop-1-yn-3-ol (**1a**) with molecular iodine (I₂) would lead to iodocyclization in a 5-exo-dig manner resulting in an indole derivative. To test the viability of the above hypothesis we treated **1a** with I₂, but unexpectedly the ¹H and ¹³C NMR data revealed quinoline derivative **2a** (Scheme 1). Herein, we report a general and highly regioselective protocol for the synthesis of substituted quinolines by 6-endo-dig iodocyclization of 2-tosylaminophenylprop-1-yn-3-ols. In 2005, although Flynn developed the synthesis of 3-iodoquinolinium salts by a related process,³³ the lengthy preparation of their starting materials, the low substrate scope, and the separation of quinolinium salts make it less attractive synthetically. We were extremely pleased to see that our 2-tosylaminophenylprop-1-yn-3-ols reacted smoothly to give 3-iodoquinolines in a single step with no quinolinium salts.

To explore suitable reaction conditions for this useful transformation, we started by using 0.2 mmol of **1a** and 1.5 equiv of I₂ in nitromethane at room temperature; to our

Scheme 1. Design of an Electrophile-Driven Cascade Cyclization




delight, the quinoline derivative **2a** was obtained in 38% yield after 2 h. Increasing the amount of iodine to 2.0 equiv, the yield slightly increased to 41%. Further increasing the amount of iodine resulted in a decrease of yield. The influence of different acids and increase of reaction temperature in nitromethane was studied but no useful results were obtained. We then discovered that the best result can be obtained by carrying out the reaction in methanol at 60 °C. Thus, changing the solvent from polar aprotic to polar protic the yield dramatically increased to 93%. The protic nature of the solvent is most probably involved in the easy removal of hydroxyl group to give intermediate **D** (see the mechanism, Scheme 2). Solvents such as dichloromethane, acetonitrile, and tetrahydrofuran were found to be less effective, whereas dimethylformamide and 1,4-dioxane were found to be ineffective (see the Supporting Information).

To investigate the scope of the cascade iodocyclization a variety of differently aliphatic and aromatic substituted 2-tosylaminophenylprop-1-yn-3-ols were subjected to the



Table 1. Cascade 6-Endo-Dig Iodocyclization of 2-Tosylaminophenylprop-1-yn-3-ols (eq 1)^a

entry	no.	alcohol			time (h)	quinoline	isolated yield (%)
		R ¹	R ²	R ³			
1	1a	H	Ph	Ph	6	2a	93
2	1b	H	Ph	<i>p</i> -MeC ₆ H ₄	6	2b	94
3	1c	H	Ph	<i>p</i> -MeOC ₆ H ₄	6	2c	96
4	1d	H	Ph	<i>p</i> -ClC ₆ H ₄	6	2d	88
5	1e	H	Ph	<i>p</i> -BrC ₆ H ₄	6	2e	81
6	1f	H	Ph	<i>m</i> -MeC ₆ H ₄	6	2f	77
7	1g	H	Ph	<i>m</i> -MeOC ₆ H ₄	6	2g	84
8	1h	H	Ph	<i>o</i> -MeC ₆ H ₄	12	2h	66
9	1i	H	Ph	<i>o</i> -ClC ₆ H ₄	12	2i	71
10	1j	H	Ph	<i>o,p</i> -Cl ₂ C ₆ H ₃	12	2j	60
11	1k	5-Cl	Ph	Ph	12	2k	88
12	1l	5-Br	Ph	Ph	12	2l	76
13	1m	H	Ph	<i>n</i> -propyl	12	2m	62
14	1n	H	<i>p</i> -MeC ₆ H ₄	Ph	6	2n	74
15	1o	H	<i>p</i> -BrC ₆ H ₄	Ph	6	2o	99
16	1p	H	Me	Ph	12	2p	54
17	1q	H	Et	Ph	12	2q	40

^a All reactions were run with 0.2 mmol of the alcohol, 2 equiv of I₂ in 2.0 mL of CH₃OH at 60 °C, followed by addition of 8.0 mL of saturated aqueous Na₂S₂O₃ to remove the excess I₂.



The mechanism of this iodocyclization involves anti-


$$\text{Ph} \quad \text{Ph}$$


- (14) For early studies on the electrophilic cyclization reaction, see: (a) Ren, X.-F.; Turos, E. *Tetrahedron Lett.* **1993**, *34*, 1575. (b) Kitamura, T.; Takachi, T.; Kawasato, H.; Taniguchi, H. *J. Chem. Soc., Perkin Trans.* **1992**, *1*, 1969. (c) Kitamura, T.; Kobayashi, S.; Taniguchi, H.; Hori, K. *J. Am. Chem. Soc.* **1991**, *113*, 6240. (d) Ten Hoedt, R. W. M.; Van Koten, G.; Noltes, J. G. *Synth. Commun.* **1977**, *7*, 61. (e) Sonoda, T.; Kawakami, M.; Ikeda, T.; Kobayashi, S.; Taniguchi, H. *J. Chem. Soc., Chem. Commun.* **1976**, 612.
- (15) (a) Larock, R. C.; Yue, D. *Tetrahedron Lett.* **2001**, *42*, 6011. (b) Yue, D.; Larock, R. C. *J. Org. Chem.* **2002**, *67*, 1905. (c) Flynn, B. L.; Verdier-Pinard, P.; Hamel, E. *Org. Lett.* **2001**, *3*, 651. (d) Hessian, K. O.; Flynn, B. L. *Org. Lett.* **2003**, *5*, 4377.
- (16) (a) Arcadi, A.; Cacchi, S.; Fabrizi, G.; Marinelli, F.; Moro, L. *Synlett* **1999**, 1432. (b) Mehta, S.; Waldo, J. P.; Larock, R. C. *J. Org. Chem.* **2009**, *74*, 1141.
- (17) (a) Knight, D. W.; Redfern, A. L.; Gilmore, J. *Chem. Commun.* **1998**, 2207. (b) Knight, D. W.; Redfern, A. L.; Gilmore, J. *J. Chem. Soc., Perkin Trans.* **2001**, *1*, 2874. (c) Knight, D. W.; Redfern, A. L.; Gilmore, J. *J. Chem. Soc., Perkin Trans.* **2002**, *1*, 622.
- (18) (a) Bew, S. P.; Knight, D. W. *Chem. Commun.* **1996**, 1007. (b) Djuardi, E.; McNelis, E. *Tetrahedron Lett.* **1999**, *40*, 7193. (c) Sniady, A.; Wheeler, K. A.; Dembinski, R. *Org. Lett.* **2005**, *7*, 1769. (d) Yao, T.; Zhang, X.; Larock, R. C. *J. Org. Chem.* **2005**, *70*, 7679. (e) Liu, Y.; Zhou, S. *Org. Lett.* **2005**, *7*, 4609. (f) Bew, S. P.; El-Taeb, G. M. M.; Jones, S.; Knight, D. W.; Tan, W.-F. *Eur. J. Org. Chem.* **2007**, 5759. (g) Arimitsu, S.; Jacobsen, J. M.; Hammond, G. B. *J. Org. Chem.* **2008**, *73*, 2886. (h) Huang, X.; Fu, W.; Miao, M. *Tetrahedron Lett.* **2008**, *49*, 2359.
- (19) (a) Mohapatra, D. K.; Das, P. P.; Pattanayak, M. R.; Yadav, J. S. *Chem.—Eur. J.* **2010**, *16*, 2072. (b) Mancuso, R.; Mehta, S.; Gabriele, B.; Salerno, G.; Jenks, W. S.; Larock, R. C. *J. Org. Chem.* **2010**, *75*, 897.
- (20) (a) Barluenga, J.; Trincado, M.; Rublio, E.; Gonzalez, J. M. *Angew. Chem., Int. Ed.* **2003**, *42*, 2406. (b) Amjad, M.; Knight, D. W. *Tetrahedron Lett.* **2004**, *45*, 539. (c) Yue, D.; Larock, R. C. *Org. Lett.* **2004**, *6*, 1037.
- (21) (a) Barluenga, J.; Vazquez-Villa, H.; Ballesteros, A.; Gonzalez, J. M. *J. Am. Chem. Soc.* **2003**, *125*, 9028. (b) Yue, D.; Della Ca, N.; Larock, R. C. *Org. Lett.* **2004**, *6*, 1581.
- (22) (a) Yao, T.; Larock, R. C. *Tetrahedron Lett.* **2002**, *43*, 7401. (b) Yao, T.; Larock, R. C. *J. Org. Chem.* **2003**, *68*, 5936. (c) Oliver, M. A.; Gandour, R. D. *J. Org. Chem.* **1984**, *49*, 558. (d) Biagetti, M.; Bellina, F.; Carpita, A.; Stabile, P.; Rossi, R. *Tetrahedron* **2002**, *58*, 5023. (e) Rossi, R.; Carpita, A.; Bellina, F.; Stabile, P.; Mannina, L. *Tetrahedron* **2003**, *59*, 2067.
- (23) (a) Huang, Q.; Hunter, J. A.; Larock, R. C. *Org. Lett.* **2001**, *3*, 2973. (b) Huang, Q.; Hunter, J. A.; Larock, R. C. *J. Org. Chem.* **2002**, *67*, 3437. (c) Barluenga, J.; Gonzalez, J. M.; Campos, P. J.; Asensio, G. *Angew. Chem., Int. Ed. Engl.* **1988**, *27*, 1546. (d) Huo, Z.; Gridnev, I. D.; Yamamoto, Y. *J. Org. Chem.* **2010**, *75*, 1266. (e) Fischer, D.; Tomeba, H.; Pahadi, N. K.; Patil, N. T.; Huo, Z.; Yamamoto, Y. *J. Am. Chem. Soc.* **2008**, *130*, 15720. (f) Ding, Q.; Wu, J. *Adv. Synth. Catal.* **2008**, *350*, 1850. (g) Ding, Q.; Wang, Z.; Wu, J. *J. Org. Chem.* **2009**, *74*, 921.
- (24) Waldo, J. P.; Larock, R. C. *Org. Lett.* **2005**, *7*, 5203.
- (25) Hashmi, A. S. K.; Weyrauch, J. P.; Frey, W.; Bats, W. *J. Org. Lett.* **2004**, *6*, 4391.

cation **D**. Finally, **D** gains full aromaticity by eliminating a tosyl group to form quinoline derivative **2** (Scheme 2).

In conclusion, we have discovered an efficient and highly regioselective protocol for preparing quinolines from readily available and simple starting materials. The reaction is metal-free and can produce substituted 3-iodoquinolines in a single step in moderate to excellent yields. The resulting 3-iodoquinolines can be further functionalized by using known organopalladium chemistry, such as Sonogashira³⁴

and Suzuki³⁵ cross-couplings and the Heck reaction³⁶ (Scheme 3). Further exploration of the reaction scope and mechanism is underway.

Acknowledgment. We thank the National Science Foundation (NSF-21072080) for financial support. We also acknowledge the National Basic Research Program of China (973 Program), 2010CB833203.

Supporting Information Available. Detailed experimental procedures, copies of ¹H NMR and ¹³C NMR spectra of all compounds, and X-ray data of **2b** in CIF format. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(26) (a) Just, Z. W.; Larock, R. C. *J. Org. Chem.* **2008**, *73*, 2662. (b) Crone, B.; Kirsch, S. F. *J. Org. Chem.* **2007**, *72*, 5435.

(27) Arcadi, A.; Cacchi, S.; Di Giuseppe, S.; Fabrizi, G.; Marinelli, F. *Org. Lett.* **2002**, *4*, 2409.

(28) (a) Zhang, X.; Larock, R. C. *J. Am. Chem. Soc.* **2005**, *127*, 12230. (b) Tang, B.-X.; Tang, D.-J.; Tang, S.; Yu, Q.-F.; Zhang, Y.-H.; Liang, Y.; Zhong, P.; Li, J.-H. *Org. Lett.* **2008**, *10*, 1063.

(29) Yao, T.; Yue, D.; Larock, R. C. *J. Org. Chem.* **2005**, *70*, 9985.

(30) Zhang, X.; Sarkar, S.; Larock, R. C. *J. Org. Chem.* **2006**, *71*, 236.

(31) Barluenga, J.; Vazquez-Villa, H.; Ballesteros, A.; Gonzalez, J. M. *Org. Lett.* **2003**, *5*, 4121.

(32) Kothandaraman, P.; Rao, W.; Foo, S. J.; Philip Wai, H. C. *Angew. Chem., Int. Ed.* **2010**, *49*, 1–6.

(33) Hessian, K. O.; Flynn, B. L. *Org. Lett.* **2005**, *8*, 243–246.

(34) (a) Sonogashira, K. In *Metal-Catalyzed Cross-Coupling Reactions*; Diederich, F., Stang, P. J., Eds.; Wiley-VCH: Weinheim, Germany, 1998; Chapter 5, pp 203–229. (b) Sonogashira, K.; Tohda, Y.; Hagihara, N. *Tetrahedron Lett.* **1975**, 4467.

(35) (a) Suzuki, A. *J. Organomet. Chem.* **1999**, *576*, 147. (b) Lloyd-Williams, P.; Giralt, E. *Chem. Soc. Rev.* **2001**, *30*, 145. (c) Suzuki, A. *Metal-Catalyzed Cross-Coupling Reactions*; Diederich, F., Stang, P. J., Eds.; Wiley-VCH: Weinheim, Germany, 1998; 49. (d) Miyaura, N. *Chem. Rev.* **1995**, *95*, 2457.

(36) (a) Crisp, G. T. *Chem. Soc. Rev.* **1998**, *27*, 427. (b) Brase, S.; De Meijere, A. *Metal-Catalyzed Cross-Coupling Reactions*; Diederich, F., Stang, P. J., Eds.; Wiley-VCH: Weinheim, Germany, 1998; p 99. (c) Heck, R. F. *Palladium Reagents in Organic Synthesis*; Academic Press: San Diego, CA, 1985; pp 276–287.