This article was downloaded by: [University of Tennessee, Knoxville] On: 13 April 2014, At: 22:29 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/gpss20</u>

On the Reaction of Triaryl Trithioarsenites, (ArS)₃As, with Iodine

Maria D. Markaki^a & Panayiotis V. Ioannou^a ^a Department of Chemistry, University of Patras, Patras, Greece Published online: 15 Aug 2006.

To cite this article: Maria D. Markaki & Panayiotis V. Ioannou (2006) On the Reaction of Triaryl Trithioarsenites, $(ArS)_3As$, with Iodine, Phosphorus, Sulfur, and Silicon and the Related Elements, 181:4, 921-929, DOI: <u>10.1080/10426500500272178</u>

To link to this article: http://dx.doi.org/10.1080/10426500500272178

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.



This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions





On the Reaction of Triaryl Trithioarsenites, $(ArS)_3As$, with lodine

Maria D. Markaki Panayiotis V. Ioannou Department of Chemistry, University of Patras, Patras, Greece

Iodine in dry nonprotic solvents oxidizes triaryl trithioarsenites, $(ArS)_3As$, to As(III) iodide, AsI_3 , and disulfides, ArSSAr. The reaction most likely involves arylsulfenyl iodide, ArSI, as an intermediate. With triphenyl and tris(4-chlorophenyl) trithioarsenites, AsI_3 is prepared in very good yields. When the disulfide, which is produced, has MeCONH- or $-NH-CMe_2-OH$ groups, then it acts as a Lewis base towards AsI_3 forming complexes with stoichiometry $2AsI_3 \cdot 3ArSSAr$. Probable coordination modes of the AsI_3 are discussed.

Keywords Arsenic triiodide; arsenic triiodide as Lewis acid; diphenyl aryldithioarsonites; disulfides; iodine; triaryl trithioarsenites

INTRODUCTION

We have recently been studying the effect of the oxidizing element octasulfur on dithioarsonites [L-As(SPh)₂, L = Ar or R]¹ and on triaryl² and trialkyl³ trithioarsenites [(ArS)₃As and (RS)₃As], spurred by the (erroneous) report⁴ that (*p*-MePhS)₃As is oxidized by octasulfur in refluxing carbon disulfide to the ester (*p*-MePhS)₃As=S. We have found that the previously mentioned As(III) esters are not oxidized to the As(V) ones probably because the As(III) is not nucleophilic towards octasulfur.

Next, we studied the air oxidation of triaryl⁵ and trialkyl³ trithioarsenites, and we found that electron-withdrawing groups at the *para* position of the phenyl group tend to favor the oxidative decomposition compared to those having electron-donating groups, according to Eq. (1):

$$2(ArS)_3As + 3/2O_2 \rightarrow As_2O_3 + 3ArSSAr$$
(1)

Aliphatic trithioarsenites bearing a terminal -COOH were decomposed by air (Eq. 1), while those bearing a $-COO^-$ group were

Address correspondence to Panayiotis V. Ioannou, University of Patras, Department of Chemistry, Patras, Greece. E-mail: ioannou@chemistry.upatras.gr



Received March 24, 2005; accepted April 12, 2005.

resistant to air oxidation. The esters $As(glutathione)_3$, $As(cysteine)_3$, and $As(penicillamine)_3$ were found to be air oxidized at a slower rate than their parent thiols. These and other⁶ oxidative decomposition results have been rationalized by assuming dioxygen binding to As(III). The adduct then decomposes to the observed products.

Stronger oxidizing elements other than octasulfur and dioxygen are the halogens. The action of iodine, dissolved in ethanol, on the ester $\mathbf{1}$ with an unspecified stoichiometry was reported⁷ to give diphenyl disulfide, but the fate of the arsenic was not determined.



FORMULAS 1-7

In this article, we report our results on the action of iodine and, in one case, bromine on the trithioarsenites **1–7**. We found that the reaction of iodine with the esters **1–7** is simpler in nonprotic solvents; arsenic triiodide is produced that either precipitates out or forms Lewis adducts when the produced disulfide has certain nitrogen and/or oxygen donor atoms.

RESULTS AND DISCUSSION

The reaction of **1** with iodine in methanol under various stoichiometries gave a complex mixture of products, among them diphenyl disulfide, as revealed by TLC analyses. However, in dry nonprotic solvents, e.g., chloroform, acetone, ether, **1** reacted cleanly with iodine giving, by TLC, only two compounds with the stoichiometry of Eq. (2):

$$2(PhS)_3As + 3I_2 \rightarrow 2AsI_3 + 3PhSSPh$$
(2)

The best solvent for the isolation of AsI₃ was ether, in which the triiodide is sparingly soluble (1:500).⁸ The yields, 85–90%, compare favorably with those reported for its preparation in aqueous media^{8a,9} using As₂O₃/HCl/KI. Because the ester **1** is easily prepared and is stable to storage,¹⁰ the reaction (2) can be useful for preparations of milligram amounts of arsenic triiodide. The ester **2** gave AsI₃ (75–80%), while



from the reaction of **3** with iodine, pure bis(4-nitrophenyl) disulfide, was isolated.

The most likely mechanism for the reaction (2) is shown in Scheme 1. It is not easy to say which of the two intermediates, the tetracoordinated $(ArS)_3AsI^+I^-$ or the pentacoordinated $(ArS)_3As(I_2)$, is preferred.



SCHEME 1

The mechanism suggests that arylsulfenyl iodide ArSI is an intermediate, and we sought a preparation of phenylsulfenyl iodide to test this hypothesis. However, arylsulfenyl iodides are only accessible in special instances,¹¹ and the use of phenylsulfenyl iodide in the literature¹² is questionable. We therefore prepared phenylsulfenyl bromide,¹³ contaminated by diphenyl disulfide, by reacting diphenyl disulfide with bromine,¹⁴ and reacted it with **1** with a 3:1 stoichiometry. The reaction was very fast, and **1** was not detected by TLC. Therefore, there are strong indications that phenylsulfenyl iodide is an intermediate in the reaction (2), and its reactivity explains the complex mixture of products obtained when the reaction (2) was done in methanol.

The oxidation of the trithioarsenite 1 with bromine gave diphenyl disulfide and AsBr₃, which could not be separated. Overall, iodine and bromine can therefore oxidize trithioarsenites to AsX₃ plus disulfides, and the use of the solid iodine is more convenient for preparative purposes.

Iodine should oxidize esters **4–7** to AsI_3 and disulfide. The gum produced from **6** could not be crystallized, while the powdery solids isolated from the reaction of **4**, **5**, and **7** with iodine were the Lewis adducts **8–10**, in which each AsI_3 acts as a Lewis acid formally coordinated to three donor atoms. Because AsI_3 is not coordinated to the -S-S- of diphenyl disulfide, it follows that it must be coordinated to N and/or O atoms in **8–10**, and it is not extracted by carbon disulfide in which it is soluble (1:18).⁸

RIGHTSLINK



FORMULAS 8-10

The first adduct of AsI₃, acting as a Lewis acid, AsI₃·3PhNH₂, was prepared by H. Schiff in 1863 and other earlier examples^{8a} are AsI₃·xpy (x = 2, 3), AsI₃·xNH₃ (x = 3, 4, 12), and AsI₃·Me₂(Ph)As. More recent work on the Lewis acid behavior of AsI₃ includes the reaction with Lewis bases having N,^{15–17} P,^{16,18} and As^{16,18} as well as O,^{17,19,20} S,²¹ and Se²¹ donor atoms. In some instances,^{18,19,21} the determination of the structure of the complexes by X-ray crystallography was possible.

Bis(4-acetamidophenyl) disulfide in the solid state must have intermolecular hydrogen bonding, forming transoid-transoid "polymer"²² because strong (at 3294 cm⁻¹) and weaker (at 3171, 3108 cm⁻¹) bands for N-H stretching and very strong amide I (at 1662 cm⁻¹) and amide II (at 1540 cm⁻¹) bands are seen in its solid phase IR spectrum. The IR spectrum of the black **8** is quite similar to that of the disulfide except that the bands are at 3304, 1654, and 1534 cm⁻¹, respectively. These data indicate that the "polymeric" structure of the bis(4-acetamidophenyl) disulfide is retained in **8**, having a C=O···H-N hydrogen-bonding network, which is slightly perturbed by AsI₃. From the composition of **8** and its IR spectrum, the AsI₃ molecule seems to be bonded to the lone pairs of three nitrogen atoms. Then, each arsenic will have a N₃I₃ environment. However, the brown 1:1 complexes of AsI₃ with acetamide and with *N*, *N*-dimethylacetamide have, by IR, arsenic coordinated to carbonyl oxygen.²⁰

The solid state IR spectrum of 9 is quite simple, does not resemble that of bis(4-aminophenyl) disulfide,²³ and does not contain decomposition products that were found with AsI₃·3PhNH₂, which in ether gave PhNH₂·HI and I-As(NHPh)₂.¹⁵ While from the mass balance and percent analysis six acetone molecules must be present, in the IR spectrum we find a very weak carbonyl peak at 1704 cm⁻¹. This fact, coupled with the absence of $-NH_2$ vibrations at 3418, 3335, and 3212 cm^{-1} of the disulfide²³ and the presence of two very strong but not too broad bands at 3444 and 2876 and a much weaker peak at 2575 cm⁻¹, indicates that the amino group of the disulfide has been added to the carbonyl group of acetone to give a geminal aminoalcohol (Nsubstituted hemiaminal), which did not dehydrate to a Schiff base most likely because the electron pair on nitrogen is not free to initiate the dehydration. Therefore, the AsI_3 must be coordinated to the nitrogen and, from the composition, each arsenic must have a N₃I₃ environment. The band, then, at 3444 cm⁻¹, must be due to intermolecularly hydrogen bonded –OH groups, and the band at 2876 cm⁻¹ must be due to a stretching vibration of an elongated and, thus, weaker N-H bond. The elongation is caused by the "donation" of the nitrogen electron pair to the arsenic.

The solid state IR spectrum of the black 10 differs from that of 2,2-dithiobis(pyridine N-oxide).²³ Of the C=C vibrations at 1463 and 1422 cm^{-1} of the free ligand, the one at 1422 cm^{-1} was less intense in 10. The N–O stretching vibrations at 1249 and 1230 cm^{-1} of the free ligand were less intense in 10, and a new vibration at 1182 cm^{-1} appeared that is attributed to N-OAs stretching. The N-O bending vibration moved from 838 to 832 cm^{-1} in **10**. The coordination of related ligands showed that, e.g., when pyridine N-oxide binds to platinum(II), the N–O stretching is shifted from 1242 to 1235 cm⁻¹.²⁴ When 2-mercaptopyridine N-oxide, which exists mainly as the tautomeric 1hydroxypyrid-2-thione with $\beta_{\rm CH}$ at 1263 cm⁻¹ and N–O stretching at 1112 and 1081 cm⁻¹, 25 is coordinated in a bidentate mode to tetraphenylantimony(V), the N–O stretching appears at 1197 cm⁻¹.²⁶ The stretching vibrations of the N-O of the free ligand, 2,2'-bipyridine N, N'dioxide (BipyO₂), at 1262 and 1255 appeared at 1265 and 1210 cm⁻¹ in the reddish-brown complex AsI₃·2BipyO₂, while the N–O bending vibrations of the free ligand at 852 and 842 cm⁻¹ appeared at 842 cm⁻¹ in the complex.¹⁷ From the composition of **10** and its IR spectrum, a bidentate coordination of the ligand to give, for example, a 9-membered ring and an $O_3I_3As^{3-}$ core with a high charge on arsenic is not probable. We propose that 2,2'-dithiobis(pyridine N-oxide) most probably acts, through its -O⁻, as a monodentate ligand, thus lowering the charge on the arsenic atom compared to the bidentate coordination, and, moreover, the positive charge on nitrogen, N⁺-O-As⁻, can be dissipated through the -S-S- bridge to the other pyridine N-oxide ring via their canonical structures.²⁷ Whether 10 is a single compound is an open question in the absence of crystallographic data.

Arsenic triodide is quite stable in water; in fact, it can be recovered unchanged from the solution⁹ and has been used in medicine for the treatment of tuberculosis in pigs, of anaemia from hereditary syphilis, of arthritis and certain dermatitises, but it is toxic.^{8a} Given the effectiveness of arsenic(III) oxide (As₂O₃, arsenox^R) in treating acute promyelocytic leukemia in man,²⁸ arsenic triiodide and its complexes, like **8–10**, should be examined in place of arsenic(III) oxide. Based on the behavior of (ArS)₃As and I₂, the reaction of iodine with thioesters of arsonous and arsinous acids, L-As(SPh)₂ and L₂As-SPh, should give L-AsI₂ and L₂As-I (L = R or Ar) and PhSSPh. From these iodides, a wide range of organoarsenic compounds can be prepared. Such studies are underway.

EXPERIMENTAL

The aromatic trithioarsenites have been prepared by literature methods as follows: $1,^{10}$ **3**, **5**, and $6,^2$ **2**, **4**, and $7.^5$ Silica gel 60 H (Merck) was used for Thin Layer Chromatography (TLC). AR grade solvents have been dried over A₄ molecular sieves and de-aerated by boiling, flushing with nitrogen, stoppering, and cooling to r.t. TLC was run on microslides using the appropriate disulfides^{2,5,10} as standards. The visualization was effected by iodine vapors followed by spraying with 35% sulfuric acid and charring. Compounds containing an ArS– group initially gave colored spots and on charring left a faint black spot. For melting points, an Electrothermal (model 9100) apparatus was used, and IR spectra were obtained on a Perkin-Elmer (model 16PC FT-IR) spectrometer. Elemental analyses were obtained through the Centre of Instrumental Analyses, University of Patras, Patras, Greece.

The Reaction of the Trithioarsenites 1–7 With lodine

Triphenyl Trithioarsenite, 1

To a solution of 1 (201 mg, 0.5 mmol) in ether (6 mL), iodine (191 mg, 0.75 mmol) was added and stirred at r.t. for 1 h. Centrifugation and washing with ether (2×2 mL) gave arsenic triiodide (204 mg, 90%) as an orange solid having m.p. 141° C (lit.^{8a} 138.6, 140.7, 146°C). IR (KBr): 3431 s, broad, 1636 w, broad, 595 m, broad. Evaporation of the dark red supernatant gave a light orange solid (169 mg), which was diphenyl disulfide (by TLC and IR) contaminated by arsenic triiodide.

Tris(4-Chlorophenyl) Trithioarsenite, 2

The reaction of **2** (101 mg, 0.2 mmol) with iodine (73 mg, 0.3 mmol) in ether (1 mL), as previously mentioned, gave arsenic triiodide (72 mg, 78%), m.p. $141-142^{\circ}$ C.

Tris(4-Nitrophenyl) Trithioarsenite, 3

To a suspension of **3** (0.25 mmol) in dichloromethane (3 mL), iodine (0.375 mmol) was added. At once, the ester **3** dissolved giving a dark brown solution from which an orange solid precipitated. Evaporation, drying in vacuum, addition of carbon disulfide (3 mL), centrifugation, and washing with carbon disulfide (1 × 2 mL) left the disulfide (99 mg, 86%), m.p. 183–184°C (lit.²⁹ 173–181°C; lit.³⁰ 191°C). Its IR (KBr) was the same as reported.²³ From the supernatant, arsenic triiodide (120 mg) contaminated by the disulfide (by TLC and IR) was obtained, m.p. 140°C.

Tris(4-Acetamidophenyl) Trithioarsenite, 4

The ester **4** (0.147 mmol) and iodine (0.221 mol) in acetone (2 mL) after 1 h of stirring and evaporation gave a cherry-red gum. The addition of carbon disulfide (3 mL) and boiling transformed the gum to a black solid. Centrifugation and washing with carbon disulfide (2 × 2 mL) left **8** as a black powder (121 mg, 88%). It is soluble in acetone (yellow solution) and in methanol (orange solution), sparingly soluble in chloroform, and insoluble in ether and carbon disulfide. M.p. 148–149°C. Calculated for $C_{48}H_{48}N_6O_6S_6I_6As_2$ (M_r 1908.58): C, 30.20; H, 2.53; N, 4.40; S, 10.08%. Found. C, 30.41; H, 2.48; N, 4.33; S, 10.55%. IR (KBr): 3303, s, sharp, 3112 m, 1654 vs, 1594 s, 1534 vs, 1490 ms, 1396 ms, 1320 m, 1117 w, 1078 w, 1045 w, 1006 m, 816 s, 672 mw, 604 mw.

Tris(4-Aminophenyl) Trithioarsenite, 5

The addition of iodine (0.375 mmol) to a solution of **5** (0.25 mmol) in acetone (3 mL) immediately precipitated a brown-orange gum. After 3 h of stirring at r.t., evaporation and drying in a vacuum gave a sticky, colored foam. It was covered with ether and stirred at r.t. overnight. Centrifugation and washing with ether $(1 \times 2 \text{ mL})$ gave **9** as a light yellow-brown solid (246 mg, 98%) insoluble in carbon disulfide and ether, sparingly soluble in chloroform (yellow), and soluble in methanol (yellow). M.p.: 73°C shrinks, 92°C turns red, 99–101°C melts to a red oil. Calculated for C₅₄H₇₂N₆O₆S₆I₆As₂ (M_r 2004.79): C, 32.35; H, 3.62; N, 4.19; S, 9.60%. Found: C, 31.97; H, 3.15; N, 4.69; S, 9.55%. IR (KBr): 3444 vs, slightly broad, 2876 vs, slightly broad, 2575 m, 1704

vw, 1618 s, 1565 m, 1488 vs, 1406 w, 1374 w, 1313 w, 1206 w, 1122 w, 1017 w, 808 s, 612 w, 495 w.

Tris(2-Pyridyl) Trithioarsenite, 6

The reaction of 6 (0.25 mmol) suspended in dichloromethane (3 mL) with iodine (0.375 mmol) gave a cherry-red oil that could not be induced to crystallize and was not studied further.

Tris(2-pyridyl N-Oxide) Trithioarsenite, 7

To a suspension of **7** (0.82 mmol) in acetone (5 mL), iodine (1.23 mol) was added. At once, the ester dissolved and a new solid formed. After 2 h of stirring, evaporation and drying in vacuum gave a black crystalline solid. The addition of carbon disulfide (3 mL), centrifugation, and washing with carbon disulfide (1 × 2 mL) gave **10** as a black-brown powder (677 mg, 99%). It is soluble in methanol (yellow) and acetone (orange), sparingly soluble in chloroform, and insoluble in ether. M.p. 105–108°C. Calculated for $C_{30}H_{24}N_6O_6S_6I_6As_2$ (M_r 1668.21): C, 21.69; H, 1.45; N, 5.04; S, 11.53%; Found: C, 21.48; H, 1.02; N, 5.36; S, 11.52%. IR (KBr): 3444 m, slightly broad, 3090 m, 1590 w, 1554 m, 1460 vs, 1418 s, 1248 m, 1230 sh, 1182 m, 1142 m, 1084 w, 1039 w, 828 m, 758 s, 704 m, 578 s.

The Reaction of Triphenyl Trithioarsenite, 1, With Bromine

The reaction of 1 (0.315 mmol) and bromine (0.473 mmol) in ether (6 mL) gave a yellow solution from which the arsenic tribromide and diphenyl disulfide could not be separated due to similar solubilities.^{8a} Hydrolysis of the mixture in acetone gave As_2O_3 (16 mg, 50% recovery) and impure diphenyl disulfide (114 mg), which was recrystallized from petroleum ether, m.p. 59–60°C (lit.³¹ 61–62°C).

The Reaction of 1 With Phenylsulfenyl Bromide

To a solution of bromine (103 mg, 0.625 mmol) in dichloromethane, diphenyl disulfide (136 mg, 0.625 mmol) was added, and the red solution was left at r.t. for 2 h. TLC (petroleum ether) showed the presence of disulfide ($R_{\rm f}$ 0.46) and a spot as $R_{\rm f}$ 0.0 (due to the hydrolysis of PhSBr on the plate). The evaporation left an oil (202 mg, expected 236 mg PhSBr). To the oil in dichloromethane (3 mL), solid 1 (0.35 mmol) was added. At once, the solution decolorized to a nearly colorless solution. TLC (petroleum ether) showed only the presence of disulfide and the absence of 1 ($R_{\rm f}$ 0.10).



REFERENCES

- [1] A. Terzis and P. V. Ioannou, Z. Anorg. Allg. Chem., 630, 278 (2004).
- [2] M. N. Haikou, G. M. Tsivgoulis, and P. V. Ioannou, Z. Anorg. Allg. Chem., 630, 1084 (2004).
- [3] G. M. Tsivgoulis, T. C. Photopoulou, and P. V. Ioannou, Phosphorus, Sulfur, and Silicon, 181, 413–425 (2006).
- [4] (a) R. Klement and R. Reuber, *Chem. Ber.*, **68**, 1761 (1935); (b) W. Herman, Methoden der Organischen Chemie, Part 2 (Houben Weyl, G. Thieme Verlag, Stuttgard, 1963) 4th ed. Vol. 6, pp. 411–416.
- [5] M. N. Haikou and P. V. Ioannou, Phosphorus, Sulfur, and Silicon, 181, 363–376 (2006).
- [6] M. A. Lala and P. V. Ioannou, J. Inorg. Biochem., 97, 331 (2003).
- [7] S. C. Chaudhry, R. K. Mahajan, S. S. Bhatt, and N. Sharma, *Indian J. Chem.*, 31A, 279 (1992).
- [8] R. Dolique, Nouveau Traité de Chimie Minérale, P. Pascal Ed. (Masson, Paris, 1958), Vol. XI (a) pp. 169–175, (b) 162–167.
- [9] J. C. Bailar, Jr., Inorg. Syntheses, 1, 103 (1939).
- [10] S. V. Serves, Y. C. Charalambidis, D. N. Sotiropoulos, and P. V. Ioannou, *Phosphorus*, *Sulfur, and Silicon*, **105**, 109 (1995).
- [11] E. Kühle, Synthesis, 561 (1970).
- [12] R. S. Dickson, G. D. Fallon, W. R. Jackson, and A. Polas, J. Organometal. Chem., 607, 156 (2000).
- [13] M. Verny and R. Vessière, Bull. Soc. Chim. France, 746 (1970).
- [14] J. Chatt, J. P. Lloyd, and R. L. Richards, J. Chem. Soc. Dalton Trans., 565 (1976).
- [15] D. Hass, Z. Anorg. Allg. Chem., 337, 43 (1965).
- [16] P. O. Dunstan, Thermochim. Acta, 279, 167 (1996).
- [17] P. O. Dunstan, Thermochim. Acta, 240, 91 (1994).
- [18] N. J. Hill, W. Levason, and G. Reid, J. Chem. Soc. Dalton Trans., 1188 (2002).
- [19] B. Borgsen, F. Weller, and K. Dehnicke, Chemiker-Zeitung, 114, 111 (1990).
- [20] P. O. Dunstan and L. C. R. Dos Santos, Thermochim. Acta, 156, 163 (1989).
- [21] N. J. Hill, W. Levason, and G. Reid, Inorg. Chem., 41, 2070 (2002).
- [22] S. F. Dyke, A. J. Floyd, M. Sainsbury, and R. S. Theobald, Organic Spectroscopy, An Introduction, (Penguin Books, Middlesex, 1971), Ch. 3, pp. 83–86.
- [23] Spectral Database of Organic Compounds, SDBS http://www. Aist. go. jp/RIODB/ SDBS
- [24] S. I. Shupack and M. Orchin, J. Am. Chem. Soc., 85, 902 (1963).
- [25] A. R. Katritzky and R. A. Jones, J. Chem. Soc., 2947 (1960).
- [26] G. M. Arvanitis, M. E. Berardini, T. B. Acton, and P. E. Dumas, *Phosphorus, Sulfur, and Silicon*, 82, 127 (1993).
- [27] R. M. Acheson, An Introduction to the Chemistry of Heterocyclic Compounds, (Wiley-Interscience, New York, 1967), 2nd ed., Ch. V, pp. 207–211.
- [28] W. H. Miller, Jr., H. M. Schipper, J. S. Lee, J. Singer, and S. Waxman, *Cancer Res.*, 62, 3893 (2002).
- [29] J. Nakabayashi, S. Kawamura, T. Horii, and M. Hamada, Chem. Letters, 869 (1976).
- [30] R. S. Atkinson and B. D. Judkins, J. Chem. Soc. Perkin Trans. I, 2615 (1981).
- [31] R. C. West, CRC Handbook of Chemistry and Physics (CRC Press, Baca Raton, FL, 1984), 65th ed.

RIGHTSLINK()

Downloaded by [University of Tennessee, Knoxville] at 22:29 13 April 2014