



Design, synthesis, and biological evaluation of novel derivatives of dithiodiglycolic acid prepared via oxidative coupling of thiols

Olga Bakulina, Anton Bannykh, Mirna Jovanović, Ilona Domračeva, Ana Podolski-Renić, Raivis Žalubovskis, Milica Pešić, Dmitry Dar'in & Mikhail Krasavin

To cite this article: Olga Bakulina, Anton Bannykh, Mirna Jovanović, Ilona Domračeva, Ana Podolski-Renić, Raivis Žalubovskis, Milica Pešić, Dmitry Dar'in & Mikhail Krasavin (2019) Design, synthesis, and biological evaluation of novel derivatives of dithiodiglycolic acid prepared via oxidative coupling of thiols, Journal of Enzyme Inhibition and Medicinal Chemistry, 34:1, 665-671, DOI: [10.1080/14756366.2019.1575372](https://doi.org/10.1080/14756366.2019.1575372)

To link to this article: <https://doi.org/10.1080/14756366.2019.1575372>



© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



View supplementary material [↗](#)



Published online: 12 Feb 2019.



Submit your article to this journal [↗](#)



Article views: 49



View Crossmark data [↗](#)

SHORT COMMUNICATION



Design, synthesis, and biological evaluation of novel derivatives of dithiodiglycolic acid prepared via oxidative coupling of thiols

Olga Bakulina^a, Anton Bannykh^a, Mirna Jovanović^b, Ilona Domračeva^c, Ana Podolski-Renić^b, Raivis Žalubovskis^{c,d}, Milica Pešić^b, Dmitry Dar'in^a and Mikhail Krasavin^a

^aSaint Petersburg State University, Saint Petersburg, Russian Federation; ^bInstitute for Biological Research “Siniša Stanković”, University of Belgrade, Belgrade, Serbia; ^cLatvian Institute of Organic Synthesis, Riga, Latvia; ^dFaculty of Materials Science and Applied Chemistry, Institute of Technology of Organic Chemistry, Riga Technical University, Riga, Latvia

ABSTRACT


Human thioredoxin reductase 1 (TrxR1) is a selenocysteine-containing enzyme which plays a crucial role in regulating numerous redox signalling pathways within the cell. While its functioning is important in all cells, levels of TrxR1 expression are higher in cancer cells, possibly as an adaptation to much higher levels of reactive oxygen species and the need for more extensive DNA synthesis. This makes TrxR1 an attractive target for cancer therapy development. Inspired by the structure of disulphide compounds which have advanced through various stages of clinical development, we designed a series of dithiodiglycolic acid derivatives. These were prepared from respective thiol synthons using an iodine- or benzotriazolyl chloride-promoted oxidative disulphide bond formation. Inhibition of TrxR present in cell lysates from human neuroblastoma cells (SH-SY5Y) and rat liver cells indicated several compounds with a potential for TrxR inhibition. Some of these compounds were also tested for growth inhibition against two human cancer cell lines and normal human keratinocytes.

ARTICLE HISTORY

Received 20 December 2018
Revised 21 January 2019
Accepted 22 January 2019

KEYWORDS

TrxR; disulphide inhibitors; dithiodiglycolic acid; anticancer activity



Introduction

Thioredoxin reductase enzymes (TrxR, EC 1.8.1.9), which are the focus of this work, belong to thioredoxin system along with NADPH and thioredoxin (Trx). The system is highly conserved among species and responsible for regulating redox processes, gene transcription, and protection against reactive oxygen species (ROS)¹. TrxR maintains Trx in its reduced bis-sulfhydryl state that, in turn, interacts with antioxidant enzymes and transcription factors². TrxR1 is a cytosolic isoform of the enzyme (along with mitochondrial TrxR2 and TrxR3 expressed primarily in the testes). TrxR enzymes are overexpressed in cancer cells contributing to their resistant phenotype characterised by higher levels of ROS³. Targeting TrxR specifically with various, mostly electrophilic, inhibitors have thus become an attractive new approach for the development of anticancer therapy⁴.

The various types of inhibitors targeting the catalytic selenocysteine residue of TrxR with varying degree of

electrophilicity – Michael acceptors, disulphide and diselenide substrate mimics, metal-based inhibitors as well as compounds of miscellaneous structure – have been recently reviewed in comprehensive accounts by Bellelli⁵ and Fang^{6,7}. Particularly relevant to the present study are the disulphide substrate analogues: e.g. alkyl 2-imidazolyl disulphides **1–3**⁸ (**2**, whose K_i value toward the enzyme was determined as 31 μM , is also known as PX-12 and has been in Phase 1 clinical trials for advanced metastatic cancer and was stopped due to safety concerns⁶), disodium 2,2'-dithio-bis-ethane sulphonate **4**⁹ (also known as TravoceptTM which advanced through phase III clinical trials in patients with non-small cell lung cancer and has a K_m value of 72 μM ⁶) as well as bicyclic disulphide-containing natural products chaetocin (**5**), gliotoxin (**6**) and chaetomin (**7**) which have been shown to possess K_m values of 4.6 μM , 16.9 μM , and 16.1 μM , respectively – and have demonstrated a substrate-competitive inhibition profile toward TrxR1 and pro-apoptotic efficacy toward cancer cells¹⁰. Other disulphide

CONTACT Mikhail Krasavin  mkrasavin@hotmail.com  Saint Petersburg State University, Saint Petersburg 199034, Russian Federation

 Supplemental data for this article can be accessed [here](#).

© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Known disulphide TrxR inhibitors (1–7) and fluorescent probes (8–10) as well as compounds explored in this work (11–13).

compounds – 5,5'-dithiobis(2-nitrobenzoic acid) (**8** also known as DTNB or Ellman's reagent)¹¹, thioredoxin reductase fluorogenic substrate (**9** or TRFS-green)¹² and mitochondria-targeted "mito-TRFS" (**10**)¹³ have not been employed as TrxR inhibitors but rather as fluorescent probes, also due to their being substrate (disulphide-linked Trx) analogues. Based on the structure of these disulphide TrxR ligands, we designed a series of symmetric (**11**) and non-symmetric (**12–13**) derivatives of dithiodiglycolic acid (Figure 1). Herein, we report the results of their synthetic exploration and biological testing for TrxR inhibition and cytotoxicity toward two human cancer cell lines and normal human keratinocytes.

Materials and methods

Chemical syntheses – general

NMR spectroscopic data were recorded with Bruker Avance 400 spectrometer (400.13 MHz for ¹H and 100.61 MHz for ¹³C) and Bruker Avance 500 spectrometer (500.03 for ¹H and 125 MHz for ¹³C) in DMSO-d₆ and in CDCl₃ and were referenced to residual solvent proton signals (δ_{H} = 2.50 and 7.26 ppm, respectively) and solvent carbon signals (δ_{C} = 39.5 and 77.0 ppm, respectively). Mass spectra were recorded with a Bruker Maxis HRMS-ESI-qTOF spectrometer (electrospray ionisation mode). Merck silica gel 60 mesh was used for column chromatography. TLC was performed with Macherey-Nagel Alugram Sil G/UV254 plates. Thioacetic acid, trityl chloride, methyl thioglycolate, amines, and 1H-benzotriazole, 2,2'-disulfanediyldiacetic acid (**14**) and 2,2'-disulfanediyldiacetic acid disodium salt (**15**) were obtained from commercial sources. 2-(Tritylthio)acetic acid (**16**) was prepared from thioacetic acid and

trityl chloride according to literature procedure¹⁴. Dimethyl 2,2'-disulfanediyldiacetate (**17**) was prepared by oxidative dimerisation of methyl thioglycolate¹⁵. 1-Chlorobenzotriazole was prepared according to known procedure¹⁶. DCM was distilled from P₂O₅ and stored over MS 4 Å. THF was distilled from sodium benzophenone under Ar. All synthesised thiols and disulphides are air-sensitive, and therefore were stored in sealed vials at –20 °C. Characterisation data are provided only for one representative compound in each group. The rest of the compounds are described in the [Supplementary Data](#).

General procedure 1: preparation of 2-(tritylthio)acetamides **18a–l**

To a solution of 2-(tritylthio)acetic acid **16** (1.5–4.5 mmol) in dry dichloromethane (DCM; 5 ml/mmol) 1,1'-carbonyldiimidazole (CDI; 1.0–1.1 equiv.) was added portionwise (gas evolution!). After stirring for 50 min, amine (1.0–1.1 equiv.) was added and the reaction mixture was left stirring at r.t. overnight. The reaction mixture was washed with citric acid (2 N aq. solution), saturated aq. NaHCO₃ and water. Organic layer was dried (over Na₂SO₄), filtered, and concentrated *in vacuo* to give pure title amides.

1-(Pyrrolidin-1-yl)-2-(tritylthio)ethan-1-one (18a**)** was prepared according to General Procedure 1 from 2-(tritylthio)acetic acid (668 mg, 2 mmol), pyrrolidine (150 mg, 2.1 mmol) and CDI (340 mg, 2.1 mmol). Yield 584 mg, 75%. White solid. ¹H NMR (400 MHz, CDCl₃) δ 7.57–7.43 (m, 6H), 7.37–7.28 (m, 6H), 7.28–7.17 (m, 3H), 3.40 (t, *J* = 6.6 Hz, 2H), 3.07 (t, *J* = 6.4 Hz, 2H), 2.93 (s, 2H), 1.85–1.77 (m, 4H). ¹³C NMR (101 MHz, CDCl₃) δ 166.8, 144.3, 129.6,

128.0, 126.8, 66.8, 46.5, 46.0, 36.1, 26.0, 24.3. HRMS (ESI), m/z calcd for $C_{25}H_{25}NOSNa$ $[M + Na]^+$ 410.1549, found 410.1553.

General procedure 2: preparation of thiols 19a–c

To a stirred solution of protected amide **18a–c** (2 mmol) in TFA (5 ml) triethylsilane (TES; 1.1 equiv.) was added at room temperature. Reaction progress was monitored by TLC and reaction mixture turned colourless from deep yellow. After 5–30 min, reaction mixture was washed, diluted with water (15 ml), and extracted with hexane (3×15 ml). The aqueous layer was then extracted with DCM (3×15 ml). The combined organic layer was dried over Na_2SO_4 , filtered and concentrated to provide pure title compounds¹⁷.

2-Mercapto-1-(pyrrolidin-1-yl)ethan-1-one (19a) was prepared according to General Procedure 2 from **18a** (774 mg, 2 mmol) and TES (254 mg, 2.2 mmol). Yield 178 mg, 61%. 1H NMR (400 MHz, $CDCl_3$) δ 3.57 (q, $J = 7.3$ Hz, 4H), 3.32 (s, 2H), 2.23 (s, 1H), 2.07 (p, $J = 6.8$ Hz, 2H), 1.96 (p, $J = 6.8$ Hz, 2H). ^{13}C NMR (101 MHz, $CDCl_3$) δ 170.7, 47.4, 47.0, 26.3, 25.9, 24.3. HRMS (ESI), m/z calcd for $C_6H_{12}NOS$ $[M + H]^+$ 146.0634, found 146.0634.

General procedure 3: preparation of thiols 19d–k

To a stirred solution of protected amide **18d–k** (1.2–2.1 mmol) in DCM (10 ml) trifluoroacetic acid (TFA; 5 equiv.) was added followed by triethylsilane (TES; 2 equiv.). Reaction progress was monitored by TLC. After 2–16 h, reaction mixture was washed with water (3×30 ml), dried and concentrated *in vacuo*. The residue was suspended in hexane (10 ml) and sonicated using ultrasonic bath, cooled and filtered to give corresponding thiol¹⁸.

N-Cyclopropyl-2-mercaptoacetamide (19d) was prepared according to General Procedure 3 from **18d** (746 mg, 2 mmol), TFA (1.27 g, 5.9 mmol) and TES (514 mg, 4.4 mmol). Yield 233 mg, 89% (85% NMR purity). 1H NMR (400 MHz, $CDCl_3$) δ 6.72 (s, 1H), 3.23 (d, $J = 9.1$ Hz, 2H), 2.76 (dq, $J = 7.1, 3.5$ Hz, 1H), 1.87 (t, $J = 9.1$ Hz, 1H), 0.88–0.79 (m, 2H), 0.60–0.52 (m, 2H). HRMS (ESI), m/z calcd for C_5H_9NOSNa $[M + Na]^+$ 154.0297, found 154.0299.

General procedure 4: preparation of symmetric disulphides 11a–k

To the solution of thiol **19a–k** (0.3–0.9 mmol) in DCM and triethylamine (TEA; 1.5–3 equiv.) solid I_2 was added portionwise until persistent deep yellow colour appeared. After stirring for 5 min, the conversion of thiol was checked by TLC. In case of compounds **11a–d, f, g, k** the reaction mixture was concentrated and the residue was crystallised from hexane-ethyl acetate to give pure disulphides. Compounds **11e, h, i, j** precipitated from the reaction mixture and were isolated by filtration and washing with DCM (2×5 ml)¹⁹.

2,2'-Disulfanediybis(1-(pyrrolidin-1-yl)ethan-1-one) (11a) was prepared according to General Procedure 4 from **19a** (90 mg, 0.31 mmol) and TEA (66 mg, 0.93 mmol). Yield 66 mg, 75%. 1H NMR (400 MHz, $CDCl_3$) δ 3.69 (s, 4H), 3.57 (t, $J = 6.8$ Hz, 4H), 3.51 (t, $J = 6.9$ Hz, 4H), 2.04–1.96 (m, 4H), 1.95–1.84 (m, 4H). ^{13}C NMR (126 MHz, $CDCl_3$) δ 166.7, 47.2, 46.2, 42.3, 26.2, 24.4. HRMS (ESI), m/z calcd for $C_{12}H_{20}N_2O_2S_2Na$ $[M + Na]^+$ 311.0858, found 311.0873.

General procedure 5: preparation of non-symmetric disulphides 12(13)a

A mixture of 1-chlorobenzotriazole (BtCl; 1.1 equiv.) and 1H-benzotriazole (BtH; 1 equiv.) was placed into a Schlenk tube, which was then evacuated and filled with argon three times followed by addition of dry DCM (10 ml) and cooling to $-78^\circ C$. First thiol (0.5–0.6 mmol) dissolved in 5 ml DCM was added dropwise during 15–20 min to form a yellow solution. Then the reaction mixture was slowly warmed to $-20^\circ C$ and the second thiol (1.1 equiv.) dissolved in 5 ml DCM was added dropwise during 15–20 min. The resulting solution was left stirring at $0^\circ C$ for 30 min. The reaction mixture was then quenched with aqueous $Na_2S_2O_3$, washed with saturated aq. $NaHCO_3$ (20 ml) and extracted with DCM (3×50 ml). Combined organic layer was dried over Na_2SO_4 , filtered and concentrated to give crude material, which was purified by silica gel column chromatography using DCM/MeOH mixture (100:1) as eluent²⁰.

2-((2-Morpholino-2-oxoethyl)disulfanyl)-N-propylacetamide (12a) was prepared according to General Procedure 5 from **19d** (84 mg, 0.63 mmol) as the first thiol, **19b** (102 mg, 0.63 mmol) as the second thiol, BtCl (106 mg, 0.69 mmol) and BtH (75 mg, 0.63 mmol). Yield 70 mg, 38%. 1H NMR (400 MHz, $CDCl_3$) δ 3.73 (dt, $J = 6.4, 3.8$ Hz, 4H), 3.69 (d, $J = 6.2$ Hz, 2H), 3.67 (s, 2H), 3.53 (t, $J = 4.9$ Hz, 2H), 3.50 (s, 2H), 3.35–3.26 (m, 2H), 1.62 (h, $J = 7.4$ Hz, 2H), 0.98 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (126 MHz, $CDCl_3$) δ 167.8, 167.0, 66.8, 66.6, 46.8, 42.9, 42.5, 41.8, 39.9, 22.8, 11.5. HRMS (ESI), m/z calcd for $C_{11}H_{20}N_2O_3S_2Na$ $[M + Na]^+$ 315.0808, found 315.0815.

Methyl 2-((2-((4-fluorophenyl)amino)-2-oxoethyl)disulfanyl)acetate (13a) was prepared according to General Procedure 5 from **19i** (110 mg, 0.59 mmol) as the first thiol, methyl thioglycolate (69 mg, 0.65 mmol) as the second thiol, BtCl (100 mg, 0.65 mmol) and BtH (71 mg, 0.6 mmol). Yield 68 mg, 40%. 1H NMR (400 MHz, $CDCl_3$) δ 8.71 (s, 1H), 7.67–7.43 (m, 2H), 7.05 (m, 2H), 3.84 (s, 3H), 3.65 (s, 2H), 3.60 (s, 2H). ^{13}C NMR (101 MHz, $CDCl_3$) δ 171.31, 166.03, 159.51 (d, $J_{CF} = 243.7$ Hz), 133.79 (d, $J_{CF} = 2.9$ Hz), 121.69 (d, $J_{CF} = 7.8$ Hz), 115.63 (d, $J_{CF} = 22.5$ Hz), 53.17, 42.85, 41.87. HRMS (ESI), m/z calcd for $C_{11}H_{12}FNO_3S_2Na$ $[M + Na]^+$ 312.0135, found 312.0139.

Cell culture

Human neuroblastoma cells (SH-SY5Y) were grown in 1:1 mixture of Eagle's Minimum Essential Medium and F12 Medium (Sigma) supplemented with 10% foetal bovine serum (Sigma). Human keratinocytes (HaCaT) were grown in DMEM, supplemented with 10% foetal bovine serum, 2 mM L-glutamine, 5000 U/ml penicillin, and 5 mg/ml streptomycin. Human glioblastoma cell (U87-MG) were grown in Minimum Essential Media (MEM), containing 10% foetal bovine serum, 2 mM L-glutamine, 5,000 U/ml penicillin, and 5 mg/ml streptomycin. All cell lines were obtained from American Type Culture Collection (ATCC, Rockville, MD). All cell lines were grown at $37^\circ C$ in a humidified 5% CO_2 atmosphere.

TrxR inhibition assay

For TrxR assay, lysate from SH-SY5Y cells and rat liver were used. SHSY5Y cells were washed twice with phosphate buffer solution (PBS) and harvested from the flask with a scraper. Cells were homogenised in ice-cold 50 mM potassium phosphate buffer (pH 7.4), containing 1 mM EDTA, with Ultra Turrax T25 (Ika-Verkerk, Janke and Kunkel Inc., Staufen, Germany). The homogenates were

centrifuged at $10,000 \times g$ for 15 min at 4°C . The protein concentrations of the supernatants were determined using a Bio-Rad protein assay kit (Bio-Rad Laboratories, USA). TrxR activity was assayed using 5,5-dithio-bis-(2-nitrobenzoic acid) (DTNB, Sigma-Aldrich, USA) as substrate²¹. The reactions were run on a 96-well plates, final volume of $100\ \mu\text{l}$ in 50 mM potassium phosphate buffer (pH 7.0), containing $50\ \mu\text{g}$ of the cell lysate proteins, 1 mM EDTA, 50 mM KCl, 0.2 mg/ml bovine serum albumin, 0.25 mM NADPH (Acros Organics, USA). Reaction mixture was incubated for 15 min at room temperature in plate shaker (PST-60HL-4, BioSan, Riga, Latvia); afterwards DTNB was added to the total concentration of 2.5 mM. Enzyme kinetics was monitored on a Tecan Infinite1000 microplate reader, by measuring the increase in absorbance at 412 nm for 20 min. The background TrxR-independent reduction of DTNB in the cell lysates, as determined in the presence of aurothiomalate ($200\ \mu\text{M}$), was subtracted from each value²². Reaction mixture with rat liver lysate contained $35\ \mu\text{g}$ of proteins, 1 mM EDTA, 50 mM KCl, 0.2 mg/ml bovine serum albumin, 0.25 mM NADPH, and 2.5 mM DTNB. Reactions with rat liver lysate were run in the same manner as cell lysates.

Cytotoxicity assay

Cells grown in 25 cm^2 tissue flasks were trypsinized, seeded into flat-bottomed adherent 96-well cell culture plates in appropriate medium (SH-SY5Y 12,000 cells/well, U87-MG 4000 cells/well, HaCaT 2000 cells/well) and incubated overnight; afterwards the cells were exposed to different concentrations of the tested compounds for 72 h. Cell viability was measured by MTT assay²³. Briefly, after incubation with the compounds, medium containing 0.2 mg/ml MTT was added. After incubation (3 h, 37°C , 5% CO_2), the MTT-containing medium was removed, and $200\ \mu\text{l}$ of dimethyl sulfoxide (DMSO) was immediately added to each sample. The samples were assessed at 540 nm on LKB 5060-006 Microplate Reader (Vienna, Austria) microplate reader. The half-maximal inhibitory concentration (IC_{50}) of each compound was calculated using Graph Pad Prism® 6.0 (GraphPad Software, Inc., USA).

Results and discussion

Chemistry

Initial attempts to elaborate commercially available dithiodiglycolic acid (**14**) into compounds **11–13** failed, since either HATU- or DCC-mediated amidation of it resulted in a difficult-to separate mixture of mono- and symmetric bis-amides of **14** (also containing **14** itself). Therefore, for the preparation of symmetric bis-amides **11a–k** we adopted the route depicted in Scheme 1.

Trityl-protected thioacetic acid (**16**)¹⁴ was activated by 1,1'-carbonyldiimidazole (CDI) to give amides **18a–k** in good to excellent yields. The trityl group was removed by the action of trifluoroacetic acid (TFA) used as a solvent (**18a–c**)¹⁸ or as a reagent (5 equiv.) in DCM (**18d–k**)¹⁹, in the presence of triethylsilane. The reaction was completed in 5–30 min in the former case and took 2–16 h – in the latter. The free thiols **19a–k** thus obtained (in moderate to excellent yield) were homo-coupled on the action of molecular iodine in the presence of triethylamine¹⁹ to give target compounds **11a–k** in moderate to excellent yield.


For the preparation of non-symmetric disulphides **12(13)a**, one of the thiols **19** was activated at -78°C by the action of a roughly equimolar mixture of *N*-chlorobenzotriazole (BtCl) and benzotriazole (BtH), and the resulting intermediate was treated with a second thiol **19** (**12a**) or methyl mercaptoacetate (**13a**) to give, on stirring at 0°C for 30 min, respective disulphides (Scheme 2).

Biological evaluation of compounds **11a–k**, **12a**, and **13a** as well as commercially available dithiodiglycolic acid (**14**), its disodium salt (**15**) and dimethyl ester (**17**) against human neuroblastoma (SHSY5Y) as well as rat liver cell lysates using DTNB assay²¹ revealed that a number of compounds (**11a–d**, **11g**, **11j–k**, **13a**, **14**, and **17**) possessed double-digit micromolar activity against liver cell lysate while their inhibitory activity measured against neuroblastoma (SHSY5Y) cell lysate was significantly lower (compounds **11a–b**, **11g**, **11j**, **14**, and **17** displaying virtually no activity). Considering that TrxR1 expression levels in hepatocytes are significantly higher compared to SHSY5Y cells^{24,25}, these results may suggest higher activity of our novel compounds towards lysates with higher TrxR1 content. Notably, compound **15** which is a close analogue (double homolog) of compound **4** (Travocept™ which entered Phase III clinical trials in 2015⁶) is completely devoid of inhibitory activity toward TrxR.


Selected compounds from the active cohort (**11b**, **11d**, **11g**) and also compound **12a** (which displayed identical inhibitory potency with IC_{50} around $45\ \mu\text{M}$ against SHSY5Y and hepatocyte lysates) were tested for cytotoxicity against neuroblastoma (SHSY5Y), glioblastoma (U87), and immortal keratinocyte (HaCaT) cells. Interestingly, compound **11b** was found to be 2–3 times more active against U87 cell line compared to the other three compounds (Table 1).

Conclusions

We have described design, synthesis, and biological evaluation of a series of dithiodiglycolic acid derivatives. Symmetric bis-amides were prepared via molecular iodine-promoted bis-homocoupling of thioglycolic acid amides, while non-symmetric diamide and



Scheme 1. Preparation of symmetric disulphides **11a–k**.



Scheme 2. Preparation of non-symmetric disulphides **12(13)a**.

Table 1. Inhibition profile of compounds prepared in this work against TrxR1 measured in neuroblastoma (SHSY5Y) and hepatocyte lysates; cytotoxicity profile of selected compounds determined against SHSY5Y, U87, and HaCaT cells.

Compound	Structure	TrxR IC ₅₀ (μM) ^a		CC ₅₀ (μM) ^a		
		SHSY5Y lysate	Hepatocyte lysate	SHSY5Y	U87	HaCaT
11a		>200	170.4 ± 54.2	nd	nd	nd
11b		>200	26.5 ± 2.9	219.7 ± 15.3	54.6 ± 3.4	58.4 ± 5.0
11c		189.5 ± 78.4	182.5 ± 32.2	nd	nd	nd
11d		63.6 ± 27.5	34.7 ± 9.1	202.3 ± 12.0	179.6 ± 12.6	69.0 ± 3.4
11e		>200	>200	nd	nd	nd
11f		>200	>200	nd	nd	nd
11g		>200	22.2 ± 3.1	174 ± 18.3	125.2 ± 15.6	43.0 ± 2.9
11h		>200	>200	nd	nd	nd

(continued)

Table 1. Continued.

Compound	Structure	TrxR IC ₅₀ (μM) ^a		CC ₅₀ (μM) ^a		
		SHSY5Y lysate	Hepatocyte lysate	SHSY5Y	U87	HaCaT
11i		>200	>200	nd	nd	nd
11j		>200	90.4 ± 18.7	nd	nd	nd
11k		>200	24.0 ± 11.5	nd	nd	nd
12a		44.4 ± 15.1	45.6 ± 6.8	176.3 ± 13.4	186.8 ± 15.9	71.8 ± 6.9
13a		186.4 ± 83.8	>200	nd	nd	nd
14		>200	>200	nd	nd	nd
15		>200	>200	nd	nd	nd
17		>200	>200	nd	nd	nd

^aThe values provided are mean ± SEM ($n = 3$).

"nd": not determined; IC: inhibitory concentration; CC: cytotoxicity concentration.

amide-ester were prepared via bis-thiol heterocoupling promoted by *N*-chorobenzotriazole/benzotriazole mixture. The compound set tested against human neuroblastoma (SHSY5Y) and rat liver cell lysates for TrxR inhibition using DTNB assay revealed a promising inhibitory profile for 6 out of 16 compounds tested, namely, higher inhibitory potency toward hepatocyte lysate TrxR1 compared to SHSY5Y lysate. Several of active compounds also showed cytotoxic effect toward human neuroblastoma (SHSY5Y) and glioblastoma (U87), as well as, immortal human keratinocyte (HaCaT) cell lines. Thus, the derivatives studied in this work represent a novel lead chemotype that could be further developed into more potent and more specific TrxR1 inhibitors valuable for anticancer therapy.

Acknowledgements

We thank the Educational Resource Centre of Chemistry, Research Centre for Magnetic Resonance and the Centre for Chemical Analysis and Materials Research of Saint Petersburg State University Research Park for obtaining the analytical data.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This work was supported by the Russian Foundation for Basic Research [project grant 18-515-76001], Ministry of Education, Science and Technological Development of the Republic Serbia

[project III41031] as well as by ERA.Net RUS plus joint programme grant RUS_ST2017-309 and State Education Development Agency of Republic of Latvia ("THIOREDIN"). This work was also supported by a Short-Term Scientific Mission (STSM) Grant from COST Action CM1407 "Challenging organic syntheses inspired by nature - from natural products chemistry to drug discovery and European Cooperation in Science and Technology.

References

- Arner ES, Holmgren A. Physiological functions of thioredoxin and thioredoxin reductase. *Eur J Biochem* 2000;267:6102–9.
- Zhong L, Arner ES, Ljung J, et al. Rat and calf thioredoxin reductase are homologous to glutathione reductase with a carboxyl-terminal elongation containing a conserved catalytically active penultimate selenocysteine residue. *J Biol Chem* 1998;273:8581–91.
- Zhang J, Li X, Han X, et al. Targeting the thioredoxin system for cancer therapy. *Trends Pharmacol Sci* 2017;38:794–808.
- Arner ES, Holmgren A. The thioredoxin system in cancer. *Semin Cancer Biol* 2006;16:420–6.
- Saccoccia F, Angelucci F, Boumris G, et al. Thioredoxin reductase and its inhibitors. *Curr Protein Pept Sci* 2014;15:621–46.
- Zhang B, Zhang J, Peng S, et al. Thioredoxin reductase inhibitors: a patent review. *Expert Opin Ther Pat* 2017;27: 547–56.
- Zhang B, Liu Y, Li X, et al. Small molecules to target the selenoprotein thioredoxin reductase. *Chem Asian J* 2018;13: 3593–600.

8. Oblong JE, Chantler EL, Gallego A, et al. Reversible inhibition of human thioredoxin reductase activity by cytotoxic alkyl 2-imidazolyl disulfide analogues. *Cancer Chemother Pharmacol* 1994;34:434–8.
9. Hausheer FH. Compositions and methods of use of compounds to increase cancer patient survival time. US Patent no. 20090232827; *Chem Abstr* 2009;151:366710.
10. Bible KC, Isham CR, Xu R, Tibodeau JD. Methods and compositions for treating cancer. PCT Int. Appl. WO2009112014; *Chem Abstr* 2009;149:347477.
11. Arner ES, Zhong L, Holmgren A. Preparation and assay of mammalian thioredoxin and thioredoxin reductase. *Meth Enzymol* 1999;300:226–39.
12. Zhang L, Duan D, Liu Y, et al. Highly selective off-on fluorescent probe for imaging thioredoxin reductase in living cells. *J Am Chem Soc* 2014;136:226–33.
13. Ma H, Zhang J, Zhang Z, et al. A fast response and red emission probe for mammalian thioredoxin reductase. *Chem Commun* 2016;52:12060–3.
14. Kimura S, Umeda IO, Moriyama N, Fujii H. Synthesis and evaluation of a novel ^{99m}Tc-labeled bioreductive probe for tumor hypoxia imaging. *Bioorg Med Chem Lett* 2011;21:7359–62.
15. Zeynizadeh B. Oxidative coupling of thiols to disulfides with iodine in wet acetonitrile. *J Chem Res Synopses* 2002;2002:564–6.
16. Rees CW, Storr RC. 1-Chlorobenzotriazole: a new oxidant. *J Chem Soc C* 1968;1474–7.
17. Yonezawa H, Nishiyama Y, Takeo T, et al. New photocleavable linker: α -Thioacetophenone-type linker. *Bioorg Med Chem Lett* 2014;24:2831–3.
18. Arabi H, Bell RA, Howard-Lock HE, et al. N^{10} -(2'-Mercaptoethanoyl)-2,2,5,5-tetramethyl-3,4-dithia-7,10-diazabicyclo[5.3.0]decane and its reaction with oxotrichlorobis(triphenylphosphine)rhenium(V). *Can J Chem* 1996;74:574–82.
19. Lv W, Zhang G, Barinka C, et al. Design and synthesis of mercaptoacetamides as potent, selective, and brain permeable histone deacetylase 6 inhibitors. *ACS Med Chem Lett* 2017;8:510–5.
20. Hunter R, Caira M, Stellenboom N. Inexpensive, one-pot synthesis of unsymmetrical disulfides using 1-chlorobenzotriazole. *J Org Chem* 2006;71:8268–71.
21. Luthman M, Holmgren A. Rat liver thioredoxin and thioredoxin reductase: purification and characterization. *Biochemistry* 1982;21:6628–33.
22. Jan Y-H, Heck DE, Gray JP, et al. Selective targeting of selenocysteine in thioredoxin reductase by the half mustard 2-chloroethyl ethyl sulfide in lung epithelial cells. *Chem Res Toxicol* 2010;23:1045–53.
23. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 1983;65:55–63.
24. Crosley LK, Méplan C, Nicol F, et al. Differential regulation of expression of cytosolic and mitochondrial thioredoxin reductase in rat liver and kidney. *Arch Biochem Biophys* 2007;459:178–88.
25. Nalvarte I, Damdimopoulos AE, Rüegg J, Spyrou G. The expression and activity of thioredoxin reductase 1 splice variants v1 and v2 regulate the expression of genes associated with differentiation and adhesion. *Biosci Rep* 2015;35:e00269.