

## Short Total Synthesis of [15N5]-Cylindrospermopsins from 15NH4Cl Enables Precise Quantification of Freshwater Cyanobacterial Contamination

Artur K. Mailyan, Joanna L. Chen, Weiwei Li, Arturo A. Keller, Shawn Sternisha, Brian G. Miller, and Armen Zakarian

*J. Am. Chem. Soc.*, **Just Accepted Manuscript** • DOI: 10.1021/jacs.8b03071 • Publication Date (Web): 19 Apr 2018

Downloaded from <http://pubs.acs.org> on April 19, 2018

### Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

# Short Total Synthesis of [ $^{15}\text{N}_5$ ]-Cylindrospermopsins from $^{15}\text{NH}_4\text{Cl}$ Enables Precise Quantification of Freshwater Cyanobacterial Contamination

Artur K. Mailyan,<sup>†</sup> Joanna L. Chen,<sup>†</sup> Weiwei Li,<sup>‡</sup> Arturo A. Keller,<sup>‡</sup> Shawn M. Sternisha,<sup>§</sup> Brian G. Miller,<sup>§</sup> Armen Zakarian<sup>\*,†</sup>

<sup>†</sup> Department of Chemistry and Biochemistry, University of California, Santa Barbara, California 93110, United States

<sup>‡</sup> Bren School of Environmental Science and Management, University of California, Santa Barbara, California 93106, United States

<sup>§</sup> Department of Chemistry and Biochemistry, Florida State University, Florida 32310, United States.

**ABSTRACT:** Fresh water cyanobacterial algal blooms represent a major health risk because these organisms produce cylindrospermopsin, a toxic, structurally complex, zwitterionic uracil-guanidine alkaloid recognized by the EPA as a dangerous drinking water contaminant. At present, the ability to detect and quantify the presence of cylindrospermopsin in water samples is severely hampered by the lack of an isotopically-labeled standard for analytical mass spectrometry. Herein, we present a concise, scaled total synthesis of  $^{15}\text{N}$  cylindrospermopsin from  $^{15}\text{N}$  ammonium chloride, which leverages a unique stereoselective intramolecular double conjugate addition step to assemble the tricyclic guanidine core. In addition to providing the first pure isotopically labeled probe for precise quantification of this potent biotoxin in fresh water sources, our results demonstrate how unique constraints associated with isotope incorporation compel novel solutions to synthesis design.

## Introduction

In recent years, cyanobacterial algal blooms have occurred with increasing frequency near population centers worldwide, causing significant socioeconomic impact and risk to human health.<sup>1</sup> Cylindrospermopsin (CYN)<sup>2</sup> and its naturally occurring 7-*epi*-<sup>3</sup> and 7-*deoxy*-<sup>4</sup> congeners are potent biotoxins released into fresh water sources during such blooms<sup>5</sup> (Figure 1a,b). In June 2015, the US Environmental Protection Agency released a drinking water health advisory on cylindrospermopsins detailing this emerging environmental threat.<sup>6</sup> In addition to the acute toxicity risks, exposure to cylindrospermopsin and its derivatives is linked to hepatotoxicity, gastrointestinal inflammation, hemorrhages, anorexia and liver failure.<sup>7</sup>


High-performance liquid chromatography coupled with mass spectrometry is the preferred method for cylindrospermopsin detection in environmental samples.<sup>8,9,10,11,12,13</sup> In particular, isotope-dilution mass spectrometry (IDMS)<sup>14</sup> provides an attractive quantification method, as it uses an isotope of the analyte as an internal reference standard to ensure high sensitivity and precision in the measurement. The application of IDMS to cylindrospermopsin analysis, however, is hampered by the lack of isotopically labeled cylindrospermopsins. To solve this important problem, we coupled recent advances in synthetic chemistry and enzymatic bio-oxidation to prepare 7-*epi*- and 7-*deoxy*-cylindrospermopsin from [ $^{15}\text{N}$ ]-ammonium chloride, achieving >99.5% isotope incorporation into all five nitrogen atoms of these complex alkaloid natural products.

To facilitate mass spectrometry data interpretation, we sought an isotope of CYN with mass difference of at least 5 atomic units relative to the most abundant natural isotope. We considered isotopes

of hydrogen, carbon, nitrogen and oxygen for this purpose. To ensure structural integrity of the analytical standard, isotope insertion at non-exchangeable positions throughout the synthesis is essential. Based on these two key requirements, we chose to pursue the total synthesis of [ $^{15}\text{N}_1,^{15}\text{N}_3,^{15}\text{N}_{16},^{15}\text{N}_{18},^{15}\text{N}_{19}$ ]-cylindrospermopsin ([ $^{15}\text{N}_5$ ]-CYN, [ $^{15}\text{N}_5$ ]-**1**, Figure 1c). The decision to incorporate  $^{15}\text{N}$  in the cylindrospermopsin scaffold required us to rethink past synthetic strategies applied to the total synthesis of CYN and structurally related alkaloids, as these approaches are incompatible with efficient label installation at non-exchangeable atomic positions. In addition, these prior strategies, while elegant in design and execution, are laborious, requiring 20 or more steps, and present uncertainties regarding scalability.<sup>15,16,17,18,19,20,21,22</sup> Similarly, preparation of labeled cylindrospermopsins via fermentation using enriched nutrients such as  $\text{Na}^{15}\text{NO}_3$  has proven untenable.<sup>23</sup> These key concerns prompted the development of a *de novo* approach, whereby we explored whether isotope incorporation at the design stage could lead to an improved synthetic strategy that is more effective than biosynthetic or synthetic strategies reported previously.

Several principal considerations factored into the synthesis planning. Clearly, the effective introduction of the  $^{15}\text{N}$  isotope becomes a major factor. Compared to a typical total synthesis effort, the payoff of a convergent versus linear, atom-by-atom assembly is now substantially enhanced because all but the most elemental sources of the  $^{15}\text{N}$  isotope ( $^{15}\text{NH}_4\text{Cl}$  or aqueous  $^{15}\text{NH}_3$ ) become impractical.<sup>24</sup> Thus, a convergent design with preassembled intermediates containing several  $^{15}\text{N}$  isotopes is essential. The common metric of synthetic efficiency, namely, minimizing the number of individual steps based on robust, reproducible chemical transformations becomes especially critical given the costs of the starting iso-

topes.<sup>25,26,27</sup> Finally, all intended transformation must ensure retention of the isotopic label.



**Figure 1.** (a) A confocal optical microscopy image of *Cyllindrospermopsis raciborskii*, (b) the structures of cyllindrospermopsins, and (c) the outline of the synthetic design for  $[^{15}\text{N}_3]\text{-CYN}$ .

## Results and Discussion

**Synthesis Design.** Following the logic outlined above, we formulated a synthesis plan depicted in Fig 1c.  $[^{15}\text{N}_3]\text{-Guanidine}$  and  $[^{15}\text{N}_2]\text{-urea}$ , each derived from  $^{15}\text{NH}_4\text{Cl}$ , serve as precursors of the tricyclic guanidine and uracil subunits of cyllindrospermopsin, respectively. The synthesis plan requires a late stage conversion of the electron-withdrawing group Z in intermediate **i** to uracil and C–H oxidation at C7. The electron-withdrawing group Z enables a key simplification to diene **ii** by disconnection of two C–N bonds in tricyclic guanidine **i** at C8 and C10. Stereoselective intramolecular addition of guanidine to diene, although lacking precedent, is expected to ensure rapid assembly of the tricyclic guanidine subunit of cyllindrospermopsins. Alkyne **iii** was intended as a precursor to diene **ii** via a transition metal-mediated rearrangement. As an essential element of the plan, homoallylic alcohol **iv** serves as the sub-

strate for a hydroxyl-directed stereoselective delivery of free  $[^{15}\text{N}_3]\text{-guanidine}$  that we previously developed for this application.<sup>28</sup>


The success of this plan rests on identifying a suitable substituent Z, which must first activate the diene for addition of guanidine, then facilitate C7 hydroxylation, and eventually be converted to uracil. Although several options can be considered, heterocyclic substituents such as 2,4-dimethoxypyrimid-6-yl appeared to be especially enticing as precursors of uracil. Various electron-deficient heterocyclic groups are known to stabilize carbanions, and activation of double bonds by substitution with heterocycles for nucleophilic addition has been reported.<sup>29,30,31,32</sup> What remained uncertain was whether *diene* activation for double addition of guanidine, a centerpiece of our strategy, was feasible. After a careful analysis supported by experimentation, we identified 2,4-dimethoxypyrimid-6-yl as group Z because of its potential to meet these criteria most effectively.

**Preparation of  $[^{15}\text{N}]\text{-labeled Precursors.}$**  While the initial validation of the entire total synthesis was carried out with the natural isotope, we performed the scaled synthesis starting with  $^{15}\text{NH}_4\text{Cl}$  as depicted in Scheme 1.  $[^{15}\text{N}]\text{-Ammonium chloride}$  proved to be the most economical source of dry liquid  $[^{15}\text{N}]\text{-ammonia}$  on the laboratory scale, affording >95% yields of the reagent reproducibly upon treatment with NaOH in solid phase as long as the scale exceeds 0.5 g.  $[^{15}\text{N}_2]\text{-2,4-Dimethoxy-6-bromopyrimidine}$   $[^{15}\text{N}_2]\text{-5}$  was obtained by the initial conversion of  $^{15}\text{NH}_3$  to urea,<sup>33</sup> its condensation with diethyl malonate to  $[^{15}\text{N}_2]\text{-barbituric acid}$   $[^{15}\text{N}_2]\text{-4}$ , treatment with POBr<sub>3</sub>, and methanolysis with 2 equiv. of NaOMe. In contrast to the synthesis of urea, the reaction of  $^{15}\text{NH}_3$  with diphenyl thiocarbonate gave ammonium thiocyanate, pyrolysis of which was required to access  $[^{15}\text{N}_2]\text{-thiourea}$ .<sup>34</sup> The preparation of  $[^{15}\text{N}_3]\text{-guanidine hydrochloride}$  was accomplished in four steps by S-methylation, treatment with Boc<sub>2</sub>O, displacement of MeS group with  $^{15}\text{NH}_3$  in chloroform, and complete hydrolysis with 12 M HCl at reflux. The resulting  $[^{15}\text{N}_3]\text{-guanidine hydrochloride}$  and  $[^{15}\text{N}_2]\text{-5}$  constitute the source of all  $^{15}\text{N}$  isotopes in the target alkaloid.


**Total synthesis of  $[^{15}\text{N}_3]\text{-Cyllindrospermopsins.}$**  Advancing the synthesis of  $[^{15}\text{N}_3]\text{-cyllindrospermopsin}$ , bromopyrimidine  $[^{15}\text{N}_2]\text{-5}$  was subjected to Sonogashira coupling with 5-hexyn-1-ol (Scheme 2). After Swern oxidation of  $[^{15}\text{N}_2]\text{-7}$  to  $[^{15}\text{N}_2]\text{-8}$ , efficient enantioselective crotylation was achieved using Leighton's reagent **935** (99% yield, dr 10:1, er 97:3). To shorten the synthesis, we explored direct conversion of  $[^{15}\text{N}_2]\text{-7}$  to  $[^{15}\text{N}_2]\text{-10}$  using a powerful method developed by Krische and co-workers.<sup>36</sup> Although the enantiocontrol was excellent (er 98:2), lower diastereoselectivity (4:1) and yield (21%) thwarted its application for the scaled synthesis of  $[^{15}\text{N}_2]\text{-10}$  (see the supporting information for additional details). The next objective was the appendage of guanidine unit to the unactivated terminal alkene in  $[^{15}\text{N}_2]\text{-10}$ . This was achieved by a two-stage approach described previously.<sup>35</sup> First, free  $[^{15}\text{N}_3]\text{-guanidine}$  was attached to the hydroxy group by a carbonyl linker in good yield using 1.2 equiv. of  $^{15}\text{N}$ -labeled reagent. Second, electrophilic cycloguanidylation was performed in the presence of *N*-iodosuccinimide, delivering  $[^{15}\text{N}_3]\text{-12}$  after treatment with 4-methoxybenzenesulfonyl chloride (MbsCl) in 71% yield and 2.5:1 dr favoring the requisite diastereomer.

Alkyne  $[^{15}\text{N}_5]\text{-12}$  served as an effective precursor of the 1,3-diene required for the key double addition of guanidine.<sup>37,38,39,40</sup>

**Scheme 1.** The preparation of [ $^{15}\text{N}_2$ ]-2,4-dimethoxy-6-bromopyrimidine and [ $^{15}\text{N}_3$ ]-guanidine hydrochloride from  $^{15}\text{NH}_4\text{Cl}$ , the key precursors constituting the source of all  $^{15}\text{N}$  isotopes in cylindrospermopsins.



**Scheme 2.** Incorporation of all five  $^{15}\text{N}$  isotopes into advanced intermediate [ $^{15}\text{N}_5$ ]-16 from [ $^{15}\text{N}_2$ ]-2,4-dimethoxy-6-bromopyrimidine [ $^{15}\text{N}_2$ ]-5 and [ $^{15}\text{N}_3$ ]-guanidine hydrochloride.




We identified  $(\text{Ph}_3\text{P})_4\text{Pd}$  and benzoic acid at reflux in dioxane as the optimal reaction system for the challenging substrate containing two heterocyclic substituents. Diene [ $^{15}\text{N}_5$ ]-13, obtained in 75% yield as a 6:1 mixture of *E* and *Z* isomers, was now poised to implement the key cyclization to the tricyclic guanidine ring system of CYNs. Simple heating [ $^{15}\text{N}_5$ ]-13 in 14% methanol in acetonitrile (v/v) in the presence of  $^t\text{Pr}_2\text{NEt}$  at reflux resulted in


rapid methanolysis to [ $^{15}\text{N}_5$ ]-14. Remarkably, the key addition of guanidine to the diene took place concurrently with methanolysis under these mild reaction conditions, and subsequent removal of methyl carbonate in [ $^{15}\text{N}_5$ ]-14 was accomplished with NaOMe *in situ*, affording [ $^{15}\text{N}_5$ ]-16. The success of this process was essential to ensure the brevity of the total synthesis, and demonstrated the

**Scheme 3.** Total synthesis of [ $^{15}\text{N}_5$ ]-CYN, [ $^{15}\text{N}_5$ ]-7-*epi*-CYN, and 7-deoxy-CYN by a) C7 bio-hydroxylation with CyrI and b) stereoselective chemical oxidation via lithiation at C7.

a. Chemoenzymatic oxidation [**most direct**]



b. Chemical oxidation [**amenable gram to scale**]



activating properties of the 2,6-dimethoxy-6-pyrimidinyl substituent for addition of nitrogen nucleophiles to alkenes.

To address one of the most challenging phases of the synthesis, the stereospecific hydroxylation of the C7 methylene group of [ $^{15}\text{N}_5$ ]-7-deoxy-CYN ([ $^{15}\text{N}_5$ ]-3), we explored enzymatic biooxidation. This conversion represents the final step of CYN biosynthesis and is catalyzed by CyrI, an  $\alpha$ -ketoglutarate-dependent non-heme iron oxygenase. We produced pure, recombinant CyrI (*Oscillatoria* strain PCC7926) in *E. coli* and applied the enzyme in the direct oxidation of ([ $^{15}\text{N}_5$ ]-3), which was readily generated from [ $^{15}\text{N}_5$ ]-16 by acidic hydrolysis (12 M HCl, reflux) and *O*-sulfonation ( $\text{SO}_3\cdot\text{Py}$ , DMF, 95% overall yield) (Scheme 3a).<sup>41</sup> The enzyme showed high selectivity and did not accept a substrate lacking the O12 sulfate, suggesting that *O*-sulfonation must precede C7 hydroxylation in the biosynthesis of CYN. Importantly, CyrI enabled a catalytic, protecting-group-free oxidation to afford the final natural product in 93% yield with >90:1 diastereoselectivity achievable with high enzyme loading. The biocatalytic synthesis produced uniformly  $^{15}\text{N}$  labeled CYN in 10 total steps with 15% overall yield,

demonstrating the advantages of combining traditional organic chemistry with enzymatic transformations.

As a comparison to the enzymatic approach, and to gain access the 7-*epi* congener of cylindrospermospin, we also investigated chemical oxidation of [ $^{15}\text{N}_5$ ]-16 (Scheme 3b). Silylation of [ $^{15}\text{N}_5$ ]-16 with  $\text{Et}_3\text{SiCl}$  (TESCl) afforded tricyclic guanidine [ $^{15}\text{N}_5$ ]-17 in 93% overall yield from diene [ $^{15}\text{N}_5$ ]-13 as a separable 3:2 mixture of diastereomers at C8, favoring the required *R* configuration. Replacement of methyl carbonate with the TES group in [ $^{15}\text{N}_5$ ]-17 facilitated two practical objectives 1) enabling a simple chromatographic separation of C8 diastereomers, and 2) compatibility with basic conditions required for the ensuing C7 metallation and hydroxylation. Although the substrate was sensitive to the nature and stoichiometry of the base, we found that rapid lithiation with LDA (5 min,  $-78\text{ }^\circ\text{C}$ ) followed by a quench with an oxaziridine afforded the diastereomeric C7 hydroxylation products in good yields along with recovered starting material. The diastereomer ratio is dependent on the oxaziridine: *rac*-19 afforded a 6:1 dr, while camphor based reagent 20 or *ent*-20 gave 25:1 and 20:1 dr in favor of [ $^{15}\text{N}_5$ ]-1, respectively. The optimized hydroxylation was scaled to 1.50 g of

[<sup>15</sup>N<sub>5</sub>]-**17**, providing [<sup>15</sup>N<sub>5</sub>]-**21** as a pure isomer in 46% yield along with 41% of starting material. Complete hydrolysis to uracil along with the removal of the Mbs group was achieved on 0.350 g scale of [<sup>15</sup>N<sub>5</sub>]-**21** by reflux in 12 M HCl for 3 h, resulting in a minor epimerization at C7 (10:1). The mixture of epimers [<sup>15</sup>N<sub>5</sub>]-**22** and [<sup>15</sup>N<sub>5</sub>]-**23** was submitted to *O*-sulfonylation with SO<sub>3</sub>•Py complex in DMF.<sup>18</sup> After optimizing the reagent loading, we observed highly selective monosulfation of the diols at the C12 OH group. The resulting mixture was readily separated by reverse-phase HPLC, affording 75 mg of [<sup>15</sup>N<sub>5</sub>]-CYN (77% yield) and 6.8 mg of [<sup>15</sup>N<sub>5</sub>]-7-*epi*-CYN (7% yield) in a single attempt. By comparison with biooxidation using CyrI, the chemical oxidation shows improved scalability, but introduces one additional step in the total synthesis with slightly diminished overall yield and lower stereoselectivity.

With the supply of [<sup>15</sup>N<sub>5</sub>]-CYN secured by total synthesis, its development as an analytical standard for the precise quantification of CYN by LC/ESI-MS/MS was undertaken. Based on reported concentrations of the cyanotoxin in environmental samples, we developed an assay for the detection of cylindrospermopsin in a broad range from 10 ng/L to 100 µg/L. While the EPA limit for cylindrospermopsin in drinking water has been proposed to be 1 µg/L,<sup>42</sup> concentrations below this level still represent an environmental hazard due to bioaccumulation and uncertain health risks of chronic sub-lethal exposure.<sup>43</sup> The limit of detection (LOD, Signal/Noise = 3) and the limit of quantification (LOQ, Signal/Noise = 10) for the method were determined to be 2.5 ng/L and 8.25 ng/L respectively, the highest sensitivity reported to date.<sup>8-13</sup> The evaluation of matrix effect for analyte quantification was performed using environmental samples free of native CYN that were spiked with 100 µg/L. In all cases the observed recoveries were nearly 100%. This method was then used to quantify cylindrospermopsin in environmental water samples in fourteen locations in Southern California where potential contamination or water resources was suspected.<sup>44</sup> Notably, availability of [<sup>15</sup>N<sub>5</sub>]-CYN afforded the reliable and precise detection of cylindrospermopsin in Lake Casitas, a previously undescribed location of cyanotoxin contamination, where we detected the concentration of 157 ng/L of the toxin.

### Concluding Remarks

In summary, we developed a strategy for the concise total synthesis of cylindrospermopsins that is directly applicable for <sup>15</sup>N isotope incorporation. We demonstrated the scalability of the synthesis by preparing 0.125 g of [<sup>15</sup>N<sub>5</sub>]-CYN in 14 steps (longest linear sequence) and 9% overall yield from [<sup>15</sup>N]-ammonium chloride, along with 7.5 mg of [<sup>15</sup>N<sub>5</sub>]-7-*epi*-CYN and 20 mg of [<sup>15</sup>N<sub>5</sub>]-7-deoxy-CYN. In the natural isotope series, the total synthesis of CYN was completed from commercially available 2,4-dimethoxy-6-bromopyrimidine (**5**) in only 10 steps via the enzymatic oxidation route (15% yield from **5**), and 11 steps by chemical C7 hydroxylation (11% yield), representing a substantial improvement in brevity and scalability compared to the previous efforts. Furthermore, the-

se efforts provided a high-quality isotopically labeled probe for the reliable detection and quantification of cyanotoxins in fresh water sources with the highest limits of detection reported to date. Our results demonstrate how atypical constraints associated with isotope incorporation empower the development of novel approaches to organic synthesis design.

### ASSOCIATED CONTENT

#### Supporting Information

Full experimental procedures and copies of <sup>1</sup>H, <sup>13</sup>C, <sup>15</sup>N NMR spectra and crystallographic information is provided in the Supporting Information. The Supporting Information is available free of charge on the ACS Publications website.

### AUTHOR INFORMATION

#### Corresponding Author

\* zakarian@chem.ucsb.edu

#### Present Addresses

†If an author's address is different than the one given in the affiliation line, this information may be included here.

#### Author Contributions

The manuscript was written through contributions of all authors. / All authors have given approval to the final version of the manuscript. / ‡These authors contributed equally. (match statement to author names with a symbol)

#### Funding Sources

Any funds used to support the research of the manuscript should be placed here (per journal style).

#### Notes

Any additional relevant notes should be placed here.

### ACKNOWLEDGMENT

This work was supported by NIH (NIGMS, R01-077379, R01-115388 and NIEHS, R03 ES025345-01). Dr. Hongjun Zhou is acknowledged for assistance with NMR spectroscopy. Dr. Dmitriy Uchenik and the UCSB mass spectroscopy facility are thanked for assistance with mass spectral analysis. Dr. Guang Wu and the UCSB X-ray analytical facility are acknowledged for assistance with and the X-ray crystallography. We are especially grateful to Anastasiia S. Minakova for assistance with enzymatic oxidation experiment. Dr. Ben Lopez and the NRI-MCDB Microscopy Facility at UCSB are acknowledged for obtaining microphotographs of *Cylindrospermopsis raciborskii*.

### REFERENCES

- (1) Cheung, M. Y.; Liang, S.; Lee, J. J. *J. Microbiol.* **2013**, *51*, 1–10.
- (2) Ohtani, I.; Moore, R. E.; Runnegar, M. T. C. *J. Am. Chem. Soc.* **1992**, *114*, 7941–7942.

- (3) Banker, R.; Teltsch, B.; Sukenik, A.; Carmeli, S. *J. Nat. Prod.* **2000**, *63*, 387–389.
- (4) Norris, R. L.; Eaglesham, G. K.; Pierens, G.; Shaw, G. R.; Smith, M. J.; Chiswell, R. K.; Seawright, A. A.; Moore M. R. *Environ. Toxicol.* **1999**, *14*, 163–165.
- (5) De la Cruz, A. A.; Hiskia, A.; Kaloudis, T.; Chernoff, N.; Hill, D.; Antoniou, M. G.; He, X.; Loftin, K.; O'Shea, K.; Zhao, C.; Pelaez, M.; Han, C.; Lynchf, T. J.; Dionysiou, D. D. *Environ. Sci.: Processes Impacts* **2013**, *15*, 1979–2003.
- (6) Drinking water health advisory for the cyanobacterial toxin cylindrospermopsin. EPA-820R15101. <https://www.epa.gov/sites/production/files/2017-06/documents/cylindrospermopsin-report-2015.pdf>.
- (7) Falconer, I. R.; Humpage, A. R. *Environ. Toxicol.* **2006**, *21*, 299–304.
- (8) Guzmán-Guillén, R.; Prieto, A. I.; González, A. G.; Soria-Díaz, M. E.; Cameán, A. M. *Environ. Toxicol. Chem.* **2012**, *31*, 2233–2238.
- (9) Graham, J. L.; Loftin, K. A.; Meyer, M. T.; Ziegler, A. C. *Environ. Sci. Technol.* **2010**, *44*, 7361–7368.
- (10) Kikuchi, S.; Kubo, T.; Kaya, K. *Anal. Chim. Acta*, **2007**, *583*, 124–127.
- (11) Bláhová, L.; Oravec, M.; Marsálek, B.; Sejnohová, L.; Simek, Z.; Bláha, L. *Toxicon*, **2009**, *53*, 519–524.
- (12) Gallo, P.; Fabbrocino, S.; Grazia Cerulo, M.; Ferranti, P.; Bruno, M.; Serpe, L. *Rapid Commun. Mass Spectrom.*, **2009**, *23*, 3279–3284.
- (13) Kubo, T.; Sano, T.; Hosoya, K.; Tanaka, N.; Kaya, K. *Toxicon*, **2005**, *46*, 104–107.
- (14) De Bièvre, P. Techniques and instrumentation in analytical chemistry **1994**, *15*, 169–183.
- (15) Xie, C.; Runnegar, M. T. C.; Snider, B. B. *J. Am. Chem. Soc.* **2000**, *122*, 5017–5024.
- (16) Heintzelman, G. R.; Fang, W.-K.; Keen, S. P.; Wallace, G. A.; Weinreb, S. M. *J. Am. Chem. Soc.*, **2001**, *123*, 8851–8853.
- (17) Heintzelman, G. R.; Fang, W.-K.; Keen, S. P.; Wallace, G. A.; Weinreb, S. M. *J. Am. Chem. Soc.* **2002**, *124*, 3939–3945.
- (18) White, J. D.; Hansen, J. D. *J. Am. Chem. Soc.* **2002**, *124*, 4950–4951.
- (19) White, J. D.; Hansen, J. D. *J. Org. Chem.*, **2005**, *70*, 1963–1977.
- (20) Looper, R. E. & Williams, R. M. *Angew. Chem. Int. Ed.* **2004**, *43*, 2930–2933.
- (21) Looper, R. E.; Runnegar, M. T. C.; Williams, R. M. *Angew. Chem. Int. Ed.*, **2005**, *44*, 3879–3881.
- (22) Looper, R. E.; Runnegar, M. T. C.; Williams, R. M. *Tetrahedron* **2006**, *62*, 4549–4562.
- (23) Kittler, K.; Hoffmann, H.; Lindemann, F.; Koch, M.; Rohn, S.; Maul, R. *Anal. Bioanal. Chem.* **2014**, *406*, 5765–5774.
- (24) Cambridge Isotope Laboratories, Inc., listed prices: NH<sub>4</sub>Cl - \$490/10g; aqueous <sup>15</sup>NH<sub>3</sub> - \$168/5g (3.3N solution); <sup>15</sup>NH<sub>3</sub> - \$454/1 L of gas; (<sup>15</sup>NH<sub>2</sub>)<sub>2</sub>CO - \$305/1g; [<sup>15</sup>N<sub>3</sub>]-guanidine – no longer available.
- (25) Trost, B. M. *Science*, **1991**, *254*, 1471–1477.
- (26) Kuttruff, C. A.; Eastgate, M. D.; Baran, P. S. *Nat. Prod. Rep.* **2014**, *31*, 419–432.
- (27) Tian, M.; Yan, M.; Baran P. S. *J. Am. Chem. Soc.* **2016**, *138*, 14234–14237.
- (28) Mailyan, A. K.; Young, K.; Chen, J. L.; Reid, B. T.; Zakarian, A. *Org. Lett.* **2016**, *18*, 5532–5535.
- (29) Gray, A. P.; Platz, R. D.; Chang, T. C. P.; Leverone, T. R.; Ferrick, D. A.; Kramer, D. N. *J. Med. Chem.* **1985**, *28*, 111–116.
- (30) Macor, J. E.; Ordway, T.; Smith, R. L.; Verhoest, P. R.; Mack, R. A. *J. Org. Chem.* **1996**, *61*, 3228–3229.
- (31) Djung, J. F.; Hart, D. J.; Young, E. R. *J. Org. Chem.* **2000**, *65*, 5668–5675.
- (32) Schroeder, A. C.; Bloch, A.; Perman, J. L.; Bobek, M. *J. Med. Chem.* **1982**, *25*, 1255–1258.
- (33) Leitch, L.C.; Davidson, W. M. *Can. J. Agr. Sci.* **1949**, *29*, 189–190.

- 1  
2  
3  
4 (34) Laxer, A.; Major, D. T.; Gottlieb, H. E.; Fischer, B. *J. Org. Chem.* **2001**, *66*, 5463–5481.  
5  
6 (35) Kim, H.; Ho, S.; Leighton, J. L. *J. Am. Chem. Soc.* **2011**, *133*, 6517–6520.  
7  
8 (36) Kim, I. S.; Han, S. B.; Krische, M. J. *J. Am. Chem. Soc.* **2009**, *131*, 2514–2520.  
9  
10 (37) Kwong, C. K.-W.; Fu, M. Y.; Lam, C. S.-L.; Toy, P. H. *Synthesis* **2008**, *15*, 2307–2317.  
11  
12 (38) Shintani, R.; Duan, W.-L.; Park S.; Hayashi, T. *Chem. Comm.* **2006**, 3646–3647.  
13  
14 (39) Kadota, I.; Shibuya, A.; Mpaka Lutete L.; Yamamoto, Y. *J. Org. Chem.* **1999**, *64*, 4570–4571.  
15  
16 (40) Patil, N. T.; Huo, Z.; Bajracharya, G. B.; Yamamoto, Y. *J. Org. Chem.* **2006**, *71*, 3612–3614.  
17  
18 (41) Mazmouz, R.; Chapuis-Hugon, F.; Pichon, V.; Méjean, A.; Ploux, O. *ChemBioChem* **2011**, *12*, 858–862.  
19  
20 (42) Humpage, A.R.; Falconer, I.R.; *Environ. Toxicol.* **2003**, *18*, 94–103.  
21  
22 (43) Kinneer, S. *Mar. Drugs* **2010**, *8*, 542–564.  
23  
24 (44) Tatters, A. O.; Howard, M. D. A.; Nagoda, C.; Busse, L.; Gellene, A. G.; Caron, D. A. *Toxins* **2017**, *9*, 95–111.

25  
26  
27  
28 TOC graphic:  
29

