This article was downloaded by: [University of Colorado at Boulder Libraries] On: 02 January 2015, At: 02:07 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides and Nucleotides

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lncn19

³¹P NMR Studies on Oxidative Transformations of Aryl Nucleoside H-Phosphonate Diesters

Annika Kers $^{\rm a}$, Inger Kers $^{\rm a}$, Jacek Cieslak $^{\rm b}$, Adam Kraszewski $^{\rm b}$ & Jacek Stawinski $^{\rm a}$

 $^{\rm a}$ Department of Organic Chemistry, Arrhenius Laboratory , Stockholm University , S-106 91, Stockholm, Sweden

^b Institute of Bioorganic Chemistry, Polish Academy of Sciences, Noskowskiego 12/14, 61-704, Poznan, Poland Published online: 04 Oct 2006.

To cite this article: Annika Kers , Inger Kers , Jacek Cieslak , Adam Kraszewski & Jacek Stawinski (1999) ³¹P NMR Studies on Oxidative Transformations of Aryl Nucleoside H-Phosphonate Diesters, Nucleosides and Nucleotides, 18:4-5, 991-992, DOI: <u>10.1080/15257779908041625</u>

To link to this article: http://dx.doi.org/10.1080/15257779908041625

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

³¹P NMR STUDIES ON OXIDATIVE TRANSFORMATIONS OF ARYL NUCLEOSIDE H-PHOSPHONATE DIESTERS

Annika Kers, Inger Kers, Jacek Cieslaka, Adam Kraszewskia* and Jacek Stawinski*

Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, S-106 91 Stockholm, Sweden; ^aInstitute of Bioorganic Chemistry, Polish Academy of Sciences, Noskowskiego 12/14, 61-704 Poznan, Poland

ABSTRACT: Various aspects of oxidative coupling of nucleoside aryl H-phosphonate diesters with nucleosides to produce important synthetic intermediates in the phosphotriester approach to oligonucleotide synthesis, dinucleoside aryl phosphotriesters, was investigated.

Aryl H-phosphonates 2, due to the pronounced electrophilic character of their phosphorus center, are useful synthetic intermediates¹⁻³. As part of our interest in this type of H-phosphonate derivatives, we investigated various oxidative transformations of 2.



Nucleoside aryl H-phosphonates 2 upon treatment with iodine (1.5 equiv.) and ethanol (1.5 - 6 equiv.) were cleanly converted to phosphotriesters 3. An analogous oxidative coupling of 2 with 3'-protected nucleosides produced dinucleoside phosphate triesters 4 (Scheme 1), but this reaction was more sensitive to spurious water as it was apparent from the formation of varying amounts of the corresponding pyrophosphates. Under optimal reaction conditions (strictly anhydrous medium, presence of N-methylimidazole) aryl dinucleoside phosphates 4 were formed almost quantitatively, with only traces of phosphate diester present (^{31}P NMR). Compounds 4 could be isolated by silica gel column chromatography in 70-80% yield.

Since aryl nucleoside H-phosphonate diesters 2 are usually not stable enough to permit their isolation, these compounds were produced *in situ* from nucleoside H-phosphonate monoester 1 and a phenol, *via* a coupling agent promoted condensation. Along with pivaloyl chloride, diphenyl phosphorochloridate and 2-chloro-5,5-dimethyl-2-oxo-1,3,2-dioxaphosphinane were also found to be useful reagents for this purpose.



We observed, however, that aryl H-phosphonates 2 produced with different condensing agents, showed different reactivity in subsequent oxidative coupling reactions (*e.g.* formation of side products of type 5-7). These, we assigned to the presence of various nucleophilic and electrophilic species derived from condensing agents that might react with nucleosidic components and/or intermediates generated during oxidation of 2 with iodine.

Acknowledgements

Financial support from the Swedish Natural Science Research Council and the State Committee for Scientific Research, Republic of Poland, is gratefully acknowledged.

REFERENCES

- 1. Jankowska, J.; Sobkowski, M.; Stawinski, J.; Kraszewski, A. Tetrahedron Lett. 1994, 35, 3355-3358.
- Kers, A.; Kers, I.; Stawinski, J.; Sobkowski, M.; Kraszewski, A. Tetrahedron 1996, 52, 9931-9944.
- Sobkowska, A.; Sobkowski, M.; Cieslak, J.; Kraszewski, A.; Kers, I.; Stawinski, J. J. Org. Chem. 1997, 62, 4791-4794.