

View Article Online View Journal

ChemComm

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: D. Zhu, D. Chang and L. Shi, *Chem. Commun.*, 2015, DOI: 10.1039/C5CC00875A.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/chemcomm

ChemComm

RSCPublishing

Journal Name

COMMUNICATION

Cite this: DOI: 10.1039/x0xx00000x

Transition-metal-free cross-coupling of thioethers with aryl(cyano)iodonium triflates: a facile and efficient method for the one-pot synthesis of thiocyanates

previous work

Received 00th January 2012 Accepted 00th January 2012 Dan Zhu, Denghu Chang and Lei Shi*

DOI: 10.1039/x0xx00000x

www.rsc.org/

A novel transition-metal-free cross-coupling method for the one-step synthesis of thiocyanates via the C-S bond cleavage of readily available thioethers with aryl(cyano)iodonium triflates as the cyanating agent is developed. This process features relatively broad substrate scopes, less-toxic hypervalent iodine reagents, mild operating conditions, excellent functional group compatibilities, and affords various thiocyanates in moderate to good yields.

Organic thiocyanates are well-known in the area of organosulfur chemistry and have attracted considerable attention, since they are not only the substructures or building blocks of bioactive natural products¹ and pesticides,² but also the versatile synthetic precursors and useful intermediates for the synthesis of sulfur-containing compounds such as thiols,³ thioethers,⁴ disulfides,⁵ trifluoromethyl sulfides⁶ and thiocarbamates.⁷ On the one hand, thiocyanates can usually be obtained by the nucleophilic or electrophilic thiocyanation of the corresponding organic molecules with stoichiometric amounts of inorganic thiocyanates (Scheme 1, eqn. (1)),⁸ which commonly requires more careful control of the reaction conditions to avoid the undesirable formation of isothiocyanates. On the other hand, the cyanation of sulfur-containing compounds, such as thiol derivatives (Scheme 1, eqn. $(2)^9$ and $(3)^{10}$) and arylsulfinate salts (Scheme 1, eqn. (4)),¹¹ provides a beneficial complement and perfection for the aforementioned transformation. However, most of these cyanation methods are confronted with various drawbacks and limitations, including the utilization of highly poisonous and notorious cyanating agents (e.g. KCN, CuCN, Zn(CN)₂ and TMSCN), the prefunctionalization or pre-activation of substrates and harsh reaction conditions.

In the past few years, great achievements have been made in the efficient introduction of C=N moieties via alternative cyano sources¹² into molecules. More recently, Wang and co-wokers¹³ reported an efficient method for iron(II)-catalyzed direct cyanation of arenes with aryl(cyano)-iodonium triflates as the cyanating agents, which were developed by Stang and co-workers.14 ⁺ Herein. we report a one-step, metal-free synthetic route to the synthesis of thiocyanates between readily available thioethers and aryl(cyano)iodonium triflates as the cyano source under mild conditions (Scheme 1, eqn.





CBX = 1-cyano-1,2-benziodoxol-3-(1H)-one



Scheme 1 Existing methods to prepare aryl thiocyanates.

(5)). This protocol offers several advantages of relatively broad substrate scopes and the utilization of less-toxic hypervalent iodine reagents as the cyanide source. To the best of our knowledge, the present study is the first example of accessing the thiocyanates via the unusual C-S bond cleavage of thioethers under metal-free condition, although the cleavage of C-S bonds has been extensively studied with a vast range of transition-metal catalysts.¹

Published on 17 March 2015. Downloaded by FLORIDA INSTITUTE OF TECHNOLOGY on 19/03/2015 15:53:25.

View Article Online DOI: 10.1039/050008/3A10

Table 1 Optimization of the reaction conditions^a



^a The reactions were conducted with 0.5 mmol of **1a** and 0.75 mmol of **2a-b** for 12 h. ^b Isolated yields. ^c Stirred at 40 °C.

We initiated our studies by investigating the reaction of thioanisole (1a) and 1.5 equiv of 4-fluorophenyl(cyano)iodonium triflate (2a) in various commonly used organic solvents for 12 h under nitrogen atmosphere. To our delight, it was found that with CH₃CN as the solvent, the reaction provided a 77% yield of the desired product (Table 1, entry 5). In contrast, none of the expected product was formed in DMF and THF (Table1, entries 1 and 2), or relatively low yields of product were obtained in toluene and DCM (Table 1, entries 3 and 4). Further improvement in the yield was achieved by replacing 4-fluorophenyl(cyano)iodonium triflate (2a) with 3,5-di(trifluoromethyl)phenyl(cyano)iodonium triflate (2b) as the cyanating agent (Table1, entry 6). Consequently, the reaction condition described in entry 6 was selected as the standard condition for investigation of the reaction scope.



^a The reactions were proceeded in the standard condition.
^b Isolated yields. ^c The yield was calculated based on one side.
Scheme 2 Diversity in C–S bond cleavage^{a,b}

Table 2 Scope of thioether derivatives^{a,b}



^a The reactions were conducted with 0.5 mmol of **1a-u** and 0.75 mmol of **2b** for 12 h. ^b Isolated yields.

To assess the ability and efficiency of Csp^3 –S bond cleavage in this protocol, various alkyl phenyl sulfides, such as ethyl, isopropyl, benzyl, cyclopropane and phenyl sulfides were subsequently reacted with 3,5-di(trifluoromethyl)phenyl(cyano)iodonium triflate (2b) under the above-optimized reaction condition, and the results are summarized in Table 2. In all cases studied, the reaction proceeded smoothly to afford the corresponding thiocyanates 3a and 3i in moderate to excellent yields under metal-free condition.

With the highly efficient condition in regard to the cleavage of $S-CH_3$ bond in hand, we next commit to explore the scope of this reaction with respect to a variety of thioether derivatives. As shown in Table 3, the thioanisole derivatives bearing electron-donating substituents, gave slightly better yields than

those bearing electron-withdrawing substituents, such as nitryl (3e), cyano (3f). Gratifyingly, diverse functional groups, including halides (3b, 3c and 3d), cyano (3f), acetyl (3g), ester (3h), nitro (3e), alkyl (3i) and alkoxy (3j), were very well tolerated under the optimal reaction condition, furnishing the desired products in generally moderate to excellent yields. However, thioanisoles containing an active hydrogen atom such as OH, NH₂, or aldehyde did not undergo the same reactions (3s, 3t and 3u). Furthermore, the position of attached substituent groups on the phenyl ring did not affect the efficiency of this transformation (3b, 3l and 3m). In addition to mono-substituted thioanisoles, bis-substituted thioanisole was also compatible with this novel reaction, generating the corresponding thiocyanate (3n) in 59% yield. It was noteworthy that a miserable yield of 6% was observed when pentachlorothioanisole (10) was employed as substrate, possibly due to its poor solubility in acetonitrile. Besides the substituted thioanisole derivatives, the reactions of 2-(methylthio)thiophene (1q), and 2-(methylthio)pyridine (1r) with 3,5-di(trifluoromethyl)phenyl(cyano)iodonium triflate (2b) under standard condition were also investigated, and the desired thiocyanate products 3q and 3r were obtained in 68% and 22% yields, respectively. Finally, we also detected this reaction with alkyl thioethers under the standard condition, and the expected thiocyanate (3p) could be isolated in a relatively low yield of 36%.



Scheme 3 Proposed mechanism for the reaction

On the basis of a previous report by Sanford and co-workers,¹⁶ a possible mechanism is depicted in Scheme 2. Initially, the oxidation of thioethers with aryl(cyano)-iodonium triflates gives rise to the corresponding sulfonium salts A and aryl iodides. After subsequent nucleophilic substitution of A with triflate (OTf), the expected thiocyanate products could be released with alkyl triflate as byproduct to finish the reaction. To prove our mechanistic hypothesis, we repeated the model reaction (Table 1, entry 6), and to our delight, the existence of MeOTf ($R^2 = Me$) and 1-iodo-3,5bis(trifluoromethyl)benzene could been observed in the crude mixture by GC-MS analysis. Furthermore, when the reaction of thioanisole (1a) and 3,5-di(trifluoromethyl)phenyl(cyano)iodonium triflate (2b) was carried out in CDCl₃, we could also observe the generation of MeOTf and 1-iodo-3,5-bis(trifluoromethyl)benzene by ¹³C NMR analysis. These observations lend very strong support to our mechanistic proposal.

In summary, we have developed a transition-metal-free transformation of readily available thioethers into the corresponding thiocyanates through the Csp^3 –S bond cleavage. This process features relatively broad substrate scopes, less-toxic hypervalent iodine reagents, mild operating conditions, excellent functional group compatibilities, and affords various thiocyanates in moderate to good yields. Notably, this protocol is not only complementary to existing methods of thiocyanate formation, but also offers an opportunity to extend the applications of less-toxic hypervalent iodine reagents in organic synthesis. Further investigations on related reactions involving hypervalent iodine reagents for the

preparation of other useful sulfur-containing compounds are still underway in our laboratory.

The work was supported by the "Hundred Talents Program" of Harbin Institute of Technology (HIT), the NSFC (21202027), the NCET (NCET-12-0145), and the "Technology Foundation for Selected Overseas Chinese Scholar" of Ministry of Human Resources and Social Security of China (MOHRSS).

Notes and references

Institute of Organic Chemistry, The Academy of Fundamental and Interdisciplinary Science, Harbin Institute of Technology, Harbin 150080, P. R. China. Tel: +86-451-86403860; E-mail: <u>lshi@hit.edu.cn</u>

† Electronic Supplementary Information (ESI) available: Detailed synthetic procedures and characterization of new compounds. See DOI: 10.1039/c000000x/

- (a) A. D. Patil, A. J. Freyer, R. Reichwein, B. Carte, L. B. Killmer, L. Faucette, R. K. Johnson and D. J. Faulkner, *Tetrahedron Lett.*, 1997, **38**, 363–364; (b) S. Dutta, H. Abe, S. Aoyagi, C. Kibayashi and K. S. Gates, *J. Am. Chem. Soc.*, 2005, **127**, 15004–15005; (c) A. T. Pham, T. Ichiba, W. Y. Yoshida, P. J. Scheuer, T. Uchida, J.-I. Tanaka and T. Higa, *Tetrahedron Lett.*, 1991, **32**, 4843–4846; (d) I. C. Piña, J. T. Gautschi, G.-Y.-S. Wang, M. L. Sanders, F. J. Schmitz, D. France, S. Cornell-Kennon, L. C. Sambucetti, S. W. Remiszewski, L. B. Perez, K. W. Bair and P. Crews, *J. Org. Chem.*, 2003, **68**, 3866–3873.
- 2 A. A. Newman, Chemistry and Biochemistry of Thiocyanic Acid and Its Derivatives, Academic Press, London, 1975.
- 3 (a) J. Houk and G. M. Whitesides, J. Am. Chem. Soc., 1987, 109, 6825–6836; (b) L. Linderoth, P. Fristrup, M. Hansen, F. Melander, R. Madsen, T. L. Andresen and G. H. Peters, J. Am. Chem. Soc., 2009, 131, 12193–12200.
- 4 (a) Z. Pakulski, D. Pierożyński and A. Zamojski, *Tetrahedron*, 1994, 50, 2975–2992; (b) I. W. J. Still and F. D. Toste, *J. Org. Chem.*, 1996, 61, 7677–7680; (c) F. Ke, Y. Qu, Z. Jiang, Z. Li, D. Wu and X. Zhou, *Org. Lett.*, 2011, 13, 454–457.
- 5 (a) K. R. Prabhu, A. R. Ramesha and S. Chandrasekaran, J. Org. Chem., 1995, 60, 7142–7143; (b) D. Sengupta and B. Basu, *Tetrahedron Lett.*, 2013, 54, 2277–2281.
- 6 T. Billard, S. Large and B. R. Langlois, *Tetrahedron Lett.*, 1997, 38, 65–68.
- 7 (a) R. Riemschneider, J. Am. Chem. Soc., 1956, 78, 844–847; (b) R. Riemschneider, F. Wojahn and G. Orlick, J. Am. Chem. Soc., 1951, 73, 5905–5907.
- For a selected review, see: (a) A. W. Erian and S. M. Sherif, 8 Tetrahedron, 1999, 55, 7957-8024; For representative examples on nucleophilic or electrophilic thiocyanation, see: (b) N. Sun, H. Zhang, W. Mo, B. Hu, Z. Shen and X. Hu, Synlett, 2013, 24, 1443-1447; (c) K. Takagi, H. Takachi and K. Sasaki, J. Org. Chem., 1995, 60, 6552-6556; (d) I. P. Beletskaya, A. S. Sigeev, A. S. Peregudov and P. V. Petrovskii, Mendeleev Communications, 2006, 5, 250-251; (e) B. Akhlaghinia, A.-R. Pourali and M. Rahmani, Synth. Commun., 2012, 42, 1184-1191; (f) M. Barbero, I. Degani, N. Diulgheroff, H. Dughera and R. Fochi, Synthesis, 2001, 4, 585-590; (g) B. D. Grishchuk, S. I. Klimnyuk, G, Ya. Zagrichuk, M. P. Kravchenyuk, T. S. Kolomiets and P. M. Gorbovoi, Pharm. Chem. J, 1999, 33, 143-144; (h) K. Takagi, H. Takachi and N. Hayama, Chem. Lett., 1992, 509-510; (i) K. Tamao, T. Kakui and M. Kumada, Tetrahedron Lett., 1980, 21, 111-114; (j) B. R. Vaddula, R. S. Varma and J. Leazer, E. J. Org. Chem., 2012, 6852-6855; (k) R. G. Guy, in The Chemistry of Cyanates and Their Thio Derivatives, ed. S. Patai, Wiley, Chichester, 1977, ch. 18; (l) Y.-F. Wang, Y. Zhou, J.-R. Wang, L. Liu and Q.-X. Guo, Chin. Chem. Lett., 2006, 17, 1283.
- For selected examples, see: (a) B. M. Trost, W. L. Schinski, F. Chen and I. B. Mantz, J. Am. Chem. Soc., 1971, 93, 676–684; (b) Y. Q. Wu, D. C. Limburg, D. E. Wilkinson and G. S. Hamilton, Org. Lett., 2000, 2, 795– 797; (c) Z. Miao and J. P. Tam, Org. Lett., 2000, 2, 3711–3713; (d) I. W. J. Still and I. D. G. Watson, Synth. Commun., 2001, 31, 1355–1359; (e) G. Westerberg and B. Långström, Acta Chemica Scandinavica, 1993, 47, 974–978; (f) G. Pipes, A. Kosky, J. Abel, Y. Zhang, M. Treuheit and G. Kleemann, Pharm. Res., 2005, 22, 1059–1068; (g) J. J. Kim, D. H. Kweon, S. D. Cho, H. K. Kim, E. Y. Jung, S. G. Lee, J. R. Falck and Y.

J. Yoon, Tetrahedron, 2005, **61**, 5889–5894; (*h*) K. Yamaguchi, K. Sakagami, Y. Miyamoto, X. Jin and N. Mizuno, *Org. Biomol. Chem.*, 2014, **12**, 9200–9206; (*i*) R. Frei, T. Courant, M. D. Wodrich and J. Waser, *Chem. Eur. J.*, 2015, **21**, 2662–2668.

- For selected examples, see: (a) F. Teng, J. T. Yu, H. T. Yang, Y. Jiang and J. Cheng, Chem. Commun., 2014, **50**, 12139–12141; (b) K. Takeda, K. Horiki, Heterocycles, 1990, **30**, 367–373; (c) H. F. Gilbert, J. Am. Chem. Soc., 1980, **102**, 7059–7065; (d) K. Fujiki, E. Yoshida, Synth. Commun., 1999, **29**, 3289–3294; (e) J. L. Kice and J. M. Anderson, J. Org. Chem., 1968, **33**, 3331–3333; (f) R. A. Gancarz and J. L. Kice, J. Org. Chem., 1981, **46**, 4899–4906; (g) T. Castanheiro, M. Gulea, M. Donnard, and J. Suffert, Eur. J. Org. Chem., **2014**, 7814–7817.
- For selected examples, see: (a) S. Kagabu, K. Sawahara, M. Maehara, S. Ichihashi and K. Saito, *Chem. Pharm. Bull.*, 1991, **39**, 784–785; (b) S. Kagabu, M. Maehara, K. Sawahara and K. Saito, *J. Chem. Soc.*, *Chem. Commun.*, 1988, 1485–1486; (c) S. Harusawa and T. Shioiri, *Tetrahedron Lett.*, 1982, **23**, 447–448.
- 12 For a review, see: (a) J. Kim, H. J. Kim and S. Chang, Angew. Chem. Int. Ed., 2012, 51, 11948-11959; Angew. Chem., 2012, 124, 12114-12125; For selected examples, see: (b) P. Anbarasan, H. Neumann and M. Beller, Chem. Eur. J., 2010, 16, 4725-4728; (c) S. Ushijima and H. Togo, Synlett, 2010, 1562-1566; (d) G. Zhang, X. Ren, J. Chen, M. Hu and J. Cheng, Org. Lett., 2011, 13, 5004-5007; (e) G. Ishii, K. Moriyama and H. Togo, Tetrahedron Lett., 2011, 52, 2404-2406; (f) S. Ushijima, K. Moriyama and H. Togo, Tetrahedron, 2011, 67, 958-964; (g) S. Kamijo, T. Hoshikawa and M. Inoue, Org. Lett., 2011, 13, 5928-5931; (h) P. Anbarasan, H. Neumann and M. Beller, Angew. Chem. Int. Ed., 2011, 50, 519-522; Angew. Chem., 2011, 123, 539-542; (i) S. Ding and N. Jiao, J. Am. Chem. Soc., 2011, 133, 12374-12377; (j) J. Kim, J. Choi, K. Shin and S. Chang, J. Am. Chem. Soc., 2012, 134, 2528-2531; (k) S. Zheng, C. Yu and Z. Shen, Org. Lett., 2012, 14, 3644-3647; (1) J. Peng, J. Zhao, Z. Hu, D. Liang, J. Huang and Q. Zhu, Org. Lett., 2012, 14, 4966-4969; (m) G. Zhang, S. Chen, H. Fei, J. Cheng and F. Chen, Synlett, 2012, 23, 2247-2250; (n) Q. Wen, J. Jin, B. Hu, P. Lu and Y. Wang, RSC Adv., 2012, 2, 6167-6169; (o) S. Xu, X. Huang, X. Hong and B. Xu, Org. Lett., 2012, 14, 4614-4617; (p) G.-Y. Zhang, J.-T. Yu, M.-L. Hu and J. Cheng, J. Org. Chem., 2013, 78, 2710-2714; (q) O. Y. Yuen, P. Y. Choy, W. K. Chow, W. T. Wong and F. Y. Kwong, J. Org. Chem., 2013, 78, 3374-3378; (r) C. Pan, H. Jin, P. Xu, X. Liu, Y. Cheng and C. Zhu, J. Org. Chem., 2013, 78, 9494-9498; (s) C. Tao, F. Liu, Y. Zhu, W. Liu and Z. Cao, Org. Biomol. Chem., 2013, 11, 3349-3354; (t) A. B. Khemnar, D. N. Sawant and B. M. Bhanage, Tetrahedron Lett., 2013, 54, 2682-2684; (u) K. Zheng, P. Yu, S. Chen, F. Chen and J. Cheng, Chin. J. Chem., 2013, 31, 449-452; (v) Y. Yan, Y. Yuan and N. Jiao, Org. Chem. Front., 2014, 1, 1176-1179; (w) Y. Yang and S. L. Buchwald, Angew. Chem., Int. Ed., 2014, 53, 8677-8681; (x) X. Hong, H. Wang, G. Qian, Q. Tan and B. Xu, J. Org. Chem., 2014, 79, 3228-3237.
- 13 Z. Shu, W. Ji, X. Wang, Y. Zhou, Y. Zhang and J. Wang, Angew. Chem. Int. Ed., 2014, 53, 2186–2189; Angew. Chem., 2014, 126, 2218–2221.
- For reviews, see: (a) P. J. Stang and V. V. Zhdankin, *Chem. Rev.*, 1996, 96, 1123–1178; (b) V. V. Zhdankin and P. J. Stang, *Chem. Rev.*, 2002, 102, 2523–2584.
- For a review, see: (a) F. Pan and Z.-J. Shi, ACS Catal., 2014, 4, 280–288; For selected examples, see: (b) D. H. Ortgies, F. Chen and P. Forgione, E. J. Org. Chem., 2014, 3917–3922; (c) T. Sugahara, K. Murakami, H. Yorimitsu, and A. Osuka, Angew. Chem. Int. Ed., 2014, 53, 9329–9333; Angew. Chem., 2014, 126, 9483–9487; (d) K. Murakami, H. Yorimitsu and A. Osuka, Angew. Chem. Int. Ed., 2014, 53, 7510–7513; Angew. Chem., 2014, 126, 7640–7643; (e) S. Otsuka, D. Fujino, K.Murakami, H. Yorimitsu and A. Osuka, Chem. Eur. J., 2014, 20, 13146–13149.
- 16 A. M. Wagner and M. S. Sanford, J. Org. Chem., 2014, 79, 2263-2267.