


This article was downloaded by: [Akdeniz Universitesi]

On: 20 December 2014, At: 00:23

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides and Nucleotides

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn19>

1,5-Anhydro-2-Deoxy-D-Altritol Oligonucleotides as Conformationally Restricted Analogues of Rna

Brigitte Allart^a, Arthur Van Aerschot^a & Piet Herdewijn^a

^a Laboratory of Medicinal Chemistry, Rega Institute for Medical Research, Katholieke Universiteit Leuven, Minderbroedersstraat 10, B-3000, Leuven, Belgium

Published online: 22 Aug 2006.

To cite this article: Brigitte Allart, Arthur Van Aerschot & Piet Herdewijn (1998) 1,5-Anhydro-2-Deoxy-D-Altritol Oligonucleotides as Conformationally Restricted Analogues of Rna, Nucleosides and Nucleotides, 17:9-11, 1523-1526, DOI: [10.1080/07328319808004682](https://doi.org/10.1080/07328319808004682)

To link to this article: <http://dx.doi.org/10.1080/07328319808004682>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>


1,5-ANHYDRO-2-DEOXY-D-ALTRITOL OLIGONUCLEOTIDES AS CONFORMATIONALLY RESTRICTED ANALOGUES OF RNA

Brigitte Allart, Arthur Van Aerschot, and Piet Herdewijn*

Laboratory of Medicinal Chemistry, Rega Institute for Medical Research, Katholieke Universiteit Leuven, Minderbroedersstraat 10, B-3000 Leuven, Belgium

Abstract : As part of a project concerning the investigation of new hexitol nucleic acids (HNA), the 1,5-anhydro-2-deoxy-D-altritol nucleoside building blocks with a uracil, cytosine, adenine and guanine base moiety were synthesized. The uracil analogue was used for the automated synthesis of corresponding oligonucleotides. Hybridization capabilities of these altritol nucleic acids (ANA) are illustrated by the T_m values obtained for the $(a^hU)_{13}/(dA)_{13}$ duplex.

Hexitol nucleic acids (HNA) built from 1,5-anhydro-2,3-dideoxy-D-arabino-hexitol nucleoside analogues **I** ($X=H$)¹ have proven to be one of the most efficient conformationally restricted oligonucleotides in terms of duplex formation with natural nucleic acids, and therefore are very promising candidates for antisense strategies². The strong hybridizing properties of HNA could be explained by the conformational preference of the hexitol moiety to adopt a ⁴C₁ conformation, which fits very well with the 3'-endo conformation adopted by furanose nucleosides in natural nucleic acids (A form). This ⁴C₁ conformation is preserved by introducing a supplementary hydroxy group on C-3' in α -position³. This result prompted us to consider the synthesis of (6' → 4') oligonucleotides built from 1,5-anhydro-2-deoxy-D-altritol nucleosides **II**.



I: X = H 1,5-anhydro-2,3-dideoxy-D-arabino-hexitol (hA, hG, hT, hC)

II: X = OH 1,5-anhydro-2-deoxy-D-altrito-hexitol (ahA, ahG, ahT, ahC)




FIG. 1

General scheme for the synthesis of the *altritrol* phosphoramidite building blocks

In this communication we describe the general synthetic route to prepare the uracil phosphoramidite building block **6a** (*a*^hU) and the synthesis of the all-uracil 13-mer (*a*^hU)₁₃. Compound **6a** was synthesized in 10 steps starting from tetraacetyl- α -D-bromoglucose (FIG. 1) as follows:

Introduction of the base moiety was achieved by the regiospecific opening of 1,5:2,3-dianhydro-4,6-*O*-benzylidene-D-allitol **1** (prepared in 5 steps from tetraacetyl bromoglucose according to the procedure of Kocienski⁴) by the sodium salt of uracil. On ribonucleosides, the tert-butyldimethylsilyl group is most commonly used for protection of the free 2'-OH position which is necessary for RNA synthesis. Because of the steric hindrance protection of the 3'-OH group of **2a** with a tert-butyldimethylsilyl group proved to be difficult. Therefore we used a benzoyl protecting group which is, however, prone to migration reaction. Treatment of **3a** with 90% aq. CF₃CO₂H (1h), careful evaporation at T < 40°C followed by precipitation in diethyl ether afforded the crude product **4a** in 70% yield. Compound **4a** was then monomethoxytritylated without further purification. Since benzoyl migration from the 3'- to the 4'- position was found to occur more quickly under the classical conditions (MMTrCl/pyridine/4,4'-dimethylaminopyridine as catalyst), the reaction was performed in DMF, using silver nitrate as activator (1 eq) in presence of a small quantity of 2,4,6-collidine (1.5 eq) [4] and led to **5a** in 74% yield. Compound **5a** was converted into the corresponding 2-cyanoethyl-*N,N*-diisopropyl phosphoramidite **6a** by the general procedure in 94% yield⁵.

TABLE I : Compared T_m values obtained for pairing of *altrito-(a^h)* and *arabino-hexitol (h)* oligomers with their DNA complement

mixture (4 μM of each strand)	T _m ^(a)	
	0.1M NaCl	1M NaCl
(<i>a^h</i> U) ₁₃ /(dA) ₁₃	30	48
(hU) ₁₃ /(dA) ₁₃	22	36-39

(a) T_m were measured in a buffer containing 0.02M KH₂PO₄, pH 7.5, 0.1 mM EDTA with 0.1 or 1M NaCl

Compounds **5b**, **5c**, and **5d** were synthesized following a similar synthetic scheme, with an additional step for protection of the base moiety (*N*⁶-benzoyl, *N*⁴-benzoyl or *N*²-isobutyryl for adenine, cytosine and guanine derivatives respectively) using conventional procedures and their conversion into the phosphoramidites **6b**, **6c** and **6d** is under way.

Oligonucleotide synthesis : Making use of compound **6a**, a (all-uracil) homopolymer was synthesized with the aim to optimize the conditions of oligomer deprotection and to evaluate the hybridization capabilities of the altritol nucleic acids.

Oligonucleotide synthesis was carried out on an automated DNA-synthesizer using the phosphoramidite approach, with some modifications for final deprotection of the oligomer. After assembly, the 2- cyanoethyl groups were selectively cleaved by a 30 min. treatment of the support with a piperidine/dioxane (1:4) solution, before final treatment with concentrated ammonia (50°C, 16h). The (all-uracil) 13-mer prepared from **6a** was found to be >95% homogeneous after purification.

Hybridization Properties : The UV-melting curves at 260 nm (and 284 nm) of the all-uracil altritol oligomer with complementary DNA (dA)₁₃ demonstrate the formation of a (*a^h*U)₁₃/(dA)₁₃ complex. According to T_m values (**TABLE I**), the (*a^h*U)₁₃/(dA)₁₃ duplex is more stable than the (hU)₁₃/(dA)₁₃ duplex. This may be attributed to a better hydration of the altritol nucleic acid/DNA duplex, which is now under investigation. Incorporation of the other phosphoramidite derivatives **6b**, **6c** and **6d** and the study of the pairing properties of this new construct with DNA and with RNA is in progress. The preliminary studies, however, demonstrate that altritol nucleic acids might form more stable duplexes with natural nucleic acids than HNA itself.

REFERENCES

1. Van Aerschot, A., Verheggen, I., Hendrix, C., and Herdewijn, P. *Angewandte Chemie Int. Ed. English*, **1995**, *34*, 1338-1339. Hendrix, C., Verheggen, I., Rosemeyer, H., Seela, F., Van Aerschot, A., Herdewijn, P., *Chemistry Eur. J.* **1997**, *3*, 110-120.

2. Herdewijn, P. *Liebigs Ann.*, **1996**, 1337-1348.
3. Pérez-Pérez, M.-J., De Clercq, E., and Herdewijn, P. *Bioorg. Med. Chem. Lett.*, **1996**, *6(13)*, 1457-1460.
4. Reddy, M. P., Rampal, J. B., and Beaucage, S. L., *Tetrahedron Lett.*, **1987**, *28(1)*, 23-26.
5. Beaucage, S. L., and Iyer, R. P. *Tetrahedron*, **1993**, *49*, 192-1963.