


Metal-Free C–H Cross-Coupling toward
Oxygenated Naphthalene-Benzene Linked
BiarylsToshifumi Dohi,[†] Motoki Ito,^{†,‡} Itsuki Itani,[†] Nobutaka Yamaoka,[†] Koji Morimoto,[†]
Hiromichi Fujioka,[‡] and Yasuyuki Kita^{*,†}College of Pharmaceutical Sciences, Ritsumeikan University, 1-1-1 Nojihigashi,
Kusatsu, Shiga 525-8577, Japan, and Graduate School of Pharmaceutical Sciences,
Osaka University, 1-6 Yamada-oka, Suita, Osaka 565-0871, Japan

kita@ph.ritsumeik.ac.jp

Received September 30, 2011

ABSTRACT



The intermolecular C–H cross-coupling between aromatic ethers has been achieved for the first time using perfluorinated hypervalent iodine(III) compounds as extreme single-electron-transfer (SET) oxidants. The demonstrations of this specific coupling could provide a direct route to valuable oxygenated mixed naphthalene-benzene biaryls **3** only, without formation of other biaryl-derived byproducts.

The oxidative coupling between arene C–H bonds is recognized as a straightforward and attractive approach that creates new C–C bonds of biaryl molecules without the use of prefunctionalized substrates, i.e., organometallic compounds or halides.¹ A recent breakthrough in this synthetic area by participation of transition metal chemistry has permitted direct access to a variety of mixed biaryl

compounds containing heteroaromatic rings² and metal-coordinating directing groups³ by C–H cross-couplings. On the other hand, oxidative cross-coupling between C–H bonds of aromatic hydrocarbons not containing directing groups, such as naphthalenes and benzenes, is still difficult in regard to the product selectivity and yield even after modern synthetic advances.⁴ Previously, we reported the unprecedented success of the cross-coupling between naphthalenes and alkylbenzenes using phenyliodine bis(trifluoroacetate) (PIFA).⁵ As a continuation of our

[†] Ritsumeikan University.[‡] Osaka University.

(1) (a) McGlacken, G. P.; Bateman, L. M. *Chem. Soc. Rev.* **2009**, 38, 2447. (b) Ashenhurst, J. A. *Chem. Soc. Rev.* **2010**, 39, 540. (c) Yeung, C. S.; Dong, V. M. *Chem. Rev.* **2011**, 111, 1215.

(2) (a) Stuart, D. R.; Fagnou, K. *Science* **2007**, 316, 1172. (b) Stuart, D. R.; Villemure, E.; Fagnou, K. *J. Am. Chem. Soc.* **2007**, 129, 12072. (c) Dwight, T. A.; Rue, N. R.; Charyk, D.; Josselyn, R.; DeBoef, B. *Org. Lett.* **2007**, 9, 3137. (d) Cho, S. H.; Hwang, S. J.; Chang, S. J. *Am. Chem. Soc.* **2008**, 130, 9254. (e) Xi, P.; Yang, F.; Qin, S.; Zhao, D.; Lan, J.; Gao, G.; Hu, C.; You, J. *J. Am. Chem. Soc.* **2010**, 132, 1822. (f) He, C.-Y.; Fan, S.; Zhang, X. *J. Am. Chem. Soc.* **2010**, 132, 12850. (g) Wang, Z.; Li, K.; Zhao, D.; Lan, J.; You, J. *Angew. Chem., Int. Ed.* **2011**, 50, 5365.

(3) (a) Hull, K. L.; Sanford, M. S. *J. Am. Chem. Soc.* **2007**, 129, 11904. (b) Li, B.-J.; Tian, S.-L.; Fang, Z.; Shi, Z.-J. *Angew. Chem., Int. Ed.* **2008**, 47, 1115. (c) Brasche, G.; Garcia-Fortanet, J.; Buchwald, S. L. *Org. Lett.* **2008**, 10, 2207. (d) Zhao, X.; Yeung, C. S.; Dong, V. M. *J. Am. Chem. Soc.* **2010**, 132, 5837.

(4) (a) Li, R.; Jiang, L.; Lu, W. *Organometallics* **2006**, 25, 5973. (b) Rong, Y.; Li, R.; Lu, W. *Organometallics* **2007**, 26, 4376. (c) Wei, Y.; Su, W. *J. Am. Chem. Soc.* **2010**, 132, 16377.

(5) Dohi, T.; Ito, M.; Morimoto, K.; Iwata, M.; Kita, Y. *Angew. Chem., Int. Ed.* **2008**, 47, 1301.

(6) Intramolecular couplings: (a) Kita, Y.; Gyoten, M.; Ohtsubo, M.; Tohma, H.; Takada, T. *Chem. Commun.* **1996**, 1481. (b) Takada, T.; Arisawa, M.; Gyoten, M.; Hamada, R.; Tohma, H.; Kita, Y. *J. Org. Chem.* **1998**, 63, 7698. (c) Hamamoto, H.; Anilkumar, G.; Tohma, H.; Kita, Y. *Chem.—Eur. J.* **2002**, 8, 5377.

(7) Intermolecular couplings: (a) Arisawa, M.; Utsumi, S.; Nakajima, M.; Ramesh, N. G.; Tohma, H.; Kita, Y. *Chem. Commun.* **1999**, 469. (b) Tohma, H.; Morioka, H.; Takizawa, S.; Arisawa, M.; Kita, Y. *Tetrahedron* **2001**, 57, 345. (c) Tohma, H.; Iwata, M.; Maegawa, T.; Kita, Y. *Tetrahedron Lett.* **2002**, 43, 5377.

research involving metal-free biaryl couplings using hypervalent iodine chemistry,^{5–8} we now report the extremely specific C–H cross-coupling for a new combination of aromatic rings, that is, between aromatic ethers, to give the corresponding valuable oxygenated mixed biaryls (see examples in eq 1).

Highly oxygenated biaryl structures, which are frequently involved in many natural products, have been extensively utilized as the core components of pharmaceuticals, dyes, liquid crystals, organic devices, and conductors and for the design of ligands and catalysts in organic synthesis.⁹ The intramolecular oxidative couplings of phenyl ether rings are widely used for the construction of the cyclic biaryl structures of these natural products and other useful compounds.¹⁰ However, some difficulties emerge when performing intermolecular cross-coupling by the methods. The first commonly known problem is the coformation of an undesired homocoupling dimer together with a mixed biaryl product.^{11,12} Another specific concern regarding the coupling of aromatic ethers is that the formed biaryls, that is, the oxidation-sensitive electron-rich compounds,¹³ would be further oxidized and produce oligomers, quinone derivatives, and other byproducts, thus decreasing the yield of the coupling product. The intermolecular coupling for obtaining oxygenated mixed biaryls is believed to be very troublesome because of the chemoselective issues of both the substrates and products.

With respect to the excellent single-electron-transfer (SET) oxidizing ability of hypervalent iodine reagents to aromatic rings,¹⁴ we first anticipated their possible use with the intermolecular cross-coupling between aromatic ethers employing PIFA. Indeed, the treatment of PIFA in the presence of the activator BF₃·Et₂O^{5–7} in an attempt at the

cross-coupling of the naphthalene and phenyl ethers **1a** and **2a** (eq 1) produced the expected naphthalene-benzene linked biaryl **3aa** in 63% yield (Table 1, entry 1). Nevertheless, PIFA caused homocoupling in this case, along

Table 1. Influence of the Structure of Hypervalent Iodine Reagents in the Cross-Coupling of **1a** and **2a** (eq 1)^a

entry	oxidant Ar-I(OCOCF ₃) ₂	mixed biaryl 3aa ^b (%)	homocoupling of 2a ^c
1	Ar = Ph (PIFA)	63	+ (10%)
2	Ar = 4-MeC ₆ H ₄	14	+
3	Ar = 4-MeOC ₆ H ₄	24	+
4	Ar = 2,4,6-MeC ₆ H ₂	26	+
5	Ar = 4-CF ₃ C ₆ H ₄	63	–
6	Ar = C ₆ F ₅ (FPIFA)	82	–
7	Ar = 4-CF ₃ C ₆ F ₄	78	–
8 ^d	(FPIFA)	>99	–
9 ^{d,e}	(FPIFA)	96	–

^a Reactions were examined using **1a** (1 equiv) and **2a** (2 equiv) with oxidants (1 equiv) in the presence of BF₃·Et₂O (2 equiv) at 0 °C.

^b Isolated yield of the pure product based on the substrate **1a** used.

^c The formation of the homodimer of **2a**, that is, 2,2'-dimethyl-4,4',6,6'-tetramethoxybiphenyl, was checked by ¹H NMR, GC, and HPLC.

^d Reaction was performed at –40 °C. ^e 4 equiv of **2a**.

with ca. 10% formation of the undesired dimer of methoxybenzene **2a**. Detailed screenings of the reaction conditions, such as the concentration, temperature, solvent, amounts of the reagent and activator, order of the reagent addition and their addition rate, were examined for the PIFA/BF₃·Et₂O system for the purpose of suppressing the homodimer, the formation of which had not yet been eliminated. An alternative use of bromotrimethylsilane as an activator⁸ was then investigated, which did not yield the mixed biaryl **3aa**. Instead, we found that a subtle change in the oxidant nature significantly affected the distribution of the coupling products, the cross-coupling biaryl **3aa** versus the homodimer of **2a**, as well as the reaction yield.

In comparison to PIFA, the relatively electron-rich aryl moieties in the oxidants Ar-I(OCOCF₃)₂, where Ar = 4-Me, 4-MeO, and 2,4,6-Me, were less positive, and the product **3aa** was no longer formed in good yield (entries 2–4). In addition, formation of the homodimer of **2a** was observed. On the contrary, the oxidant having the CF₃ substituent as the electron-deficient group showed a degree of product formation similar to that of PIFA, while the homodimer was not detected (entry 5). With this perspective, we further evaluated the perfluorinated PIFA^{15a–c}

(8) Cross-couplings: (a) Kita, Y.; Morimoto, K.; Ito, M.; Ogawa, C.; Goto, A.; Dohi, T. *J. Am. Chem. Soc.* **2009**, *131*, 1668. (b) Morimoto, K.; Yamaoka, N.; Ogawa, C.; Nakae, T.; Fujioka, H.; Dohi, T.; Kita, Y. *Org. Lett.* **2010**, *12*, 3804.

(9) (a) Bringmann, G.; Gulder, T.; Gulder, T. A. M.; Breuning, M. *Chem. Rev.* **2011**, *111*, 563. (b) Hertweck, C. *Angew. Chem., Int. Ed.* **2009**, *48*, 4688. (c) Bringmann, G.; Gunther, C.; Ochse, M.; Schupp, O.; Tasler, S. *Prog. Chem. Org. Nat. Prod.* **2001**, *82*, 1. (d) van Leeuwen, P. W. N. M.; Kamer, P. C. J.; Claver, C.; Pamies, O.; Dieguez, M. *Chem. Rev.* **2011**, *111*, 2077. (e) Terada, M. *Chem. Commun.* **2008**, 4097.

(10) (a) Taylor, E. C.; Andrade, J. G.; Rall, G. J. H.; McKillop, A. *J. Am. Chem. Soc.* **1980**, *102*, 6513. (b) Planchenault, D.; Dhal, R.; Robin, J. P. *Tetrahedron* **1993**, *49*, 5823. (c) Tanaka, M.; Mukaiyama, C.; Mitsuhashi, H.; Maruno, M.; Wakamatsu, T. *J. Org. Chem.* **1995**, *60*, 4339 and references therein. (d) Zhai, L.; Shukla, R.; Rathore, R. *Org. Lett.* **2009**, *11*, 3474. See also our methods in ref 6.

(11) (a) Nyberg, K. *Acta. Chem. Scand.* **1973**, *27*, 503. (b) Nyberg, K. *Chem. Scr.* **1974**, *5*, 120.

(12) The oxidative cross-couplings of naphthols were reported, which are not applicable to the naphthyl ether couplings: (a) Hovorka, M.; Gunterova, J.; Zavada, J. *Tetrahedron Lett.* **1990**, *31*, 413. (b) Vyskocil, S.; Smrcina, M.; Lorenc, M.; Hanus, V.; Polasek, M.; Kocovsky, P. *Chem. Commun.* **1998**, 585. (c) Li, X.; Hewgley, J. B.; Mulrooney, C. I. A.; Yang, J.; Kozlowski, M. C. *J. Org. Chem.* **2003**, *68*, 5500.

(13) Oxygenated aryl naphthalenes are sensitive to oxidations. For examples of quinone formations, see: (a) Lumb, J.-P.; Trauner, D. *Org. Lett.* **2005**, *7*, 5865. (b) Larsen, D. S.; O'Shea, M. D. *J. Org. Chem.* **1996**, *61*, 5681. (c) Schäfer, H. J. In *Organic Electrochemistry*; Lund, H., Baizer, M. M., Eds.; Marcel Dekker: New York, 1991; Chapter 23.

(14) (a) Kita, Y.; Tohma, H.; Hatanaka, K.; Takada, T.; Fujita, S.; Mitoh, S.; Sakurai, H.; Oka, S. *J. Am. Chem. Soc.* **1994**, *116*, 3684. Accounts: (b) Kita, Y.; Takada, T.; Tohma, H. *Pure Appl. Chem.* **1996**, *68*, 627. (c) Dohi, T.; Ito, M.; Yamaoka, N.; Morimoto, K.; Fujioka, H.; Kita, Y. *Tetrahedron* **2009**, *65*, 10797.

(15) For the reactions using FPIFA and related compound, see: (a) Moriarty, R. M.; Prakash, I.; Penmasta, R. *J. Chem. Soc., Chem. Commun.* **1987**, 202. (b) Moriarty, R. M.; Penmasta, R.; Awasthi, A. K.; Prakash, I. *J. Org. Chem.* **1988**, *53*, 6124. (c) Harayama, Y.; Yoshida, M.; Kamimura, D.; Wada, Y.; Kita, Y. *Chem.—Eur. J.* **2006**, *12*, 4893. (d) Schaefer, S.; Wirth, T. *Angew. Chem., Int. Ed.* **2010**, *49*, 2786.

(16) (a) Tamura, Y.; Yakura, T.; Haruta, J.; Kita, Y. *J. Org. Chem.* **1987**, *52*, 3927. (b) Moriarty, R. M.; Prakash, O. *Org. React.* **2001**, *5*, 327. (c) Jean, A.; Cantat, J.; Berard, D.; Bouchu, D.; Canesi, S. *Org. Lett.* **2007**, *9*, 2553.

(FPIFA) and its CF₃-substituted derivative.^{15d} Gratefully, both oxidants worked well as more effective promoters for the cross-coupling (entries 6 and 7). A more pronounced advantage of the perfluorinated oxidants was no formation of the homodimer, which is in clear contrast to PIFA. Further improvement of the product yield was achieved after optimization of the conditions, and the use of commercial FPIFA gave a perfect result (entry 8). It should be noted that the perfluorinated oxidant, FPIFA, did not induce the homodimer formation even in the presence of a greater excess amount of the methoxybenzene **2a** (entry 9). None of other typical chemical SET oxidants¹⁰ [specifically, FeCl₃, MoCl₅, Mn(OAc)₃, Ce(NH₄)₂(NO₃)₆ (CAN), and 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ)] or anodic oxidations^{11a} produced the cross-coupled **3aa** as the major product.

It became evident that selection of the oxidant is very important for successful cross-coupling. The consumption of all of the naphthalene ether **1a** and the nearly perfect yield of **3aa** as the sole coupling product suggested the increased SET oxidation efficiency of the perfluorinated hypervalent iodine compounds over PIFA to quantitatively produce the naphthalene cation radical^{14a} of **1a** (Figure 1, left), which was then trapped by the neutral **2a** to




Figure 1. Selective C-T complex formation between FPIFA and **1a**.

give the mixed biaryl **3aa**. The generation of the cation radical of **2a** that led to the homodimer did not occur. Hence, the Lewis acidity of the iodine atom in the FPIFA/BF₃·Et₂O adduct should just match the formation of the C-T complex with the naphthalene ether **1a** that was involved before starting the SET oxidation process. Similarly, overoxidation of the formed biaryl **3** can be ruled out. The reaction of the substrates **1a** (0.200 mmol) and **2a** (0.40 mmol) or **2b** (R' = OMe, 0.40 mmol) in the presence of the prepared **3aa** (0.200 mmol) or **3ab** (0.200 mmol) under the standard coupling conditions resulted in the recovery of 0.364 mmol (for **3aa**) and >0.399 mmol (for **3ab**) of the biaryls after isolation (see Supporting Information), which revealed the revival of at least 82%

of the initially added biaryl **3aa** and over 99% of the biaryl **3ab** during the control experiments. This is also rationalized by considering the steric factor in the formed C-T complex (Figure 1, right). Apparently, the existence of the angular methoxybenzene ring in the products **3aa** and **3ab** forces the complexation with FPIFA to be unfavorable.

For these reasons, the extended series of the phenyl ethers **2b–i** could also be used as the coupling partners of the naphthalene ether **1a** with the optimized oxidant (Table 2). All coupling reactions proceeded to satisfacto-

Table 2. Examples of the Cross-Coupling Partners **2b–i**^a

entry	arene 2	mixed biaryl 3 , yield ^b
	 2b–f	 3ab–af
1	R = Me, R' = OMe; 2b	R = Me, R' = OMe; 3ab , >99%
2	R = Bn, R' = OBn; 2c	R = Bn, R' = OBn; 3ac , 87%
3	R = Me, R' = CH ₂ OAc; 2d	R = Me, R' = CH ₂ OAc; 3ad , 80%
4 ^c	R = Me, R' = C≡C–Bu; 2e	R = Me, R' = C≡C–Bu; 3ae , 60%
5 ^c	R = Me, R' = H; 2f	R = Me, R' = H; 3af , 65%
	 2g–i	 3ag–ai
6	R' = OBn; 2g	R' = OBn; 3ag , 58%, 3'ag , 41%
7	R' = OH; 2h	R' = OH; 3ah , nd, 3'ah , 68%
8	R' = NHTs; 2i	R' = NHTs; 3ai , 4%, 3'ai , 88%


^a Reactions were performed using naphthalene ether **1a** (1 equiv) and **2b–d,g–i** (2 equiv) under the optimized conditions in Table 1 unless otherwise noted. ^b Isolated yield of the pure products based on the substrate **1a** used. ^c 4 equiv of **2e** and **2f** was used. nd = not determined (less than 3% yield formation). Ts = *p*-tosyl.

rily give the corresponding oxygenated mixed biaryls **3** without formation of the homodimer in these cases. Several protecting groups (benzyl, acetyl, etc.) and functionalities (alkyne, phenol, amine, and others) were acceptable as the coupling partners **2**. The phenyl ethers **2g–i** produced two separable cross-coupling products with preference of the biaryls, **3ag**, **3'ah**, and **3'ai**, by the *para*-reactions of the

methoxy group or acidic substituents of **2h** and **2i** (entries 6–8). To our surprise, the oxidations of the phenol and aniline rings¹⁶ were relatively slower than the cross-coupling with FPIFA (entries 7 and 8).

Regarding the naphthalene ring, the use of similar protecting groups and functionalities was also possible (Scheme 1). In **1c**, the couplings preferentially occurred


Scheme 1. Reactions of Naphthalene Ethers **1b–d**^a



^a Reactions were performed using 2 equiv of **2b**.

at the less hindered position of the naphthalene ring. Curiously, the naphthalene ether **1d** selectively reacted at the *ortho* position of the methyl group to give only the biaryl **3'db**. On the other hand, interesting remote regioselectivity controls at the naphthalene rings were observed (Scheme 2). Particularly, the OTBDPS and halogen in **1f**

Scheme 2. Remote Regioselectivity Control in Naphthalenes^a



^a TBDPS = *tert*-butyldiphenylsilyl


and **1g** showed different regioselectivities to the OAc group in **1e**, which even overwhelmed the steric influence of the isobutyl group in substrate **1h**.

The oxygenated aryl naphthalenes **3** belong to a class of valuable synthetic intermediates that are the core structures of many useful compounds.¹⁷ Their further oxidized aryl naphthoquinones are also versatile for divergent access to naturally occurring types of compounds. Accordingly,

(17) For recent synthetic studies toward oxygenated aryl naphthalenes, see: (a) James, C. A.; Snieckus, V. *J. Org. Chem.* **2009**, *74*, 4080 and references therein. (b) Mori, K.; Ohmori, K.; Suzuki, K. *Angew. Chem., Int. Ed.* **2009**, *48*, 5633. (c) Martinez, A.; Barcia, J. C.; Estevez, A. M.; Fernandez, F.; Gonzalez, L.; Estevez, J. C.; Estevez, R. *J. Tetrahedron Lett.* **2007**, *48*, 2147. (d) James, C. A.; Snieckus, V. *Tetrahedron Lett.* **1997**, *38*, 8149.

elaboration of the obtained biaryl in our method toward the benz[*b*]phenanthridine structure¹⁸ was demonstrated (Scheme 3). The biaryl **3ad** was thus converted into the aryl naphthoquinone **4** by treatment of CAN^{13b} followed by the installation of an amino group, and its cyclization could afford the target tetracyclic compound **6**. Considering the necessity of sacrificial functional groups, such as halogens, in order to construct the phenanthrine structures via other synthetic routes, our metal-free C–H coupling method is capable of providing unique access toward molecules retaining such functional groups.

Scheme 3. Elaboration of the Biaryl Structure of **3ad**



In summary, we have reported for the first time the intermolecular C–H cross-coupling between the aromatic ethers **1** and **2**. The key point of the reaction success is the utilization of the excellent ability of hypervalent iodine reagents, especially the perfluorinated PIFAs, in the SET oxidation process. The remarkable high affinity of the reagents to the naphthalene ethers **1** over the phenyl ethers **2** as well as the formed mixed biaryl products **3** in the reactions can definitively exclude the undesired homocoupling and overoxidation of the biaryl products **3**, thus realizing the specific cross-coupling.

Acknowledgment. This work was partially supported by Grant-in-Aid for Scientific Research (A) from the Japan Society for the Promotion of Science (JSPS) and Ritsumeikan Global Innovation Research Organization (R-GIRO) project. T.D. is thankful for support from The Sumitomo Foundation and the Industrial Technology Research Grant Program from the New Energy and Industrial Technology Development Organization (NEDO) of Japan.

Supporting Information Available. Representative experimental procedures and detailed spectroscopic data of the products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(18) (a) Gore, M. P.; Gould, S. J.; Weller, D. D. *J. Org. Chem.* **1992**, *57*, 2774. (b) Echavarren, A. M.; Tamayo, N.; Cardenas, D. J. *J. Org. Chem.* **1994**, *59*, 6075. (c) Mohri, S.-I.; Stefinovic, M.; Snieckus, V. *J. Org. Chem.* **1997**, *62*, 7072. (d) De Frutos, O.; Atienza, C.; Echavarren, A. M. *Eur. J. Org. Chem.* **2001**, 163. (e) Akagi, Y.; Yamada, S.-I.; Etomi, N.; Kumamoto, T.; Nakanishi, W.; Ishikawa, T. *Tetrahedron Lett.* **2010**, *51*, 1338. (f) Shan, M.; Sharif, E. U.; O'Doherty, G. A. *Angew. Chem., Int. Ed.* **2010**, *49*, 9492.