

# Oxidative Chlorination of Various Iodoarenes to (Dichloroiodo)arenes with Chromium(VI) Oxide as the Oxidant†

*J. Chem. Research (S)*,  
1999, 64–65†

Pawet Kaźmierczak, Lech Skulski\* and Nicolas Obeid

Chair and Laboratory of Organic Chemistry, Faculty of Pharmacy, Medical Academy of Warsaw,  
1 Banacha Street, PL 02-097 Warsaw, Poland

Chromium(VI) oxide dissolved in a mixture of acetic acid with concentrated hydrochloric acid converts, at or near room temperature, iodoarenes to (dichloroiodo)arenes, in a very simple and efficient procedure.


Organic hypervalent iodine reagents play an important role in organic synthesis, as demonstrated by a number of recent reviews.<sup>1–6</sup> The more stable, solid (dichloroiodo)arenes, ArICl<sub>2</sub>, *e.g.* (dichloroiodo)benzene, are widely used as potent and fairly selective halogenating and/or oxidizing agents. They have a practical advantage over elemental chlorine, due to their easy and safe handling. Moreover, they may be readily converted to the respective iodosylarenes, (diacyloxyiodo)arenes, (difluoroiodo)arenes, and iodylarenes, used as important reagents in organic syntheses.<sup>1–6</sup> The most common procedure<sup>1–3,7,8</sup> for the synthesis of ArICl<sub>2</sub> is by reacting an iodoarene, ArI, with dichlorine in chloroform at 0 °C; the yields are generally excellent when this method is applicable;<sup>1–3,7</sup> the anhydrous conditions previously demanded<sup>8</sup> are not necessary.<sup>9</sup> Quite recently,<sup>10</sup> the repeated preparations of PhICl<sub>2</sub> on a 20 kg scale (94% yield, mp 113–117 °C with decomp.) have been conducted by the direct chlorination of PhI at –3 to +4 °C; the substitution of CHCl<sub>3</sub> and CCl<sub>4</sub> as solvents for the less toxic CH<sub>2</sub>Cl<sub>2</sub> is worth mentioning. PhICl<sub>2</sub> could be safely handled without major concerns for its thermal decomposition; under cooled conditions, it was possible to chlorinate 4-aminoacetophenone on a large scale, in 87% yield.

The inconvenient use of gaseous Cl<sub>2</sub> to afford ArICl<sub>2</sub> from ArI may be avoided as follows: (i) by using liquid SO<sub>2</sub>Cl<sub>2</sub> added to a solution of ArI either in wet diethyl ether,<sup>11</sup> or (better) in 98% acetic acid;<sup>12</sup> (ii) by the action of hydrochloric acid on either iodosylarenes or (diacyloxy-

iodo)arenes previously otherwise obtained from ArI;<sup>1–3,7</sup> (iii) by the action of a liquid mixture: cobalt(III) acetate–KCl–67% aq. CF<sub>3</sub>COOH on PhI;<sup>13</sup> (iv) by the use of hydrochloric acid oxidized *in situ* with sodium perborate in either acetonitrile or carbon tetrachloride containing a dissolved ArI.<sup>14</sup>

ArICl<sub>2</sub>, yellow crystalline compounds, are light- and heat-sensitive and often unstable to storage. They do not usually give satisfactory microanalyses, and mostly should be used immediately after preparation. Due to their thermal lability, their melting points are rather uncertain, depend upon the rate of heating, and may be those of the parent ArI.<sup>1,2</sup>

We report in this paper a new, very simple, and efficient laboratory method for the conversion of some iodoarenes to (dichloroiodo)arenes, including (dichloroiodo)benzene obtained from iodobenzene in 95% crude yield (reaction time: 1 h; 95% purity). Our method avoids gaseous Cl<sub>2</sub> and, in our opinion, is relatively safe and inexpensive. We used concentrated (36%) hydrochloric acid oxidized *in situ* with different amounts (see Table 1) of chromium(VI) oxide, a common and easily handled oxidant, dissolved in aqueous acetic acid, and acting upon a dissolved (or suspended) iodoarene. The reactions proceeded according to the equation:



**Table 1** Preparative details, crude yields, and melting points (uncorrected) of (dichloroiodo)arenes

Substrate ArI	CrO <sub>3</sub> /g	t/h	Product ArICl <sub>2</sub>	Yield (%)	Mp/°C (decomp.) (Lit., mp)
C <sub>6</sub> H <sub>5</sub> I	1.0	1.0	C <sub>6</sub> H <sub>5</sub> ICl <sub>2</sub>	95 <sup>a</sup>	111–112 (121), <sup>14</sup> (110–136; 115–127) <sup>18</sup>
4-MeOC <sub>6</sub> H <sub>4</sub> I	1.0	0.5	4-MeOC <sub>6</sub> H <sub>4</sub> ICl <sub>2</sub>	62	75–76 (75–78) <sup>14</sup>
3-MeC <sub>6</sub> H <sub>4</sub> I	1.0	1.0	3-MeC <sub>6</sub> H <sub>4</sub> ICl <sub>2</sub>	80	97–98 (100) <sup>14</sup>
4-MeC <sub>6</sub> H <sub>4</sub> I	1.0	1.0	4-MeC <sub>6</sub> H <sub>4</sub> ICl <sub>2</sub>	81	108–111 (97) <sup>14</sup>
4-FC <sub>6</sub> H <sub>4</sub> I	1.0	2.0	4-FC <sub>6</sub> H <sub>4</sub> ICl <sub>2</sub>	95	106–107 (105) <sup>14</sup>
2-ClC <sub>6</sub> H <sub>4</sub> I	1.0	2.0	2-ClC <sub>6</sub> H <sub>4</sub> ICl <sub>2</sub>	78	96–98 (96–97) <sup>18</sup>
3-ClC <sub>6</sub> H <sub>4</sub> I	1.0	2.0	3-ClC <sub>6</sub> H <sub>4</sub> ICl <sub>2</sub>	79	96–98 (99–100) <sup>18</sup>
4-ClC <sub>6</sub> H <sub>4</sub> I	1.0	1.0	4-ClC <sub>6</sub> H <sub>4</sub> ICl <sub>2</sub>	86	110–112 (113) <sup>14</sup>
2,4-Cl <sub>2</sub> C <sub>6</sub> H <sub>3</sub> I	3.0	5.0	2,4-Cl <sub>2</sub> C <sub>6</sub> H <sub>3</sub> ICl <sub>2</sub>	62	96–98 (107) <sup>18</sup>
2-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> I	3.0	3.0	2-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> ICl <sub>2</sub>	77	92–94 (96) <sup>18</sup>
3-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> I	3.0 <sup>b</sup>	2.0	3-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> ICl <sub>2</sub>	74	94–95 (90) <sup>14</sup>
3-HO <sub>2</sub> CC <sub>6</sub> H <sub>4</sub> I	3.0	3.0	3-HO <sub>2</sub> CC <sub>6</sub> H <sub>4</sub> ICl <sub>2</sub>	94	183–185 (185) <sup>14</sup>
3-MeO <sub>2</sub> CC <sub>6</sub> H <sub>4</sub> I	1.0	1.0	3-MeO <sub>2</sub> CC <sub>6</sub> H <sub>4</sub> ICl <sub>2</sub>	84	108–110
3-EtO <sub>2</sub> CC <sub>6</sub> H <sub>4</sub> I	1.0	1.0	3-EtO <sub>2</sub> CC <sub>6</sub> H <sub>4</sub> ICl <sub>2</sub>	83	98–100
5-Cl-2-MeOC <sub>6</sub> H <sub>3</sub> I	1.0	1.0	5-Cl-2-MeOC <sub>6</sub> H <sub>3</sub> ICl <sub>2</sub>	91	100–102 ( <i>ca.</i> 76) <sup>19</sup>
2-MeO-5-NO <sub>2</sub> C <sub>6</sub> H <sub>3</sub> I	3.0	3.0	2-MeO-5-NO <sub>2</sub> C <sub>6</sub> H <sub>3</sub> ICl <sub>2</sub>	74	105–106
4-MeO-3-NO <sub>2</sub> C <sub>6</sub> H <sub>3</sub> I	1.0	1.0	4-MeO-3-NO <sub>2</sub> C <sub>6</sub> H <sub>3</sub> ICl <sub>2</sub>	94	112–113

<sup>a</sup>When we enlarged the preparative scale twentyfold, the final yields for PhICl<sub>2</sub> were 92–94%. <sup>b</sup>On adding alternatively: (a) 1.0 g; (b) 2.0 g; (c) 3.0 g of CrO<sub>3</sub> (2.0 h, room temp.) to the reaction mixtures, the respective crude yields for 3-NO<sub>2</sub>C<sub>6</sub>H<sub>4</sub>ICl<sub>2</sub> were as follows: 48, 59 and 74%.

\*To receive any correspondence.

†This is a **Short Paper** as defined in the Instructions for Authors, Section 5.0 [see *J. Chem. Research (S)*, 1999, Issue 1]; there is therefore no corresponding material in *J. Chem. Research (M)*.

The crude yellow products precipitated by water were collected by filtration, washed well on the filter with ice-cold water to remove CrCl<sub>3</sub>, HCl, and AcOH, then with carbon tetrachloride to remove most of the unreacted ArI, and

air-dried in the dark. Iodometric titrations,<sup>7</sup> carried out immediately, showed that they were of 92–98% purity. Hence, they may be used as such in subsequent reactions. So far, we have failed to chlorinate oxidatively 4-iodonitrobenzene, 4-iodobenzonitrile and 2,4,6-trichloroidobenzene.

The same procedure is effective for the aromatic chlorination of activated arenes. For example, we chlorinated 4-nitroanisole, 4-nitroacetanilide and 4-methoxybenzoic acid to obtain the respective pure chlorinated products: 2-chloro-4-nitroanisole (64%), mp 94–95 °C (from ethanol), lit.,<sup>15</sup> mp 93–94 °C; 2-chloro-4-nitroacetanilide (58%), mp 138–139 °C (from ethanol), lit.,<sup>16</sup> mp 139–140 °C; 3,5-dichloro-4-methoxybenzoic acid (42%), mp 202–204 °C (from ethanol), lit.,<sup>17</sup> mp 202–202.5 °C. Their purity was supported by correct elemental analyses (C, H, Cl).

## Experimental

The starting iodoarenes, ArI, were either commercial or obtained by reported methods. They were purified prior to use, and analyzed (C, H, I). All the chlorination reactions were carried out **under a fume hood**. The toxic residues containing chromium salts were disposed of according to the local safety measures. Working on a small scale, we did not recover or remove chromium salts from the wastes. This should be taken into account, when preparations of larger quantities are required.

The melting points with decomposition (see Table 1) were uncorrected and were measured as follows: after an approximate mp had been taken in a capillary tube, a new sample was introduced about 10 °C below this point, and the temperature was raised at a rate of 10 °C min<sup>-1</sup>.

**General Procedure.**—Powdered CrO<sub>3</sub> [either (see Table 1): (a) 1.0 g, 10 mmol, 50% excess; (b) 2.0 g, 20 mmol, 200% excess; (c) 3.0 g, 30 mmol, 350% excess] was dissolved in water (ca. 3–5 ml), this was added to glacial acetic acid (15 ml), then an appropriate iodoarene (10 mmol) was either dissolved or suspended, keeping the temperature below 30 °C. Concentrated (36%) hydrochloric acid (14 ml, 160 mmol, 300% excess) was added with vigorous stirring at or near room temp. The reaction mixture turned from orange to deep green, and a precipitate appeared either immediately or within several minutes. The resulting slurry was stirred for 0.5–5.0 h (see Table 1), then the whole was poured into ice-water (100 ml). The yellow precipitate was collected by filtration, washed well with ice-cold water until the washings were colourless and neutral, then with CCl<sub>4</sub>. The crude product was air dried‡ in the dark. See Table 1 for the crude yields

and melting points (with decomposition). Note: easily oxidizable iodoarenes, e.g. 4-iodoanisole and iodotoluenes, should be added to the reaction mixture after the addition of the whole portion of conc. HCl. Otherwise, the final yields of the corresponding ArICl<sub>2</sub> are very low.

Received, 1st July 1998; Accepted, 9th October 1998

Paper E/8/05063E

## References

- 1 A. Varvoglis, *The Organic Chemistry of Polycordinated Iodine*, VCH, New York, Weinheim, Cambridge, 1992.
- 2 A. Varvoglis, *Hypervalent Iodine in Organic Synthesis*, Academic Press, San Diego, London, 1997.
- 3 P. J. Stang and V. V. Zhdankin, *Chem. Rev.*, 1996, **96**, 1123.
- 4 A. Varvoglis, *Tetrahedron*, 1997, **53**, 1179.
- 5 R. M. Moriarty and O. Prakash, *Adv. Heterocycl. Chem.*, 1998, **69**, 1.
- 6 T. Kitamura and Y. Fujiwara, *Org. Prep. Proced. Int.*, 1995, **3**, 261.
- 7 C. Willgerodt, *Die organischen Verbindungen mit mehrwertigem Jod*, Enke, Stuttgart, 1914.
- 8 H. J. Lucas and E. R. Kennedy, *Org. Synth., Coll. Vol. III*, Wiley, New York, 1955, p. 482.
- 9 R. T. Taylor and T. A. Stevenson, *Tetrahedron Lett.*, 1988, **29**, 2033, and references therein.
- 10 A. Zanka, H. Takeuchi and A. Kubota, *Org. Process Res. Dev.*, 1998, **2**, 270.
- 11 A. Töhl, *Ber. Dtsch. Chem. Ges.*, 1893, **26**, 2949.
- 12 B. Ya. Karele and O. Ya. Neiland, *Latv. PSR Zinat. Akad. Vestis, Kim. Ser.*, 1970, 587; *Chem. Abstr.* 1971, **74**, 42033. They failed to chlorinate three iodonitrobenzenes, 4-iodobenzoic acid, and 2-iodoacetophenone; for ten appropriate iodobenzenes the crude yields for ArICl<sub>2</sub> were 73–96%.
- 13 D. I. Makhon'kov, A. V. Cheprakov and I. P. Beletskaya, *Zh. Org. Khim.*, 1986, **22**, 681; *Chem. Abstr.*, 1987, **106**, 155951. Only iodobenzene was chlorinated there to give PhICl<sub>2</sub> (75%; 94% purity).
- 14 D. Koyunçu, A. McKillop and L. McLaren, *J. Chem. Res. (S)*, 1990, 21. They failed to chlorinate 2-iodonitrobenzene; for fifteen appropriate iodobenzenes the crude yields for ArICl<sub>2</sub> were 60–98%.
- 15 H. Fischli, *Ber. Dtsch. Chem. Ges.*, 1878, **11**, 1461.
- 16 A. Mannino and L. di Donato, *Gazz. Chim. Ital.*, 1908, **38**, **II**, 20.
- 17 T. H. Durrans, *J. Chem. Soc.*, 1923, **123**, 1424.
- 18 F. M. Beringer and E. M. Gindler, *Iodine Abstr. Rev.*, 1956, **3**, 1.
- 19 C. K. Ingold, E. W. Smith and C. C. N. Vass, *J. Chem. Soc.*, 1927, 1245.

‡By drying the crude ArICl<sub>2</sub> in a vacuum desiccator, the chlorine percentage was lowered: ArICl<sub>2</sub> → ArI + Cl<sub>2</sub>.