

+ H<sub>2</sub>O

+ 2 H<sub>2</sub>O

R<sup>1</sup>

26 examples (up to 91% yield)

OH

8 examples

(up to 89% yield)

# Nickel-Catalyzed Guerbet Type Reaction: C-Alkylation of Secondary Alcohols *via* Double (de)Hydrogenation

Reshma Babu, Murugan Subaramanian, Siba P. Midya, and Ekambaram Balaraman\*



R, R<sup>1</sup> = Aromatic, aliphatic,

acyclic, etc.

2 Ar OH

OH

**ABSTRACT:** Acceptorless double dehydrogenative cross-coupling of secondary and primary alcohols under nickel catalysis is reported. This Guerbet type reaction provides an atom- and a stepeconomical method for the C-alkylation of secondary alcohols under mild, benign conditions. A broad range of substrates including aromatic, cyclic, acyclic, and aliphatic alcohols was well tolerated. Interestingly, the C-alkylation of cholesterol derivatives and the double C-alkylation of cyclopentanol with various alcohols were also demonstrated.

he construction of functionalized diverse molecules from simple abundant chemicals by C-C bond-forming reactions is one of the fundamental reactions in organic synthesis.<sup>1</sup> Traditionally, the C-C bond formation of carbonyl compounds (*i.e.*,  $C(\alpha)$ -alkylation) has been achieved by nucleophilic substitution reactions with alkyl halides or other activated derivatives.<sup>2</sup> Thus, classical methods require highly reactive, toxic, and expensive reagents, which generate stoichiometric amounts of hazardous waste.<sup>3</sup> In this regard, the development of a green and sustainable method by utilizing abundantly available feedstock alcohols for selective C-C bond formation is highly desirable.<sup>4</sup> Recently, transition metalcatalyzed alcohol activation through a borrowing hydrogen strategy (BH) has been seen to play a significant role in the direct C-C and C-N bond forming reactions, which often overcome limitations of classical methods and produce water as the byproduct.<sup>5</sup> The BH reaction relies on catalytic dehydrogenation of alcohols to carbonyl compounds, condensation reaction with a suitable C-, or N-nucleophile to form an  $\alpha_{\beta}$ -unsaturated compound and an imine intermediate, respectively, and finally, selective hydrogenation of the  $\alpha$ ,  $\beta$ unsaturated or imine intermediate leads to the saturated product by utilizing the hydrogen gas liberated in the initial dehydrogenation step.<sup>6</sup> Thus, the net reaction eliminates water as a sole byproduct.

Of late, C–C bond formation *via* the acceptorless dehydrogenative cross-coupling reaction of different alcohols is very interesting. This Guerbet type reaction provides an atom- and a step-economical method for the C-alkylation of secondary alcohols under mild, benign conditions (Scheme 1). Notably, the  $\beta$ -alkylation of secondary alcohols with primary alcohols has been widely investigated under precious metal (Ru, Rh, Ir, and Pd) catalysis.<sup>7–10</sup> In addition to this, seminal examples based on heterogeneous catalytic systems have been established for the  $\beta$ -alkylation of secondary alcohols.<sup>11</sup> In



cat. NiBra

cat.TMEDA

кон

\* Broad Subsrate Scope

\* ß-Alkylation of Cholesterol \* Water as the sole by-product

Double C-alkylation of cyclopentanol



recent times, judiciously designed earth-abundant 3d-transition metal catalysts for sustainable chemical synthesis *via* the C–C cross-coupling of alcohols has gained much interest due to economic and environmental benefits.<sup>12</sup>

In 2017, a seminal work on the PNP-cobalt pincer complex catalyzed C–C cross-coupling of secondary alcohols using primary alcohols has been reported by Kempe and co-workers.<sup>13</sup> Of late, the research group of Yu has reported the manganese-catalyzed cross-coupling of secondary alcohols with

Received:March 9, 2021Published:April 21, 2021



primary alcohols to access  $\beta$ -alkylated secondary alcohol products under dehydrogenative conditions.<sup>14</sup> Recently, notable progress has been made in the development of homogeneous catalysts used for the Guerbet-type reactions.<sup>15</sup> Notably, the first enantioselective Guerbet reaction at room temperature has been achieved using a commercially available chiral ruthenium complex developed by Zhao and coworkers.<sup>15</sup>

Lang and co-workers have studied the homogeneous Ni(II) N-heterocycle thiolate cluster mediated coupling of secondary and primary alcohols to yield  $\alpha,\beta$ -unsaturated ketones,  $\alpha$ alkylated ketones, and  $\beta$ -alkylated secondary alcohols by tuning the reaction conditions.<sup>16</sup> However, this hexanuclear nickel thiolate cluster catalyzed cross-coupling reaction of alcohols requires high catalyst loading, multistep synthesis of [Ni- $(dmpymt)_2_6$  cluster, and limited substrate scope. Thus, the development of a simple, elegant catalytic system for C-C bond formation via the Guerbet type strategy is highly desirable and demanding in contemporary science. Herein, we have reported the selective  $\beta$ -alkylation of secondary alcohols with primary alcohols, using a commercially available, inexpensive NiBr<sub>2</sub>/TMEDA as an efficient catalytic system. Interestingly, more challenging substrate conversions were achieved under benign conditions using primary alcohols as potential alkylating agents with various secondary alcohols. A broad range of substrates including aromatic, cyclic, acyclic, and aliphatic alcohols was well tolerated. Gratifyingly, the double C-alkylation of cyclopentanol with various alcohols was also demonstrated.

For the nickel-catalyzed dehydrogenative C-alkylation of secondary alcohols with primary alcohols, we have chosen 1phenyl ethanol (1a) and benzyl alcohol (2a) as the benchmark substrates. Initially, a model reaction was performed using NiBr<sub>2</sub> (5 mol %) as nickel source,  $N_1, N_1, N'_2$ -tetramethylethylenediamine (TMEDA, 5 mol %) as a ligand, and KO<sup>t</sup>Bu (1 equiv) as a base in toluene solvent at 130 °C for 18 h (Table 1, entry1). To our delight, the  $\beta$ -alkylated product 3a was obtained in 72% of isolated yield. It was found that n-octane could be the optimal solvent achieving the desired product 3a in excellent yield with selectivity (Table 1, entry 2). Under identical conditions, the effect of various bases such as KOH, LiO<sup>t</sup>Bu, NaO<sup>t</sup>Bu, NaOH, LiOH $\cdot$ H<sub>2</sub>O, and K<sub>2</sub>CO<sub>3</sub> was examined (Table 1, entries 3-8). Notably, KOH was found to be efficient and gave the desired product 3a in excellent yield (Table 1, entry 3). The variation of solvent indicated that the reaction proceeded efficiently in *n*-octane (Table 1, entries 1,3 and 9–11).

Other nickel catalysts were not effective for the present catalytic transformation and yielded the product 3a in moderate yields (Table1, entries 3 and 12–15). Also, attempts to decrease the mole percentage of base ended with low yields of the desired product 3a. Next, we have examined the effect of various commercially available nitrogen-based bidentate ligands (L1–L6) (Table 1, entries 19–23). Gratifyingly, the ligand L1 was found to be more effective for this reaction (Table1, entry 3).

A control experiment in the absence of a base, ligand, as well as nickel catalyst led to no product formation (Table 1, entries 16–18). These results confirmed that the combination of the base, ligand, and nickel catalyst is necessary to access the desired  $\beta$ -alkylated alcohols. By lowering the reaction temperature from 130 to 100 °C, the yield of the product **3a** was reduced to 70% (Table 1, entries 3 and 24–25).

#### Table 1. Optimization Studies<sup>c</sup>

$\bigcirc$	OH + ( 1a	он 2а	[Ni]/Ligand Base, Solvent Temp. (°C), 18 I	• ()	OH J J J J J J	+ H <sub>2</sub> O
Entry	[Ni] source	Base	Solvent	Ligand	Temp. (°C)	Yield(%) <sup>a</sup>
1	NiBr <sub>2</sub>	KO <sup>t</sup> Bu	Toluene	L1	130	72
2	NiBr <sub>2</sub>	KO <sup>t</sup> Bu	n-Octane	L1	130	75
3	NiBr <sub>2</sub>	КОН	n-Octane	L1	130	91(83) <sup>b</sup>
4	NiBr <sub>2</sub>	LiO <sup>t</sup> Bu	n-Octane	L1	130	19
5	NiBr <sub>2</sub>	NaO <sup>t</sup> Bu	n-Octane	L1	130	47
6	NiBr <sub>2</sub>	NaOH	n-Octane	L1	130	65
7	NiBr <sub>2</sub>	LiOH.H <sub>2</sub> O	n-Octane	L1	130	NR
8	NiBr <sub>2</sub>	K <sub>2</sub> CO <sub>3</sub>	n-Octane	L1	130	25
9	NiBr <sub>2</sub>	кон	<i>m</i> -Xylene	L1	130	44
10	NiBr <sub>2</sub>	КОН	1,4-dioxane	L1	130	58
11	NiBr <sub>2</sub>	KOH	DMF	L1	130	10
12	NiCl <sub>2</sub>	KOH	n-Octane	L1	130	68
13	Ni(acac) <sub>2</sub>	кон	n-Octane	L1	130	69
14	Ni(OTf) <sub>2</sub>	кон	n-Octane	L1	130	NR
15	NiCl <sub>2</sub> (PMe <sub>3</sub> ) <sub>2</sub>	КОН	n-Octane	L1	130	68
16		KOH	n-Octane	L1	130	NR
17	NiBr <sub>2</sub>		n-Octane	L1	130	NR
18	NiBr <sub>2</sub>	КОН	n-Octane		130	NR
19	NiBr <sub>2</sub>	КОН	n-Octane	L2	130	75
20	NiBr <sub>2</sub>	кон	n-Octane	L3	130	72
21	NiBr <sub>2</sub>	KOH	n-Octane	L4	130	55
22	NiBr <sub>2</sub>	КОН	n-Octane	L5	130	78
23	NiBr <sub>2</sub>	KOH	n-Octane	L6	130	NR
24	NiBr <sub>2</sub>	KOH	n-Octane	L1	110	81
25	NiBr <sub>2</sub>	КОН	n-Octane	L1	100	,
Me <sub>\N</sub> He	Me N <sub>Me</sub> ( L <sub>1</sub>			N=		
		I=/		,	Me L <sub>6</sub>	Me

<sup>*a*</sup>Yield calculated by <sup>1</sup>H NMR using dibromomethane as an internal standard. <sup>*b*</sup>Isolated yield. <sup>*c*</sup>Reaction conditions: **1a** (0.5 mmol), **2a** (0.5 mmol), [Ni] source (5 mol %), Ligand ( $L_1-L_6$ ) (5 mol %), base (0.5 mmol), and 1 mL of solvent heated in an oil-bath for 18 h.

With the optimized reaction conditions in hand, we next explored the applicability of the present nickel catalysis for the  $\beta$ -alkylation of secondary alcohols using diverse primary alcohols as an alkylating agent. At first, the scope of primary alcohols with secondary alcohols under standard reaction conditions was investigated (Table 2). As shown in Table 2, the electron-donating substituents such as methyl, ethyl, isopropyl, t-butyl, methoxy, and thioether groups containing benzyl alcohols afforded good to excellent yields of the  $\beta$ alkylated product (products 3b-3l in 74-87% isolated yields). Notably, halide substituted benzyl alcohols resulted in the products 3m-3o in 66-79% of isolated yields. In particular, a fused aromatic compound, 1-naphthyl methanol, produced the corresponding product 3p in 71% yield under the optimized reaction conditions. Moreover, aliphatic alcohols such as 1hexanol and 1-pentanol resulted in the corresponding products 3q in 75%, and 3r in 72% of isolated yields, respectively.

Subsequently, we have explored the scope of secondary alcohol for the selective  $\beta$ -alkylation reaction using 4-methoxy benzyl alcohol (**2e**) as the benchmark substrate (Table 3). Pleasingly, the dehydrogenative C–C coupling reactions proceeded smoothly for a variety of secondary alcohols with electron-neutral, electron-deficient, and electron-rich substituents led to the corresponding  $\beta$ -alkylated product in good to excellent yields. Under the optimized reaction conditions, the





<sup>*a*</sup>Reaction conditions: 1 (0.5 mmol, 2 (0.5 mmol), NiBr<sub>2</sub> (5 mol %), TMEDA (5 mol %), and KOH (0.5 mmol) in 1 mL of *n*-octane heated at 130 °C (oil-bath temperature) for 18 h. <sup>*b*</sup>Isolated yields.



<sup>*a*</sup>Reaction conditions: **1** (0.5 mmol, **2e** (0.5 mmol), NiBr<sub>2</sub> (5 mol %), TMEDA (5 mol %), and KO'Bu (0.5 mmol) in 1 mL of *n*-octane heated at 130 °C (oil-bath temperature) for 18 h. <sup>*b*</sup>Isolated yields.

electron-donating substituents such as -Me, and -OMe groups on secondary alcohols afforded the  $\beta$ -alkylated products in good yields (products **4b**-**4c**, 79-84%). In particular, the halide (-Cl, and -Br) group containing secondary alcohols successfully underwent  $\beta$ -alkylation reaction and gave the products **4d** to **4f** in 76-75% of isolated yields under the optimized reaction conditions.

Interestingly, fused-aromatic secondary alcohol such as 1-(2-naphthyl)ethanol was efficiently  $C(\beta)$ -alkylated with 4-methyl benzyl alcohol under the present nickel-catalyzed conditions and produced the product **4g** in 69% yield. Unfortunately, secondary alcohols bearing a strong electron-withdrawing group (*e.g.*,  $-NO_2$ ) failed to yield the desired product under our reaction conditions. Furthermore, 1-cyclopropylethanol

resulted in the expected product 4i in 71% of isolated yield under the present nickel-catalyzed conditions.

Interestingly, we extended our present catalytic system for tandem double alkylation of cyclopentanol **5a** with primary alcohols to access di( $\beta$ -alkylated) secondary alcohol **6** products (Table 4). Thus, the benzyl alcohol bearing electron-donating

# Table 4. Substrate Scope of Double C-Alkylation of Cyclopentanol $^{a,b}$



"Reaction conditions: **5a** (0.5 mmol, **2** (1 mmol), NiBr<sub>2</sub> (5 mol %), TMEDA (5 mol %), and KOH (0.5 mmol) in 1 mL of *n*-octane heated at 130 °C (oil-bath temperature) for 18 h. <sup>*b*</sup>Isolated yields.

group, for example, methyl group at the *ortho*, *meta*, and *para* positions, and halide substituted benzyl alcohols (-Cl, and -Br) underwent  $\beta$ -alkylation reaction and gave the di( $\beta$ -alkylated) products **6b**-**6g** in 79–89% of isolated yields under the optimized reaction conditions. Notably, other cyclic secondary alcohols such as cyclohexanol and cycloheptanol did not afford the desired products under standard reaction conditions.

The present nickel-catalyzed Guerbet type  $C(\beta)$ -alkylation strategy was successfully applied for the derivatization of cholesterol derivatives. As a result, the reaction of cholesterol derivative 7a with 2.5 equiv of benzyl alcohol under standard reaction conditions yielded the corresponding  $C(\beta)$ -alkylated product 8a in 78% isolated yield, wherein hydrogenation of C==C bond was observed (Scheme 2). Similarly, with an excess of benzyl alcohol compound 7b yielded bis- $\beta$ -alkylated product 8b in 87% yield with the hydrogenation of olefinic and carbonyl moieties of 7b.

To gain preliminary insights into the present nickelcatalyzed  $\beta$ -alkylation of secondary alcohols with primary alcohols, various control experiments were performed. Evaluation of H<sub>2</sub> gas from the benchmark reaction was detected by gas chromatography (GC) analysis (Scheme 3a). Next, two independent reactions of acetophenone 1a' with benzyl alcohol 2a and 1-phenyl ethanol 1a with benzaldehyde 2a' gave a mixture of C-C bonded products such as 1,3diphenylpropan-1-one (3a', major), (E)-chalcone (3a", minor), and trace amounts of 1,3-diphenylpropan-1-ol (3a) (Scheme 3b). Then, the reaction of (E)-chalcone 3a" in the presence of 1-phenyl ethanol 1a or benzyl alcohol 2a resulted in 1,3-diphenylpropan-1-one 3a' as a major product and 1,3diphenylpropan-1-ol 3a as a minor product (Scheme 3c). From



# Scheme 2. $\beta$ -Alkylation of Cholesterol Derivatives

#### Scheme 3. Control Experiments



these results, it was observed that the reaction proceeds via the initial acceptorless dehydrogenation of alcohols followed by condensation and subsequent transfer hydrogenation of intermediate chalcone 3a". Moreover, the deuterium labeling experiments were performed using deuterated benzyl alcohol 2a-D and deuterated 1-phenyl ethanol 1a-D under the present nickel-catalyzed conditions and yielded the deuteriumincorporated  $\beta$ -alkylated secondary alcohols 3a-D and 3a'-D. The presence of deuterium atom at  $\alpha$ ,  $\beta$ , and  $\gamma$  positions of **3a**-D and 3a'-D has further confirmed the dehydrogenation of an alcohol followed by (transfer) hydrogenation of intermediate 3a" (Scheme 3d). Notably, in the presence of radical scavengers, such as BHT and TEMPO, the product formation was not affected, and indeed, the product 3a was obtained in excellent yield (Scheme 3e). This result illustrates that the present nickel catalysis does not follow the SET (single electron transfer) mechanism.

On the basis of the control experiments and previous reports,  $^{17-20}$  a plausible mechanism for the nickel-catalyzed

cross dehydrogenative coupling of secondary alcohols with primary alcohols is proposed in Scheme 4. Initially, both the





primary and secondary alcohols undergo dehydrogenation under nickel catalysis leads to the corresponding carbonyls derivatives (ketone (1') and aldehyde (2'), respectively) with the generation of hydrogen gas. Subsequently, a base-mediated cross-aldol type condensation of 1' and 2' generates an intermediate  $\alpha,\beta$ -unsaturated ketone (3"). Finally, in the presence of a nickel catalyst transfer hydrogenation of  $\alpha,\beta$ unsaturated ketone (by the liberated hydrogen from the foremost dehydrogenation step) to access the desired  $\beta$ alkylated secondary alcohol product (3a). Overall, the process is redox neutral and produces water as a sole byproduct.

In summary, we have reported an efficient  $\beta$ -alkylation of secondary alcohols with primary alcohols under benign conditions. This elegant approach exhibits a broad substrate scope providing the C–C coupled products of secondary alcohols and the double alkylation of cyclopentanol using aryl, fused aryl, and aliphatic primary alcohols. This Guerbet type protocol extended for the functionalization of cholesterol derivatives. Preliminary mechanistic studies and isotopic labeling experiments suggest that the C–C coupling reaction proceeds *via* the borrowing hydrogenation with the formation of water as a byproduct.

# ASSOCIATED CONTENT

#### **1** Supporting Information

The Supporting Information is available free of charge at https://pubs.acs.org/doi/10.1021/acs.orglett.1c00782.

- Experimental section; <sup>1</sup>H and <sup>13</sup>C NMR (PDF)
- FAIR data, including the primary NMR FID files, for compounds 3a-3r; 3a'; 3a''; 3a-D; 3b-D; 4a-4i; 6a-6h; 8a; 8b (ZIP)

### AUTHOR INFORMATION

#### **Corresponding Author**

Ekambaram Balaraman – Department of Chemistry, Indian Institute of Science Education and Research (IISER) Tirupati, Tirupati 517507, India; orcid.org/0000-0001-9248-2809; Email: eb.raman@iisertirupati.ac.in

# Authors

Reshma Babu – Department of Chemistry, Indian Institute of Science Education and Research (IISER) Tirupati, Tirupati 517507, India Murugan Subaramanian – Department of Chemistry, Indian Institute of Science Education and Research (IISER) Tirupati, Tirupati 517507, India

Siba P. Midya – Department of Chemistry, Indian Institute of Science Education and Research (IISER) Tirupati, Tirupati 517507, India

Complete contact information is available at: https://pubs.acs.org/10.1021/acs.orglett.1c00782

#### Notes

The authors declare no competing financial interest.

# ACKNOWLEDGMENTS

This research work was supported by IISER Tirupati, and SERB, India (grant no. CRG/2018/002480/OC). E.B. acknowledges SwarnaJayanti Fellowship grant (DST/SJF/CSA-04/2019-2020 and SERB/F/5892/2020-2021). R.B. thanks SERB-PMRF for the research fellowship. M.S. thanks the UGC for the research fellowship.

### REFERENCES

(1) Casiraghi, G.; Battistini, L.; Curti, C.; Rassu, G.; Zanardi, F. The vinylogous aldol and related addition reactions: Ten years of progress. *Chem. Rev.* **2011**, *111*, 3076–3154. (b) Kawahara, R.; Fujita, K.; Yamaguchi, R. Dehydrogenative oxidation of alcohols in aqueous media using water-soluble and reusable Cp\*Ir catalysts bearing a functional bipyridine ligand. J. Am. Chem. Soc. **2012**, *134*, 3643–3646. (c) Kawahara, R.; Fujita, K.; Yamaguchi, R. Cooperative catalysis by Ir complexes with a bipyridonate ligand: Versatile dehydrogenative oxidation of alcohols and reversible dehydrogenation-hydrogenation between 2-propanol and acetone. *Angew. Chem., Int. Ed.* **2012**, *51*, 12790–12794. (d) Rygus, J. P. G.; Crudden, C. M. Enantiospecific and iterative Suzuki-Miyaura cross-couplings. J. Am. Chem. Soc. **2017**, *139*, 18124–18347.

(2) (a) Yi, H.; Zhang, G. T.; Wang, H. M.; Huang, Z. Y.; Wang, J.; Singh, A. K.; Lei, A. W. T. Recent advances in radical C-H activation/ radical cross-coupling. *Chem. Rev.* **2017**, *117*, 9016–9085. (b) Akhtar, W. M.; Armstrong, R. J.; Frost, J. R.; Stevenson, N. G.; Donohoe, T. J. Stereoselective synthesis of cyclohexanes *via* an Ir catalyzed (5 + 1) annulation strategy. *J. Am. Chem. Soc.* **2018**, *140*, 11916–11920.

(3) (a) Frisch, A. C.; Beller, M. Catalysts for cross-coupling reactions with non-activated alkyl halides. *Angew. Chem., Int. Ed.* **2005**, *44*, 674–688. (b) Zhang, Y. F.; Shi, Z. J. Upgrading cross-coupling reactions for biaryl synthesis. *Acc. Chem. Res.* **2019**, *52*, 161–169.

(4) (a) Hamid, M. H. S. A.; Slatford, P. A.; Williams, J. M. J. Borrowing hydrogen in the activation of alcohols. *Adv. Synth. Catal.* **2007**, 349, 1555–1575. (b) Gunanathan, C.; Milstein, D. Applications of acceptorlesss dehydrogenation and related transformations in chemical synthesis. *Science* **2013**, 341, 1229712. (c) Irrgang, T.; Kempe, R. 3d-Metal catalyzed N- and C-alkylation reactions *via* borrowing hydrogen or hydrogen autotransfer. *Chem. Rev.* **2019**, *119*, 2524–2549.

(5) (a) Iuchi, Y.; Obora, Y.; Ishii, Y. Ir-catalyzed $\alpha$ -alkylation of acetates with primary alcohols and diols. *J. Am. Chem. Soc.* **2010**, *132*, 2536–2537. (b) Cai, Y.; Li, F.; Li, Y. Q.; Zhang, W. B.; Liu, F. H.; Shi, S. L. Base metal-catalyzed alcohol C-C couplings under hydrogen transfer conditions. *Tetrahedron Lett.* **2018**, *59*, 1073–1079. (c) Li, Y.; Li, H. Q.; Junge, H.; Beller, M. Selective ruthenium-catalyzed methylation of 2-arylethanols using methanol as C1 feedstock. *Chem. Commun.* **2014**, *50*, 14991–14994.

(6) (a) Kwon, M. S.; Kim, N.; Seo, S. H.; Park, I. S.; Cheedrala, R. K.; Park, J. Recyclable palladium catalyst for highly selective  $\alpha$ -alkylation of ketones with alcohols. *Angew. Chem., Int. Ed.* **2005**, *44*, 6913–6915. (b) Liu, X.; Ding, R. S.; He, L.; Liu, Y. M.; Cao, Y.; He, H. Y.; Fan, K. N. C-C cross coupling of primary and secondary

benzylic alcohols using supported gold based bimetallic catalysts. *ChemSusChem* **2013**, *6*, 604–608.

(7) (a) Ruiz-Botella, S.; Peris, E. Unveiling the importance of  $\pi$ stacking in borrowing hydrogen processes catalyzed by Ir complexes with pyrene tags. Chem. - Eur. J. 2015, 21, 15263-15271. (b) Jiménez, M. V.; Fernández-Tornos, J.; Modrego, F. J.; Pérez-Torrente, J. J.; Oro, L. A. Oxidation and  $\beta$ -alkylation of alcohols catalyzed by Ir(I) complexes with functionalised N-Heterocyclic carbene ligands. Chem. - Eur. J. 2015, 21, 17877-17889. (c) Xu, C.; Goh, L. Y.; Pullarkat, S. A. Efficient Ir-thioether-dithiolate catalyst for  $\beta$ -alkylation of alcohols and selective imine formation via N-alkylation reactions. Organometallics 2011, 30, 6499-6502. (d) Segarra, C.; Mas-Marzá, E.; Mata, J. A.; Peris, E. Shvo's catalyst and [IrCp\*Cl<sub>2</sub>(amidine)] effectively catalyze the formation of tertiary amines from the reaction of primary alcohols and ammonium salts. Adv. Synth. Catal. 2011, 353, 2078-2084. (e) Gong, X.; Zhang, H.; Li, X. Ir phosphine abnormal Nheterocyclic carbene complexes in catalytic hydrogen transfer reactions. Tetrahedron Lett. 2011, 52, 5596-5600. (f) Gnanamgari, D.; Sauer, E. L. O.; Schley, N. D.; Butler, C.; Incarvito, C. D.; Crabtree, R. H. Ir and Ru complexes with chelating N-heterocyclic carbenes: efficient catalysts for transfer hydrogenation, and Nalkylation of amines. Organometallics 2009, 28, 321-325. (g) da Costa, A. P.; Sanaú, M.; Peris, E.; Royo, B. Easy preparation of Cp\*functionalized N-heterocyclic carbenes and their coordination to Rh and Ir. Dalton Trans. 2009, 35, 6960-6966. (h) da Costa, A. P.; Viciano, M.; Sanaú, M.; Merino, S.; Tejeda, J.; Peris, E.; Royo, B. First Cp\*-functionalized N-heterocyclic carbene and its coordination to Ir: study of the catalytic properties. Organometallics 2008, 27, 1305-1309. (i) Gnanamgari, D.; Leung, C. H.; Schley, N. D.; Hilton, S. T.; Crabtree, R. H. Alcohol cross-coupling reactions catalyzed by Ru and Ir terpyridine complexes. Org. Biomol. Chem. 2008, 6, 4442-4445. (j) Fujita, K.; Asai, C.; Yamaguchi, T.; Hanasaka, F.; Yamaguchi, R. Direct  $\beta$ -alkylation of secondary alcohols with primary alcohols catalyzed by a Cp\*Ir complex. Org. Lett. 2005, 7, 4017-4019. (k) Genc, S.; Arslan, B.; Gulcemal, S.; Gunnaz, S.; Cetinkaya, B.; Gulcemal, D. Ir(I)-catalyzed C-C and C-N bond formation reactions via the borrowing hydrogen strategy. J. Org. Chem. 2019, 84, 6286-6297.

(8) (a) Zhang, C.; Zhao, J. P.; Hu, B.; Shi, J.; Chen, D. Ru catalyzed  $\beta$ -alkylation of secondary alcohols and  $\alpha$ -alkylation of ketones via borrowing hydrogen: Dramatic influence of the pendant N-Heterocycle. Organometallics 2019, 38, 654-664. (b) Roy, B. C.; Debnath, S.; Chakrabarti, K.; Paul, B.; Maji, M.; Kundu, S. Ortho -amino group functionalized 2,2'-bipyridine based Ru(II) complex catalyzed alkylation of secondary alcohols, nitriles and amines using alcohols. Org. Chem. Front. 2018, 5, 1008-1018. (c) Shee, S.; Paul, B.; Panja, D.; Roy, B. C.; Chakrabarti, K.; Ganguli, K.; Das, A.; Das, G. K.; Kundu, S. Tandem cross coupling reaction of alcohols for sustainable synthesis of  $\beta$ -alkylated secondary alcohols and flavan derivatives. Adv. Synth. Catal. 2017, 359, 3888-3893. (d) Roy, B. C.; Chakrabarti, K.; Shee, S.; Paul, S.; Kundu, S. Bifunctional Ru<sup>II</sup>-complex catalyzed tandem C-C bond formation: Efficient and atom economical strategy for the utilization of alcohols as alkylating agents. Chem. - Eur. J. 2016, 22, 18147-18155. (e) Chakrabarti, K.; Paul, B.; Maji, M.; Roy, B. C.; Shee, S.; Kundu, S. Bifunctional Ru(II) complex catalysed C-C bond formation: An eco-friendly hydrogen borrowing strategy. Org. Biomol. Chem. 2016, 14, 10988-10997. (f) Wang, Q.; Wu, K.; Yu, Z. Ru(III)catalyzed  $\beta$ -alkylation of secondary alcohols with primary alcohols. Organometallics 2016, 35, 1251-1256. (g) Cho, C. S.; Kim, B. T.; Kim, H.-S.; Kim, T. J.; Shim, S. C. Ru-catalyzed one pot  $\beta$ -alkylation of secondary alcohols with primary alcohols. Organometallics 2003, 22, 3608-3610. (h) Prades, A.; Viciano, M.; Sanaú, M.; Peris, E. Preparation of a series of "Ru(p-cymene)" complexes with different N-Heterocyclic carbene ligands for the catalytic  $\beta$ -alkylation of secondary alcohols and dimerization of phenylacetylene. Organometallics 2008, 27, 4254-4259.

(9) Satyanarayana, P.; Reddy, G. M.; Maheswaran, H.; Kantam, M. L. Tris(acetylacetonato)rhodium(III)-catalyzed  $\alpha$ -alkylation of ke-

tones,  $\beta$ -alkylation of secondary alcohols and alkylation of amines with primary alcohols. *Adv. Synth. Catal.* **2013**, 355, 1859–1867.

(10) Kose, O.; Saito, S. Cross-coupling reaction of alcohols for C-C bond formation using pincer type NHC/palladium catalysts. *Org. Biomol. Chem.* **2010**, *8*, 896–900.

(11) (a) Shimizu, K.; Sato, R.; Satsuma, A. Direct C-C cross coupling of secondary and primary alcohols catalyzed by a gammaalumina-supported silver subnanocluster. *Angew. Chem., Int. Ed.* **2009**, *48*, 3982–3986. (b) Liu, X.; Ding, R. S.; He, L.; Liu, Y. M.; Cao, Y.; He, H. Y.; Fan, K. N. C-C Cross coupling of primary and secondary benzylic alcohols using supported gold-based bimettalic catalysts. *ChemSusChem* **2013**, *6*, 604–608. (c) Zhou, M.; Zhang, L.; Miller, J. T.; Yang, X.; Liu, X.; Wang, A.; Zhang, T. Hydrogen auto-transfer under aerobic oxidative conditions: Efficient synthesis of saturated ketones by aerobic C-C cross coupling of primary and secondary alcohols catalyzed by a Au<sub>6</sub>Pd/resin catalyst. *Chin. J. Catal.* **2016**, *37*, 1764–1770. (d) Xu, J.; Yue, H.; Liu, S.; Wang, H.; Du, Y.; Xu, C.; Dong, W.; Liu, C. Cu-Ag/hydrotalcite catalysts for dehydrogenative cross-coupling of primary and secondary benzylic alcohols. *RSC Adv.* **2016**, *6*, 24164–24174.

(12) El-Sepelgy, O.; Matador, E.; Brzozowska, A.; Rueping, M. Calkylation of secondary alcohols by primary alcohols through Mncatalyzed double hydrogen autotransfer. *ChemSusChem* **2019**, *12*, 3099–3102.

(13) Freitag, F.; Irrgang, T.; Kempe, R. Cobalt-catalysed alkylation of secondary alcohols with primary alcohols *via* borrowing hydrogen/hydrogen autotransfer. *Chem. - Eur. J.* **201**7, *23*, 12110–12113.

(14) Liu, T.; Wang, L.; Wu, K.; Yu, Z. Manganese-catalyzed  $\beta$ -alkylation of secondary alcohols with primary alcohols under phosphine-free conditions. *ACS Catal.* **2018**, *8*, 7201–7207.

(15) (a) Fu, S.; Shao, Z.; Wang, Y.; Liu, Q. Manganese-catalyzed upgrading of ethanol into 1-butanol. J. Am. Chem. Soc. 2017, 139, 11941-11948. (b) Liu, Y.; Shao, Z.; Wang, Y.; Xu, L.; Yu, Z.; Liu, Q. Manganese-catalyzed selective upgrading of ethanol with methanol into isobutanol. ChemSusChem 2019, 12, 3069-3072. (c) kaithal, A.; Bonn, P. V.; Hölscher, M.; Leitner, W. Manganese(I)-catalyzed  $\beta$ methylation of alcohols using methanol as C1 source. Angew. Chem., Int. Ed. 2020, 59, 215-220. (d) Schlagbauer, M.; Kallmeier, F.; Irrgang, T.; Kempe, R. Manganese-catalyzed  $\beta$ -methylation of alcohols using methanol. Angew. Chem., Int. Ed. 2020, 59, 1485-1490. (e) kaithal, A.; Gracia, L. L.; Camp, C.; Quadrelli, E. A.; Leitner, W. Direct synthesis of cycloalkanes from diols or ketones using a homogeneous Mn catalyst. J. Am. Chem. Soc. 2019, 141, 17487-17492. (f) Ng, W. T.; Liao, G.; Lau, K.; Pan, H. J.; Zhao, Y. Roomtemperature Guerbet reaction with unprecedented catalytic efficiency and enantioselectivity. Angew. Chem., Int. Ed. 2020, 59, 11384-11389. (16) Zhang, M.; Li, H.; Young, D. J.; Li, H. Y.; Lang, J. P. Reaction

condition controlled nickel(II)-catalyzed C-C cross coupling of alcohols. Org. Biomol. Chem. 2019, 17, 3567–3574.

(17) Das, J.; Singh, K.; Vellakkaran, M.; Banerjee, D. Nickelcatalysed hydrogen borrowing strategy for  $\alpha$ -alkylation of ketones with alcohols: A new route to branched gem-bis(alkyl)ketones. *Org. Lett.* **2018**, *20*, 5587–5591.

(18) Das, J.; Vellakkaran, M.; Banerjee, D. Nickel-catalyzed alkylation of ketone enolates: Synthesis of monoselective linear ketones. *J. Org. Chem.* **2019**, *84*, 769–779.

(19) (a) Yang, J.; Liu, X.; Meng, D. L.; Chen, H. Y.; Zong, Z. H.; Feng, T. T.; Sun, K. Efficient Fe-catalyzed direct  $\beta$ -alkylation of secondary alcohols with primary alcohols. *Adv. Synth. Catal.* **2012**, 354, 328–334. (b) Liao, S.; Yu, K.; Li, Q.; Tian, H.; Zhang, Z.; Yu, X.; Xu, Q. Copper-catalyzed C-alkylation of secondary alcohols and methyl ketones with alcohols employing the aerobic relay race methodology. *Org. Biomol. Chem.* **2012**, *10*, 2973–2978.

(20) (a) Alonso, F.; Riente, P.; Yus, M. The  $\alpha$ -alkyltion of ketones with primary alcohols promoted by nickel nanoparticles under mild and ligandless conditions. *Synlett* **2007**, 2007, 1877–1880. (b) Alonso, F.; Riente, P.; Yus, M. Alcohols for the  $\alpha$ -alkylation of methyl ketones and indirect Aza-Wittig reaction promoted by nickel nanoparticles. *Eur. J. Org. Chem.* **2008**, 2008, 4908–4914.

Letter