

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1993). Data reduction: *MolEN* (Fair, 1990). Program(s) used to solve structure: *MolEN*. Program(s) used to refine structure: *MolEN*. Molecular graphics: *MolEN*. Software used to prepare material for publication: *MolEN*.

The authors wish to acknowledge the purchase of the CAD-4 diffractometer under grant DPT/TBAG1 of the Scientific and Technical Research Council of Turkey.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: AB1518). Services for accessing these data are described at the back of the journal.

References

- Allen, F. H., Kennard, O., Watson, D., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
- Awouters, F. H. L., Niemegeers, C. J. E. & Janssen, P. A. J. (1983). *Arzneim. Forsch. Drug Res.* **33**, 381–388.
- Brandstrom, A., Lindberg, P. & Junggren, U. (1985). *Scand. J. Gastroenterol.* **20** (Suppl. 108), 15–22.
- Enraf–Nonius (1993). *CAD-4 EXPRESS*. Version 1.1. Enraf–Nonius, Delft, The Netherlands.
- Escande, A. & Galigne, J. G. (1974). *Acta Cryst.* **B30**, 1647–1648.
- Fair, C. K. (1990). *MolEN. An Interactive Intelligent System for Crystal Structure Analysis*. Enraf–Nonius, Delft, The Netherlands.
- Özbe, S., İde, S. & Kendi, E. (1998). *J. Mol. Struct.* In the press.
- Saito, T., Hagihara, A., Igarashi, N., Matsuda, N., Yamashita, A., Ito, K., Mio, M. & Tasaka, K. (1993). *Jpn J. Pharmacol.* **62**, 137–143.
- Sakai, T., Hamada, T., Awata, N. & Watanabe, J. (1989). *J. Pharm. Dyn.* **12**, 530–536.
- Vasudevan, K. T., Puttaraja & Kulkarni, M. V. (1994). *Acta Cryst.* **C50**, 1286–1288.

Acta Cryst. (1998). **C54**, 856–859

2-Amino-5-nitrobenzophenone and 2'-Fluoro-2-methylamino-5-nitrobenzophenone

PHILIP J. COX,^a ABU T. MD. ANISUZZAMAN,^b R. HOWARD PRYCE-JONES,^b GRAHAM G. SKELLERN,^b ALASTAIR J. FLORENCE^b AND NORMAN SHANKLAND^b

^aSchool of Pharmacy, The Robert Gordon University, Schoolhill, Aberdeen AB10 1FR, Scotland, and ^bDepartment of Pharmaceutical Sciences, University of Strathclyde, 204 George Street, Glasgow G1 1XW, Scotland. E-mail: paspc@pharmacy.rgu.ac.uk

(Received 19 November 1997; accepted 11 December 1997)


Abstract

The title compounds, C₁₃H₁₀N₂O₃ and C₁₄H₁₁FN₂O₃, were prepared by acid degradation of the 1,4-benzodiazepine drugs nitrazepam and flunitrazepam, respec-

tively. The structure of 2-amino-5-nitrobenzophenone reported here is a polymorphic form of a known structure.


Comment

The 1,4-benzodiazepine drugs nitrazepam and flunitrazepam are prescribed for the short-term treatment of insomnia. Both drugs are hydrolysed in aqueous acid solution to produce substituted benzophenone products *via* a ring-opened intermediate. Thus nitrazepam, (I) (see scheme below), hydrolyses to 2-amino-5-nitrobenzophenone, (II) (Han *et al.*, 1977; Broxton & Morrison, 1985; Davidson & Smail, 1991; Anisuzzaman, 1995), and flunitrazepam, (III), to 2-methylamino-2'-fluoro-5-nitrobenzophenone, (IV) (Debruyne *et al.*, 1984; Moro *et al.*, 1991; Anisuzzaman, 1995). Crystal structures have been reported for (I) (Gilli *et al.*, 1977), (II) (Dvorkin *et al.*, 1985) and (III) (Butcher *et al.*, 1983), but not for (IV). We report here two new structures, namely, that of (IV) and that of a polymorph of the previously reported compound (II).



(I) $R^1 = \text{H}, R^2 = \text{H}$

(III) $R^1 = \text{CH}_3, R^2 = \text{F}$




(II) $R^1 = \text{H}, R^2 = \text{H}$

(IV) $R^1 = \text{CH}_3, R^2 = \text{F}$

Benzophenones almost always crystallize in the propeller conformation, with the torsion angles φ_1 and φ_2 (see scheme below) taking the same sign (Rappoport *et al.*, 1990). This is indeed the case for compounds (II) [$\varphi_1 = 21.7(3)$ and $\varphi_2 = 45.9(3)^\circ$] and (IV) [$\varphi_1 = -8.5(4)$ and $\varphi_2 = -57.7(4)^\circ$]. The previously reported polymorph of (II) has $\varphi_1 = 24.6$ and $\varphi_2 = 49.1^\circ$. By way of comparison, the corresponding angles observed in energy-optimized structures of benzophenone are $\varphi_1 = \varphi_2 = 26.2^\circ$ (Kendrick, 1990) and $\varphi_1 = \varphi_2 =$

30° (Rappoport *et al.*, 1990), while in the crystal structure of benzophenone $\varphi_1 = 29.4$ and $\varphi_2 = 30.9^\circ$ (Fleischer *et al.*, 1968).



In both compounds, an amino H atom forms an intramolecular hydrogen bond with the carbonyl O atom, O1 [in compound (II), $\text{H1B} \cdots \text{O1} = 2.03 \text{ \AA}$, and in compound (IV), $\text{H1} \cdots \text{O1} = 1.98 \text{ \AA}$]. Another feature common to both structures is the presence of a single intermolecular hydrogen bond between an amino H atom and an O atom [$\text{H1A} \cdots \text{O3}(-1 + x, \frac{1}{2} - y, -\frac{1}{2} + z) = 2.14 \text{ \AA}$ in (II) and $\text{H1} \cdots \text{O1}(-x, -y, -z) = 2.42 \text{ \AA}$ in (IV)]. These features are shown in Figs. 1 and 2.




Fig. 1. The atomic arrangement and hydrogen bonding in molecules of (II). Displacement ellipsoids are shown at the 50% probability level. H atoms have been set artificially small for clarity. O3' is related to O3 by the symmetry operation $-1 + x, \frac{1}{2} - y, -\frac{1}{2} + z$.




Fig. 2. The atomic arrangement and hydrogen bonding in molecules of (IV). Displacement ellipsoids are shown at the 50% probability level. H atoms have been set artificially small for clarity. Primed (') atoms are generated from their unprimed counterparts by inversion through the origin.

Atom H1 in compound (IV) therefore acts as a donor for two hydrogen bonds in a bifurcated system. The previously published polymorph of 2-amino-5-nitrobenzophenone, (IIP) (Dvorkin *et al.*, 1985), is also monoclinic ($P2_1/b$, $a = 7.851$, $b = 12.686$, $c = 11.121 \text{ \AA}$ and $\gamma = 95.58^\circ$) and possesses a similar hydrogen-bonding network to that of the structure reported here. The amino H-atom bond geometries for these three compounds are given in Table 3, where the O-atom coordinates of (IIP) are transposed by $-x, \frac{1}{2} - y, -\frac{1}{2} + z$.

A scan of the close contacts in the crystal structure of (IV) also reveals a short C—H $\cdots$ O contact: $\text{H14C} \cdots \text{O2}(\frac{3}{2} - x, -\frac{1}{2} + y, \frac{1}{2} - z) = 2.38 \text{ \AA}$.

Experimental

Nitrazepam was hydrolysed with aqueous HCl to give (II), and flunitrazepam was hydrolysed with aqueous HCl to give (IV). Crystals of (II) were obtained from aqueous HCl solution, while (IV) was recrystallized by slow evaporation of an ethanol solution.

Compound (II)

Crystal data

$\text{C}_{13}\text{H}_{10}\text{N}_2\text{O}_3$
 $M_r = 242.23$

Mo $K\alpha$ radiation
 $\lambda = 0.7107 \text{ \AA}$

Monoclinic

 $P2_1/c$ $a = 5.7358 (15) \text{ \AA}$ $b = 14.693 (2) \text{ \AA}$ $c = 13.1120 (12) \text{ \AA}$ $\beta = 98.87 (4)^\circ$ $V = 1091.8 (3) \text{ \AA}^3$ $Z = 4$ $D_x = 1.474 \text{ Mg m}^{-3}$ D_m not measured

Data collection

Delft Instruments FAST
diffractometer

Area-detector scans

Absorption correction: none

4705 measured reflections

1705 independent reflections

Refinement

Refinement on F^2 $R[F^2 > 2\sigma(F^2)] = 0.038$ $wR(F^2) = 0.077$ $S = 0.757$

1705 reflections

164 parameters

H atoms: see below

 $w = 1/[\sigma^2(F_o^2) + (0.0128P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ Cell parameters from 250
reflections $\theta = 2.10\text{--}25.09^\circ$ $\mu = 0.107 \text{ mm}^{-1}$ $T = 150 (2) \text{ K}$

Needle

 $0.20 \times 0.15 \times 0.15 \text{ mm}$

Yellow

1004 reflections with

 $I > 2\sigma(I)$ $R_{\text{int}} = 0.082$ $\theta_{\text{max}} = 25.09^\circ$ $h = -6 \rightarrow 6$ $k = -17 \rightarrow 17$ $l = -15 \rightarrow 11$ $(\Delta/\sigma)_{\text{max}} = 0.037$ $\Delta\rho_{\text{max}} = 0.192 \text{ e \AA}^{-3}$ $\Delta\rho_{\text{min}} = -0.143 \text{ e \AA}^{-3}$

Extinction correction: none

Scattering factors from

*International Tables for
Crystallography* (Vol. C)

Refinement

Refinement on F^2 $R[F^2 > 2\sigma(F^2)] = 0.033$ $wR(F^2) = 0.068$ $S = 0.496$

1946 reflections

184 parameters

H atoms: see below

 $w = 1/\sigma^2(F_o^2)$ $(\Delta/\sigma)_{\text{max}} < 0.001$ $\Delta\rho_{\text{max}} = 0.156 \text{ e \AA}^{-3}$ $\Delta\rho_{\text{min}} = -0.139 \text{ e \AA}^{-3}$

Extinction correction: none

Scattering factors from

*International Tables for
Crystallography* (Vol. C)Table 2. Selected geometric parameters ($\text{\AA}$, $^\circ$) for (IV)

F1—C9	1.365 (3)	O3—N2	1.236 (3)
O1—C7	1.230 (3)	N1—C2	1.343 (3)
O2—N2	1.236 (3)	N2—C5	1.447 (4)
C6—C1—C7—C8	−8.5 (4)	C1—C7—C8—C9	−57.7 (4)

Table 3. Hydrogen-bond geometry ($\text{\AA}$, $^\circ$)

	N—H	O...H	N...O	N—H...O
(II) (intra)	0.86	2.03	2.664 (3)	123
(II) (inter)	0.86	2.14	2.972 (3)	164
(IV) (intra)	0.88	1.97	2.658 (4)	134
(IV) (inter)	0.88	2.41	3.112 (4)	137
(IIP) (intra)	1.00	2.01	2.698	124
(IIP) (inter)	0.92	2.18	3.083	175

The unit-cell and intensity data were obtained using the routines *ENDEX*, *REFINE* and *MADONL* in the *MADNES* suite of software (Pflugrath & Messerschmidt, 1989) and processed using *ABSMAD* (Karaulov, 1992); detailed procedures are described by Darr *et al.* (1993). All non-H atoms were refined with anisotropic displacement parameters. The H atoms were initially placed in calculated positions and thereafter allowed to ride on their attached atoms with a common isotropic displacement parameter, which converged to 0.027 (2) $\text{\AA}^2$ for (II), and to 0.026 (3) (non-methyl) and 0.070 (6) $\text{\AA}^2$ (methyl) for (IV). We attribute the poor value of S for structure (IV) to the effects of uncertainties on F^2 in a very weak data set.

For both compounds, program(s) used to solve structures: *SIR92* (Altomare *et al.*, 1994). Program(s) used to refine structures: *SHELXL93* (Sheldrick, 1993) for (II); *SHELXL97* (Sheldrick, 1997) for (IV). For both compounds, molecular graphics: *ZORTEP* (Zsolnai, 1997).

The use of the EPSRC X-ray Crystallographic Service at the University of Wales, Cardiff, Wales, is gratefully acknowledged.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: BM1214). Services for accessing these data are described at the back of the journal.

References

- Altomare, A., Cascarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
 Anisuzzaman, A. T. Md. (1995). PhD thesis, University of Strathclyde, Glasgow, Scotland.
 Broxton, T. J. & Morrison, S. R. (1985). *Aust. J. Chem.* **38**, 1037–1043.
 Butcher, H., Hamor, T. A. & Martin, I. L. (1983). *Acta Cryst.* **C39**, 1469–1472.
 Darr, J. A., Drake, S. R., Hursthouse, M. B. & Malik, K. M. A. (1993). *Inorg. Chem.* **32**, 5704–5708.

Table 1. Selected geometric parameters ($\text{\AA}$, $^\circ$) for (II)

O3—N2	1.243 (2)	N1—C2	1.336 (3)
O2—N2	1.225 (2)	N2—C5	1.443 (3)
O1—C7	1.236 (2)		
C6—C1—C7—C8	21.7 (3)	C1—C7—C8—C9	45.9 (3)

Compound (IV)

Crystal data

C₁₄H₁₁FN₂O₃ $M_r = 274.25$

Monoclinic

 $P2_1/n$ $a = 4.2150 (17) \text{ \AA}$ $b = 13.6020 (18) \text{ \AA}$ $c = 21.547 (7) \text{ \AA}$ $\beta = 94.7480 (11)^\circ$ $V = 1231.1 (7) \text{ \AA}^3$ $Z = 4$ $D_x = 1.480 \text{ Mg m}^{-3}$

Data collection

Delft Instruments FAST
diffractometer

Area-detector scans

Absorption correction: none

5315 measured reflections

1946 independent reflections

Mo $K\alpha$ radiation $\lambda = 0.7107 \text{ \AA}$ Cell parameters from 250
reflections $\theta = 2.42\text{--}24.98^\circ$ $\mu = 0.116 \text{ mm}^{-1}$ $T = 150 (2) \text{ K}$

Needle

 $0.22 \times 0.18 \times 0.12 \text{ mm}$

Yellow

736 reflections with

 $I > 2\sigma(I)$ $R_{\text{int}} = 0.094$ $\theta_{\text{max}} = 24.98^\circ$ $h = -4 \rightarrow 3$ $k = -15 \rightarrow 14$ $l = -25 \rightarrow 24$

- Davidson, A. G. & Smail, G. A. (1991). *Int. J. Pharm.* **69**, 1–3.
- Debruyne, M. M. A., Sinnema, A. & Verweij, A. M. A. (1984). *Forensic Sci. Int.* **24**, 125–135.
- Dvorkin, A. A., Andronati, S. A., Gifeisman, T. S., Simonov, Y. A., Yavorsky, A. S. & Pavlovsky, V. I. (1985). *Dokl. Akad. Nauk Ukr. RSR Ser. B Geol. Khim. Biol. Nauk*, **8**, 34–37.
- Fleischer, E. B., Sung, N. & Hawkinson, S. (1968). *J. Phys. Chem.* **72**, 4311–4312.
- Gilli, G., Bertolasi, V., Sacerdoti, M. & Borea, P. A. (1977). *Acta Cryst.* **B33**, 2664–2667.
- Han, W. W., Yakatan, G. J. & Maness, D. D. (1977). *J. Pharm. Sci.* **66**, 795–798.
- Karaulov, A. I. (1992). *ABSMAD. Program for FAST Data Processing*. University of Wales, Cardiff, Wales.
- Kendrick, J. (1990). *J. Chem. Soc. Faraday Trans.* **86**, 3995–4000.
- Moro, M. E., Novilofertrell, J., Velazquez, M. M. & Rodriguez, L. J. (1991). *J. Pharm. Sci.* **80**, 459–468.
- Pflugrath, J. W. & Messerschmidt, A. (1989). *MADNES*. Version of 11 September 1989. Distributed by Delft Instruments, Delft, The Netherlands.
- Rappoport, Z., Biali, S. E. & Kaftory, M. (1990). *J. Am. Chem. Soc.* **112**, 7742–7748.
- Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997). *SHELXL97. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Zsolnai, L. (1997). *ZORTEP. An Interactive ORTEP Program*. University of Heidelberg, Germany.

Acta Cryst. (1998). **C54**, 859–860

3-*tert*-Butyl-2-hydroxy-5-methoxyazobenzene†

ŞAMIL IŞIK,^a MUHITTİN AYĞÜN,^b HASAN KOCAOKUTGEN,^c M. NAWAZ TAHİR,^d ORHAN BÜYÜKGÜNGÖR^a AND AHMET ERDÖNMEZ^a

^aDepartment of Physics, Ondokuz Mayıs University, TR-55139, Samsun, Turkey, ^bBuca Eğitim Fakültesi, Fizik Eğitimi Bölümü, Buca, 35160 İzmir, Turkey, ^cDepartment of Chemistry, Ondokuz Mayıs University, TR-55139, Samsun, Turkey, and ^dDepartment of Physics and Engineering, Hacettepe University, Beytepe 06532, Ankara, Turkey. E-mail: samili@samsun.omu.edu.tr

(Received 2 January 1997; accepted 6 January 1998)


Abstract

The structure of the title compound, C₁₇H₂₀N₂O₂, shows features characteristic of azobenzene derivatives and is perfectly planar in the solid state with a *trans* configuration. The molecule has an intramolecular O1—H1···N1 hydrogen bond.

† IUPAC name: 2-*tert*-butyl-4-methoxy-6-(phenyldiazenyl)phenol.

Comment

Azo compounds are widely used in the textile industry as synthetic colouring materials. The structure of the title compound, (I), is very similar to the structures of azo compounds studied previously (Işık *et al.*, 1997; Glowka & Olubek, 1994; Rodrigues *et al.*, 1996). The two phenyl rings are in a *trans* configuration. The lengths of the two C—N bonds are almost identical and have an average value of 1.410(3) Å. The average value of the C—N—N angles is 115.7(2)°. The N=N bond length is 1.274(3) Å, in accordance with the expected value for aromatic azo compounds. The whole molecule, except for two methyls of the *tert*-butyl group, lies on a crystallographic mirror plane, so that the molecule is perfectly planar.



The planarity of the molecule facilitates an O1—H1···N1 intramolecular hydrogen bond, in which the O1 hydroxyl group adjacent to the N=N bond serves as the donor to the N1 atom. The O1—H1···N1 angle is 149.6(2)° and the length of the hydrogen bond (H1···N1) is 1.598(2) Å. The length of the O1···N1 bond is 2.534(2) Å, which is shorter than a typical hydrogen bond of this type [N···O = 2.78(10) Å; Vinogradov & Linnell, 1971]. This shows the presence of a very strong intramolecular interaction in the molecule. There are no intermolecular interactions other




Fig. 1. ORTEP (Johnson, 1976) view of the title molecule and the atomic numbering. The displacement ellipsoids are at the 50% probability level. H atoms are shown as spheres of arbitrary size. [Symmetry code: (i) $x, \frac{1}{2} - y, z$.]