

Tetrahedron Letters, Vol. 37, No. 43, pp. 7801-7804, 1996 Copyright © 1996 Elsevier Science Ltd Printed in Great Britain. All rights reserved 0040-4039/96 \$15.00 + 0.00

PII: S0040-4039(96)01782-0

Synthesis and DNA Binding Properties of Amide Bond-Modified Analogues Related to Distamycin

Yasuo Yamamoto, Tetsutaro Kimachi*, Yoshitomo Kanaoka, Satoshi Kato, and Kiyoshi Bessho*

Faculty of Pharmaceutical Sciences, Kyoto University, Sakyo-ku Kyoto 606-01, Japan Takuyuki Matsumoto, Tetsuya Kusakabe, and Yukio Sugiura

Institute for Chemical Research, Kyoto University, Uji, Kyoto 611, Japan

Abstract : Novel nitro analogues of distamycin which have trans olefin or α -diketo moiety instead of amide bond were synthesized. Their DNA binding properties studied by ethidium displacement assay and MPE footprinting experiments were also described. Copyright © 1996 Elsevier Science Ltd

There is considerable interest in the development of agents which recognize and bind to DNA in a sequence specific manner. Interest is particularly focused on the minor groove binding agents. The naturally occurring tripeptide antibiotic distamycin (1) originally isolated from *Streptomyces Distallicus* is a nonintercalative minor groove binder with high selectivity for AT-rich sequences^{1, 2}. Recently several derivatives of 1 have been prepared and examined for their biological activities including DNA binding properties³. Among the structural modifications, it has been shown that replacement of pyrrole rings with other heterocyclic moieties (e.g. imidazole or thiazole) creates compounds with increased tolerance for GC base pairs at their binding sites⁴.





As shown in Scheme 1, we focus on the variation of the N-terminal formamide and the interpyrrole amide function of 1. Herein we report the synthesis of novel nitro derivatives of distamycin which have

trans-olefin or α -diketo interpyrrole connection instead of amide and also report their DNA binding properties. The compounds 2 and 3, which have formamide and nitro function in the N-terminal respectively, were prepared according to the known procedure^{3b, 5}. Synthesis of novel nitro analogues 4 and 5 which one of the interpyrrole linkage is replaced by *trans*-olefin and α -diketone respectively are shown in Scheme 2. In every case dimethylamino group was introduced in the C-terminal of the analogues instead of the amidine present in 1 (replacement of the amidinium group by a dimethylamino group has been shown to afford similar selectivity with respect to DNA sequences⁶).



Reagents and Conditions: i) HNO₃ / Ac₂O / -40°C \rightarrow rt. ii) MePPh₃Br / tBuOK / THF/ rt. iii) NBS / THF / -10°C \rightarrow rt. iv) MeONa / MeOH / rt. v) Pd(OAc)₂ / Et₃N / P(o-tol)₃ / DMF / 120 °C. vi) NaOH aq. / EtOH / 30°C. vii) WSCI / HOBt / DMF / rt. viii) HCI-Et₂O / MeOH. ix) KMnO₄ / Ac₂O / -5°C \rightarrow 10°C.

The olefinic compound 8 was obtained from commercially available aldehyde 6 via 4-nitro derivative 7 prepared by nitration with nitric acid-acetic anhydride mixture⁷, and following Wittig reaction. After several trials, it was found that Heck reaction⁸ between olefin 8 and bromopyrrole 9 afforded the coupling product 10 in 62 % yield. The coupling reaction of carboxylic acid 11 derived from 10 (by alkaline hydrolysis) with a freshly prepared amine 12 (by catalytic hydrogenation of the corresponding nitro compound) afforded the target molecule 4 as a hydrochloride salt⁹. The special interest of the olefin intermediate 10 resides in its useful application to introduce α -diketo- function in the molecule by KMnO4 oxidation¹⁰. Alkaline hydrolysis of 13 followed by the same procedure for 11 gave α -diketone product 5 as a hydrochloride salt¹¹.

Binding properties of compounds (1-5) to DNA were evaluated with the ethidium displacement assay¹²). The calculated apparent binding constant to poly (dA-dT) and poly (dG-dC) are listed in Table 1. These data show that all compounds can bind to DNAs used. It is noteworthy that the nitro analogue 3 has weaker binding ability $[8.7x10^5 \text{ M}^{-1}]$ to poly (dA-dT) than 1 and 2 $[345x10^5, 150.8x10^5 \text{ M}^{-1}]$, but, binds stronger to poly (dG-dC) $[3.5x10^5 \text{ M}^{-1}]$ than 1 and 2 $[1.3x10^5, 0.9x10^5 \text{ M}^{-1}]$, indicating that the N-terminal nitro group affected for the affinity toward poly (dG-dC). The α -diketone linked analogue 5 has markedly improved acceptance of GC base pairs. The Kapp value of compound 5 for poly (dG-dC) $[20.9x 10^5 \text{ M}^{-1}]$ is close to that for poly (dA-dT) $[26.6x10^5 \text{ M}^{-1}]$, while the Kapp value of compound 1 and 2 for poly (dG-dC) are about 2 orders of magnitude lower than those for poly (dA-dT).

Table 1. The apparent binding constants $Kapp(x10^5M^{-1})$ determined by an ethidium displacement assay

	1	2	3	4	5
poly(dA-dT)	345(348) ^{4d}	150.8	8.7	16.9	26.6
poly(dG-dC)	1.3(2.0) ^{4d}	0.9	3.5	7.4	20.9

To further confirm the DNA binding interactions, MPE-Fe(II) footprinting experiments were carried out for 1, 4 and 5 on the *Eco* RI / *Ssp* I restriction fragment of pBR 322 DNA¹³. The protection patterns from cleavage on bp 4301-4350 are shown in Figure 1.





a) Bar heights are proportional to the protection from cleavage at each band.

These results, in agreement with the data from the ethidium displacement assay, suggest the high selectivity of 1 for AT-rich sequences, the low acceptance of 4 for GC base pairs and the great tolerance of 5 for GC regions in their binding sites.

In conclusion, we demonstrated the synthesis of novel analogues of distamycin *via* intermolecular Heck reaction and KMnO4 olefin oxidation as key reactions. It is also demonstrated that amide bond modifications of distamycin could lead to compounds with quite different DNA recognition patterns.

REFERENCES AND NOTES

- 1. Di Marco, A.; Gaetani, M.; Orezzi, P.; Scotti, T.; Arcamone, F. Cancer Chemother. Rep. 1962, 18, 15.
- 2. Kopka, M. L.; Yoon, C.; Goodsell, D.; Pijura, P.; Dickerson, R. E. Proc. Natl. Acad. Sci. USA 1985, 82, 1376.
- Recent references; (a) Hartley, J. A.; Webber, J.; Wyatt, M. D.; Bordenic, N.; Lee, M. Bioorg. & Med. Chem. 1995, 3, 623. (b) Chen, Y.-H.; Lown, J. W. J. Am. Chem. Soc. 1994, 116, 6995. (c) Yamamoto, Y.; Cai, J.; Nakamura, H.; Sadayori, N.; Asao, N.; Nemoto, H. J. Org. Chem. 1995, 60, 3352. (d) Mrksich, M.; Dervan, P. B. J. Am. Chem. Soc. 1995, 117, 3325.
- (a) Lown, J. W.; Krowicki, K.; Bhat, U. G.; Skorobogaty, A.; Ward, B.; Dabrowiak, J. C. Biochemistry 1986, 25, 7408. (b) Kissinger, K.; Krowicki, K.; Dabrowiak, J. C.; Lown, J. W. Biochemistry 1987, 26, 5590. (c) Lee, M.; Chang, D. K.; Hartley, J. A.; Pon, R. T.; Krowicki, K.; Lown, J. W. Biochemistry 1988, 27, 445. (d) Lee, M.; Chang, D. K.; Hartley, J. A. Biochemistry 1993, 32, 4237.
- 5. Nishiwaki, E.; Tanaka, S.; Lee, H.; Shibuya, M. Heterocycles 1988, 27, 1945.
- 6. Taylor, J. S.; Schults, P. G.; Dervan, P. B. Tetrahedron 1984, 40, 457.
- 7. Special care must be paid to avoid the drastic elevation of reaction temperature when it bacomes higher than -5°C.
- (a) Frank, W. C.; Kim, Y. C.; Heck, R. F. J. Org. Chem. 1978, 43, 2947. (b) de Meijere, A.; Meyer, F. E. Angew. Chem. Int. Ed. Engl. 1994, 33, 2379. (c) Heck, R. F. Organic Reactions, published by John Wiley & Sons Inc., 1982, 27, 345.
- 9. Spectral data for 4 ¹H-NMR (270MHz, DMSO d-6) δ 1.78 (m, 2H), 2.56 (s, 6H), 2.79 (m, 2H), 3.24 (m, 2H), 3.73 (s, 3H), 3.81 (s, 3H), 3.89 (s, 3H), 6.65 (d, 1H, J = 16.2Hz), 6.90 (d, 1H, J = 1.7Hz), 6.94 (d, 1H, J = 1.7Hz), 6.98 (d, 1H, J = 16.2Hz), 7.15 (d, 1H, J = 1.7Hz), 7.18 (d, 1H, J = 1.7Hz), 7.20 (d, 1H, J = 1.7Hz), 7.97 (d, 1H, J = 1.7Hz), 8.16 (t, 1H, J = 5.3Hz), 9.98 (s, 1H), 10.36 (bs, 1H), FABMS (m/z) 482 (M-HCl+H), HRFABMS (m/z) 482.2522 (M-HCl+H, 482.2519 calcd for C_{24H32N7O4}).
- (a) Jensen, H. P.; Sharpless, K. B. J. Org. Chem. 1974, 39, 2314. (b) Sharpless, K. B.; Lauer, R. F.; Oljan, R.; Teranishi, A. Y.; Williams, D. R. J. Am. Chem. Soc. 1971, 93, 3303.
- Spectral data for 5 ¹H-NMR (270 MHz, DMSO-d 6) δ 1.87 (m, 2H), 2.74 (s, 3H), 2.76 (s, 3H), 3.03 (m, 2H), 3.24 (m, 2H), 3.82 (s, 3H), 3.94 (s, 3H), 4.07 (s, 3H), 6.92 (d, 1H, J = 2.0Hz), 7.22 (d, 1H, J = 1.7Hz), 7.38 (d, 1H, J = 2.0 Hz), 7.51 (d, 1H, J = 2.0Hz), 7.89 (d, 1H, J = 1.7Hz), 8.23 (bs, 1H), 8.52 (d, 1H, J = 1.7Hz), 10.12 (bs, 1H), 10.27 (s, 1H), FABMS (m/z) 512 (M-HCl+H), HRFABMS (m/z) 512.2238 (M-HCl+H), 512.2260 calcd for C₂₄H₃₀N₇O₆)
- Morgan, A. R.; Lee, J. S.; Pulleyblank, D. F.; Murray, N. L.; Evans, D. H. Nucl. Acids. Res. 1979, 7, 547. These experiments followed conditions used in ref. 4 (d). Buffer solution (pH = 7.0) containing Tris-HCl (10 mM) and sodium chloride (10mM) was used.
- 13. A 50 μM MPE-Fe(II) solution was prepared by mixing a 100 μM MPE aqueous solution (20 μL) with a freshly prepared 100 μM ferrous ammonium sulfate aqueous solution (20 μL). Reaction mixture (16 μL) containing each compound, the labeled restriction fragment, sonicated calf thymus DNA (0.4 μg), sodium chloride, and Tris-HCl (pH = 7.5) was incubated at 20 °C for 30 min. To each tube was added 2 μL of a 50 μM MPE- Fe(II) solution followed by 2 μL of 100 mM dithiothreitol. Final concentrations were 5μM [MPE-Fe(II)], 10mM (dithiothreitol), 50 mM (sodium chloride), and 20 mM (Tris-HCl). The solutions were incubated at 20 °C for 10 min, lyophilized and electrophopresed.

(Received in Japan 9 August 1996; revised 6 September 1996; accepted 10 September 1996)