

Synthesis of Chiral Flavanones from Tricarbonyl (η^6 -Arylbenzaldehyde)chromium(0)

GUI-XIANG LIU¹, YU WAN², LING-LING ZHAO¹, HAI-YING WANG¹, ZHOU XU¹, JIN-LONG QI¹ and HUI WU^{1,2,*}

¹School of Chemistry and Chemical Engineering, Jiangsu Normal University, Jiangsu Key Laboratory of Green Synthesis of Functional Material, Xuzhou 221116, P.R. China

²Jiangsu Key Laboratory of Biotechnology on Medical Plant, Xuzhou 221116, P.R. China

*Corresponding author: Fax: +86 516 83500164; Tel: +86 516 83403163; E-mail: wuhui72@yahoo.com.cn

(Received: 29 September 2012;

Accepted: 18 July 2013)

AJC-13819

A synthesis of chiral flavanones *via* the condensation of tricarbonyl(η^6 -arylbenzaldehyde)chromium(0) and *o*-hydroxyacetophenone in a shorter time at room temperature have been developed. The tricarbonylchromium(0) group was removed by virtue of light and the enantioenriched flavanones was formed with highly enantioselectivity (> 95 % ee).

Key Words: Synthesis, Chiral flavanones, Chiral tricarbonylchromium(0) aromatic aldehydes.

INTRODUCTION

Flavanones constitute a large number of natural products with many medicinal applications¹. Because of their broad range of biological activities such as anti HIV², anticancer³, antioxidant⁴, gastro-protective⁵, antibacterial and antifungal^{6,7}, a variety of approaches have been developed to synthesize flavanones. The most commonly one was the intramolecular cyclodehydration of 1-(2-hydroxyphenyl)-1,3-diketones⁸. Although the diastereoselective flavanones has been shown to proceed, in some cases, with high to excellent diastereoselectivities, the enantioselective variation of this reaction is less developed. A limited number of strategies have been developed to obtain chiral flavanones, such as resolution of the related alcohols⁹ or substitution reactions¹⁰. Due to its difficult synthesis and harsh experimental conditions, it is needed to find efficient method for the synthesis of chiral flavanones.


Herein we report an efficient synthesis of chiral flavanones *via* the reaction of tricarbonylchromium(0) and *o*-hydroxyacetophenone.

EXPERIMENTAL

Solvents were dried and liquid aromatic aldehydes were distilled prior to use. NMR spectra were recorded on a Bruker Avance DMX400 spectrometer in DMSO-*d*₆ with TMS as an internal standard. Infrared spectra were recorded on Bruker FTIR-Tensor27 spectrometer expressed in cm⁻¹ (KBr). Mass was determined by using a Bruker Micro TOF-MS high resolution mass spectrometer. TLC analysis was performed with

glass backed plates precoated with silica gel and examined under UV (254 nm).

Experimental process: Synthesis of racemic flavanones (**Scheme-I**): Compound **3** (2-hydroxychalcones) was prepared through condensation of aromatic aldehyde **1** (5 mmol) and *o*-hydroxyacetophenone **2** (5 mmol) in 20 % NaOH aqueous solution (2 mL), using TBAB (tetrabutylammonium bromide, 0.15 mmol, 0.05 g) as phase transfer catalyst under microwave irradiation for 5 min. Then the reaction mixture was poured into ethyl acetate (20 mL) and washed with water for 2-3 times. The combined organic layers were dried over anhydrous Na₂SO₄, concentrated in vacuo and purified by chromatography on silica gel (petroleum ether:ethyl acetate = 3:1) to give **3**.




Scheme-I: Synthesis of enantiomers 2-phenychroman-4-ones

The mixture of **3** (2 mmol), hexahydropyridine (1 mL) and water (10 mL) was taken in a 50 mL round bottom flask and stirred at room temperature for 5-6 h. Then by filtering flavanones **4** was obtained (Table-1).

TABLE-1
SYNTHETIC RESULTS OF RACE-FLAVANONE **4**

Entry	R	Formula	m.p. (°C)	Time	Yield (%)
4a	2 -OCH ₃	C ₁₆ H ₁₄ O ₃	106-107	3.0	85
4b	2,3(-OCH ₃) ₂	C ₁₇ H ₁₆ O ₄	92-93	3.0	80
4c	3,4(-OCH ₃) ₂	C ₁₇ H ₁₆ O ₄	73-74	3.5	85
4d	2-Br	C ₁₅ H ₁₁ O ₂ Br	97-98	3.0	89
4e	3-Br	C ₁₅ H ₁₁ O ₂ Br	95-96	3.0	90
4f	4-Cl	C ₁₅ H ₁₁ O ₂ Cl	81-82	2.0	92
4g	—	C ₁₅ H ₁₂ O ₂	70-72	2.0	90

Synthesis of chiral flavanones (Scheme-II): In N₂ atmosphere the chiral tricarbonylchromium(0) aromatic aldehydes **5** (based on our previous work¹¹) (5 mmol) and *o*-hydroxyacetophenone **2** (5 mmol) were stirred with 20 % NaOH solution (2 mL) in 95 % ethanol (10 mL) and THF (tetrahydrofuran 10 mL) at room temperature for 1 h. Then the reaction mixture was poured into ethyl acetate (20 mL) and washed with water for 2-3 times. The combined organic layers were dried over anhydrous Na₂SO₄, concentrated *in vacuo* and purified by chromatography on silica gel (petroleum ether:ethyl acetate = 3:1) to give chiral tricarbonylchromium(0) 2-hydroxychalcones **6**.



Scheme-II: Synthesis of chiral 2-phenylchroman-4-ones


The mixture of **6** (2 mmol), sodium acetate and 95 % ethanol (10 mL) was taken in a 50 mL round bottom flask and stirred at room temperature for 5-6 h. Then by filtering we got chiral tricarbonylchromium(0) flavanones **7**. By virtue of light, the tricarbonylchromium(0) group was removed and obtained the chiral flavanones **8** (Table-2).

TABLE-2
SYNTHETIC RESULTS OF CHIRAL FLAVANONES

Entry	Time	Yield (%) ^a	e.e (%) ^b	Config.
+8a	2.0	65	96	R
-8a	2.0	65	98	S
+8b	2.0	60	95	R
-8b	2.0	60	94	S

^aIsolated yields after chromatography. ^bDetermined by chiral-HPLC using AD-H column.

Probable mechanism (Scheme-III): Firstly, the starting chalcone was accessed *via* Knoevenagel condensation, the chiral tricarbonylchromium(0) aromatic aldehydes reacted with *o*-hydroxyacetophenone in alkaling condition. Then, based on acid catalyzed the flavanones was formed upon intramolecular cyclization. At last the tricarbonylchromium(0) group was removed by virtue of light and the enantioenriched flavanones was formed (Scheme-III).



Scheme-III: Probable mechanism

Spectral data for new compounds

2,3-Dihydro-2-(2-methoxyphenyl)chromen-4-one (4a): colourless crystals, IR (KBr, $\nu_{\max}$, cm⁻¹): 3030, 2960, 2920, 2870, 1680, 1600, 1500, 1460, 1380, 1260. ¹H NMR (δ ppm, DMSO-*d*₆): 2.86 (dd, $J_1 = 2.4$ Hz, $J_2 = 16.8$ Hz, C₃-H), 3.33 (t, $J = 13.6$ Hz, C₃-H), 3.89 (s, 3H, CH₃O), 5.84 (dd, $J_1 = 2.8$ Hz, $J_2 = 13.2$ Hz, 1H, C₂-H), 7.70-7.13 (m, 8H, ArH). Anal. calcd. (%) for C₁₆H₁₄O₃: C 63.0, H 5.5; found (%): C 63.2, H 5.6.

2,3-Dihydro-2-(2,3-dimethoxyphenyl)chromen-4-one (4b): IR (KBr, $\nu_{\max}$, cm⁻¹): 3030, 2960, 2920, 2860, 1680, 1600, 1500, 1450, 1380, 1260, 1120. ¹H NMR (δ ppm, DMSO-*d*₆): 2.86 (dd, $J_1 = 2.4$ Hz, $J_2 = 16.8$ Hz, C₃-H), 3.32 (t, $J = 13.6$ Hz, C₃-H), 3.83 (s, 3H, CH₃O), 3.89 (s, 3H, CH₃O), 5.84 (dd, $J_1 = 2.8$ Hz, $J_2 = 13.2$ Hz, 1H, C₂-H), 7.70-7.13 (m, 7H, ArH). Anal. calcd. (%) for C₁₇H₁₆O₄: C 71.8, H 5.6; found (%): C 71.9, H 5.7.

2,3-Dihydro-2-(3,4-dimethoxyphenyl)chromen-4-one (4c): IR (KBr, $\nu_{\max}$, cm⁻¹): 3030, 2960, 2920, 2870, 1680, 1600, 1500, 1450, 1380, 1260, 1130. ¹H NMR (δ ppm, DMSO-*d*₆): 2.85 (dd, $J_1 = 2.4$ Hz, $J_2 = 16.8$ Hz, C₃-H), 3.32 (t, $J = 13.6$ Hz, C₃-H), 3.83 (s, 3H, CH₃O), 3.89 (s, 3H, CH₃O), 5.84 (dd, $J_1 = 2.8$ Hz, $J_2 = 13.2$ Hz, 1H, C₂-H), 7.70-7.13 (m, 7H, ArH). Anal. calcd. (%) for C₁₇H₁₆O₄: C 71.8, H 5.6; found (%): C 71.8, H 5.7.

2,3-Dihydro-2-(2-bromo)chromen-4-one (4d): IR (KBr, $\nu_{\max}$, cm⁻¹): 3030, 2920, 1680, 1600, 1500, 1450, 1260. ¹H NMR (δ ppm, DMSO-*d*₆): 3.33 (t, $J = 13.6$ Hz, C₃-H), 2.84 (dd, $J_1 = 2.4$ Hz, $J_2 = 16.8$ Hz, C₃-H), 5.86 (dd, $J_1 = 2.8$, $J_2 = 13.2$ Hz, 1H, C₂-H), 8.27-7.00 (m, 8H, ArH). Anal. calcd. (%) for C₁₅H₁₁O₂Br: C 59.4, H 3.6; found (%): C 59.5, H 3.4.

2,3-Dihydro-2-(2-chloro)chromen-4-one (4e): IR (KBr, $\nu_{\max}$, cm⁻¹): 3030, 2920, 1680, 1600, 1500, 1450, 1260. ¹H NMR (δ ppm, DMSO-*d*₆): 3.33 (t, $J = 13.6$ Hz, C₃-H), 2.84 (dd, $J_1 = 2.4$ Hz, $J_2 = 16.8$ Hz, C₃-H), 5.86 (dd, $J_1 = 2.8$, $J_2 = 13.2$ Hz, 1H, C₂-H), 8.27-7.00 (m, 8H, ArH). Anal. calcd. (%) for C₁₅H₁₁O₂Cl: C 59.4, H 3.6; found (%): C 59.5, H 3.4.

2,3-Dihydro-2-(4-chloro)chromen-4-one (4f): IR (KBr, $\nu_{\max}$, cm⁻¹): 3030, 2920, 1680, 1600, 1500, 1450, 1260, 1130. ¹H NMR (δ ppm, DMSO-*d*₆): 2.83 (dd, $J_1 = 2.4$ Hz, $J_2 = 16.8$ Hz, C₃-H), 3.35 (t, $J = 13.6$ Hz, C₃-H), 5.82 (dd, $J_1 = 2.8$ Hz, $J_2 = 13.2$ Hz, 1H, C₂-H), 7.70-7.13 (m, 8H, ArH). Anal. calcd. (%) for C₁₅H₁₁O₂Cl: C 69.6, H 4.3; found (%): C 69.7, H 4.4.

RESULTS AND DISCUSSION

In this work, it was found that the ee % of products were high when chiral tricarbonylchromium(0) aromatic aldehydes were used as chiral reactant. The results indicated that both (-)-tricarbonylchromium(0) aromatic aldehydes and (+)-

tricarbonylchromium(0) aromatic aldehydes played the same activity and stereoselective role in the synthesis of chiral flavanones.

Conclusion

In summary, we have developed an efficient method for the synthesis of chiral flavanones. This is a novel application of chiral tricarbonylchromium(0) aromatic aldehydes. Shorter total times (5-6 h) were taken with highly enantioselective (> 95 % ee) at room temperature comparing with previous methods. Our method for the synthesis of these natural products provided an efficient way and made significant improvement.

ACKNOWLEDGEMENTS

The authors are grateful to the foundation of the Priority Academic Program Development of Jiangsu Higher Education Institutions, Key Basic Research Project in Jiangsu University (No. 10KJA430050, 11KJB150017), the Graduate Innovation Project in Jiangsu Province (No. CXZZ12_0979), The Project of Outstanding Scientific and Technological Innovation Team

for Higher Education Institutions in Jiangsu Province (Pre-development of medicinal microbial resources) for financial support.

REFERENCES

1. D.O. Bennardi, G.P. Romanelli, J.C. Autino, G.T. Baronettic and H.J. Thomas, *Appl. Catal. A*, **404**, 68 (2011).
2. J. Wu, X. Wang, Y. Yi and K. Lee, *Bioorg. Med. Chem. Lett.*, **13**, 1813 (2003).
3. S. Martens and A. Mithofer, *Phytochemistry*, **66**, 2399 (2005).
4. H. Chu, H. Wu and Y. Lee, *Tetrahedron*, **60**, 2647 (2004).
5. J. Ares, P. Outt, J. Randall, P. Murray, P. Weisshaar, L. Obrien, B. Ems, S. Kakodkar, G. Kelm, W. Kershaw, K. Werchowski and A. Parkinson, *J. Med. Chem.*, **38**, 4937 (1995).
6. S. Alam, *J. Chem. Sci.*, **116**, 325 (2004).
7. H. Goker, D. Boykin and S. Yildiz, *Bioorg. Med. Chem.*, **13**, 1707 (2005).
8. G. Kabalka and A. Mereddy, *Tetrahedron Lett.*, **46**, 6315 (2005).
9. S. Ramadas and G.L.D. Krupadanam, *Tetrahedron: Asymm.*, **15**, 3381 (2004).
10. K.J. Hodgetts, *Tetrahedron*, **61**, 6860 (2005).
11. J.-J. Shi, L.-L. Zhao, X.-X. Zhang, C. Wang, Y. Wan, Z. Xu and H. Wu, *Asian J. Chem*, **23**, 5319 (2011).