

Sodium Hexamethyldisilazide: Using ^{15}N – ^{29}Si Scalar Coupling to Determine Aggregation and Solvation States

Ryan A. Woltornist and David B. Collum*

Cite This: <https://dx.doi.org/10.1021/jacs.0c00331>

Read Online

ACCESS |



Metrics & More



Article Recommendations



Supporting Information

ABSTRACT: ^{29}Si NMR spectroscopy, the method of continuous variations, and density functional theory computations show that sodium hexamethyldisilazide (NaHMDS) is a disolvated dimer in toluene, a mixture of disolvated dimer and tetrasolvated monomer in THF/toluene, and exclusively monomer in neat THF. The dioxane-solvated NaHMDS only partially deaggregates to monomer even in neat dioxane. ^{15}N – ^{29}Si coupling constants and ^{29}Si chemical shifts show a high and dependable correlation with the aggregation state. Monitoring either chemical shift or coupling constant versus THF concentration even in the high-temperature, rapid-exchange limit affords the solvation numbers consistent with DFT computations. The preparation of ^{15}N -labeled NaHMDS has been improved.

A select few groups have been hailing the organic chemistry community to pay more attention to organosodium




Figure 1. DFT-computed, toluene-complexed dimer **1a**.

chemistry. We joined in only recently and possibly may not be the last to do so. To this end, sodium hexamethyldisilazide (NaHMDS) is arguably *the* preeminent organosodium reagent in both academic and industrial laboratories.¹ Despite its prominence, studies of its properties in solution are restricted to a couple of NMR spectra^{2,3} and a handful of computations.⁴

We have embarked on an extensive study using a combination of methods to determine the structure of NaHMDS in over 30 commonly employed organic solvents to begin to study how aggregation and solvation influence reactivity and selectivity.⁵ An unexpectedly important protocol revolves around the interrogation of structure using ^{15}N – ^{29}Si coupling observed in ^{15}N -labeled NaHMDS. Lukevics and co-workers used natural abundance to examine $^1\text{J}_{\text{N-Si}}$ coupling in a series of disilazanes including NaHMDS in benzene.³ Unaided by additional data, their suggestion that NaHMDS is tetrameric in benzene was suspect,^{2a,b} but the tactic had merit.

We find a highly predictable correlation of ^{15}N – ^{29}Si coupling with aggregation state. The high sensitivity, resolution, and quantitation offered by ^{29}Si NMR spectroscopy and the low cost of the ^{15}N label (7% the cost of an NMR tube) render this of potential interest to those studying M–




Figure 2. ^{29}Si NMR spectra of (a) ^{15}N NaHMDS (0.20 M) in toluene at $-80\text{ }^{\circ}\text{C}$ showing homodimer **1a**, (b) NaTMDS (0.20 M) in toluene at $-80\text{ }^{\circ}\text{C}$ showing homodimer **2a**, and (c) a 1:1 mixture of ^{15}N NaHMDS and NaTMDS (0.20 M total titer) in toluene recorded at $-80\text{ }^{\circ}\text{C}$ showing homo- and heterodimers **1a** and **3a**. (d) ^{29}Si NMR spectrum of 0.15 M ^{15}N NaHMDS in 0.75 M THF with Me_2NEt (DMEA) as cosolvent recorded at $-120\text{ }^{\circ}\text{C}$ shows dimer **1b** and monomer **4**.

Received: January 10, 2020




Figure 3. Job plot showing relative integrations of the ^{29}Si resonances of NaHMDs homodimer **1a** (red), NaTMDs-derived homodimer **2a** (blue), and heterodimer **3a** (green; eq 1) versus the measured¹⁶ mole fraction of NaHMDs (X_{NaHMDs}) at 0.20 total molarity¹⁷ in neat toluene- d_8 at -80°C .




Figure 4. DFT-computed, THF-complexed dimer **1a** and monomer **4**.

$\text{N}(\text{SiR}_3)_2$ and $\text{M}-\text{N}(\text{SiR}_3)(\text{R})$ derivatives. We present herein preliminary studies that focus on three prominent solvents: toluene, THF, and dioxane.

A $^{29}\text{Si}\{^1\text{H}\}$ INEPT experiment circumvented the problems posed by background signal from glass NMR tubes and NOE effects,⁶ allowing the relative integrations to be ascertained. ^{15}N NaHMDs was prepared in 45% overall yield from $^{15}\text{NH}_4\text{Cl}$ by an optimized protocol.⁷ ^1H and ^{13}C NMR spectra are archived [Supporting Information](#). Density functional theory (DFT) computations were carried out at the M06-2X level of theory^{8,9} for geometry optimizations and single-point calculations.¹⁰ The standard Def2-SVP basis set was used for geometry optimizations, and the expanded basis set Def2-TZVP for single point calculations.^{11,12}

The high solubility of NaHMDs in toluene and insolubility in hexane implicated an explicit π complexation, as observed




Figure 5. ^{29}Si chemical shift (green) and $^{15}\text{N}-^{29}\text{Si}$ coupling constants (black) plotted versus $[\text{THF}]$ in 2:1 pentane/toluene as cosolvent measured at -20°C . The curves are fits to a model based on an $\text{A}_2\text{S}_2-\text{AS}_4$ equilibrium ([Supporting Information](#)).




Figure 6. $^{15}\text{N}-^{29}\text{Si}$ coupling constants plotted versus $[\text{THF}]$ in 2:1 pentane/toluene as cosolvent measured at -20°C (black), 20°C (green), and 50°C (blue). The functions are fits to a model based on an $\text{A}_2\text{S}_2-\text{AS}_4$ equilibrium ([Supporting Information](#)).

crystallographically for many metals, including sodium.¹³ The ^{29}Si NMR spectrum shows a doublet with $^1J_{\text{N}-\text{Si}} = 7.9$ Hz akin to that for the benzene solvate noted by Lukevics, which proves to be characteristic of dimeric NaHMDs ([Figure 1](#)).

The dimer assignment was secured using the method of continuous variations (MCV).¹⁴ Mixtures of NaHMDs and the structurally similar sodium tetramethyldisilazide $[\text{NaTMDs}; (\text{HMe}_2\text{Si})_2\text{NNA}]$ ¹⁵ in toluene show two homodimers along with a heterodimer manifesting a ^{29}Si singlet corresponding to the TMDs fragment and a doublet with coupling characteristic of dimers ($^1J_{\text{N}-\text{Si}} = 8.9$ Hz; [Figure 2c](#)). Plotting the proportions of homo- and heteroaggregates versus measured¹⁶ mole fraction of NaHMDs (X_{NaHMDs}) affords the Job plot in [Figure 3](#). Quantitative heterodimerization is supported computationally (eq 1) and presumably derives from steric relief in the NaHMDs homodimer **1**. An analogous Job plot is obtained in toluene with 5.0 equiv of THF, conditions in which THF quantitatively displaces toluene to




Figure 7. ^{15}N – ^{29}Si coupling constants plotted versus [THF] (black) and [1,4-dioxane] (green) in 2:1 pentane/toluene as cosolvent measured at 20 °C. The functions are fit to a model based on an A_2S_2 – AS_4 equilibrium (Supporting Information).

form **1b** as shown by titrations.¹³ Dimer **1b** has been characterized crystallographically.^{2c}

Titration of solutions of [^{15}N]NaHMDS (0.15 M) in DMEA to record ^{29}Si NMR spectra with added THF at –120 °C reveals a markedly upfield-shifted resonance displaying a large coupling ($^1J_{\text{N-Si}} = 13.4$ Hz, Figure 2d) characteristic of NaHMDS monomers (Figure 4). The monomer becomes the sole observable form by 10 equiv (1.50 M).

Exchange of free and sodium-bound THF is rapid even at –120 °C. Although computations are supportive of both di- and tetrasolvated dimer, a host of other monodentate solvents show desolvation to be the norm.⁵ Couplings and chemical shifts are proxies for aggregation even in the high-temperature, rapid-exchange limit as illustrated in Figure 5. Figure 6 shows the temperature dependence of the deaggregation, revealing the anticipated preference for monomer at lower temperatures. Moreover, the curves in Figures 5 and 6 result from fits according to the equilibrium in Figure 4 with tetrasolvated monomer **4**. Analogous fits assuming a trisolvated monomer are decidedly inferior.

The correlation of coupling constant to aggregation state can also be used to assign dimer–monomer ratios in solvents that eluded assignment at lower temperatures. The prominent ethereal solvent 1,4-dioxane produced a highly insoluble white crystalline material at low temperature probably owing to a polymeric network of monomers characterized crystallographically,¹⁸ precluding solution structural studies. ^{29}Si spectra in the rapid exchange limit at 20 °C show dioxane-concentration-dependent coupling consistent with partial deaggregation of dimer **5** to monomer **6** at 20 °C. In neat dioxane, the 10.7 Hz coupling indicates that approximately

50% of the titer derives from monomer **6** (Figure 7). DFT computations indicate monomer **6** is only trisolvated.

We have shown that chemical shift and ^{15}N – ^{29}Si coupling for the dimers (7.5–8.5 Hz) and monomers (13.0–13.5 Hz), in conjunction with results from a much more broadly based study,⁵ are highly diagnostic of aggregation state. From a single spectrum, even at ambient temperatures, one can assess the relative proportions of monomers and dimers. Ironically, in over a dozen papers describing the structure of lithium hexamethyldisilazide using ^{15}N -labeled substrate in >100 solvents, we did not record a single ^{29}Si NMR spectrum: we did not need them. Belatedly, we find the analogous [^{15}N]LiHMDS/THF dimer and monomer ^{15}N – ^{29}Si couplings are 7.0 and 11.7 Hz, respectively. We suspect that other organometallic complexes with silazide-based ligands are likely to show diagnostic trends as well.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.0c00331>.

Spectroscopic, computational, and MCV data (PDF)

■ AUTHOR INFORMATION

Corresponding Author

David B. Collum – Baker Laboratory, Cornell University, Ithaca, New York 14853-1301, United States; orcid.org/0000-0001-6065-1655; Email: dbc6@cornell.edu

Author

Ryan A. Woltornist – Baker Laboratory, Cornell University, Ithaca, New York 14853-1301, United States

Complete contact information is available at:

<https://pubs.acs.org/10.1021/jacs.0c00331>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the National Institutes of Health (GM131713) for support.

■ REFERENCES

- (1) Watson, B. T.; Lebel, H. Sodium hexamethyldisilazide. In *e-EROS Encyclopedia of Reagents for Organic Synthesis*; John Wiley & Sons: New York, 2005; pp 1–10.
- (2) (a) Knizek, J.; Krossing, I.; Noth, H.; Schwenk, H.; Seifert, T. Synthesis and Structures of Sodium Phenylhydrazides. *Chem. Ber.* **1997**, *130*, 1053. (b) Driess, M.; Pritzkow, H.; Skipinski, M.; Winkler, U. Synthesis and Solid State Structures of Sterically Congested Sodium and Cesium Silyl(Fluorosilyl)Phosphanide Aggregates and Structural Characterization of the Trimeric Sodium Bis-(Trimethylsilyl)Amide. *Organometallics* **1997**, *16*, S108. (c) Karl, M.; Seybert, G.; Massa, W.; Harms, K.; Agarwal, S.; Maleika, R.; Stelter, W.; Greiner, A.; Neumüller, W. H.; Dehnicke, K. Amidometallate von Seltenerdelementen. Synthese Und Kristallstrukturen von $[\text{Na}(12\text{-Krone-4})_2][\text{M}\{\text{N}(\text{SiMe}_3)_2\}_3(\text{OSiMe}_3)]$ ($\text{M} = \text{Sm}, \text{Yb}$), $[\text{Na}(\text{THF})_3\text{Sm}\{\text{N}(\text{SiMe}_3)_2\}_3(\text{C}\equiv\text{C-Ph})]$, $[\text{Na}(\text{THF})_6][\text{Lu}_2(\mu\text{-NH}_2)(\mu\text{-NSiMe}_3)\{\text{N}(\text{SiMe}_3)_2\}_4]$ sowie von $[\text{NaN}(\text{SiMe}_3)_2(\text{THF})]_2$. *Anwendungen Der. Z. Anorg. Allg. Chem.* **1999**, *625*, 1301. (d) Ojeda-Amador, A. I.; Martínez-Martínez, A. J.; Kennedy, A. R.; Armstrong, D. R.; O'Hara, C. T. Monodentate Coordination of the Normally Chelating Chiral Diamine (*R,R*)-TMCD. *Chem. Commun.* **2017**, *53*, 324. (e) Forbes, G. C.; Kennedy, A. R.; Mulvey, R. E.; Rodger, P. J. A.

TEMPO: A Novel Chameleonic Ligand for s-Block Metal Amide Chemistry. *Chem. Commun.* **2001**, 1, 1400. (f) Neufeld, R.; Michel, R.; Herbst-Irmer, R.; Schöne, R.; Stalke, D. Introducing a Hydrogen-Bond Donor into a Weakly Nucleophilic Brønsted Base: Alkali Metal Hexamethyldisilazides (MHMDS, M = Li, Na, K, Rb and Cs) with Ammonia. *Chem. - Eur. J.* **2016**, 22, 12340. (g) Goodwin, C. A. P.; Smith, A.; Ortu, F.; Vitorica-Yrezabal, I. J.; Mills, D. P. Salt Metathesis versus Protonolysis Routes for the Synthesis of Silylamide Hauser Base (R_2NMgX ; X = Halogen) and Amido-Grignard (R_2NMgR) Complexes. *Dalt. Trans.* **2016**, 45, 6004. (h) Schüler, P.; Görls, H.; Westerhausen, M.; Kriek, S. Bis(trimethylsilyl)amide Complexes of s-Block Metals with Bidentate Ether and Amine Ligands. *Dalton Trans* **2019**, 48, 8966. (i) A molecular weight measurement of NaHMDS in Et_2O implicated dimer: Wannagat, U. N-Metallated Silicon-Nitrogen Derivatives: Preparation, Structure, and Reactions. *Pure Appl. Chem.* **1969**, 19, 329.

(3) Kupce, E.; Lukevics, E. Silicon-29-Nitrogen-15 Spin-Spin Coupling Constants in Silazanes. *Organometallics* **1988**, 7, 1649.

(4) Luo, G.; Luo, Y.; Qu, J. Direct Nucleophilic Trifluoromethylation Using Fluoroform: A Theoretical Mechanistic Investigation and Insight into the Effect of Alkali Metal Cations. *New J. Chem.* **2013**, 37, 3274.

(5) Woltornist, R. A.; Collum, D. B. Unpublished.

(6) Uhlig, F. D.; Marzmann, H. C. Silicon-29 NMR Some Practical Aspects in *Gelest Catalog Silicon Compounds: Silanes & Silicones Aufl.* (2003) 208. Morrisville, PA, USA, self-published.

(7) (a) [^{15}N]Hexamethyldisilazane was prepared from [^{15}N]NH $_3$ ¹⁹ by mixing [^{15}N]ammonium chloride (3.0 g, 55.0 mmol, >99% ^{15}N isotopic purity) with 6.00 g (0.15 mol) of granular NaOH in a 25 mL in a one-neck, round-bottom flask cooled to $-78\text{ }^{\circ}C$. The mixture was warmed with a heat gun for approximately 20 min. After the transfer of ammonia was complete, 1-(trimethylsilyl)imidazole (14.7 g, 15.3 mL, 105 mmol, 98% purity) was added to ammonia at $-78\text{ }^{\circ}C$ and stirred. HCl begins to off-gas, and imidazole precipitates immediately. Anhydrous diethyl ether (20 mL) is then added to the flask, and the mixture is held at $0\text{ }^{\circ}C$ for 40 min. Cholesterol (3.0 g) was added to the [^{15}N]hexamethyldisilazane with stirring for 45 min to remove excess 1-(trimethylsilyl)imidazole. Short-path distillation at atmospheric pressure removed the diethyl ether. Vacuum distillation (40 mm Hg, $20\text{ }^{\circ}C$) afforded 4.75 mL (47% yield) of $(Me_3Si)_2^{15}NH$. (b) [^{15}N]-Sodium hexamethyldisilazide (**1**) was prepared by a known procedure as follows.²⁰ Sliced sodium metal (1.20 g, 52.4 mmol) was added to a flame-dried, fine-mesh swivel frit setup (Supporting Information) in the glovebox. The apparatus was moved to a Schlenk line for the remainder of the procedure. Under an argon atmosphere, [^{15}N]HMDS (7.31 g, 9.45 mL, 45.0 mmol) and 40 mL of DMEA were added to the reaction flask at room temperature. Isoprene (2.62 mL, 26.2 mmol) dissolved in 8 mL of dry DMEA was then added over 1 h via syringe pump to the mixture. After addition of isoprene, the reaction was stirred at rt for an additional 2 h. The solution was then filtered through the frit and canula transferred to a second coarse swivel frit setup, and the solution was evaporated to dryness under vacuum for at least 10 h to yield a white powder. The powder was then suspended in dry pentane ($\sim 20\text{ mL}$), stirred for 1 h, and filtered. Washing with an additional 20 mL of pentane yielded 6.70 g (91% yield) of [^{15}N]NaHMDS, which was transferred to a glovebox and stored at room temperature. It should be noted that NaHMDS can be recrystallized as described previously²⁰ but with no detectable improvement. 1H NMR spectrum (500 MHz, toluene- d_8) δ 0.160 ppm; ^{13}C NMR spectrum (125.72 MHz, toluene- d_8) δ 7.28 ppm; ^{29}Si NMR spectrum (99.36 MHz, toluene- d_8) δ -14.6 ppm. (c) NaTMDs (**2**) was prepared using the same method described for NaHMDS. ^{13}C NMR spectrum (125.72 MHz, toluene- d_8) δ 5.93 ppm; ^{29}Si NMR spectrum (99.36 MHz, toluene- d_8) δ -25.57 ppm.

(8) Zhao, Y.; Truhlar, D. G. The M06 Suite of Density Functionals for Main Group Thermochemistry, Thermochemical Kinetics, Noncovalent Interactions, Excited States, and Transition Elements: Two New Functionals and Systematic Testing of Four M06-Class

Functionals and 12 Other Functionals. *Theor. Chem. Acc.* **2008**, 120, 215.

(9) In a number of organosodium studies we have noted that computations at the MP2 level of theory can fail to afford minima or single-point energies on what seem like pedestrian structures. This has not been a problem using the M06 functionals.

(10) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Zakrzewski, V. G.; Montgomery, J. A., Jr.; Stratmann, R. E.; Burant, J. C.; Dapprich, S.; Millam, J. M.; Daniels, A. D.; Kudin, K. N.; Strain, M. C.; Farkas, O.; Tomasi, J.; Barone, V.; Cossi, M.; Cammi, R.; Mennucci, B.; Pomelli, C.; Adamo, C.; Clifford, S.; Ochterski, J.; Petersson, G. A.; Ayala, P. Y.; Cui, Q.; Morokuma, K.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Cioslowski, J.; Ortiz, J. V.; Baboul, A. G.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Gomperts, R.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Gill, A.; Nanayakkara, C.; Gonzalez, M.; Challacombe, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Andres, J. L.; Gonzalez, C.; Head-Gordon, M.; Replogle, E. S.; Pople, J. A. *Gaussian 09*, Revision A.02; Gaussian, Inc.: Wallingford, CT, 2009.

(11) Weigend, F.; Ahlrichs, R. Balanced Basis Sets of Split Valence, Triple Zeta Valence and Quadruple Zeta Valence Quality for H to Rn: Design and Assessment of Accuracy. *Phys. Chem. Chem. Phys.* **2005**, 7, 3297.

(12) Legault, C. Y. CYLview, 1.0b; Université de Sherbrooke: 2009 (<http://www.cylview.org>).

(13) Yamada, S. Cation- π Interactions in Organic Synthesis. *Chem. Rev.* **2018**, 118, 11353.

(14) Renny, J. S.; Tomasevich, L. L.; Tallmadge, E. H.; Collum, D. B. Method of Continuous Variations: Applications of Job Plots to the Study of Molecular Associations in Organometallic Chemistry. *Angew. Chem., Int. Ed.* **2013**, 52, 11998.

(15) (a) Eppinger, J.; Herdtweck, E.; Anwender, R. Synthesis and Characterisation of Alkali Metal Bis(Dimethylsilyl) Amides: Infinite All-Planar Laddering in the Unsolvated Sodium Derivative. *Polyhedron* **1998**, 17, 1195. (b) Schneider, J.; Popowski, E.; Reinke, H. Darstellung, Charakterisierung und Reaktionsverhalten von Natrium- und Kaliumhydrosilylamiden $R_2(H)Si-N(M)R'$ (M = Na, K) - Kristallstruktur von $[(Me_3C)_2(H)Si-N(K)SiMe_3]_2 \cdot THF$. *Z. Anorg. Allg. Chem.* **2003**, 629, 55.

(16) The measured mole fraction—the mole fraction within only the ensemble of interest—eliminates the distorting effects of impurities. This problem has been highlighted: Hibbert, D. B.; Thordarson, P. The Death of the Job Plot, Transparency, Open Science and Online Tools, Uncertainty Estimation Methods and Other Developments in Supramolecular Chemistry Data Analysis. *Chem. Commun.* **2016**, 52, 12792.

(17) The concentration of NaHMDS, although expressed in units of molarity, refers to the concentration of the monomer subunit (normality).

(18) (a) Edelmann, F. T.; Pauer, F.; Wedler, M.; Stalke, D. Preparation and Structural Characterization of Dioxane Coordinated Alkali Metal Bis(Trimethylsilyl)Amides. *Inorg. Chem.* **1992**, 31, 4143.

(b) For an extensive review on the chemistry of the alkali metal amides, see: Mulvey, R. E.; Robertson, S. D. Synthetically Important Alkali-Metal Utility Amides: Lithium, Sodium, and Potassium Hexamethyldisilazides, Diisopropylamides, and Tetramethylpiperidides. *Angew. Chem., Int. Ed.* **2013**, 52, 11470.

(19) Mailyan, A. K.; Chen, J. L.; Li, W.; Keller, A. A.; Sternisha, S. M.; Miller, B. G.; Zakarian, A. Short Total Synthesis of [$^{15}N_5$]-Cylindrospermopsins from $^{15}NH_4Cl$ Enables Precise Quantification of Freshwater Cyanobacterial Contamination. *J. Am. Chem. Soc.* **2018**, 140, 6027.

(20) Tomasevich, L. L.; Collum, D. B. Method of Continuous Variation: Characterization of Alkali Metal Enolates Using 1H and ^{19}F -NMR Spectroscopies. *J. Am. Chem. Soc.* **2014**, 136, 9710.