NOVEL SYNTHESIS OF ENOL ESTERS THROUGH SILVER-CATALYZED REACTION OF ACETYLENIC COMPOUNDS WITH CARBOXYLIC ACIDS

Yoshio ISHINO*, Ikuzo NISHIGUCHI, Satoshi NAKAO, and Tsuneaki HIRASHIMA Osaka Municipal Technical Research Institute, 2-1-1, Ogimachi, Kita-ku Osaka 530

It was found that the reactions of 1,4-diacetoxy-2-butyne and related acetylenic compounds with carboxylic acids are efficiently catalyzed by silver salts to give the corresponding enol acetates in good yield.

Much attention has been focused on useful application of 2-butyne-1,4-diol and its derivatives to organic synthesis, and on extensive utilization in industry.¹⁾ For example, we have already reported a facile synthesis of 2,3-diaryl-1,3-butadienes from the reaction of 1,4-dimethoxy-2-butyne with Grignard reagents in the presence of copper(I) salts.²⁾

In this communication, we wish to describe a versatile method for synthesis of enol esters through silver-catalyzed addition of carboxylic acids to 1,4-diacetoxy-2-butyne and related acetylenic compounds.

The hitherto known methods³⁻⁶⁾ for the addition of carboxylic acid to a carboncarbon triple bond have not been so useful in organic synthesis owing to low yield, troublesome procedure, or employment of poisonous reagents.

The present method for transformation of acetylenic alcohols to the correspond-

ing enol esters is characterized by simple procedure, unique pattern of the reaction and good yield. A typical procedure is as follows: To a solution of 0.01 mole of 1,4-diacetoxy-2-butyne(<u>1a</u>) in 10ml of acetic acid(2a) was added 0.05 g(0.00018 mole) of silver carbonate in smallportions during 1/2 hr at 80 °C with stirring under nitrogen atmosphere. After the addition, the mixture was stirred at 110 °C for 6 hr. Then the solvent was evaporated out under reduced pressure, and distillation or/and column chromatography of the organic residue gave 1,2,4-triacetoxy-2-butene (3a) as a single product in a 80 % yield: bp. 160 °C/27 mmHg. NMR(CCl₄, ppm) **§** =2.00 [d.(J=4.0 Hz), 6H], 2.18[s. 3H], 4.53-4.75[m. 4H], 5.73 and 5.69[t. (J=7.2 Hz), 1H]. MASS: m/e=170(M⁺-HOAc). IR(neat, cm⁻¹) 1739.

Catalytic activities of a variety of metal salts in the reaction <u>la</u> with <u>2a</u> are summarized in Table 1. It is interesting that silver carbonate shows the best reactivity among them. While mercury acetate is known as common catalyst for usual electrophilic reactions of ethynyl bond³⁻⁶⁾, treatment of <u>la</u> with mercury acetate under the similar conditions as reported in the literature gave <u>3a</u> only in a poor yield. Also as is shown in Table I, 0.05 g of silver carbonate(0.0181 eq. mole based on <u>la</u>) was enough to promote the present reaction.

Metal Salt	Amount(g)*	Yield of <u>3a</u> (%)
none	_	0
Ag ₂ CO ₃	0.01(0.0036)	65
Ag ₂ CO ₃	0.05(0.0181)	80
Ag ₂ CO ₃	0.10(0.0363)	73
Ag_2CO_3	0.50(0.1813)	72
AgOAc	0.05(0.0300)	53
AgNO ₃	0.05(0.0294)	49
Ag ₂ SO ₄	0.50(0.1604)	3
Hg(OAc) ₂	0.50(0.1569)	26
CuC1	0.50(0.0551)	0

Table 1Catalytic Activities of a Variety of Metal Salts in
the Reaction of lawith 2a

* The numbers in the parentheses show equivalent mole of each metal salt based on <u>la</u>. Under the similar conditions, the addition of a variety of carboxylic acids to <u>la</u> and related acetylenic compounds easily took place to give the corresponding enol esters in good yields(Table 2). It may be quite noteworthy that the silver-catalyzed reaction also successfully proceeded with dimethyl acetylenedicarboxylate in which

Chemistry Letters, 1981

the ethynyl bond were substituted by typical electron-withdrawing groups. The mixture of enol acetates $\underline{4}$ and $\underline{5}$ (ratio of $\underline{4}/\underline{5}$ = ca. 2.0) were obtained from the reaction of $\underline{1d}$ with acetic acid, indicating some regioselectivity of the reaction.

HC≡CCH ₂ OH	+ сн ₃ соон	$\xrightarrow{110 \ ^{\circ}C}_{Ag} H_{3}COOCH=CHCH_{2}OCOCH_{3}$	+ CH ₂ =CCH ₂ OCOCH ₃
<u>1d</u>	<u>2a</u>	<u>4</u>	5

	R ¹ -	$C \equiv C - R^2 (1) R^2$	к ³ соон к ³	(<u>2</u>)	(R ¹ -(isolat	COR^3 C=CH-R ² (<u>3</u>) ted yield(%)	Stereoisomeric ratio ^{*1}
<u>1a</u>	H ₃ CCOOCH ₂	H ₃ CCOOCH ₂	CH ₃	(<u>2a</u>)		80	8/2
			C ₂ H ₅	(<u>2b</u>)		83	8/2
			n-C ₃ H ₅	(<u>2c</u>)		94	7/3
			C ₆ H ₅	(<u>2d</u>)		53	6/4
<u>1b</u>	H ₃ COCH ₂	H ₃ COCH ₂	CH ₃	(<u>2a</u>)		49	9/1
			C ₂ H ₅	(<u>2b</u>)		55	9/1
			$C_{6}^{H_{5}}$	(<u>2d</u>)		69	9/1
<u>1c</u>	н ₃ соос	H ₃ COOC	CH ₃	(<u>2a</u>)		60	9/1
<u>1d</u>	Н	HOCH ₂	CH ₃	(<u>2a</u>)		63 ^{*2} , ³	6/4 ^{*7}
<u>le</u>	C ₆ H ₅	HOCH ₂	CH ₃	(<u>2a</u>)		54 ^{*2,4}	9/1
$\underline{1f}$	C ₆ H ₅	Н	CH ₃	(<u>2a</u>)		28 ^{*5}	-
<u>1g</u>	C ₆ H ₅	с ₂ н ₅	CH ₃	(<u>2a</u>)		0 ^{*6}	-

Table 2The Silver-Catalyzed Reaction of Acetylenic Compounds withCarboxylic Acids

*1 The stereoisomeric ratio for each product was determined by gas chromatographic technique.⁷⁾

- *2 All the hydroxyl groups of 3 were acetylated.
- *3 The isomeric mixture of 4 and 5.
- *4 Based on the comsumed starting acetylene(conversion: 53 %).
- *5 Acetophenone was also formed in a 23 % yield.
- *6 Propiophenone was formed in a 60 % yield.
- *7 The ratio for 4.

It has been reported that an acyloxyallene $\underline{7}$ was obtained from the similar silver-catalyzed reaction of an acetylenic tertiary alcohols $\underline{6}$ with acetic acid⁹⁾, which may give a distinct contrast with the present result.¹⁰⁾

$$HC=C-C-OH + CH_{3}COOH \xrightarrow{reflux} CH_{3}COOCH=C=C < \binom{R'}{R''}$$

$$\frac{6}{2}$$

Although the detailed mechanism of the present reaction does not seem simple,

it may be proposed as one of the most plausible pathways that initial electrophilic addition of a silver cation to a ethynyl bond gives a π -complex 8, which is successively transformed to the corresponding σ -complex 9 by electrophilic attack of a carboxylate anion. The silver complex 9 may be so reactive as to be decomposed to an enol ester 3 by substitution of the silver atom with a proton, while the corresponding mercury¹¹) or thallium complex¹²) was isolated. Therefore, a catalytic amount of a silver salt may be enough effective to promote the esterification of acetylenic compounds, which may be quite significant in organic synthesis.

 $R^{1}-C=C-R^{2} \xrightarrow{Ag^{+}} R^{1}-C=C-R^{2} \xrightarrow{R^{3}COO^{-}} R^{1}-C=C-R^{2} \xrightarrow{H^{+}} R^{1}-C=CH-R^{2}$ $\frac{1}{4g} \xrightarrow{8} \frac{9}{2} \xrightarrow{3}$

References and Notes

- 1) K.Weissermel, and H.J.Arpe, "Industrielle Organische Chemie", Verlarg Chemie GmbH, Weinheim, Germany, 1976, Charpter 6.
- 2) Yoshio Ishino, Ikuzo Nishiguchi, Fugio Takihira, and Tsuneaki Hirashima, Tetrahedron Lett., <u>1980</u>, 1527.
- 3) P.F.Hudrlik and A.M.Hudrlik, J.Org.Chem., <u>38</u>, 4254(1973).
- 4) H.Lemaire and H.J.Lucas, J.Am.Chem.Soc., <u>77</u>, 939(1955).
- 5) P.E.Peterson and J.E.Duddey, ibid., 88, 4990(1966).
- 6) K.Fujita, K.Yamamoto, and T.Shono, Tetrahedron Lett., 1973, 3865.
- 7) While the vinyl proton cis to the acetoxy function of enol acetates was reported to have its NMR signal at approximately 0.2 to 0.3 p.p.m. lower field,⁸⁾ the differences in the chemical shift(0.05-0.10 p.p.m.) of the vinyl proton in the NMR spectra of both stereoisomers of each product is not enough to make definite stereochemical assignment for either (E)- or (Z)-isomer.
- 8) H.O.House and V.Kramar, J.Org.Chem., <u>28</u>, 3362(1963).
- 9) G.Saucy, R.Marbet, H.Lindlar, and O.Isler, Helv.Chem.Acta, 42, 1945(1973).
- 10)Further study on these contrasting behaviours is in progress.
- 11)S.Uemura, H.Miyoshi, K.Sohma, and M.Okano, J.Chem.Soc., Chem.Comm., 1975, 548.

12)S.Uemura, K.Sohma, H.Tara, and M.Okano, Chem.Lett., 1973, 545.

(Received February 26, 1981)