

## Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

### AN IMPROVED PREPARATION OF 4-AMINO-3-MERCAPTOBENZOIC ACID

Robert C. Lang<sup>a</sup>, Craig M. Williams<sup>a</sup> & Mary J. Garson<sup>a</sup>

<sup>a</sup> Chemistry Department, School of Molecular and Microbial Sciences, The University of Queensland, St. Lucia, 4072, Queensland, AUSTRALIA E-mail:

Version of record first published: 11 Feb 2009

To cite this article: Robert C. Lang, Craig M. Williams & Mary J. Garson (2003): AN IMPROVED PREPARATION OF 4-AMINO-3-MERCAPTOBENZOIC ACID, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 35:5, 520-524

To link to this article: <http://dx.doi.org/10.1080/00304940309355864>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

10. K. Undheim and B. P. Nilsen, *Acta Chem. Scand. B*, **29**, 503 (1975).
11. S. K. Ghorai, *Ph.D Thesis, I.I.T. Kharagpur, (2001)*.
12. B. M. Choudary, M. L. Kantam, B. Bharathi and C. V. Reddy, *Synlett*, 1203 (1998).
13. N. K. Hamer, *J. Chem. Soc. Perkin I*, 1285 (1979).
14. G. B. Payne, *J. Org. Chem.*, **25**, 275 (1960).
15. K. L. Reed, J. T. Gupton and T. L. Solarz, *Synth. Commun.*, **19**, 3579 (1989).
16. J. Katsuhara, *J. Org. Chem.*, **32**, 797 (1967).
17. T. S. Straub, *Tetrahedron Lett.*, **36**, 663 (1995).
18. R. B. Garland, M. Miyano, D. Pireh, M. Clare, P. M. Finnegan and L. Swenton, *J. Org. Chem.*, **55**, 5854 (1990).

\*\*\*\*\*

### AN IMPROVED PREPARATION OF 4-AMINO-3-MERCAPTobenzoic ACID

*Submitted by*  
(04/04/03)

Robert C. Lang, Craig M. Williams\* and Mary J. Garson

*Chemistry Department, School of Molecular and Microbial Sciences  
The University of Queensland, St. Lucia, 4072, Queensland, AUSTRALIA  
E-Mail: c.williams3@mailbox.uq.edu.au*

4-Amino-3-mercaptobenzoic acid (**4**) is commonly used as a precursor of 6-carboxy-benzothiazole<sup>1</sup> which is heavily utilized by materials chemists<sup>2</sup> and to a lesser extent by medicinal chemists.<sup>3</sup> Two procedures have been reported for the synthesis of **4**; the first involves reaction of sodium sulfide<sup>3b,4</sup> with 4-amino-3-thiocyanatobenzoic acid (**2**), and the second<sup>1a</sup> requires hydrolysis of 1-amino-6-carboxybenzothiazole (**3**), derived from 4-aminobenzoic acid (**1**). Although, the later procedure<sup>1a</sup> can be conducted on large scale and has successfully been repeated by others (*Scheme 1*),<sup>2d,2e,3a</sup> it requires either catalytic<sup>3a</sup> or stoichiometric<sup>1a</sup> amounts of



Downloaded by [University of Saskatchewan Library] at 02:13 01 August 2012

Downloaded by [University of Saskatchewan Library] at 02:13 01 August 2012



Downloaded by [University of Saskatchewan Library] at 02:13 01 August 2012

Downloaded by [University of Saskatchewan Library] at 02:13 01 August 2012

## Downloaded by [University of Saskatchewan Library] at 02:13 01 August 2012

Downloaded by [University of Saskatchewan Library] at 02:13 01 August 2012

on a Hewlett-Packard 5890A chromatograph fitted with a DBS EC-5 Alltech column (30 m x 0.25 mm), coupled to a 5970 series mass spectrometer. Melting points were determined on a Fischer Johns Melting Point apparatus and are uncorrected. Column chromatography (Flash Silica gel 230–400 mesh, Merck) on silica gel was performed using distilled solvents.

**2-Aminobenzothiazole-6-carboxylic Acid Hydrochloride (3).**– The published procedure<sup>3a</sup> was followed. To a stirred suspension of sodium thiocyanate (13 g, 0.160 mol) and 4-aminobenzoic acid **3** (20 g, 0.146 mol) in MeOH (75 mL) was slowly added bromine (7.5 mL, 0.146 mol) while the temperature was maintained below –5°C. After addition, the mixture was stirred at –5°C for 2 h. The precipitated solid was collected and washed with water. The isolated solid was suspended in hydrochloric acid (1M, 70 mL), refluxed for 30 mins and then filtered while hot. Conc. hydrochloric acid (30 mL) was added to the filtrate producing a white solid. The solid was collected and dried under vacuum to afford **3** (11.5 g, 34%). The crude product was used without further purification. A sample was recrystallized from methanol, mp. >290°C (dec.). *Lit.*<sup>2d</sup> mp. 288–290°C (dec.).

<sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ 7.54 (d, 1H, H4, *J* = 8.5 Hz), 7.94 (dd, 1H, H5, *J* = 8.5, 1.6 Hz), 8.45 (d, 1H, H7, *J* = 1.6 Hz). <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>): δ 114.6, 124.4, 125.86, 125.91, 128.5, 145.2, 166.6, 170.1. Mass spectrum (EI) *m/z* (%): 194(M<sup>+</sup> -HCl, 100), 177 (82), 149 (27), 122 (29).

HRMS (ESI): Calcd for C<sub>8</sub>H<sub>7</sub>N<sub>2</sub>O<sub>2</sub>S [M+H]: 195.0228; Found: 195.0228.

**4-Amino-3-mercaptopbenzoic Acid Hydrochloride (4).**– A suspension of **3** (6.0 g, 0.026 mol) and potassium hydroxide (26.3 g, 0.469 mol) in rigorously degassed demineralized water (50 mL) was shielded from light (aluminium foil) and then refluxed under nitrogen (or argon) for 4 h. After cooling to room temperature, conc. hydrochloric acid (45 mL) was added dropwise under nitrogen. The resulting mixture was cooled in a cold-water bath (~–5°C) and stirred for 30 mins. The precipitate was collected under a nitrogen blanket and dried under vacuum (0.01 mmHg) to give the desired product as a white solid (4.2 g, 79%). Light- and oxygen-free storage (–5°C) is recommended.

<sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ 6.69 (d, *J* = 8.4 Hz), 6.74 (d, 1H, H5, *J* = 8.4 Hz), 7.45 (d, 1H, H2, *J* = 1.6 Hz), 7.52 (dd, *J* = 8.4, 1.6 Hz), 7.61 (dd, 1H, H6, *J* = 8.4, 1.6 Hz), 7.76 (d, *J* = 1.6 Hz).

<sup>1</sup>H NMR [D<sub>2</sub>O (5%)/DMSO-*d*<sub>6</sub> (95%)]: δ 6.75 (d, 1H, H5, *J* = 8.6 Hz), 7.41 (d, 1H, H2, *J* = 2.1 Hz), 7.61 (dd, 1H, H6, *J* = 8.6, 2.1 Hz). <sup>13</sup>C NMR [D<sub>2</sub>O (5%)/DMSO-*d*<sub>6</sub> (95%)]: δ 114.3, 115.2, 117.8, 133.2, 138.5, 153.8, 167.0.

All attempts to recrystallize the title compound resulted in the formation of *bis*-(2-amino-3-mercaptopbenzoic acid) disulfide hydrochloride, mp. 284–286°C (dec.) (methanol). *Lit.*<sup>7</sup> mp. 285°C (dec.).

<sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ 6.28 (s, 2H, NH), 6.74 (d, 1H, H5, *J* = 8.6 Hz), 7.47 (d, 1H, H2, *J* = 2.0 Hz), 7.62 (dd, 1H, H6, *J* = 8.6, 2.0 Hz), 12.15 (bs, 1H, CO<sub>2</sub>H). <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>): δ 113.8 (C-5), 114.8 (C-3), 117.5 (C-1), 132.7 (C-6), 138.0 (C-2), 153.6 (C-4), 166.5 (–CO<sub>2</sub>H). Mass spectrum (EI) *m/z* (%): 336 (M<sup>+</sup>-2HCl, 28), 194 (36), 177 (18), 169 (55), 168 (100), 152 (28), 124 (46).

HRMS (ESI): Calcd. for C<sub>14</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>S<sub>2</sub>Na [M+Na]: 359.0136; Found: 359.0132.

**Methyl 4-Amino-3-methylsulfanybenzoate (5).**- To a solution of **1** (15 mg, 0.073 mmol) in MeOH (0.3 mL) was added a solution of diazomethane in diethyl ether at room temperature until effervescence ceased (performed in fumehood). The unreacted diazomethane was removed by flushing the solution with nitrogen gas, and the solvents were evaporated under reduced pressure affording a yellow colored oily residue. Column chromatography (ethyl acetate/hexane; 1:4) on silica gel gave the title compound (12mg, 83%) as a white solid, mp. 58-60°C.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  2.36 (s, 3H,  $\text{SCH}_3$ ), 3.84 (s, 3H,  $\text{CO}_2\text{CH}_3$ ), 6.66 (d, 1H, H5,  $J = 8.4$  Hz), 7.75 (dd, 1H, H6,  $J = 8.4, 2.0$  Hz), 8.04 (d, 1H, H2,  $J = 2.0$  Hz).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  17.8, 51.7, 113.6, 119.5, 120.1, 130.8, 135.5, 151.0, 166.7. GC/MS (EI)  $m/z$  (%): 197 ( $\text{M}^+$ , 100), 182 (53), 166 (87), 151 (16), 138 (12), 123 (14).

*Anal.* Calcd. for  $\text{C}_9\text{H}_{11}\text{NO}_2\text{S}$ : C, 54.80; H, 5.62; N, 7.10;  $\text{M}^+$  197.0510. Found: C, 54.77; H, 5.64; N, 7.01; 197.0503.

**bis-(Methyl 4-amino-3-mercaptopbenzoate) Disulfide (6).**- The same procedure to that directly above afforded the title compound as a yellow solid in near quantitative yield, mp. 166-168°C. All attempts at obtaining an analytical sample failed.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  3.77 (s, 3H,  $\text{CO}_2\text{CH}_3$ ), 6.68 (d, 1H, H5,  $J = 8.4$  Hz), 7.74 (d, 1H, H2,  $J = 2.0$  Hz), 7.82 (dd, 1H, H6,  $J = 8.4, 2.0$  Hz).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  51.7, 114.3, 117.3, 119.7, 133.4, 139.2, 152.2, 166.2. Mass spectrum (EI)  $m/z$  (%): 364 ( $\text{M}^+$ , 28), 197 (6), 183 (39), 182 (100), 152 (33), 124 (14).

HRMS (ESI): Calcd. for  $\text{C}_{16}\text{H}_{17}\text{N}_2\text{O}_4\text{S}_2$  [ $\text{M}+\text{H}$ ]: 365.0630; Found: 365.0633.

## REFERENCES

1. a) M. Schubert, *Ann.*, **558**, 10 (1947); b) K. R. Nivalkar and S. H. Mashraqui, *Synth. Commun.*, **26**, 3535 (1996); c) S. H. Mashraqui, S. Kumar and K. R. Nivalkar, *Heterocyclic Commun.*, **7**, 73 (2001).
2. See for example: a) D. J. Perettie, P. E. Pierini and T. McCarthy, PCT Int. Appl. WO 9319461 and 9319462 (1993) (*Chem. Abstr.*, **120**, 234558 and 234557); b) T. T. Tasi, F. E. Arnold and W. F. Hwang, *J. Polym. Sci., Part A: Polym. Chem.*, **27**, 2839 (1989); c) S. Kitamura, H. Kitano and F. Tanimoto, Jpn. Kokai Tokkyo Koho JP 01040530 (1989) (*Chem. Abstr.*, **111**, 174918); d) A. W. Chow, S. P. Bitler, P. E. Penwell and J. F. Wolfe, *Macromolecules*, **22**, 3514 (1989) and references therein; e) P. M. Hergenrother and H. H. Levine, *J. Polym. Sci., Part A-1*, **6**, 2939 (1968).
3. a) H. Matsuoka, N. Ohi, M. Mihara, H. Suzuki, K. Miyamoto, N. Maruyama, K. Tsuji, N. Kato, T. Akimoto, Y. Takeda, K. Yano and T. Kuroki, *J. Med. Chem.*, **40**, 105 (1997); b) T. Kosaka, K. Ochiai, S. Ohba, T. Wakabayashi and S.-i. Murota, *Yakugaku Zasshi*, **117**, 611 (1997) (*Chem. Abstr.*, **127**, 326160); c) S. Singh, R. S. Narang, M. Singh and H. Singh, *Indian J. Chem.*, **6**, 12 (1968).
4. A. I. Kiprianov and I. L. Mushkalo, *J. Gen. Chem. USSR*, **32**, 3967 (1962).

5. It should be noted that both procedures (Chow *et al.*<sup>2d</sup> and Matsuoka *et al.*<sup>3a</sup>) are identical.
6. A. Adeeko, D. Li, D. S. Forsyth, V. Cassey, G. M. Cooke, J. Barthelemy, D. G. Cyr, J. M. Trasler, B. Robaire and B. F. Hales, *Toxicol. Sci.*, **74**, 407 (2003).
7. H. Wojahn and I. Lerch, *Arzneim. Forsch.*, **2**, 455 (1952).

\*\*\*\*\*


### IMPROVED SYNTHESIS OF 1,3-DIAZA-6-ADAMANTANONE

Submitted by Timothy A. Lewis\* and Gurmit Grewal  
(08/11/03)

UCB Research, Inc.

840 Memorial Drive, Cambridge, MA 02139

1,3-Diaza-6-adamantanone was first synthesized by oxidation of 1,3-diaza-6-adamantanol.<sup>1</sup> The alcohol synthesis<sup>2</sup> (6 steps, 10%) was later shortened (3 steps, 8%) and the oxidation yield improved.<sup>3</sup> We have synthesized the ketone directly, improving the overall yield (3 steps, 35%) while avoiding the lachrymators and thiols used previously.



a) Benzylamine,  $(\text{CH}_2\text{O})_n$ , HOAc/HCl/MeOH (69%), ref. 4; b)  $\text{H}_2$ ,  $\text{Pd}(\text{OH})_2/\text{C}$ , HOAc/MeOH;  
(c)  $\text{CH}_2\text{O}$  (aq) (51 %, two steps)

Mannich condensation of benzylamine, paraformaldehyde, and *N*-benzyl-4-piperidone was performed using a literature procedure.<sup>4</sup> Hydrogenolysis of the *bis*-benzylamine product proceeded slowly under atmospheric pressure. Cyclization of the crude product with aqueous formaldehyde gave 1,3-diaza-6-adamantanone, which was isolated by hot filtration.

### EXPERIMENTAL SECTION

Mps were determined on a Mel-temp apparatus and are uncorrected. All reactions were run under an argon atmosphere except where noted. All solvents and reagents were purchased from Aldrich or VWR and used without purification, except for benzylamine and *N*-benzyl-4-piperi-