

# Preparation of 2-(*N,N*-dialkylamino)-2,4,6-cycloheptatrien-1-one derivatives from an isocyanide, a primary amine, propionaldehyde, and tropolone via Ugi-Smiles coupling reaction in the presence of silica nanoparticles

Abdolhossain MASSOUDI<sup>1</sup>, Issa AMINI<sup>1</sup>, Ali RAMAZANI<sup>2,\*</sup>,  
Fatemeh ZEINALI NASRABADI<sup>3</sup>

<sup>1</sup>*Department of Chemistry, Payame Noor University, P.O. Box 19395-3697, Tehran-IRAN*

<sup>2</sup>*Department of Chemistry, Zanzan Branch, Islamic Azad University,  
P.O. Box 49195-467, Zanzan-IRAN*

*e-mail: aliramazani@gmail.com*

<sup>3</sup>*Department of Chemistry, University of Zanzan, P.O. Box 45195-313, Zanzan-IRAN*

Received: 03.02.2012

The use of Smiles rearrangement in Ugi-type couplings with tropolone allows very straightforward multicomponent formation of 2-(*N,N*-dialkylamino)-2,4,6-cycloheptatrien-1-one derivatives. The 4-component reaction of isocyanides, tropolone (2-hydroxy-2,4,6-cycloheptatrien-1-one), primary amines, and propionaldehyde in the presence of silica nanoparticles (silica NPs, ca. 42 nm) proceeds easily in methanol to form the title compounds in a new Ugi-Smiles-type reaction.

**Key Words:** Ugi-type coupling, tropolone, isocyanides, primary amines, silica (NP), Ugi-Smiles-type reaction

## Introduction

Multicomponent reactions (MCRs) allow more than 2 simple and flexible building blocks to be combined in practical, timesaving one-pot operations. Due to their valued features such as atom economy, inherent simple

---

\*Corresponding author

experimental procedures, and their one-pot character, they are perfectly suited for automated synthesis.<sup>1</sup> Therefore, MCRs have attracted much attention because of their exceptional synthetic efficiency.<sup>2–5</sup> Since all the organic reagents employed are used and moved toward the target compound the purification of products resulting from MCRs is simple.<sup>6,7</sup> Isocyanide-based multicomponent reactions (abbreviated to IMCRs by Ugi and Dömling) have an advantageous position. The special features of IMCRs including unique synthetic potential, high atom economy, convergent nature, ease of implementation, and the molecular diversity generation are considered acceptable factors in the relative advantage of the reactions.<sup>6–13</sup>

In recent years, nanoparticles (NPs) have attracted tremendous attention in catalysis because of their improved efficiency under mild and environmentally benign conditions in the context of ecological (green) synthesis.<sup>14,15</sup> Due to their enormously large and highly reactive surface area, NPs have potential to exhibit superior catalytic activity in comparison to their bulk counterparts.<sup>16,17</sup> In the past, we established a one-pot method for the preparation of organic compounds.<sup>18–24</sup> As part of our ongoing program to develop efficient and robust methods for the synthesis of heteroatom-containing compounds,<sup>25–32</sup> we wish to report the preparation of a new class of 2-(*N,N*-dialkylamino)-2,4,6-cycloheptatrien-1-one derivatives **5a–h** by a novel 4-component condensation reaction of propionaldehyde **1**, primary amines **2**, isocyanides **3**, and tropolone **4** in the presence of silica NPs in excellent yields (Scheme 1).

## Experimental

The starting materials and solvents were obtained from Merck (Germany) and Fluka (Switzerland) and were used without further purification. The methods used to follow the reactions were TLC and NMR. TLC and NMR indicated that there were no side products. IR spectra were measured on a Jasco 6300 FTIR spectrometer. <sup>1</sup>H- and <sup>13</sup>C-NMR spectra (CDCl<sub>3</sub>) were recorded on a BRUKER DRX-400 AVANCE spectrometer at 400.22 and 100.63 MHz, respectively. Elemental analyses were performed using a Heraeus CHN-O-Rapid analyzer. Mass spectra were recorded on a FINNIGAN-MATT 8430 mass spectrometer operating at an ionization potential of 70 eV. Preparative layer chromatography (PLC) plates were prepared from Merck silica gel (F<sub>254</sub>) powder.

### General procedure for the preparation of compounds 5a-5h

To magnetically stirred silica NPs (0.2 g) and solution of propionaldehyde **1** (1 mmol), primary amine **2** (1 mmol), and tropolone **4** (1 mmol) in CH<sub>3</sub>OH (7 mL) was added dropwise a solution of isocyanide **3** (1 mmol) in CH<sub>3</sub>OH (2 mL) at 60 °C over 5 min. The reaction mixture was refluxed for 6 h. The solvent was removed under reduced pressure and the viscous residue was purified by preparative layer chromatography (silica gel (F<sub>254</sub>); petroleum ether–ethyl acetate (4:1)). The solvent was removed under reduced pressure and the products (**5a–h**) were obtained. The characterization data of the compounds are given below.

#### 2-[Benzyl(7-oxo-1,3,5-cycloheptatrienyl)amino]-N<sup>1</sup>-cyclohexylbutanamide (5a)

Yellow oil, (Yield: 92%). IR (neat):  $\nu = 3440, 2952, 1745, 1651, 1600, 1466, 1227, 1077 \text{ cm}^{-1}$ . <sup>1</sup>H-NMR (400.22 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 0.79 (t, 3H,  $J = 7.4 \text{ Hz}$ , CH<sub>3</sub>), 0.86–2.1 (m, 10H, 5CH<sub>2</sub> of cyclohexyl and m, 2H, CH<sub>2</sub> of propyl), 3.80–4.02 (m, 1H, CH–N of cyclohexyl), 4.50 (AB, 2H,  $J = 16.8 \text{ Hz}$ , PhCH<sub>2</sub>), 4.55 (t,

1H,  $J = 7.0$  Hz, CH), 6.53 (d, 1H,  $J = 10.4$  Hz, CH of tropone), 6.63 (t, 1H,  $J = 9.0$  Hz, CH of tropone), 6.84 (t, 1H,  $J = 10.2$  Hz, CH of tropone); 7.07-7.35 (m, 7H, CH of tropone and ph); 8.69 (d, 1H,  $J = 8.0$  Hz, NH exchanged by D<sub>2</sub>O addition). <sup>13</sup>C-NMR (100.63 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 11.74 (CH<sub>3</sub>), 22.35 (CH<sub>2</sub> of propyl), 24.81, 25.66, 33.11 (3CH<sub>2</sub> of cyclohexyl), 47.84 (CH of cyclohexyl), 49.56 (PhCH<sub>2</sub>), 63.55 (CH), 121.51, 126.58, 126.98, 127.22, 128.55, 133.83, 134.56, 136.56 (8 CH of tropone and Ph), 135.93 (C<sub>ip</sub> of Ph), 158.12 (C of tropone), 169.80 (C=O of amide), 184.38 (C=O of tropone). Analysis of C<sub>24</sub>H<sub>30</sub>N<sub>2</sub>O<sub>2</sub> (378.23). (% calculation/found): C: 76.16/76.12, H: 7.99/7.96, N: 7.40/7.45. MS,  $m/z$ (%): 378 (4), 287 (14), 252 (95), 210 (63), 188 (21), 160 (33), 148 (44), 132 (20), 106 (20), 105 (13), 91 (100), 77 (17), 65 (18), 57 (15), 55 (34), 41 (32).

### 2-[Benzyl(7-oxo-1,3,5-cycloheptatrienyl)amino]-N<sup>1</sup>- (tert-butyl)butanamide (5b)

Yellow oil, (Yield: 90%). IR (neat):  $\nu = 3423, 2922, 1742, 1636, 1620, 1461, 1234, 1030$  cm<sup>-1</sup>. <sup>1</sup>H-NMR (400.22 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 0.78 (t, 3H,  $J = 7.4$  Hz, CH<sub>3</sub>), 1.37 (s, 9H, CMe<sub>3</sub>), 1.70-2.10 (m, 2H, CH<sub>2</sub> of propyl), 4.50 (AB, 2H,  $J = 16.8$  Hz, PhCH<sub>2</sub>), 4.51 (t, 1H,  $J = 6.4$  Hz, CH), 6.51 (d, 1H,  $J = 10.4$  Hz, CH of tropone), 6.66 (t, 1H,  $J = 9.0$  Hz, CH of tropone), 6.85 (t, 1H,  $J = 10.2$  Hz, CH of tropone), 7.07-7.38 (m, 7H, CH of tropone and Ph), 8.65 (s, 1H, NH exchanged by D<sub>2</sub>O addition). <sup>13</sup>C-NMR (100.63 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 11.77 (CH<sub>3</sub>), 22.20 (CH<sub>2</sub> of propyl), 28.72 (3CH<sub>3</sub>), 49.43 (PhCH<sub>2</sub>), 50.67 (CMe<sub>3</sub>), 64.01 (CH), 121.25, 126.47, 126.91, 127.24, 128.57, 133.85, 134.37, 136.57 (8 CH of tropone and Ph), 135.91 (C<sub>ip</sub> of Ph), 158.18 (C of tropone), 170.02 (C=O of amide), 184.46 (C=O of tropone). Analysis of C<sub>22</sub>H<sub>28</sub>N<sub>2</sub>O<sub>2</sub> (352.22). (% calculation/found): C: 74.97/74.94, H: 8.01/8.05, N: 7.95/7.92. MS,  $m/z$ (%): 352 (14), 280 (10), 261 (25), 252 (100), 210 (67), 188 (24), 160 (56), 148 (14), 132 (41), 106 (15), 105 (20), 91 (89), 77 (36), 65 (26), 57 (48), 41 (37).

### N<sup>1</sup>-(tert-butyl)-2-[(7-oxo-1,3,5-cycloheptatrienyl)(propyl)amino]butanamide (5c)

Yellow oil, (Yield: 83%). IR (neat):  $\nu = 3444, 2922, 1746, 1650, 1608, 1469, 1256, 1084$  cm<sup>-1</sup>. <sup>1</sup>H-NMR (400.22 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 0.70 (t, 3H,  $J = 7.4$  Hz, CH<sub>3</sub> of (CH<sub>2</sub>)<sub>2</sub>CH<sub>3</sub>), 0.88 (t, 3H,  $J = 7.4$  Hz, CH<sub>3</sub> of CH<sub>2</sub>CH<sub>3</sub>), 1.39 (s, 9H, CMe<sub>3</sub>), 1.23-2.20 (m, 4H, 2CH<sub>2</sub>), 3.07-3.37 (m, 2H, NCH<sub>2</sub>), 4.35 (t, 1H,  $J = 7.0$  Hz, H-C), 6.66 (d, 1H,  $J = 10.4$  Hz, CH of tropone), 6.72 (t, 1H,  $J = 9.2$  Hz, CH of tropone), 7.00-7.30 (m, 3H, CH of tropone), 8.39 (s, 1H, NH exchanged by D<sub>2</sub>O addition). <sup>13</sup>C-NMR (100.63 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 11.46 and 11.68 (2 CH<sub>3</sub>), 19.06 and 21.85 (2CH<sub>2</sub>), 28.69 (3CH<sub>3</sub>), 47.01 (CH<sub>2</sub>N); 50.51 (CMe<sub>3</sub>), 64.07 (CH), 119.92, 125.98, 133.93, 134.12, 136.43 (5 CH of tropone), 158.61 (C of tropone), 170.21 (C=O of amide), 184.42 (C=O of tropone). Analysis of C<sub>18</sub>H<sub>28</sub>N<sub>2</sub>O<sub>2</sub> (304.22). (% calculation/found): C: 71.02/71.05, H: 9.27/9.25, N: 9.20/9.23. MS,  $m/z$ (%): 304 (28), 261 (10), 252 (91), 236 (15), 220 (42), 210 (31), 205 (98), 204 (100), 188 (13), 162 (85), 160 (33), 132 (28), 105 (13), 91 (24), 77 (14), 57 (10), 41 (8).

### N<sup>1</sup>-cyclohexyl-2-[(7-oxo-1,3,5-cycloheptatrienyl)(propyl)amino]butanamide (5d)

Yellow oil, (Yield: 80%). IR (neat):  $\nu = 3445, 2927, 1740, 1654, 1616, 1465, 1260, 1018$  cm<sup>-1</sup>. <sup>1</sup>H-NMR (400.22 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 0.71 (t, 3H,  $J = 7.4$  Hz, CH<sub>3</sub> of (CH<sub>2</sub>)<sub>2</sub>CH<sub>3</sub>), 0.85 (t, 3H,  $J = 7.4$  Hz, CH<sub>3</sub> of CH<sub>2</sub>CH<sub>3</sub>), 0.96-2.39 (m, 14H, 5 CH<sub>2</sub> of cyclohexyl and 2 CH<sub>2</sub> of acyclic), 3.07-3.41 (m, 2H, NCH<sub>2</sub>),

3.70-3.91 (m, 1H, CH-N of cyclohexyl), 4.40 (t, 1H,  $J = 7.0$  Hz, CH), 6.66 (d, 1H,  $J = 10.4$  Hz, CH of tropone), 6.72 (t, 1H,  $J = 9.2$  Hz, CH of tropone), 7.04-7.30 (m, 3H, CH of tropone), 8.43 (d, 1H,  $J = 8.0$  Hz, NH exchanged by D<sub>2</sub>O addition). <sup>13</sup>C-NMR (100.63 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 11.41 and 11.65 (2 CH<sub>3</sub>), 19.00 and 21.99 (2 CH<sub>2</sub>), 24.79, 26.65, 32.96 (2 CH<sub>2</sub> of cyclohexyl), 47.03 (CH<sub>2</sub>N); 47.79 (CH of cyclohexyl), 63.60 (CH), 120.02, 126.07, 133.94, 136.44 (5 CH of tropone), 158.57 (C of tropone), 170.03 (C=O of amide), 184.28 (C=O of tropone). Analysis of C<sub>20</sub>H<sub>30</sub>N<sub>2</sub>O<sub>2</sub> (330.23). (% calculation/found): C: 72.69/72.65, H: 9.15/9.18, N: 8.48/8.44.

**2-[benzyl (7-oxo-1,3,5-cycloheptatrienyl) amino]-N<sup>1</sup>-(2,4,4-trimethyl sec. pentyl) butanamide (5e)**

Yellow oil, (Yield: 85%). IR (neat):  $\nu = 3429, 2923, 1748, 1651, 1619, 1417, 1221, 1020$  cm<sup>-1</sup>. <sup>1</sup>H-NMR (400.22 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 0.79 (t, 3H,  $J = 7.4$  Hz, CH<sub>3</sub>), 1.05 (s, 9H, CMe<sub>3</sub>), 1.43 (s, 6H, CMe<sub>2</sub>NH), 1.62-2.20 (m, 2H, CH<sub>2</sub> of propyl and s, 2H, CH<sub>2</sub>CMe<sub>3</sub>), 4.49 (t, 1H,  $J = 7.0$  Hz, CH), 4.51 (AB, 2H,  $J = 16.8$  Hz, PhCH<sub>2</sub>), 6.51 (d, 1H,  $J = 10.4$  Hz, CH of tropone), 6.63 (t, 1H,  $J = 9.0$  Hz, CH of tropone), 6.85 (t, 1H,  $J = 10.4$  Hz, CH of tropone), 7.08-7.38 (m, 7H, CH of tropone and ph), 8.50 (s, 1H, NH exchanged by D<sub>2</sub>O addition). <sup>13</sup>C-NMR (100.63 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 11.80 (CH<sub>3</sub>), 22.26 (CH<sub>2</sub> of propyl), 29.27 (CMe<sub>2</sub>NH), 31.51 (CMe<sub>3</sub>), 31.64 (CMe<sub>3</sub>), 49.48 (PhCH<sub>2</sub>), 51.16 (CH<sub>2</sub>CMe<sub>3</sub>), 54.63 (CMe<sub>2</sub>NH), 64.08 (CH), 121.10, 126.35, 126.91, 127.18, 128.53, 133.75, 134.44, 136.37 (8 CH of tropone and Ph), 135.92 (C<sub>ip</sub> of Ph), 158.02 (C of tropone), 169.71 (C=O of amide), 184.27 (C=O of tropone). Analysis of C<sub>26</sub>H<sub>36</sub>N<sub>2</sub>O<sub>2</sub> (408.28). (% calculation/found): C: 76.43/76.45, H: 8.88/8.91, N: 6.86/6.82.

**N<sup>1</sup>-(tert-butyl)-2-[cyclohexyl(7-oxo-1,3,5-cycloheptatrienyl)amino]butanamide (5f)**

Yellow oil, (Yield: 75%). IR (neat):  $\nu = 3424, 2921, 1743, 1652, 1616, 1473, 1220, 1032$  cm<sup>-1</sup>. <sup>1</sup>H-NMR (400.22 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 0.88 (t, 3H,  $J = 7.4$  Hz, CH<sub>3</sub>), 1.33 (s, 9H, CMe<sub>3</sub>), 0.89-2.20 (m, 10H, CH<sub>2</sub> of cyclohexyl and m, 2H, CH<sub>2</sub> of propyl), 3.30 (t, 1H,  $J = 11.4$  Hz, CH-N of cyclohexyl), 4.00 (t, 1H,  $J = 7.0$  Hz, H-C), 6.68 (t, 1H,  $J = 9.0$  Hz, CH of tropone), 6.95-7.30 (m, 4H, CH of tropone), 8.23 (s, 1H, NH, exchanged by D<sub>2</sub>O addition). <sup>13</sup>C-NMR (100.63 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 12.53 (CH<sub>3</sub>), 24.00 (CH<sub>2</sub> of propyl), 25.73, 26.23, 26.47, 30.62, 33.32 (5 CH<sub>2</sub> of cyclohexyl), 28.58 (3CH<sub>3</sub>), 50.49 (CMe<sub>3</sub>), 60.98 (CH of cyclohexyl), 65.20 (CH), 120.01, 126.44, 133.15, 133.91, 135.92 (5 CH of tropone), 158.42 (C of tropone), 170.88 (C=O of amide), 185.61 (C=O of tropone). Analysis of C<sub>21</sub>H<sub>32</sub>N<sub>2</sub>O<sub>2</sub> (344.25). (% calculation/found): C: 73.22/73.18, H: 9.36/9.33, N: 8.13/8.09.

**2-[cyclohexyl (7-oxo-1,3,5-cycloheptatrienyl) amino]-N<sup>1</sup>-(2,4,4-trimethyl sec. pentyl) butanamide (5g)**

Yellow oil, (Yield: 70%). IR (neat):  $\nu = 3291, 2933, 1741, 1659, 1608, 1461, 1250, 1071$  cm<sup>-1</sup>. <sup>1</sup>H-NMR (400.22 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 0.90 (t, 3H,  $J = 7.4$  Hz, CH<sub>3</sub>), 1.01 (s, 9H, CMe<sub>3</sub>), 1.42 (s, 6H, CMe<sub>2</sub>NH), 1.07-2.30 (m, 10H, 5 CH<sub>2</sub> of cyclohexyl and m, 2H, CH<sub>2</sub> of propyl and s, 2H, CH<sub>2</sub>CMe<sub>3</sub>), 3.37 (t, 1H,  $J = 11.6$  Hz, CH-N of cyclohexyl), 3.98 (t, 1H,  $J = 7.0$  Hz, H-C), 6.68 (t, 1H,  $J = 9.0$  Hz, CH of tropone), 6.95-7.30 (m, 4H, CH of tropone), 8.04 (s, 1H, NH, exchanged by D<sub>2</sub>O addition). <sup>13</sup>C-NMR (100.63 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm):

12.78 (CH<sub>3</sub>), 24.18 (CH<sub>2</sub> of propyl), 25.75, 26.23, 26.49, 30.64, 33.35 (5 CH<sub>2</sub> of cyclohexyl), 28.71 (CMe<sub>2</sub>), 31.48 (CMe<sub>3</sub>), 31.56 (CMe<sub>3</sub>), 51.36 (CH<sub>2</sub> CMe<sub>3</sub>), 54.54 (CMe<sub>2</sub>NH), 60.88 (CH of cyclohexyl), 65.11 (CH), 121.57, 126.22, 133.06, 133.87, 135.72 (5 CH of tropone), 158.23 (C of tropone), 170.44 (C=O of amide), 185.33 (C=O of tropone). Analysis of C<sub>25</sub>H<sub>40</sub>N<sub>2</sub>O<sub>2</sub> (400.31). (% calculation/found): C: 74.95/74.99, H: 10.06/10.10, N: 6.99/6.94.

#### N<sup>1</sup>-cyclohexyl-2-[cyclohexyl (7-oxo-1,3,5-cycloheptatrienyl)amino] butanamide (5h)


Yellow oil, (Yield: 79%). IR (neat):  $\nu = 3340, 2971, 1740, 1653, 1612, 1478, 1222, 1080 \text{ cm}^{-1}$ . <sup>1</sup>H-NMR (400.22 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 0.88 (t, 3H,  $J = 7.4 \text{ Hz}$ , CH<sub>3</sub>), 0.91-2.20 (m, 20H, 10 CH<sub>2</sub> of 2 cyclohexyl and m, 2H, CH<sub>2</sub> of propyl), 3.20-3.80 (m, 2H, 2 CH of 2 cyclohexyl), 4.01 (t, 1H,  $J = 7.0 \text{ Hz}$ , H-C), 6.72 (t, 1H,  $J = 9.0 \text{ Hz}$ , CH of tropone), 6.96-7.30 (m, 4H, CH of tropone), 8.31 (d, 1H,  $J = 8.0 \text{ Hz}$ , NH, exchanged by D<sub>2</sub>O addition). <sup>13</sup>C-NMR (100.63 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 12.67 (CH<sub>3</sub>), 24.09 (CH<sub>2</sub> of propyl), 24.72, 24.80, 25.60, 25.74, 26.22, 26.37, 30.74, 32.81, 32.95, 33.03 (10 CH<sub>2</sub> of 2 cyclohexyl), 64.77 (CH), 122.94, 126.92, 133.09, 134.39, 135.95 (5 CH of tropone), 158.38 (C of tropone), 170.99 (C=O of amide), 185.83 (C=O of tropone). Analysis of C<sub>23</sub>H<sub>34</sub>N<sub>2</sub>O<sub>2</sub> (370.26). (% calculation/found): C: 74.55/74.59, H: 9.25/9.22, N: 7.56/7.59.

## Results and discussion

The 1:1 imine intermediate generated by the addition of primary amine **2** to propionaldehyde **1** is trapped by isocyanide **3** and tropolone **4** in the presence of silica NPs, leading to the formation of 2-(*N,N*-dialkylamino)-2,4,6-cycloheptatrien-1-one derivatives **5** (Scheme 1 and Table 1). The reaction proceeds smoothly and cleanly under mild conditions, and no side reactions are observed.

Silica NPs were found to catalyze the synthesis of 2-(*N,N*-dialkylamino)-2,4,6-cycloheptatrien-1-one derivatives from propionaldehyde **1**, primary amines **2**, isocyanides **3**, and tropolone **4** in methanol at 60 °C with high efficiency (Scheme 1 and Table 1). We also used silica gel powder (Merck) instead of silica NPs in this reaction, but increasing reaction times and decreasing 2-(*N,N*-dialkylamino)-2,4,6-cycloheptatrien-1-one yields were observed (Scheme 1 and Table 1). The use of just 0.2 g of silica NPs per mmol of reactants is sufficient to push the reaction forward. Higher amounts of silica NPs (0.3 g) did not improve the result to a great extent (Table 2, entries 4-7). In order to investigate the effects of other reaction media in this reaction, we also carried out the described condensation in MeOH, H<sub>2</sub>O and solvent-free (neat) systems (Table 2, entries 1-3).

A possible mechanism for the present reaction is shown in Scheme 2, which envisages a tandem sequence. On the basis of the chemistry of isocyanides, it is reasonable to assume that the first step may involve the formation of imine **6** by the condensation reaction of primary amine **2** with propionaldehyde **1**; the next step may involve nucleophilic addition of the isocyanide **3** to the imine intermediate **7**, leading to nitrilium intermediate **8**. This intermediate may be attacked by the conjugate base of the tropolone **4** to form 1:1:1 adduct **9**. The intermediate **9** may undergo Smiles rearrangements to afford the isolated 2-(*N,N*-dialkylamino)-2,4,6-cycloheptatrien-1-one derivatives **5** via intermediate **10**. It may be speculated that the polar amphoteric surface (hydroxyl groups of the silica NPs) facilitates the interaction of adsorbed weak acidic and basic components due to stabilization of the corresponding transition states and intermediates by H-bonding. This interaction with



**5a:**  $\text{R}^1 = \text{Benzyl}$ ,  $\text{R}^2 = \text{Cyclohexyl}$ ; **5b:**  $\text{R}^1 = \text{Benzyl}$ ,  $\text{R}^2 = \text{tert-Butyl}$ ; **5c:**  $\text{R}^1 = n\text{-Propyl}$ ,  $\text{R}^2 = \text{tert-Butyl}$ ; **5d:**  $\text{R}^1 = n\text{-Propyl}$ ,  $\text{R}^2 = \text{Cyclohexyl}$ ; **5e:**  $\text{R}^1 = \text{Benzyl}$ ,  $\text{R}^2 = 1,1,3,3\text{-tetramethylbutyl}$ ; **5f:**  $\text{R}^1 = \text{Cyclohexyl}$ ,  $\text{R}^2 = \text{tert-Butyl}$ ; **5g:**  $\text{R}^1 = \text{Cyclohexyl}$ ,  $\text{R}^2 = 1,1,3,3\text{-tetramethylbutyl}$ ; **5h:**  $\text{R}^1 = \text{Cyclohexyl}$ ,  $\text{R}^2 = \text{Cyclohexyl}$ .

**Scheme 1.** Four-component synthesis of 2-(*N,N*-dialkylamino)-2,4,6-cycloheptatrien-1-one derivatives **5** in the presence of silica NPs (see Table 1).

**Table 1.** SiO<sub>2</sub> NP-promoted synthesis of 2-(*N,N*-dialkylamino)-2,4,6-cycloheptatrien-1-one derivatives **5**<sup>a</sup>.

<b>5</b>	R <sup>1</sup>	R <sup>2</sup>	Yield (%) <sup>b</sup>
<b>a</b>	Benzyl	Cyclohexyl	92
<b>b</b>	Benzyl	<i>tert</i> -Butyl	90
<b>c</b>	<i>n</i> -Propyl	<i>tert</i> -Butyl	83
<b>d</b>	<i>n</i> -Propyl	Cyclohexyl	80
<b>e</b>	Benzyl	1,1,3,3-Tetramethylbutyl	85
<b>f</b>	Cyclohexyl	<i>tert</i> -Butyl	75
<b>g</b>	Cyclohexyl	1,1,3,3-Tetramethylbutyl	70
<b>h</b>	Cyclohexyl	Cyclohexyl	79


<sup>a</sup> See Scheme 1; 0.2 g SiO<sub>2</sub> NP/mmol reactants were applied. <sup>b</sup> Yield of isolated **5**.

**Table 2.** Synthesis of 2-(*N,N*-dialkylamino)-2,4,6-cycloheptatrien-1-one derivatives **5a** from the reaction of propionaldehyde, benzylamine, cyclohexyl isocyanide, and tropolone under various conditions.

Entry	Catalyst <sup>a</sup> or solvent	Temp (°C)	Time (h)	Yield (%) <sup>b</sup>
1	MeOH	60	24	65
2	H <sub>2</sub> O	90	24	57
3	neat	90	24	35
4	MeOH/Silica gel powder (Merck) (0.2 g)	60	24	77
5	MeOH/SiO <sub>2</sub> NP (0.1 g)	60	6	83
6	MeOH/SiO <sub>2</sub> NP (0.2 g)	60	6	92
7	MeOH/SiO <sub>2</sub> NP (0.3 g)	60	6	92

<sup>a</sup> Amount of SiO<sub>2</sub> catalyst per mmol of reactants. <sup>b</sup> Yields of isolated **5a**.

the neighboring silanol groups is shown in Scheme 2 for the first reaction step. Participation of 2 proximate silanol groups (one as H-bond donor and the other as H-bond acceptor) in the reaction mechanism also seems plausible.



**Scheme 2.** Proposed mechanism for the formation of 2-(*N,N*-dialkylamino)-2,4,6-cycloheptatrien-1-one derivatives **5** in the presence of silica NPs.

The structures of the products were deduced from their IR,  $^1\text{H-NMR}$ , and  $^{13}\text{C}$  NMR-spectra and mass spectrometry. For example, the  $^1\text{H-NMR}$  spectrum of **5a** consisted of a triplet for the  $\text{CH}_3$  of *n*-propyl ( $\delta = 0.79$  ppm,  $J = 7.4$  Hz), several multiplets for the cyclohexyl and *n*-propyl ( $\delta = 0.86$ -2.10 ppm) moieties, a multiplet for the NCH ( $\delta = 3.80$ -4.02 ppm) of cyclohexyl moiety, an AB-quartet for  $\text{CH}_2$  of  $\text{PhCH}_2$  ( $\delta = 4.50$  ppm,  $J = 16.8$  Hz), a triplet for the NCH ( $\delta = 4.55$  ppm,  $J = 7.0$  Hz) of the acyclic part, a doublet for

the tropolone hydrogen ( $\delta = 6.53$  ppm,  $J = 10.4$  Hz), a triplet for the tropolone hydrogen ( $\delta = 6.63$  ppm,  $J = 9.0$  Hz), a triplet for the tropolone hydrogen ( $\delta = 6.84$  ppm,  $J = 10.2$  Hz), a multiplet for a tropolone hydrogen and phenyl group ( $\delta = 7.07$ - $7.35$  ppm), and a doublet for the NH that was exchangeable with D<sub>2</sub>O ( $\delta = 8.69$  ppm,  $J = 8.0$  Hz). The <sup>13</sup>C-NMR spectrum of **5a** showed 20 distinct signals. Partial assignment of these signals is given in the experimental section. The <sup>1</sup>H- and <sup>13</sup>C-NMR spectra of compounds **5b–h** were similar to those of **5a**, except for the aromatic or aliphatic moieties, which exhibited characteristic signals with appropriate chemical shifts.

## Conclusions

In summary, the reported method offers a mild, simple, and efficient route for the preparation of 2-(*N,N*-dialkylamino)-2,4,6-cycloheptatrien-1-one derivatives **5** from propionaldehyde **1**, primary amine **2**, isocyanide **3**, and tropolone **4** in the presence of silica NPs. Its ease of work-up, high yields, and fairly mild reaction conditions make it a useful addition to modern synthetic methodologies.

## References

1. Zhu, J.; Bienaymé, H. *Multicomponent Reactions*; Wiley-VCH, Weinheim, **2005**.
2. Heshmati-Gonbari, M.; Ramazani, A.; Souldozi, A. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2009**, *184*, 309-314.
3. Ramazani, A.; Kazemizadeh, A. R. *Curr. Org. Chem.* **2011**, *15*, 3986-4020.
4. Ramazani, A.; Shajari, N.; Mahyari, A.; Ahmadi, Y. *Mol. Divers.* **2011**, *15*, 521.
5. Ramazani, A.; Zeinali Nasrabadi, F.; Mashhadi Malek Zadeh, A.; Ahmadi, Y. *Monatsh. Chem.* **2011**, *142*, 625.
6. Souldozi, A.; Ramazani, A.; Bouslimani, N.; Welter, R. *Tetrahedron Lett.* **2007**, *48*, 2617-2620.
7. Domling, A. *Chem. Rev.* **2006**, *106*, 17.
8. Ugi, I.; Werner, B.; Dömling, A. *Molecules* **2003**, *8*, 53.
9. Waller, R. W.; Diorazio, L. J.; Taylor, B. A.; Motherwell, W. B.; Sheppard, T. D.
10. *Tetrahedron* **2010**, *66*, 6496.
11. Ramazani, A.; Ahmadi, Y.; Rouhani, M.; Shajari, N.; Souldozi, A. *Heteroatom Chem.* **2010**, *21*, 368.
12. Ramazani, A.; Ahmadi, Y.; Tarasi, R. *Heteroatom Chem.* **2011**, *22*, 79.
13. Nagaraja G. K.; Prakash G. K.; Kumaraswamy M. N.; Vaidya V. P.; Mahadevan K. M. *Arkivoc.* **2006**, *vx*, 160
14. Zeinali Nasrabadi F.; Ramazani, A.; Ahmadi, Y. *Mol. Divers.* **2011**, *15*, 791.
15. Ramazani, A.; Ahmadi, Y.; Mahyari, A. *Synthetic Communications* **2011**, *41*, 2273-2282.
16. Ramazani, A.; Ahmadi, Y.; Zeinali Nasrabadi, F. *Z. Naturforsch.* **2011**, *66b*, 184-190.
17. Lewis, L. N. *Chem. Rev.* **1993**, *93*, 2693.
18. Banerjee, S.; Santra, S. *Tetrahedron Lett.* **2009**, *50*, 2037.
19. Ramazani, A.; Azizian, A.; Bandpey, M.; Noshiranzadeh, N. *Phosphorus, Sulfur, and Silicon Relat. Elem* **2006**, *181*, 2731.

20. Yavari, I.; Ramazani, A. *Phosphorus, Sulfur, and Silicon*. **1997**, *130*, 73.
21. Ramazani, A.; Bodaghi, A. *Tetrahedron Lett.* **2000**, *41*, 567-568.
22. Ramazani, A.; Mahyari, A. *Helv. Chim. Acta* **2010**, *93*, 2203.
23. Souldozi, A.; Ramazani, A. *Tetrahedron Lett.* **2007**, *48*, 1549.
24. Ramazani, A.; Souldozi, A. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2009**, *184*, 3191.
25. Ramazani, A.; Souldozi, A. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2009**, *84*, 2350.
26. a) Ramazani, A.; Rezaei, A. *Org. Lett.* **2010**, *12*, 2852-2855; b) Ramazani, A.; Karimi, Z.; Souldozi, A.; Ahmadi, Y. *Turk. J. Chem.* **2012**, *36*, 81-91; c) Valizadeh Holagh, M.; Maharramov, A. M.; Allahverdiyev, M. A.; Ramazani, A.; Ahmadi, Y.; Souldozi, A. *Turk. J. Chem.* **2012**, *36*, 179-188; d) Souldozi, A.; Ramazani, A. *Arkivoc* **2008**, *xvi*, 235-242; e) Ramazani, A.; Rouhani, M.; Rezaei, A.; Shajari, N.; Souldozi, A. *Helv. Chim. Acta* **2011**, *94*, 282-288; f) Shajari, N.; Ramazani, A.; Ahmadi, Y. *Bull. Chem. Soc. Ethiop.* **2011**, *25*, 1-6; g) Ramazani, A.; Nasrabadi, F. Z.; Karimi, Z.; Rouhani, M. *Bull. Korean Chem. Soc.* **2011**, *32*, 2700-2704; h) Ramazani, A.; Farshadi, A.; Mahyari, A.; Slepokura, K.; Lis, T.; Rouhani, M.; *J. Chem. Crystallogr.* **2011**, *41*, 1376-1385; i) Massoudi, A.; Amini, I.; Ramazani, A.; Nasrabadi, F. Z.; Ahmadi, Y. *Bull. Korean Chem. Soc.* **2012**, *33*, 39-42; j) Ramazani, A.; Rezaei, A.; Ahmadi, Y. *Phosphorus, Sulfur, Silicon Relat. Elem* **2012**, *187*, 22-31; k) Ramazani, A.; Momeni-Movahhed, A.; Gouranlou, F. *Phosphorus, Sulfur, Silicon Relat. Elem* **2002**, *177*, 903-907; l) Yavari, I.; Ramazani, A.; Yahyazadeh, A. *Synth. Commun.* **1996**, *26*, 4495-4499; m) Pakravan, P.; Ramazani, A.; Noshiranzadeh, N.; Sedrpoushan, A. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2007**, *182*, 545-549; n) Ramazani, A.; Rahimifard, M.; Souldozi, A. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2007**, *182*, 1-5; o) Ramazani, A.; Rahimifard, M.; Noshiranzadeh, N.; Souldozi, A. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2007**, *182*, 413-417; p) Ramazani, A.; Souldozi, A. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2003**, *178*, 2663-2666; q) Ramazani, A.; Shajari, N.; Gouranlou, F. *Phosphorus, Sulfur, Silicon Relat. Elem* **2001** *174*, 223-227; r) Ramazani, A.; Mahyari, A. T.; Rouhani, M.; Rezaei, A. *Tetrahedron Lett.* **2009**, *50*, 5625-5627; s) Kazemizadeh, A. R.; Ramazani, A. *J. Braz. Chem. Soc.* **2009**, *20*, 309-312; t) Ramazani, A.; Amini, I.; Massoudi, A. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2006**, *181*, 2225-2229; u) Ramazani, A.; Souldozi, A. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2003**, *178*, 1325-1328; v) Ramazani, A.; Souldozi, A. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2003**, *178*, 1329-1332; w) Souldozi, A.; Ramazani, A. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2009**, *184*, 2344-2350; x) Ramazani, A.; Souldozi, A. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2004**, *179*, 529-534; y) Ramazani, A.; Bodaghi, A. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2004**, *179*, 1615-1620; z) Kazemizadeh, A. R.; Ramazani, A. *Arkivoc* **2008**, *xv*, 159-165.
27. Souldozi, A.; Slepokura, K.; Lis, T.; Ramazani, A. *Z. Naturforsch.* **2007**, *62b*, 835.
28. Astruc, D.; Lu, F.; Aranzas, J. R. *Angew. Chem. Int. Ed.* **2005**, *44*, 7852.
29. Beletskaya, I. P.; Cheprakov, A. V. *Chem. Rev.* **2000**, *100*, 3009.
30. Ramazani, A.; Rezaei, A.; Mahyari, A. T.; Rouhani, M.; Khoobi, M. *Helv. Chim. Acta* **2010**, *93*, 2033.
31. a) Ramazani, A.; Zeinali Nasrabadi, F.; Ahmadi, Y. *Helv. Chim. Acta* **2011**, *94*, 1024; b) Ramazani, A.; Noshiranzadeh, N.; Ghamkhari, A.; Slepokura, K.; Lis, T. *Helv. Chim. Acta* **2008**, *91*, 2252-2261.
32. Ramazani, A.; Mahyari, A.; Lashgari, H.; Slepokura, K.; Lis, T. *Helv. Chim. Acta* **2011**, *94*, 611-622.
33. Ramazani, A.; Mahyari, A.; Farshadi, A.; Rouhani, M. *Helv. Chim. Acta* **2011** *94*, 1831-1837

Copyright of Turkish Journal of Chemistry is the property of Scientific and Technical Research Council of Turkey and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.