Kinetics and Mechanism of Oxidation of Dimethyl Sulphoxide by Mono- and Di-Substituted N,N-Dichlorobenzenesulphonamides in Aqueous Acetic Acid

B. Thimme Gowda, K. L. Jayalakshmi, and K. Jyothi

Department of Post-Graduate Studies and Research in Chemistry, Mangalore University, Mangalagangothri-574 199, Mangalore, India

Reprint requests to Prof. B. Thimme Gowda. Fax: 91 824 2287 367. E-mail: gowdabt@yahoo.com

Z. Naturforsch. 58b, 787-794 (2003); received February 6, 2003

In an effort to introduce N,N-dichloroarylsulphonamides of different oxidising strengths, four mono- and five di-substituted N,N-dichlorobenzenesulphonamides are prepared, characterised and employed as oxidants for studying the kinetics of oxidation of dimethyl sulphoxide (DMSO) in 50% aqueous acetic acid. The reactions show first order kinetics in [oxidant], fractional to first order in [DMSO] and nearly zero order in [H⁺]. Increase in ionic strength of the medium slightly increases the rates, while decrease in dielectric constant of the medium decreases the rates. The results along with those of the oxidation of DMSO by N,N-dichlorobenzenesulphonamide and N,N-dichloro-4methylbenzenesulphonamide have been analysed. Effective oxidising species of the oxidants employed in the present oxidations is Cl⁺ in different forms, released from the oxidants. Therefore the introduction of different substituent groups into the benzene ring of the oxidant is expected to affect the ability of the reagent to release Cl⁺ and hence its capacity to oxidise the substrate. Significant changes in the kinetic and thermodynamic data are observed in the present investigations with change of substituent in the benzene ring. The electron releasing groups such as CH3 inhibit the ease with which Cl⁺ is released from the oxidant, while electron-withdrawing groups such as Cl enhance this ability. The Hammett equation, $\log k_{obs} = -3.19 + 1.05 \sigma$, is found to be valid for oxidations by all the p-substituted N,N-dichlorobenzenesulphonamides. The substituent effect on the energy of activation, E_a and log A for the oxidations is also analysed. The enthalpies and free energies of activation correlate with an isokinetic temperature of 320 K.

Key words: Kinetics, Oxidation, Dimethyl Sulphoxide, N,N-Dichloroarylsulphonamides

Introduction

Many sulphonamides and their N-halo compounds are of fundamental chemical interest as they show distinct physical, chemical and biological properties. They also exhibit pharmacological activity. Further, many of these compounds exhibit fungicidal and herbicidal activities, due to their oxidising action in aqueous, partial aqueous and non-aqueous media [1-6]. This has stimulated recent interest in their chemistry. Therefore an understanding of the formation, properties and reactions of arylsulphonamides, their N-chloro and N,N-dichloro compounds is of interest in medicinal and redox chemistry, and a great deal of work on the spectroscopic, structural and kinetic aspects of these compounds needs to be done for correlating their reactivities with the chemical bond parameters. We are interested in the chemistry of these compounds in particular and amides in general [6-14] and have recently

reported the spectroscopic studies on a number of arylsulphonamides [12], the sodium salts of *N*-chloro compounds [13] and *N*,*N*-dichloroarylsulphonamides [14].

Dimethyl sulphoxide is a versatile solvent in industry and chemical laboratories as medium for carrying out reactions. It is also a popular medicinal compound because of its low toxicity and high skin permeability. Hence the reactions of dimethyl sulphoxide are of interest. The kinetics of oxidation of DMSO by a number of oxidants have been reported [15-20].

We report herein the kinetics of oxidation of dimethyl sulphoxide by four mono- and five disubstituted *N*,*N*-dichlorobenzenesulphonamides and analyse the results along with those by *N*,*N*-dichlorobenzenesulphonamide and *N*,*N*-dichloro-4methylbenzenesulphonamide. The oxidants or *N*,*N*-dichloro compounds employed were of the configuration ArSO₂NCl₂, where Ar = C₆H₅ (1); 4-CH₃-C₆H₄ (2); 4-C₂H₅-C₆H₄ (3); 4-F-C₆H₄ (4); 4-Cl-C₆H₄ (5); 4-Br-

0932-0776 / 03 / 0800-0787 \$ 06.00 © 2003 Verlag der Zeitschrift für Naturforschung, Tübingen · http://znaturforsch.com

<u>788</u>

B. Thimme Gowda et al. · Kinetics and Mechanism of Oxidation of Dimethyl Sulphoxide

10 ³ [DCSB] _o	10[DMSO] _o	10[HClO ₄]	10	$0^4 k_{\rm obs} ({\rm s}^{-1})^4 k_{\rm obs}$	l) for i-X-C	C ₆ H ₄ SC	D_2NCl_2 ,	i-X =
$(mol dm^{-3})$	$(mol dm^{-3})$	$(\text{mol } \text{dm}^{-3})$	4-H	4-CH ₃	$4-C_2H_5$	4-F	4-Cl	4-Br
Effect of var	ying [DCSB] _o							
0.5	1.0	1.0	11.9	9.0	11.6	14.8	20.0	20.4
1.0	1.0	1.0	12.1	9.4	11.8	15.2	20.7	21.8
2.0	1.0	1.0	12.3	9.9	12.2	15.3	21.1	22.3
4.0	1.0	1.0	13.0	10.5	12.9	15.8	21.6	22.5
Effect of var	ying [DMSO] _c	•						
1.0	0.1	1.0	1.9	1.4	2.0	2.0	2.8	2.9
1.0	0.5	1.0	7.2	5.4	5.8	8.4	10.5	11.2
1.0	1.0	1.0	12.1	9.4	11.8	15.2	20.6	21.8
1.0	2.0	1.0	26.6	18.4	21.5	29.1	40.9	43.7
Effect of var	ying [HClO ₄]							
1.0	1.0	0.1	12.1	8.6	10.5	15.3	20.0	21.3
1.0	1.0	0.5	12.1	9.1	10.5	15.3	20.3	21.3
1.0	1.0	1.0	12.2	9.4	11.8	15.3	20.6	21.7
1.0	1.0	2.0	12.9	9.9	11.9	15.4	20.6	21.7
10 ³ [DCSB] _o	10[DMSO] _o	10[HClO ₄]	$10^4 k_{\rm obs}$	(s^{-1}) for i	i-X-j-Y-C ₆ I	H_3SO_2	NCl ₂ , i-Z	K-j-Y =
$(mol dm^{-3})$	$(mol dm^{-3})$	(mol dm^{-3}) $\overline{2,3}$	3-(CH ₃) ₂	2,4-(CH	3)2 2-CH3	-4-Cl	2,4-Cl ₂	3,4-Cl ₂
Effect of vary	ing [DCSB]							
0.5	1.0	1.0	8.9	20.2	24	.2	118.1	41.0
1.0	1.0	1.0	9.5	21.0	24	.9	119.6	41.9
2.0	1.0	1.0	9.7	22.0	25	.5	119.8	43.4
4.0	1.0	1.0	10.2	22.2	26	.2	120.2	44.0
Effect of vary	ing [DMSO] _o							
1.0	0.1	1.0	1.5	4.2	4	.4	12.4	4.8
1.0	0.5	1.0	5.2	14.0	15	.7	60.0	22.6
1.0	1.0	1.0	9.5	21.0	24	.9	119.6	41.9
1.0	2.0	1.0	17.5	42.2	50	.3	_	73.1
Effect of vary	ing [HClO ₄]							
1.0	1.0	0.1	9.6	20.8	24	.7	119.6	41.7
1.0	1.0	0.5	9.6	20.9	24	.7	119.6	41.7
1.0	1.0	1.0	9.7	21.0	24	.9	119.7	41.8
1.0	1.0	2.0	9.9	21.0	25	.1	120.4	41.8
10[<i>I</i>]		10	$^{4}k_{obs}$ (s ⁻	¹) for				Table 3
$(mol dm^{-3})$	i-X-C ₆ H ₄ SO	$NCl_2, i-X =$	i-X-j -	Y-C ₆ H ₃ S	02NCl2, i-2	X-j-Y =	=	strength
	4-H 4-C ₂ H	- 4-Cl	2 4-(0	(H ₂) ₂ 2-($\overline{CH_2}$ 4-Cl	2 4-	Cla	the react
1.0	11.8 0.0	5 4 Cl	2,4 (0	7	22.5	2,4	12	idation of
1.0	11.8 9.9	19.4	20	./	22.3	107	1.2	by the
2.0	12.0 9.9	19.8	21	.0	23.8	10	1.2	dichloro
5.0	12.1 11.8	20.7	21	.0	24.9	115	7.0 C O	in 1:1 (v
5.0	15.5 11.9	21.1	21	.2	23.4	120).9	cept duri
% AcOH		10	k_{obs} (s ⁻	¹) for	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~		_	-
(v/v)	1-X-C ₆ H ₄ S	O_2NCl_2 , i-X =	i-X-	· J-Y-C ₆ H ₃	$_{3}$ SO ₂ NCl ₂ ,	1-X-j-Y	(=	10^{3} [DCS
	4-H 4-C ₂ H	5 4-Cl	2,4-(C	$(H_3)_2$ 2-	CH_3 -4-Cl	2,4-	Cl_2	1.0 mol di
20	22.5 19.6	44.6	52	.8	60.3	-		
30	18.7 16.1	33.6	41	.4	45.7	156	5.9	
50	12.1 11.8	20.7	21	.0	24.9	119	9.6	

Table 1. Pseudo-first order rate constants (k_{obs}) for the oxidation of dimethyl sulphoxide (DMSO) by monosubstituted *N*,*N*-dichlorobenzenesulphonamides (DCSB) in 1:1 (v/v) water-acetic acid medium at 303 K (I = 0.30 mol dm⁻³).

Table 2. Pseudo-first order rate constants (k_{obs}) for the oxidation of dimethyl sulphoxide (DMSO) by disubstituted *N*,*N*-dichlorobenzenesulphonamides (DCSB) in 1:1 (v/v) water-acetic acid medium at 303 K (I = 0.30 mol dm⁻³).

Table 3. Effect of variation in ionic
strength (I) and solvent composition of
the reaction medium on the rate of ox-
idation of dimethyl sulphoxide (DMSO)
by the mono- and di-substituted N,N-
dichlorobenzenesulphonamides (DCSB)
in 1:1 (v/v) water-acetic acid medium (ex-
cept during its variations) at 303 K.

 $10^3 \text{ [DCSB]}_0 = 10 \text{ [DMSO]}_0 = 10[\text{HClO}_4] = 1.0 \text{ mol dm}^{-3}.$

 C_6H_4 (6); 2,3-(CH₃)₂- C_6H_3 (7); 2,4-(CH₃)₂- C_6H_3 (8); 2-CH₃-4-Cl- C_6H_3 (9); 2,4-Cl₂- C_6H_3 (10) and 3,4-Cl₂- C_6H_3 (11).

11.2

12.9

13.5

70.1

Results and Discussion

7.4

7.3

70

Kinetic measurements: The kinetic runs were made in glass stoppered pyrex boiling tubes under pseudofirst order conditions with [substrate] \gg [oxidant] (5 to 50 times). The reactions were initiated by the rapid addition of requisite amounts of oxidant solution (0.0005–0.004 mol dm⁻³), pre-equilibrated at a desired temperature, to solution mixtures containing known amounts of dimethyl sulphoxide (0.01–0.2 mol dm⁻³), perchloric acid (0.01–0.2 mol dm⁻³), acetic

Table 4. Kinetic data for the oxidation of dimethyl sulphoxide (DMSO) by the mono- and di-substituted N,N-dichlorobenzenesulphonamides (DCSB) in 1:1(v/v) wateracetic acid medium at 303 K.

Orders		i-X	$-C_6H_4SO_2I$	NCl ₂ , i	-X =	
in	4-H	4-CH ₃	$4-C_2H_5$	4-F	4-Cl	4-Br
[DCSB]	1.0	1.0	1.0	1.0	1.0	1.0
[DMSO]	0.9	0.8	0.7	1.0	1.0	1.0
[HClO ₄]	0	0	0	0	0	0
		i-X-j-Y-0	C ₆ H ₃ SO ₂ N	Cl ₂ , i-X	K-j-Y =	
	2,3-(CH ₃)	2,4-(CH	3) ₂ 2-CH ₃	-4-Cl	2,4-Cl ₂	3,4-Cl ₂
[DCSB]	1.0	1.0	1.	0	1.0	1.0
[DMSO]	0.8	0.7	1.	0	1.0	1.0
[HClO ₄]	0	0	C)	0	0



Fig. 1. Plot of k_{obs} vs. [DMSO]₀; 10³ [DCSB]₀ = 10 [HClO₄] = 3.33 I = 1.0 mol dm⁻³; Temp. 303 K. (DCSB: i-XC₆H₄SO₂NCl₂; i-X = 4-CH₃ or 4-C₂H₅).

acid and water, thermostated at the same temperature. The progress of the reactions was monitored for two half-lives by the iodometric estimation of unreacted oxidant at regular intervals of time. The pseudo-first order rate constants were computed by the graphical methods and the values were reproducible within \pm 3% error.

The kinetic data on the oxidation of dimethyl sulphoxide by the substituted N,N-dichlorobenzene-sulphonamides (DCSB) **3**–**11**, under varying [DCSB], [DMSO], [HClO₄], ionic strength and solvent composition of the medium, are shown in Tables 1–4



Fig. 2. Plot of k_{obs} vs. [DMSO]₀; 10³ [DCSB]₀ = 10 [HClO₄] = 3.33 I = 1.0 mol dm⁻³; Temp. 303 K. (DCSB: i-X-j-YC₆H₃SO₂NCl₂; i-X-j-Y = 2,3-(CH₃)₂; 2,4-(CH₃)₂).



Fig. 3. Plot of k_{obs} vs. % acetic acid; 10^3 [DCSB]₀ = 10 [HClO₄] = 3.33 I = 1.0 mol dm⁻³; Temp. 303 K. (DCSB: i-XC₆H₄SO₂NCl₂; i-X: 4-H, 4-C₂H₅ or 4-Cl and i-X-j-YC₆H₃SO₂NCl₂ = 2,4-(CH₃)₂; 2-CH₃-4-Cl; 2,4-Cl₂).

and Figures 1-4. The kinetics of oxidation of DMSO by the oxidants 1 and 2 have also been carried out

Activation		i	-X-C ₆ H ₄ SC	D_2NCl_2 , i-X	ζ =	
parameters	4-H	4-CH ₃	$4-C_2H_5$	4-F	4-Cl	4-Br
$E_{\rm a}$ (kJ mol ⁻¹)	65.5	58.3	59.0	67.3	65.1	62.2
log A	8.4	7.0	7.2	8.8	8.5	8.1
ΔH^{\neq} (kJ mol ⁻¹)	64.0	56.3	55.2	60.2	62.4	58.6
ΔS^{\neq} (JK ⁻¹ mol ⁻¹)	-89.8	-117.3	-119.0	-100.2	-90.4	-102.7
ΔG^{\neq} (kJ mol ⁻¹)	91.2	91.8	91.2	90.6	89.8	89.7
Optimised values	with refe	erence to lo	A of the p	parent oxid	ant	
$E_{\rm a}$ (kJ mol ⁻¹)	65.7	66.3	65.7	65.1	64.3	64.2
ΔH^{\neq} (kJ mol ⁻¹)	64.0	64.6	64.0	63.4	62.7	62.5
Optimised values	with refe	erence to E	a of the pare	ent oxidant		
log A	8.4	8.3	8.4	8.5	8.6	8.7
ΔS^{\neq} (JK ⁻¹ mol ⁻¹)	-89.7	-91.8	-89.9	-87.8	-85.3	-84.8



Fig. 4. Plot of (i) log k_{obs} vs. $\sigma\rho$ and (ii) ΔH^{\neq} vs. ΔS^{\neq} ; 10³ [DCSB]₀ = 10 [HClO₄] = 3.33 I = 1.0 mol dm⁻³; Temp. 303 K. (DCSB: i-XC₆H₄SO₂NCl₂; i-X: 4-H, 4-CH₃, 4-C₂H₅, 4-F, 4-Cl or 4-Br and i-X-j-Y-C₆H₃SO₂NCl₂; i-X-j-Y=2,3-(CH₃)₂; 2,4-(CH₃)₂; 2-CH₃-4-Cl; 2,4-Cl₂ or 3,4-Cl₂).

under identical conditions and included for completeness.

Effect of varying $[DCSB]_0$: At constant [DMSO](5–50 fold excess over [DCSB]) and $[H^+]$, first order plots of log[DCSB] versus time were linear up to at least 75% completion of the reaction. The pseudo-first order rate constants (k_{obs}) computed from the plots remained unaffected by the changes in [DCSB], establishing first order dependence of the rate on [DCSB], in all the cases (Tables 3 and 4).

Effect of varying $[DMSO]_0$: At constant $[DCSB]_0$ and $[HClO_4]$, the rates increased with increase in

Table 5. Activation parameters for the oxidation of dimethyl sulphoxide by mono-substituted N,N-dichlorobenzenesulphonamides in 1:1 (v/v) water-acetic acid in presence of perchloric acid.

Table 6. Activation parameters for the oxidation of dimethyl sulphoxide by di-substituted N,N-dichlorobenzenesulphonamides in 1:1 (v/v) water-acetic acid in presence of perchloric acid.

i-X	-j-Y-C ₆ H ₃ S	O ₂ NCl ₂ , i-X	K-j-Y =	
2,3-(CH ₃) ₂	2,4-(CH ₃) ₂	2-CH ₃ -4-Cl	2,4-Cl ₂	3,4-Cl ₂
61.6	65.7	58.4	53.5	58.0
7.6	8.6	7.5	7.3	7.6
) 59.7	58.7	55.6	52.0	56.5
$^{-1}) - 106.0$	-102.5	-111.4	-110.3	-104.1
) 91.8	89.8	89.4	85.4	88.0
ues with ref	erence to lo	A of the particular of the p	arent ox	idant
66.2	64.3	63.8	59.9	62.5
) 64.6	62.6	62.2	58.2	60.9
ues with ref	erence to E	of the paren	nt oxida	nt
8.3	8.6	8.7	9.4	8.9
$^{-1}$) -91.7	-85.1	-83.7	-70.7	-79.4
	$\begin{array}{c} \text{i-X}\\ 2,3-(\text{CH}_3)_2\\ \hline 61.6\\ 7.6\\ 7.6\\ 9.59.7\\ ^{-1})-106.0\\ 9.18\\ \text{lues with ref}\\ 66.2\\ 0.64.6\\ \text{lues with ref}\\ 8.3\\ ^{-1})-91.7\\ \end{array}$	$\begin{array}{r} \text{i-X-j-Y-C_6H_3S}\\ 2,3-(\text{CH}_3)_2 2,4-(\text{CH}_3)_2\\ \hline 61.6 & 65.7\\ 7.6 & 8.6\\ \text{)} & 59.7 & 58.7\\ \hline ^{-1}) -106.0 & -102.5\\ \text{)} & 91.8 & 89.8\\ \text{lues with reference to Ic}\\ 66.2 & 64.3\\ \text{)} & 64.6 & 62.6\\ \text{lues with reference to } E\\ 8.3 & 8.6\\ \hline ^{-1}) -91.7 & -85.1\\ \end{array}$	$\begin{array}{c c} \text{i-X-j-Y-C}_6\text{H}_3\text{SO}_2\text{NCl}_2, \text{i-X}\\ 2,3-(\text{CH}_3)_2\ 2,4-(\text{CH}_3)_2\ 2-\text{CH}_3-4-\text{Cl}\\\hline\hline 61.6 & 65.7 & 58.4\\ 7.6 & 8.6 & 7.5\\ 9.7 & 58.7 & 55.6\\ -^1)\ -106.0 & -102.5 & -111.4\\ 9.91.8 & 89.8 & 89.4\\ \text{lues with reference to } \log A \text{ of the pr}\\ 66.2 & 64.3 & 63.8\\ 9. & 64.6 & 62.6 & 62.2\\ \text{lues with reference to } E_a \text{ of the parent}\\ 8.3 & 8.6 & 8.7\\ -^1)\ -91.7 & -85.1 & -83.7\\ \end{array}$	$\begin{array}{c c} \text{i-X-j-Y-C}_{6}\text{H}_{3}\text{SO}_{2}\text{NCl}_{2}, \text{i-X-j-Y} = \\ 2,3-(\text{CH}_{3})_{2} 2,4-(\text{CH}_{3})_{2} 2-\text{CH}_{3}-4-\text{Cl} 2,4-\text{Cl}_{2} \\ \hline 61.6 & 65.7 & 58.4 & 53.5 \\ 7.6 & 8.6 & 7.5 & 7.3 \\ 0, 59.7 & 58.7 & 55.6 & 52.0 \\ \hline ^{-1}) -106.0 & -102.5 & -111.4 - 110.3 \\ 0, 91.8 & 89.8 & 89.4 & 85.4 \\ \text{lues with reference to } \log A \text{ of the parent ox} \\ 66.2 & 64.3 & 63.8 & 59.9 \\ 0, 64.6 & 62.6 & 62.2 & 58.2 \\ \text{lues with reference to } E_{a} \text{ of the parent oxidal} \\ 8.3 & 8.6 & 8.7 & 9.4 \\ \hline ^{-1}) -91.7 & -85.1 & -83.7 & -70.7 \end{array}$

[DMSO] with fractional to first order dependence of the rate on [DMSO] (Tables 1, 2, and 4). The k_{obs} versus [DMSO] plots were linear passing through the origin with the oxidants 4-6 and 9-11. The plots were also linear with the oxidants 1-3, 7, 8 but with intercepts on the ordinates (Fig. 1 and 2).

Effect of varying $[HClO_4]$: At constant $[DCSB]_0$ and $[DMSO]_0$, the rates remained almost unaffected by the changes in $[HClO_4]$, indicating nearly zero order dependence of rates on $[H^+]$ (Tables 1 and 2).

Effect of varying other parameters: The rates slightly increased with increase in ionic strength of the medium, while decrease in dielectric constant of the medium by increasing the acetic acid composition of the solvent decreased the rates (Table 3 and Fig. 3). The rates were measured at different temperatures and the activation parameters have been computed from the Arrhenius and Eyring plots (Tables 5 and 6).

N,*N*-dichloroarylsulphonamides furnish the following reactive species in its aqueous acetic acid solutions [2, 4, 6]: ArSO₂NCl₂, ArSO₂NHCl and HOCl

Table 7. The calculated rate constants k_1 and k'_2 at different temperatures for the oxidation of dimethyl sulphoxide by mono and di-substituted *N*,*N*-dichlorobenzenesulphonamides in 1:1 (ν/ν) water-acetic acid medium in the presence of perchloric acid.

Temp.	$10^3 k_1 (\mathrm{dm}^3 \mathrm{mol}^{-1} \mathrm{s}^{-1})$						
(K)	i-X	K-C ₆ H ₄ SO	$_2$ NCl ₂ ,	i-X-j-Y-C ₆ H ₃ SO ₂ NCl ₂ ,			
		i-X =		i	-X-j-Y =		
	4-H	$4-CH_3$	$4-C_2H_5$	2,4-Cl ₂	3,4-Cl ₂		
293	6.2	4.0	3.9	3.2	12.8		
298	8.9	5.5	5.3	5.0	16.6		
303	12.9	9.0	10.5	8.9	19.6		
308	16.0	13.8	14.1	10.7	30.6		
			10 ⁵ 1	$k_{2}^{'}(s^{-1})$			
293	2.0	2.5	7.5	4.0	0		
298	4.0	5.0	10.0	6.0	10.0		
303	6.0	7.5	10.0	6.0	30.0		
308	8.0	7.5	17.5	8.0	30.0		

(where $Ar = C_6H_5$, 4- CH_3 - C_6H_4 , 4- C_2H_5 - C_6H_4 , 4-F- C_6H_4 , 4-Cl- C_6H_4 , 4-Br- C_6H_4 , 2,3-(CH_3)₂- C_6H_3 , 2,4-(CH_3)₂- C_6H_3 , 2,2- C_6H_3 , 2,4- Cl_2 - C_6H_3 and 3,4- Cl_2 - C_6H_3).

Mechanism of oxidation: The kinetics of first order each in [DCSB] and [DMSO], and nearly zero order in $[H^+]$ observed for the oxidation of dimethyl sulphoxide by 4-fluoro, 4-chloro, 4-bromo, 2-methyl-4-chloro, 2,4-dichloro and 3,4-dichloro substituted *N*,*N*-dichlorobenzene-sulphonamides in aqueous acetic acid medium may be explained by the simple scheme

$$ArSO_2NCl_2 + DMSO \xrightarrow{\kappa_1} Products (slow) (1)$$

The related rate law is

$$-d[\text{oxidant}]/dt = k_1[\text{ArSO}_2\text{NCl}_2] \qquad (2)$$

or
$$k_{\rm obs} = k_1 [\rm DMSO]$$
 (3)

The plots of k_{obs} versus [DMSO] gave straight lines passing through the origin in conformity with the above rate law (eq. (3)).

The kinetic behaviour for the oxidation of DMSO by the parent, 4-methyl, 4-ethyl, 2,3-dimethyl and 2,4-dimethyl substituted *N*,*N*-dichlorobenzenesul-phonamides was slightly different. Oxidations showed first order kinetics each in [DCSB], fractional order in [DMSO] and zero order in [H⁺]. This is also reflected in the magnitudes of the rate constants. These results may be explained either by a two pathway mechanism (eq. (1) and Scheme 1) or a Michaelis-Menten type mechanism (Scheme 2).

Table 8. Activation parameters for path 1 of the mechanism for the oxidation of dimethyl sulphoxide by monoand di-substituted *N*,*N*-dichlorobenzenesulphonamides in 1:1 (ν/ν) water-acetic acid medium in the presence of perchloric acid.

Activation	i-X-C	5H4SO2	NCl ₂ ,	i-X-j-Y-C ₆ H	$H_3SO_2NCl_2$,
parameters		i-X =		i-X-	j-Y =
	4-H	4-CH ₃	$4-C_2H_5$	$2,3-(CH_3)_2$	$2,4-(CH_3)_2$
$E_{\rm a}$ (kJ mol ⁻¹)	55.4	64.4	68.9	65.6	43.8
log A	7.7	9.1	9.9	9.3	5.8
ΔH^{\neq} (kJ mol ⁻¹)	54.2	61.3	66.4	62.7	45.1
ΔS^{\neq} (JK ⁻¹ mol ⁻¹) -	-102.2	-82.0	-63.9	-77.5	-129.1
ΔG^{\neq} (kJ mol ⁻¹)	85.2	86.1	85.7	86.1	84.2
Optimised values	with re	ference	to log A	of the parer	nt oxidant
$E_{\rm a}$ (kJ mol ⁻¹)	55.6	56.5	56.2	56.6	54.6
ΔH^{\neq} (kJ mol ⁻¹)	54.2	55.2	54.8	55.2	53.2
Optimised values	with re	ference	to E _a of	the parent of	xidant
log A	7.7	7.6	7.6	7.6	7.9
ΔS^{\neq} (JK ⁻¹ mol ⁻¹) -	-102.2	-105.2	-104.0	-105.3	-98.8

Table 9. Activation parameters for path 2 of the mechanism for the oxidation of dimethyl sulphoxide by mono and di-substituted *N*,*N*-dichlorobenzenesulphonamides in 1:1 (v/v) water-acetic acid medium in the presence of perchloric acid.

Activation	i-X-C ₆	H ₄ SO ₂	NCl ₂ , i-X =	2,3-(CH ₃) ₂ C ₆
parameters	4-H	4-CH ₃	$4 - C_2 H_5$	$H_3SO_2NCl_2$
$E_{\rm a}$ (kJ mol ⁻¹)	82.5	68.9	38.3	34.6
log A	10.0	7.8	2.6	1.8
ΔH^{\neq} (kJ mol ⁻¹)	82.1	79.8	39.9	32.4
ΔS^{\neq} (JK ⁻¹ mol ⁻¹)	-55.1	-60.8	-190.0	-219.0
ΔG^{\neq} (kJ mol ⁻¹)	98.7	98.2	97.5	98.7
Optimised values	s with re	ference	to log A of the	e parent oxidant
$E_{\rm a}$ (kJ mol ⁻¹)	82.5	81.9	81.2	82.5
ΔH^{\neq} (kJ mol ⁻¹)	82.1	81.5	80.8	82.1
Optimised values	s with re	ference	to E_a of the pa	rent oxidant
log A	10.0	10.1	10.2	10.0
ΔS^{\neq} (JK ⁻¹ mol ⁻¹)	-55.1	-53.2	-50.8	-55.1

The combined rate law based on equation (1) and Scheme 1 is given by

$$-d[\text{oxidant}]/dt = k_1[\text{oxidant}][\text{DMSO}] + k_2[\text{oxidant}][\text{H}_2\text{O}]$$
(4)

or
$$k_{obs} = k_1 [DMSO] + k_2 [H_2O] = k_1 [DMSO] + k'_2$$
, where $k'_2 = k_2 [H_2O]$ (5)

The plots of k_{obs} versus [DMSO] were linear with finite intercepts on the ordinates in accordance with the rate law (eq. (5)) (Fig. 1 and 2). The constants k_1 and

$$\begin{array}{rcl} ArSO_2NCl_2 + H_2O & \xrightarrow{k_1} & HOCl + ArSO_2NHCl & (slow) \\ HOCl + DMSO & \rightarrow & Products & (fast) \\ Scheme 1. \end{array}$$

 $\stackrel{K_3}{\rightleftharpoons}$ $ArSO_2NCl_2 + DMSO$ Intermediate Х k_4 Products Scheme 2.

 k'_{2} were calculated from the slopes and intercepts of the plots. The substrate concentrations were varied at different temperatures. The constants k_1 and k'_2 were calculated at each temperature (Table 7). Thus two sets of activation parameters have been computed from the plots of log k versus 1/T and log (k/T) versus 1/T(Tables 8 and 9).

Based on Scheme 2, the rate laws (eq. (6-8)) have been obtained,

$$-d[\text{oxidant}]/dt = K_3 k_4 [\text{oxidant}]$$
$$[\text{DMSO}]/1 + K_3 [\text{DMSO}]$$
(6)

$$k_{\rm obs} = K_3 k_4 [\rm DMSO] / 1 + K_3 [\rm DMSO]$$
(7)

$$1/k_{\rm obs} = 1/K_3 k_4 [{\rm DMSO}]/1 + k_4$$
 (8)

The plots of $1/k_{obs}$ versus 1/[DMSO] were also linear in conformity with the rate law in eq. (8). But the former direct plots had better correlations than the latter double reciprocal plots, favouring the two pathway mechanism over the Michaelis-Menten type mechanism.

The following form of the Hammett equation is found to be valid for the oxidation of dimethyl sulphoxide by p-substituted-N,N-dichlorobenzenesulphonamides (Fig. 4).

$$\log k_{\rm obs} = -3.19 + 1.05\sigma \tag{9}$$

The enthalpies and the free energies of activation for reactions of all the N,N-dichloro-arylsulphonamides have been correlated. The plot of ΔH^{\neq} versus ΔS^{\neq} (Fig. 4) was reasonably linear with an isokinetic temperature of 320 K. Further, to see the effect of substituent in benzene ring of oxidant on the energy of activation, Ea values of N,N-dichloroaryl-sulphonamides were optimised with reference to the $\log A$ value of the parent oxidant, N,N-dichlorobenzenesulphonamide, through the equation, $E_a = 2.303RT(\log A - 10^{-1})$ $\log k_{\rm obs}$). The energies of activation for the oxidants with electron withdrawing groups in the benzene ring are slightly lower than those of the parent oxidant, N,N-dichlorobenzenesulphonamide, while the effect of electron releasing groups on E_a is negligible. Enthalpies of activations have the same trend. Similarly,

log A values were optimised with reference to the E_{a} of the parent oxidant through the equation $\log A =$ $\log k_{\rm obs} + E_a/2.303RT$. Log A values were little higher for oxidants with the electron withdrawing groups, while the effect of electron releasing groups on log A is negligible. The free energies of activation remain almost the same in both the optimisations indicating the operation of similar mechanisms in all the cases.

Conclusions

Effective oxidising species of the oxidants employed in the present oxidations is Cl⁺ in different forms, released from the oxidants. The oxidative strengths of N,N-dichloroaryl-sulphonamides depend on the ease with which Cl⁺ is released from the reagents. Therefore the introduction of different substituent groups into the benzene ring of the oxidant is expected to affect the ability of the reagent to release Cl⁺ and hence its capacity to oxidise the substrate. Significant changes in kinetic and the thermodynamic data were observed in the present investigations with change of substituent in the benzene ring of the oxidant. The electron releasing groups such as CH₃ inhibit the ease with which Cl⁺ is released from the oxidant, while electron-withdrawing groups such as Cl enhance this ability.

Experimental Section

Materials and methods

The parent, mono- and di-substituted N,N-dichlorobenzenesulphonamides employed in the present studies were prepared as follows in three steps [12-14, 21]. In the first step, the corresponding arylsulphonamides of the configuration ArSO₂NH₂ were obtained by the reaction of chloroosulphonic acid on the respective substituted benzenes dissolved in chloroform at 0 °C and subsequent treatment with liquid ammonia. Then the corresponding N-chloroarylsulphonamides with constitution ArSO₂NaNCl·xH₂O were prepared by bubbling pure chlorine gas through clear aqueous solutions of arylsulphonamides in 4 M NaOH at 70 °C for about 1 h. The precipitated sodium salts of N-chloroarylsulphonamides were purified and dissolved in water and further chlorinated by passing pure chlorine gas through the solution.

The precipitated N,N-dichloroarylsulphonamides were filtered, washed, dried and recrystallised from chloroform. The purity of all the reagents was checked by determining their melting points and by estimating iodometrically the amounts of active chlorine present in them. All the arylsulphonamides, N-chloroarylsulphonamides and their N,Ndichloro compounds were further characterised by recording

B. Thi	mme Gowda et al.	· Kinetics and	Mechanism of	Oxidation	of Dimethy	yl Sul	phoxide
--------	------------------	----------------	--------------	-----------	------------	--------	---------

Aryl group (Ar)	Arylsulphonamides (ArSO ₂ NH ₂) obsd. [12-14]	Sodium salts of N-chloro- arylsulphonamides (ArSO ₂ NaNCl·xH ₂ O) obsd. [12–14]	N,N-Dichloro- arylsulphonamides (ArSO ₂ NCl ₂) obsd.
$4 - C_2 H_5 - C_6 H_4$	99-101	194	58
4-F-C ₆ H ₄	125 (124-125)	198	55 - 56
4-Cl-C ₆ H ₄	143 (142-143)	191 (190)	81
4-Br-C ₆ H ₄	162 (161.5)	179 (178)	102
2,3-(CH ₃) ₂ -C ₆ H ₃	138 - 140	167	58
2,4-(CH ₃) ₂ -C ₆ H ₃	140 - 142	154	63
2-CH3-4-Cl-C6H3	180-182 (184-185)	172	70
2,4-Cl ₂ -C ₆ H ₃	178-180 (179-180)	210	67
3,4-Cl ₂ -C ₆ H ₃	141–143 (134-135)	192	55

their infrared spectra [12-14]. The melting points of the prepared compounds are given in Table 10.

The stock solutions $(0.1 \text{ mol } \text{dm}^{-3})$ of the *N*,*N*-dichloroarylsulphonamides were prepared in distilled acetic acid and stored in amber coloured bottles. The solutions were standardised by the iodometric method. A stock solution $(1.0 \text{ mol } \text{dm}^{-3})$ of analytical grade dimethyl sulphoxide (E. Merck, Germany) in doubly distilled water was prepared as and when required. Ionic strength of the medium was maintained at $(0.30 \text{ mol } \text{dm}^{-3})$ by employing concentrated aqueous solution of sodium nitrate. All other reagents used were of analytical grades of purity.

Stoichiometry and product analysis

Reaction mixtures containing excess [oxidant] over [DMSO] were kept at 27 ± 2 °C in the presence of perchloric acid for 24 h. In a typical experiment, 4-ethylbenzenesulphonamide among the reaction products in *N*,*N*-dichloro-4-ethylbenzenesulphonamide oxidation was detected by

- [1] S.R. Sandler, W. Karo, Organic Functional Group Preparations, Vol. 3, Academic Press, London (1972).
- [2] M. M. Campbell, G. Johnson, Chem. Rev. 78, 65 (1978).
- [3] B.T. Gowda, D.S. Mahadevappa, Talanta **30**, 359 (1983).
- [4] B.T. Gowda, D.S. Mahadevappa, J. Chem. Soc., Perkin Trans. 2 (1983) 323.
- [5] B.T. Gowda, B.S. Sherigara, D.S. Mahadevappa, Microchem. J. 34, 103 (1986).
- [6] M. C. Agrawal, S. K. Upadhyay, J. Sci. Ind. Res. 49, 13 (1990).
- [7] B.T. Gowda, A. Weiss, Z. Naturforsch. 49a, 695 (1994).
- [8] B.T. Gowda, S. Dou, A. Weiss, Z. Naturforsch. 51a, 627 (1996).
- [9] B. T. Gowda, D. K. Bhat, H. Fuess, A. Weiss, Z. Naturforsch. 54a, 261, 679 (1999).

paper chromatography using benzyl alcohol saturated with water as the solvent with 5% vanillin in 1% HCl solution in ethanol as spraying reagent ($R_f = 0.95$). 2,4-Dimethyl-benzenesulphonamide among the reaction products in *N*,*N*-dichloro-2,4-dimethylbenzene-sulphonamide oxidation was detected by TLC [22] using petroleum ether-chloroform-butanol (2:2:1 v/v) as the solvent system and iodine as spray reagent. The R_f value of the reduced 2,4-dimethylbenzenesulphonamide was 0.97 (compared with the value of 0.96 for the pure compound). The presence of dimethyl sulphone among the reaction products was confirmed by TLC using petroleum ether-chloroform-butanol (2:2:1, v/v) as the solvent and iodine as the detecting reagent. The observed stoichiometry may be represented by

$$rSO_2NCl_2 + 2(CH_3)_2SO + 2H_2O$$

 $\rightarrow ArSO_2NH_2 + 2(CH_3)_2SO_2 + 2H^+ + 2Cl^-$

As = Arylgroups 1-11.

A

- [10] B. T. Gowda, H. Paulus, H. Fuess, Z. Naturforsch. 55a, 711,791(2000); 56a, 386 (2001).
- [11] B. T. Gowda, B. H. A. Kumar, H. Fuess, Z. Naturforsch. 55a, 721 (2000).
- [12] B. T. Gowda, K. Jyothi, J. D. D'Souza, Z. Naturforsch. 57a, 967 (2002).
- [13] B. T. Gowda, J. D. D'Souza, B. H. A. Kumar, Z. Naturforsch. 58a, 51 (2003).
- [14] B. T. Gowda, K. Jyothi, N. Damodara, Z. Naturforsch. 58a, xxx (2003).
- [15] R. K. Mohanty, M. Das, A. K. Das, Trans. Met. Chem., Chapman & Hall, London, 22, 487 (1997).
- [16] D. Oliveria, A. Lewis, E. H. Jona, Inorg. Nucl. Chem. Lett., 12, 195 (1976).
- [17] S. V. Rao, V. Jagannadhan, Indian J. Chem. 27A, 252 (1988).
- [18] Y. Ogata, T. Kazushige, Can. J. Chem. 59, 718 (1981).

(10)

Table 10. Melting points (°C) of arylsulphonamides and their *N*-chloro and *N*,*N*-dichloro compounds.

- [19] B. T. Gowda, P. V. V. Nambiar, J. D. D'Souza, J. Indian Chem. Soc. 78, 273 (2001).
- [20] N. Damodara, K. Jyothi, B. T. Gowda, Oxid. Commun. 27, (2003).
- [21] B. T. Gowda, D. S. Mahadevappa, Microchem. J. 28, 374 (1983).
- [22] E. Stahl, Thin Layer Chromatography. A Laboratory Hand Book, Springer, New York (1969).